

NORTHWEST ATLANTIC FISHERIES ORGANIZATION



Scientific Council Studies Number 48

Annual Temperature Curves in Twelve Regions of the Gulf of Maine 1985–2013



You are free to copy and distribute the work and to make derivative works under the following conditions:

Attribution. You must attribute the work in the manner specified by the author or licensor.

Noncommercial. You may not use this work for commercial purposes.

Any of these conditions can be waived if you get permission from the copyright holder. Your fair dealing and other rights are in no way affected by the above.

<http://creativecommons.org/licenses/by/2.5/ca/legalcode.en>

ISSN-0250-6432

Annual Temperature Curves in Twelve Regions of the Gulf of Maine 1985–2013

Ernest D. True

Department of Mathematics, Norwich University
Northfield, Vermont 05663, USA

TRUE, E. 2017. Annual Temperature Curves in Twelve Regions of the Gulf of Maine 1985–2013. *Scientific Council Studies*, **48**: 1–11. doi:10.2960/S.v48.m1

Abstract

Annual temperature functions have been constructed from historical ocean station data (OSD), expendable bathythermograph (XBT) and conductivity-temperature-depth (CTD) temperature data collected in the Gulf of Maine during the years 1985–2013. The data, from WOD13, was compiled by the Ocean Climate Laboratory (OCL) of the National Oceanographic Data Center/NOAA (NODC). The Gulf of Maine is divided into 12 regions where annual temperature variability may be compared among coastal regions, banks and basins. For each of the 12 regions, annual temperature functions, which are truncated Fourier series, are constructed for a surface layer of 0–30 m and an intermediate layer of 30–70 m. These annual temperature functions are used as benchmarks over this 28 year period so that annual temperature fluctuations can be compared with the historical data of each region. Tables are presented from which the temperature functions may be reproduced, and include the days of the year at which the temperatures obtained an annual maximum and minimum so that annual warming and cooling cycles can be compared among the regions. This work is a continuation of an earlier article in which the same temperature functions were constructed from the NODC data during the period 1912–1983. In most of the 12 regions, the mean annual temperatures for the period 1985–2013 are slightly higher than those of the period 1912–1983.

Keywords: annual temperature curves, banks, basins, Gulf of Maine, water masses

Introduction

The Gulf of Maine (GOM) is a partially enclosed portion of the continental shelf of the northwest Atlantic Ocean. Its temperature and salinity are partially influenced by exterior sources. The colder, fresher, nutrient poor Scotian Shelf Water (SSW) enters as surface water near Cape Sable, Nova Scotia in Canada, and can also flow across Brown's Bank and enter through the east edge of the Northeast Channel. Warm Slope Water (WSW) from the northwest edge of the Gulf Stream enters the Gulf through the east edge of the Northeast Channel at depth, as does the deeper Labrador Slope Water (LSW). The LSW is cooler and slightly fresher than WSW.

The exterior water sources exhibit major interannual variability in temperature, salinity, nutrients and volumes and play a significant role in the makeup of Gulf of Maine waters (Townsend, 2015). For more details on the effect of these water masses, see Li *et al.* (2014a,b), Smith *et al.* (2001), Colbourne (2004), and their references. Locally, seasonal and interannual variability occurs due to changes in precipitation and evaporation, surface wind forcing, heat flux, and freshwater river runoff.

Throughout the 1900s and early 2000s, a number of articles examined the seasonal and long-term temperature variability in the Gulf of Maine and surrounding waters. One data set of surface temperature readings taken twice daily since 1921 at St. Andrews in the Bay of Fundy, New Brunswick, Canada was used by Lauzier (1965) to show a warming period from the 1940s to the 1950s, followed by a cooling trend from the 1950s to 1965. The same data set was later used by Drinkwater and Gilbert (2004), to show another cooling trend in the late 1980s, followed by a temperature increase through the 1990s. They also used the data set of temperatures taken twice daily at Halifax, Nova Scotia since 1926, where surface temperatures rose during the 1990s. On the Scotian Shelf, they also found that near-surface temperatures in 1999 reached its second highest value in 55 years, while salinity values were at a record low value. By 2000, salinity values had returned to normal.

Ji *et al.* (2008) demonstrate how cold low salinity surface water from the Scotian Shelf into the Gulf of Maine reduces vertical mixing, reducing nutrients from rising to the surface to enhance photosynthesis, causing a reduction in net primary production. In late February 1999, during

a cruise across the central Gulf of Maine, Durbin *et al.* (2003) observed a winter phytoplankton bloom which was not present during a similar cruise in February 2000. A large amount of colder, low-salinity SSW was recorded in February 1999 in the Central Gulf, which helped to stabilize the water column and enhance early phytoplankton growth. This, in turn, supported high zooplankton populations. But during the same period in 2000, with no significant presence of Scotian Shelf waters, phytoplankton and zooplankton populations were typically low.

There has also been considerable discussion regarding the influence of external water types entering the GOM and subsequent changes in primary production for the future. In the Gulf of St. Lawrence, Drinkwater and Gilbert (2004) report that the river discharge from the St. Lawrence, Ottawa and Saguenay rivers during 1980–2000 was at its highest in 90 years. Near-surface temperatures in the southern gulf of St. Lawrence have increased since the 1970s, reaching its highest value in 1999 since the 1950s. But subsurface temperatures were generally colder in the 1990s. These waters make up part of the SSW entering the GOM. Townsend *et al.* (2015) report that in recent years Arctic melting has led to increased fresher, colder, less-nutrient fluxes into the Gulf of Maine through the Scotian Shelf current. A larger volume of this nutrient-poor water prevents the inflow through the northern section of the Northeast Channel of the deeper nutrient-rich water masses from the LSW and WSW. This reduction in nutrient rich injections to the GOM leads to a reduction in biological productivity.

The purpose of this study is to use a large accumulation of historical data to construct annual temperature curves in 12 important regions of the GOM from 1985–2013. These annual temperature curves can then be used as benchmarks to compare fluctuations during shorter time periods and in the near future.

Methods

Description of Data

The data used in this study is the historical ocean station data(OSD), XBT and CTD data available from the World Ocean Database 2013 (Boyer *et al.*, 2013). The data is managed by the Ocean Climate Laboratory(OCL), created by the National Oceanographic Data Center (NODC). The data in WOD13 includes work by numerous oceanographers around the world over the years whose data collecting surveys have been conducted with many different objectives. The OCL has processed this data into the same format for convenient access to the public. The quality control and processing of this data is described

in “The World Ocean Database 2013 User’s Manual” (Johnson *et al.*, 2013). The data includes optional quality control flags, which were used in this study to include only clean casts. Also, the XBT data was corrected following the Gourestki bias correction for XBT data (Gourestki and Kolterman, 2007). The OSD, XBT and CTD temperature data were extracted from the database for the period 1985–2013.

Description of Regions

The Gulf of Maine is partitioned into 12 regions (Fig. 1), based on the topography, basins and banks of importance in biological studies. The New England coastal areas are covered in regions 1 through 3 and 12. Regions 4 and 5 cover the Bay of Fundy and the coastal waters off Nova Scotia from Yarmouth to Barrington Bay and includes Cape Sable Island and Lurcher Shoals. Region 5 is bordered by the offshore areas on the west by Jordan Basin (region 6) and on the south by the north edge of Browns Bank. The regions 7, 9 and 10 cover Georges, Rodgers and Wilkinson Basins, respectively. Region 8 covers Georges Bank within the 80 meter isobath. The offshore area, region 11, which includes Jeffreys Bank, is made up of small basins, ridges and ledges between Jordan and Wilkinson Basins. This particular partition of the GOM is meant to examine temperature variability separately over basins, banks and strong tidal mixing regions along the coast. The seasonal amplitude in temperature differs considerably over the various regions due in part to topography, tidal mixing, river runoff, and atmospheric forcing. The flux of external water masses also plays an important role in some regions. In particular, Jordan and Georges Basin experience temperature changes as a consequence of intrusions of Scotian Shelf waters and warm slope waters (Townsend *et al.*, 2015).

The total number of OSD, XBT and CTD stations used in this study by month for each of the 12 regions is given in Table 1. The data set contains a total of 26,171 temperature stations compared with the 17,499 stations that were used in the 1912–1983 study (True and Wiitala, 1990).

Annual Temperature Curves

For each of the 12 regions, annual temperature curves for the surface layer (0–30 m) and an intermediate layer (30–70 m) were constructed. The procedure is the same as the curves presented in the earlier 1912–1983 study, so that comparisons can be made. For each station in a given region, the temperatures were integrated over a given depth interval to obtain a weighted mean temperature. Only those stations in the data set with three or more temperature readings in the given depth interval were




Fig 1. Twelve regions in the Gulf of Maine in which annual temperature curves are presented.

Table 1. The number of ODS, XBT and CTD stations for each month in each Region

Region	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Total
1	30	54	42	291	125	78	220	255	183	298	181	50	1 807
2	9	11	0	131	70	45	68	131	49	76	26	4	620
3	2	11	1	73	9	31	74	106	7	87	118	2	521
4	74	116	100	155	291	252	641	566	254	374	624	267	3 714
5	55	58	61	196	76	137	162	294	101	427	406	43	2 016
6	37	64	16	219	18	141	125	170	49	189	132	9	1 169
7	86	130	90	400	84	263	336	209	280	393	230	55	2 556
8	376	752	522	874	1001	681	390	982	286	832	763	286	7 745
9	36	67	38	136	82	69	35	55	138	151	72	16	895
10	57	103	65	204	82	144	85	136	201	237	149	34	1 497
11	46	97	39	207	67	95	111	204	81	207	141	29	1 324
12	128	169	94	340	79	144	112	240	156	396	293	156	2 307
Total	936	1 632	1 068	3 226	1 984	2 080	2 359	3 348	1 785	3 667	3 135	951	26 171

used. This eliminates stations like those in which only surface temperatures were taken. Next, for each day of the year in which more than one set of station readings occurred, the integrated temperatures were averaged to produce a single mean temperature for that day. This distributes the data more evenly throughout the year and reduces the weighting for the curve fitting scheme which followed. The resulting mean temperature could then be

plotted for the day of the year in which the station data was obtained. As an example, Figure 2 shows a scatter plot of the mean temperatures over the upper 30 m for the 241 days of the year in which at least one temperature station was recorded in region 1. The annual temperature curves to be explained next for both the earlier 1912–1983 period of study and this 1985–2013 study are also shown in Figure 2.

The annual temperature curves are constructed by using a regression scheme to fit a continuous annual temperature function to the integrated temperature data using the method of least squares. The final temperature functions are all of the form:

$$T(x) = M + A_1 \sin(0.0172x + B_1) + A_2 \sin(0.0344x + B_2) + A_3 \sin(0.0516x + B_3) \quad (1)$$

where x is the day number of the year and $M, A_1, A_2, A_3, B_1, B_2, B_3$ are the parameters which were computed from the regression scheme. The three sine functions are calculated in radians and the temperature $T(x)$ gives the temperature in degrees centigrade. This temperature function was simplified from the first seven terms of the truncated Fourier series:

$$T(x) = M + \sum_{n=1}^{n=3} \left[a_n \sin\left(\frac{n\pi x}{p}\right) + b_n \cos\left(\frac{n\pi x}{p}\right) \right] \quad (2)$$

with period $2p=365$ days. Equation (1) is derived from (2) by using the trigonometric identity

$$C_1 \sin(\theta) + C_2 \cos(\theta) = \sqrt{C_1^2 + C_2^2} \sin(\theta + k) \quad (3)$$

where $\arctan(k) = C_2/C_1$.

The coefficients and lag angles are computed by the least squares method rather than through the use of the Fourier integral, partly due to the fact that the temperature data is not evenly spaced in time and can lead to poor approximations in the integration for determining some of the coefficients. The use of least squares in place of computing Fourier coefficients is further explained by a well known result that Fourier coefficients give the best least-squares fit for the orthogonal expansion of the type of function described here. Hence, Fourier coefficients and least squares are closely bound.

The decision to express $T(x)$ in terms of the first three harmonics on a Fourier series is based on several factors. For the purpose of comparisons among regions and ease of applications, such as in computer modeling, it is desirable to have a single function type which produces accurate results for each season in each of the 12 regions. The temperature should accurately describe early spring data, when vernal surface warming initiates vertical mixing and surface turnover, as well as the autumn data when surface and intermediate layers begin to cool. The length of the warming and cooling cycles, which vary among regions, should be reflected in the temperature function as well. To achieve these objectives, the temperature data for a given region was first regressed using only the sine term of the first harmonic,

$$T(x) = M + C_1 \sin(0.0172x) \quad (4)$$




Fig 2. Integrated temperature data over the upper 30 m for each of the 241 acceptable days of the year. The corresponding fitted annual temperature curve over the upper 30 m in Region 1 is shown for 1985–2013. For comparison, the fitted curve from the 1912–1983 data is also shown.

The coefficient C_1 was then examined using the null hypothesis that $C_1=0$ against the hypothesis that $C_1 \neq 0$. At the 95% confidence interval, C_1 was found to be significant and the cosine term was then added and the same data was regressed onto the function

$$T(x) = M + C_1 \sin(0.0172x) + C_2 \cos(0.0172x) \quad (5)$$

Again, C_2 was found to be significant and included in the temperature function. The identity in equation (3) is applied to express equation (5) in the form

$$T(x) = M + A_1 \sin(0.0172x + B_1) \quad (6)$$

This first harmonic with a period of 365 days is the dominant harmonic in equation (1). It may be regarded as a first approximation to the annual fluctuation in temperature about the mean, M . But equation 6 would show the length of the warming and cooling cycles to be the same 6 months in each of the 12 regions. Moreover, equation (6) provides a less accurate fit in the spring and autumn data. The addition of the terms in the second harmonic to equation (6) were also found to be significant at the 95% confidence level, and provided a better fit overall. However, the sum of squares due to the error for just these two harmonics was most prevalent in spring and autumn. The addition of the third harmonic reduced the error and provided a

better overall fit, especially in the coastal regions where the amplitude of the third harmonic, A3, can exceed that of the second harmonic, A2. Although the third harmonic contributes less than 0.5°C in the computation of any of the temperature functions, it is statistically significant in some regions, and modestly improves the fit in others. When additional harmonic terms were added to the first three harmonics, an analysis of their coefficients lead to their rejection at the 95% confidence interval. Also, it should be mentioned that there can be more than one combination of these functions when each function is added to the regression scheme in different orders. The functions chosen here were among the most statistically acceptable. As an illustration, Figure 2 shows the fitted temperature function over the upper 30 m corresponding to the averaged temperature data for region 1. The temperature function for the period 1985–2013 shown in Figure 2 is:

$$T(x) = 8.48 - 4.57 \sin(0.0172x + 0.43) - 0.25 \sin(0.0344x + 0.02) + 0.49 \sin(0.0516x + 0.57) \quad (7)$$

Figure 2 also shows the temperature function that was fit to the 1912–1983 data set in True and Wiitala (1990) which is:

$$T(x) = 7.92 - 4.97 \sin(0.0172x + 0.47) - 0.25 \sin(0.0344x - 0.91) + 0.25 \sin(0.0516x + 1.31) \quad (8)$$

The mean temperature in region 1 for 1985–2013 is about 0.56°C warmer than in 1912–1983.

The information in Table 2 gives the values of the parameters M, A1, A2, A3, B1, B2 and B3 which are used in the temperature function $T(x)$ for each of the 12 regions for the upper 30 m. The table also includes the number of days in the year at which at least one suitable temperature station was recorded, the standard deviation about the mean, M, the correlation coefficient and the days when the temperature function shows its annual minimum and maximum temperature in the given region. Table 3 gives the same results for temperature functions in the intermediate layer 30–70 m. The last row in Tables 2 and 3 give the temperature functions for region SS, the Scotian Shelf, which will be discussed later.

Results and Discussion

The primary purpose of this work has been to offer a process for comparing future warming and cooling trends in various regions of the Gulf of Maine against standard or benchmark annual temperature curves based on historical data. These temperature functions for different regions may be plotted and compared to illustrate seasonal and geographical variability. In particular, they can be used

Table 2. Annual temperature functions integrated over the depth interval 0–30 m. The function is: $T(x) = M + A1 \sin(0.0172x + B1) + A2 \sin(0.0344x + B2) + A3 \sin(0.0516x + B3)$ where $x = \text{day number}$.

Region	No. of days	Standard deviation (SD)	Correlation coefficient (R)	M	A1	B1	A2	B2	A3	B3	Minimum		Maximum	
											temp (°C)	Day number	temp (°C)	Day number
1	241	0.79	0.97	8.48	-4.57	0.43	-0.25	0.02	0.49	0.57	3.32	70	13.35	259
2	151	0.86	0.96	7.76	-4.57	0.34	-0.17	-0.97	0.42	0.28	2.66	77	12.53	261
3	124	0.80	0.95	7.52	-3.77	0.19	-0.26	-0.77	0.36	0.82	3.14	76	11.39	262
4	312	0.65	0.98	7.24	-4.82	0.37	-0.17	0.38	0.14	1.49	2.21	65	12.13	254
5	266	1.22	0.93	6.79	-4.37	0.32	-0.23	0.44	0.29	-1.14	2.50	77	11.15	272
6	194	1.03	0.96	8.36	-4.63	0.43	0.34	0.11	0.05	0.53	3.88	75	13.28	245
7	291	0.95	0.97	8.76	-5.31	0.41	0.41	-0.28	0.10	0.93	3.70	73	14.54	248
8	338	0.61	0.99	9.70	-5.43	0.36	0.46	-1.48	0.16	-0.78	4.68	69	15.52	261
9	202	0.90	0.97	9.66	-5.32	0.45	0.51	-0.26	0.28	0.25	4.56	77	15.58	250
10	266	0.80	0.98	9.61	-5.30	0.50	0.74	-0.23	0.30	0.54	4.67	76	15.78	246
11	243	0.95	0.97	8.87	-4.79	0.43	0.42	0.16	0.31	0.33	4.01	79	14.11	250
12	287	1.04	0.96	8.51	-4.76	0.50	-0.56	0.61	0.49	0.16	3.41	66	13.63	264
SS	334	0.91	0.98	7.01	-6.80	0.37	-0.57	1.45	0.23	-0.23	0.52	68	14.37	262

for comparison with future temperature data surveys to determine cooling and warming anomalies in various topographic areas of the Gulf. Hopkins and Garfield (1979) selected the upper 50 m as the Maine Surface Water, with a temperature-salinity envelope which includes that of the Maine Intermediate Water having a depth range 50–120 m. For this study, the surface layer has been further divided into the 0–30 m and 30–70 m depth ranges. This division was partly influenced by biologists who were interested in temperature variability in a shallow upper layer. For con-

Table 3. Annual temperature functions integrated over the depth interval 30–70 m. The function is:
 $T(x) = M + A1 \sin(0.0172x + B1) + A2 \sin(0.0344x + B2) + A3 \sin(0.0516x + B3)$ where x = day number.

Region	No. of days	Standard deviation		Correlation coefficient (R)	M	A1	B1	A2	B2	A3	B3	Minimum temp (°C)		Maximum temp (°C)	
		(SD)	(R)									Day number	temp (°C)	Day number	temp (°C)
1	215	0.85	0.92	6.70	-2.72	-0.23	-0.68	-0.10	0.07	1.12	3.83	77	9.79	303	303
2	133	0.94	0.94	7.21	-3.70	0.11	-0.21	0.76	0.34	-0.52	3.37	89	11.34	277	277
3	122	0.76	0.95	7.26	-3.34	0.06	-0.28	-0.13	0.16	1.27	3.67	76	10.60	272	272
4	296	0.67	0.97	6.77	-4.13	0.15	-0.40	0.10	0.05	0.31	2.46	74	10.95	276	276
5	252	1.02	0.93	6.28	-3.43	0.19	-0.49	0.52	0.18	-0.94	2.88	72	10.03	280	280
6	190	0.81	0.93	7.03	-2.77	-0.05	-0.44	0.17	-0.04	1.08	4.28	77	10.03	292	292
7	292	0.83	0.92	7.19	-2.67	-0.08	-0.41	0.19	-0.12	1.45	4.61	84	10.17	294	294
8	335	0.72	0.98	9.26	-4.84	0.29	-0.59	1.01	0.31	-0.82	4.70	72	14.60	273	273
9	197	1.02	0.88	7.21	-2.33	-0.23	-0.62	0.18	0.09	-0.40	4.90	79	10.08	298	298
10	261	0.66	0.92	6.69	-2.09	-0.47	-0.67	-0.02	-0.08	0.51	4.90	81	9.45	313	313
11	239	0.74	0.94	6.90	-2.59	-0.21	-0.58	0.05	0.10	-1.31	4.35	84	9.99	300	300
12	258	0.85	0.92	6.21	-2.45	-0.26	-0.94	0.23	-0.21	1.20	3.97	65	9.68	305	305
SS	329	0.92	0.91	3.78	-2.49	-0.33	-1.01	-0.19	-0.25	-0.23	1.12	66	7.15	319	319

venience, the annual temperature functions summarized in Tables 2 and 3 will be referred to as the standard when compared with temperatures from different time periods.

The results of this work as summarized in Tables 2 and 3 include high correlation coefficients for fitting the temperature functions to the WOD13 data. The results for goodness-of-fit were encouraging and residuals were generally distributed evenly about the mean. The standard

deviation among the twelve regions varied from 0.61°C to 1.22°C in the upper 30 m, and 0.66°C to 1.02°C in the 30–70 m layer. The column labeled M gives the annual mean temperature for each region. In the 0–30 m layer, region 5 south of Yarmouth, Nova Scotia showed the coldest mean temperature, 6.79°C followed by region 4 for the Bay of Fundy at 7.24°C. Coastal regions 2 and 3 were only slightly warmer. The warmest regions in the upper layer were the offshore regions 8, Georges Bank at 9.70°C and region 9 over Roger's Basin at 9.66°C. In the 30–70 m layer, region 5 was again the coldest at 6.28°C, while the warmest was on Georges Bank, region 8 again at 9.26°C.

The columns in Tables 2 and 3 for the minimum and maximum temperatures show the amplitude of variability among the regions. Generally, the lowest temperatures occur in March and the highest temperatures occur in September. In the 0–30 m layer, the annual temperature increased 8.25°C in coastal region 3, and 11.11°C over Wilkinson Basin, region 10. In the 30–70 m layer, annual temperature increased 4.55°C in region 10 over Wilkinson Basin, and 9.9°C over Georges Bank, region 8. During the warming period, temperatures in all regions generally increase at about

0.055°C/day in the 0–30 m layer, and about 0.031°C/day in the 30–70 m layer.

The Day numbers in Tables 2 and 3 can also be used to examine the length of the warming and cooling cycles for each of the regions. The warming cycles in the 0–30 m layer varied from 170 days in regions 6 and 10 over Jordan and Wilkinson Basins, to 198 days in Cape Cod Bay, region 12. For the 30–70 m layer, the warming cycle varied from 188 days in coastal region 2 to 240 days again in Cape Cod Bay. Generally, except for Georges Bank, the warming cycle in the surface layer was shorter offshore and over basins, where tidal mixing is reduced.

The temperature functions described here show an increase in the annual mean temperatures for most of the regions that were developed for the 1912–1983 period by True and Wiitala (1990). The differences in mean temperature for the surface layer 0–30 m and intermediate layer 30–70 m are shown in Table 4. The greatest increases occurred along the Maine coast, regions 1 and 2, and on Georges Bank, region 8 for both depth layers. On the other hand, region 5, where SSW enters the GOM between Cape Sable in Nova Scotia and Browns Bank, shows an overall decrease in mean temperatures in both the surface and intermediate layers. This temperature decrease corresponds with that of Smith *et al.* (2012) where, during the early 2000s, there was roughly a 40% increase of Scotian Shelf Water entering region 5 around Cape Sable into the GOM

Table 4. The difference in mean temperatures from 1912–1983 to 1985–2013 temperature functions for each region at the two layers 0–30 m and 30–70 m. Positive entries indicate an increase in mean temperature from 1912–1983 to 1985–2013.

Depths	Region											
	1	2	3	4	5	6	7	8	9	10	11	12
0–30 m	0.56	0.69	0.43	0.44	-0.24	0.21	0.01	0.48	-0.01	0.15	0.34	0.27
30–70 m	0.37	0.45	0.22	-0.06	-0.54	0.11	-0.02	0.41	0.08	0.12	0.18	0.07

compared with the 90s. Such an increase of colder SSW into region 5 would be reflected in the annual temperature function that was created for that region.

As an illustration, the mean annual temperature functions might be used to compare temperatures from a hydrographic survey or satellite data over a short time period relative to the 1985–2013 time period. For example, during 2003–2005, temperatures on the Scotian Shelf were cold relative to the period 1971–2000. The Department of Fisheries and Oceans Canada (DFO, 2004) reported that sea ice cover east of Cabot Strait during the first part of 2003 was twice the normal cover relative to the 1971–2000 period. The first part of 2003 marks the beginning of a decrease in temperatures on the Scotian Shelf, where the water was slightly colder and saltier than normal. The St. Andrews, N.B. annual surface temperature was 0.6°C below normal and 1.8°C colder than normal at Halifax, N.S. For 2004, Petrie *et al.* (2005) report a continuation of below-normal temperatures at St. Andrews and Halifax. Also, the cold intermediate layer on the Scotian Shelf was colder and thicker than normal from 50 m to the bottom. The northern edge of the Gulf Stream current moved about 20 km south of its mean position, leaving room for a stronger southern flow of the Labrador current.

Tracking water types through the GOM certainly requires more parameters than temperature alone. But we can start with the eastern GOM and search through adjacent regions to compare water temperatures during 2003–2005 with the 1985–2013 annual temperature functions. In Jordan basin, region 6, there is a moored buoy, buoy M01, whose data is collected and managed by the University of Maine Ocean Observing System (UMOOS, www.umoos.org). This buoy provides hourly temperature and salinity data at the depths 1, 20, 50, 100, 150, 200 and 250 m, beginning in July, 2003. Also, UMOOS buoy N01 on the east edge of the Northeast Channel, and the eastern part of region 7, has been recording temperature and salinity at similar depths beginning in June 2004. The data from these two buoys were cleaned and integrated to obtain a daily mean temperature for each day of each month of each year. The plots in Figure 3 compare the 30–70 m annual temperature functions for Jordan’s and George’s Basin (regions 6 and 7) with the 50 m temperature recordings from the two buoys during the period 2003–2005 for Jordan’s and 2004–2005 for Georges Basin. For Jordan Basin, the buoy data for 2003–2005 shows a mean annual

temperature of 1.17°C below the mean of the annual temperature function for 1985–2013. Similarly, the buoy data in Georges Basin is 1.21°C below the 1985–2013 annual temperature mean. In both cases, the buoy data at 50 m show the temperatures to be considerably colder during the summer warming season. In addition to comparing the buoy data available in regions 6 and 7 with the annual temperature curves, the WOD13 data for 2003–2005 from these regions were also compared (not shown). The results are the same, with every 30–70 m integrated temperature plot lying below the annual temperature curves. The coastal regions 2 and 3 for 2004–2005 also showed annual mean temperatures slightly more than 1°C below the standard at both the 0–30 m and 30–70 m depths. Also, the annual mean temperatures for 2004–2005 in region 11 were below the standard by slightly more than 1°C. This may be caused by transport out of adjacent region 2 by the western Maine coastal current. Region 9, Roger’s Basin, just west of George’s Basin, was not affected by the colder temperatures in adjacent regions. The remaining regions did not show significant temperature decreases for 2003–2005.

As another illustration, Townsend *et al.* (2015) have examined the data from buoy M01 at the lower depths of 100, 150, 200 and 250 m. Their study uses temperature, salinity, nitrates and silicates to try to determine the sources of water that enter Jordan Basin. They find that the proportions of the three water masses into the GOM lead to considerable T/S variability in Jordan Basin at these lower depths. By analyzing the data from the UMOOS buoy N01 on the east edge of the Northeast Channel, they argue that much of the deeper water in Jordan Basin is of Northeast Channel origin. In particular, during the period 2003 to 2014, they observed numerous fluctuations in temperature and salinity over time periods of 6 months to years. But from mid 2009 to 2014 the data showed a period of increasing temperature and salinity, with a slight decrease in 2011. Water masses can enter Jordan Basin as surface water from SSW along the south edge of Nova Scotia, and at various depths from the Northeast Channel and Georges Basin, region 7. Here, we look at the temperatures in the surface and intermediate layers for 2009–2013 and compare them with the standard for Jordan Basin by using the WOD13 data, which is sufficient to construct temperature functions like those in equation (1). Unlike the data from the M01




Fig 3. Upper figure: the annual temperature curve in region 6 (Jordan's Basin) in the intermediate layer 30–70 m is compared with the 50 m temperature data from Buoy N for 2003–2005. Lower figure: similar to upper figure for region 7 (Georges Basin) with temperature data from Buoy M for 2004–2005.




Fig 4. Upper figure: the annual temperature curve in region 6 (Jordan's Basin) in the surface layer 0–30 m for 1985–2013 is compared with the annual temperature curve for the warmer period 2009–2013. Lower figure: similar to upper figure at the intermediate layer 30–70 m.




Fig 5. Upper figure: the annual temperature curve on the Scotian Shelf in the surface layer 0–30 m for 1985–2013 is compared with the annual temperature curve for the warmer period 2009–2013. Lower figure: similar to upper figure at the intermediate layer 30–70 m.

buoy, the WOD13 data set includes temperature data from stations all around Jordan Basin. The graphs in Figure 4 for Jordan Basin, region 6, show that temperatures in the upper layers were also generally higher throughout the period 2009–2013, with an annual mean increase of 1°C in the depth interval 0–30 m, and 0.8°C at 30–70 m relative to the 1985–2013 standard. Also, the same temperature curves were constructed for Region 5 east of Jordan Basin (not shown). Again, the temperatures for the period 2009–2013 were higher throughout the year than the standard, showing an annual mean increase of 1.5 °C in the 0–30 m depth layer, and 1.4°C in the 30–70 m layer. These results agree with Hebert (2014), who reported that temperatures at Lurcher Shoals in region 5 in 2012 (2013) were about 3.4°C (1°C) warmer at 50m than the 1981–2010 reference period. Hebert also reports that temperatures throughout the Scotian Shelf in 2013 were higher than the 1981–2010 means, and that 2012 was generally the warmest in 43 years.

In addition to the 12 regions used here for the GOM, the WOD13 data for the Scotian Shelf inside the 1000 m isobath for 1985–2013 was also processed the same way as the 12 GOM regions. The annual temperature functions appear at the end of Tables 2 and 3, labeled SS. These functions were derived from the rectangular region of

latitude 43.0 to 44.5N, and longitude 63.5 to 65.5W off Halifax, Nova Scotia. During 1985–2013, 7,617 stations were recorded in the data set. For comparison, Figure 5 compares the mean annual temperature functions for the Scotian Shelf from tables 2 and 3 with temperature functions that were constructed from the WOD13 data on the shelf during the period 2009–2013. The temperatures for 2009–2013 at the 0–30 m layer are 0.89°C higher than the 1985–2013 standard and 0.92°C higher at the 30–70 m layer. These comparisons enforce the suggestions that cooling and warming trends on the southwestern Scotian Shelf lead to similar results at the surface and intermediate layers in the eastern GOM. Temperatures in the coastal regions 2 and 3 and Georges Bank, region 8 during 2009–2013 were also greater than the 1985–2013 standard by more than 1°C, while changes in the other regions were insignificant.

It must be emphasized that the sources of water masses in basins and on coastal areas of the GOM certainly can't be determined from surface temperatures alone. For example, the water masses passing through Jordan Basin at different levels remains a difficult problem, even when temperature, salinity, nitrates, silicates and other parameters are known. The results of a numerical model may be necessary to do a thorough analysis of the pathways and volume transports

of SSW, WSW and LSW moving in and out of the various regions of the GOM. But the comparisons shown here, along with similar testing in other time periods, illustrate that the annual temperature functions in Tables 2 and 3 are reasonably effective indicators of typical or standard annual temperature curves for the 12 regions represented in the Gulf of Maine.

References

- BOYER, T. P., J. I. ANTONOV, O. K. BARANOVA, C. COLEMAN, H. E. GARCIA, A. GRODSKY, D. R. JOHNSON, R. A. LOCARNINI, A. V. MISHONOV, T. D. O'BRIEN, C. R. PAVER, J. R. REAGAN, D. SEIDOV, I. V. SMOLYAR, and M. M. ZWENG. 2013. World Ocean Database 2013, NOAA Atlas NESDIS 72, S. Levitus, Ed., A. Mishonov, Technical Ed.; Silver Spring, MD, 209 p., doi.org/10.7289/V5NZ85MT. Available at www.nodc.noaa.gov/about/oceanclimate.html
- COLBOURNE, E. B. 2004. Decadal Changes in the Ocean Climate in Newfoundland and Labrador Waters from the 1950s to the 1990s. *Journal of Northwest Atlantic Fisheries Science*, **Vol. 34**: 41–59. doi.org/10.2960/J.v34.m478
- DFO, 2004. State of the Ocean 2003: Physical Oceanographic Conditions on the Scotian Shelf, Bay of Fundy and Gulf of Maine. *DFO Can. Sci. Advis. Sec. Ecosystem Status Rep.* 2004/004.
- DRINKWATER, K. F., and D. GILBERT. 2004. Hydrographic Variability in the Waters of the Gulf of St. Lawrence, the Scotian Shelf and the Eastern Gulf of Maine. (NAFO Subarea 4) During 1991–2000. *J. Northw. Atlan. Fish. Sci.*, **34**: 83–99. doi.org/10.2960/J.v34.m545.
- DURBIN, E. G., R. G. CAMPBELL, and M. C. CASAS, *et al.* 2003. Interannual variation in phytoplankton blooms and zooplankton productivity and abundance in the Gulf of Maine during winter. *Mar. Ecol. Prog. Ser.*, **254**, 81–100.
- GOURETSKI, V., and K. P. KOLTERMANN. 2007. How much is the ocean really warming?, *Geophys. Res. Lett.*, **34**, L01610, doi.org/10.1029/2006GL027834.
- HEBERT, D., R. PETTIPAS, D. BRICKMAN, and M. DEVER. 2014. Meteorological, Sea Ice and Physical Oceanographic Conditions on the Scotian Shelf and in the Gulf of Maine during 2013. *DFO Can. Sci. Advis. Sec. Res. Doc.* 2014/070. v + 40 p.
- HOPKINS, T. S. and N. GARFIELD. 1979. Gulf of Maine Intermediate Water. *J. Mar. Res.*, **37(1)**: 103–139.
- JI, R., C. DAVIS, C. CHEN, D. TOWNSEND, D. MOUNTAIN, and R. BEARDSLEY. 2008. Modeling the influence of low-salinity water inflow on winter-spring phytoplankton dynamics in the Nova Scotian Shelf-Gulf of Maine region. *Journal of Plankton Research*, **Vol 30**: Num 12, 1399–1416 p.
- JOHNSON, D. R., T. P. BOYER, H. E. GARCIA, R. A. LOCARNINI, O. K. BARANOVA, and M. M. ZWENG. 2013. World Ocean Database 2013 User's Manual. Sydney Levitus, Ed.; Alexey Mishonov, TECHNICAL ED.; NODC Internal Report 22, NOAA Printing Office, Silver Spring, MD, 172 p. Available at www.nodc.noaa.gov/OC5/WOD13/docwod13.html.
- LAUZIER, L. M. 1965. Long term temperature variations in the Scotian shelf area. *ICNAF Spec. Publ.*, **Vol 6**: 807–815.
- LI, Y., R. HE, J. D. MCGILLICUDDY. 2014a. Seasonal and Interannual Variability in Gulf of Maine Hydrodynamics: 2002–2011. *Deep Sea Res Part 2 Top Stud Oceanogr.* 2014 **May**: 103–222. doi.org/10.1016/j.dsr2.2013.03.001
- LI, Y., R. JI, P. S. FRATANTONI, C. CHEN, J. A. HARE, C. S. DAVIS, and R. C. BEARDSLEY. 2014b. Wind-induced interannual variability of sea level slope, along-shelf flow, and surface salinity on the Northwest Atlantic shelf. *J. Geophys. Res. Oceans*, **119**: 2462–2479.
- PETRIE, B., R. G. PETTIPAS, W. M. PETRIE, and V. SOUKHOVTSEV. 2005. Physical oceanographic conditions on the Scotian Shelf and in the Gulf of Maine during 2004. *Cdn. Sci. Adv. Sec. Res. Doc.* 2005/21, 44 p.
- SMITH, P. C., N. R. HOUGHTON, R. G. FAIRBANKS, and D. G. MOUNTAIN. 2001. Interannual variability of boundary fluxes and water mass properties in the Gulf of Maine and on Georges Bank: 1993–1997. *Deep-Sea Research*, **II 48**: 37–70. [doi.org/10.1016/S0967-0645\(00\)00081-3](https://doi.org/10.1016/S0967-0645(00)00081-3)
- SMITH, P. C., N. R. PETTIGREW, P. YEATS, D. W. TOWNSEND and G. HAN. 2012. Regime Shift in the Gulf of Maine. In: *Advancing an Ecosystem Approach in the Gulf of Maine*. 415 p. Stephenson, R. L., J. H. Annala, J. A. Runge and M. Hall-Arber, (Eds). American Fisheries Society, Symposium 79, Bethesda, MD. p. 185–203
- TOWNSEND, D. W., N. R. PETTIGREW, M. A. THOMAS, M. G. NEARY, D. J. MCGILLICUDDY, and J. O'DONNELL. 2015. Water Masses and Nutrient Fluxes to the Gulf of Maine. *J. Marine Research*, **Vol 73**, Numbers 3–4, May & July 2015, 93–122 p.(30)
- TRUE, E. D., and S. A. WIITALA. 1990. Annual temperature curves in twelve regions of the Gulf of Maine. *NAFO Sci. Coun. Stud.*, **14**: 21–27.

Scientific Publications of the Northwest Atlantic Fisheries Organization

In efforts to reduce paper usage and ensure publications are accessible to all,
most publications are available FREE electronically at www.nafo.int or <http://journal.nafo.int>

The NAFO publications listed below are available through the NAFO Secretariat.
Prices include postage and handling.

Please note: Pricing for volumes shipped overseas is higher due to increased postage costs.

	Price North America (CAD \$)	Price Overseas (CAD \$)
Early Stages of Fishes in the Western North Atlantic Ocean (Davis Strait, Southern Greenland and Flemish Cap to Cape Hatteras)		
This comprehensive scientific publication is the only up-to-date textbook providing detailed descriptions and accurate drawings of the early life-history stages of the fishes from the Northwest Atlantic Ocean north of 35°N and west of 40°W. The region covers the world's most famous fishing grounds and includes the Davis Strait, southern Greenland, Flemish Cap, Georges Bank, northern Sargasso Sea and Middle Atlantic Bight to Cape Hatteras.		
Hardcover 2-Volume Edition by Michael. P. Fahay.....	120.00	135.00
Identification Guides		
<i>Coral, Sponge, and Other Vulnerable Marine Ecosystem Indicator Identification Guide, NAFO Area</i> (A pictorial Identification Guide for some species of Coral, Sponge, and other indicator species produced in full-colour on water-resistant paper and bound in a water-proof binder, with tab separators. Additions can be easily added).....	30.00	35.00
<i>Sponge Identification Guide NAFO Area</i> (A pictorial Identification Guide for some species of Sponge found on the Grand Banks of Newfoundland and Flemish Cap produced in full-colour on waterproof/tearproof stock)	30.00	35.00
<i>Coral Identification Guide NAFO Area</i> (A pictorial Identification Guide for some species of Coral found on the Grand Banks of Newfoundland and Flemish Cap produced in full-colour on waterproof/tearproof stock).....	25.00	30.00
<i>Identification Guide of Wolffish, Hake and Rockling in the Northwest Atlantic</i> (1-page Identification Guide produced in full-colour on laminated stock)	10.00	15.00
Journal of Northwest Atlantic Fishery Science - Available FREE online at http://journal.nafo.int		
This publication provides an international forum for the primary publication of original research papers on fisheries science in the Northwest Atlantic, with emphasis on environmental, biological, ecological and fishery aspects of the living marine resources and ecosystems.		
Please note: Scientific publications during ICNAF times (1949–1979) are available at the Secretariat in print format and also available online at www.nafo.int		
Vol. 48, 2016 (Regular Issue).....	40.00	45.00
Vol. 47, 2015 (Regular Issue).....	40.00	45.00
Vol. 46, 2014 (Regular Issue).....	40.00	45.00
Vol. 45, 2013 (Regular Issue).....	40.00	45.00
Vol. 44, 2012 (Regular Issue).....	40.00	45.00
Vol. 43, 2010-2011 (Regular Issue)	40.00	45.00
Vol. 42, 2009-2010 (The Role of Marine Mammals in the Ecosystem in the 21st Century).....	40.00	45.00
Vol. 41, 2008-2009 (Reproductive and Recruitment Processes of Exploited Marine Fish Stocks)	40.00	45.00
Vol. 40, 2008 (Regular issue)	40.00	45.00
Vol. 39, 2007-2008 (Environmental and Ecosystem Histories in the Northwest Atlantic – What Influences Living Marine Resources?)	40.00	45.00
Vol. 38, 2007 (Regular issue).....	40.00	45.00
Vol. 37, 2005-2007 (Flemish Cap Symposium).....	40.00	45.00

Vol. 36, 2005-2006 (Regular issue)	36.00	41.00
Vol. 35, 2005 (Symposium on Elasmobranch Fisheries: Managing for Sustainable Use and Biodiversity Conservation)	46.00	51.00
Vol. 34, 2004 (Mini-symposium on Hydrographic Variability in NAFO Waters for the Decade 1991-2000 in Relation to Past Decades)	31.00	36.00
Vol. 33, 2003 (Reproductive Potential of Fish Populations of the North Atlantic)	31.00	36.00
Vol. 32, 2003 (Regular issue).....	31.00	36.00
Vol. 31, 2003 (Joint NAFO/ICES/CSIRO Symposium entitled "Deep-Sea Fisheries").....	41.00	46.00
Vol. 30, 2002 (Regular issue).....	31.00	36.00
Vol. 29, 2001 (Regular issue).....	31.00	36.00
Vol. 28, 2000 (Special Issue on A Review of the Cod Fisheries at Greenland, 1910-1995).....	31.00	36.00
Vol. 27, 2000 (NAFO/ICES/PICES Symposium on Pandalid Shrimp Fisheries – Science and Management at the Millennium)	41.00	46.00
Vol. 26, 2000 (Regular issue).....	31.00	36.00
Vol. 25, 1999 (Variations in Maturation, Growth, Condition and Spawning Stock Biomass Production in Groundfish).....	Out of Print	
Vol. 24, 1998 (Regular issue).....	26.00	31.00
Vol. 23, 1998 (What Future for Capture Fisheries).....	36.00	41.00
Vol. 22, 1997 (NAFO/ICES Symposium on The Role of Marine Mammals in the Ecosystem).....	31.00	36.00
Vol. 21, 1997 (Regular issue).....	26.00	31.00
Vol. 20, 1996 (Special Issue on North Atlantic Fishery Management Systems: A Comparison of Management Methods and Resource Trends)	26.00	31.00
Vol. 19, 1996 (Gear Selectivity/Technical Interactions in Mixed Species Fisheries Symposium)	26.00	31.00
Vol. 18, 1996 (Regular issue)	23.00	28.00
Vol. 17, 1994 (Special Issue on Report of the Canada-USSR Capelin Symposium)	21.00	26.00
Vol. 16, 1994 (Regular issue)	21.00	26.00
Vol. 15, 1993 (Special Issue on Decapod Crustacean Larvae from Ungava Bay)	21.00	26.00
Vol. 14, 1992 (Special Session on Changes in Biomass, Production and Species Composition).....	21.00	26.00
Vol. 13, 1992 (Regular issue).....	16.00	21.00
Vol. 12, 1992 (Regular issue).....	16.00	21.00
Vol. 11, 1991 (Regular issue).....	16.00	21.00
Vol. 10, 1990 (Special issue on The Delimitation of Fishing Areas in the Northwest Atlantic)	16.00	21.00
Vol. 9, No. 2, 1989 (Regular issue).....	16.00	21.00
Vol. 9, No. 1, 1989 (Regular issue).....	16.00	21.00
Vol. 8, 1988 (Regular issue).....	16.00	21.00
Vol. 7, No. 2, 1987 (Regular issue).....	16.00	21.00
Vol. 7, No. 1, 1986 (Regular issue).....	16.00	21.00
Vol. 6, No. 2, 1985 (Regular issue).....	16.00	21.00
Vol. 6, No. 1, 1985 (Regular issue).....	16.00	21.00
Vol. 5, No. 2, 1984 (Regular issue).....	16.00	21.00
Vol. 5, No. 1, 1984 (Regular issue).....	16.00	21.00
Vol. 4, 1983 (Guide to the Early Stages of Marine Fishes in the Western North Atlantic Ocean, Cape Hatteras to the Southern Scotian Shelf)	26.00	31.00
Vol. 3, No. 2, 1982 (Regular issue).....	14.00	19.00
Vol. 3, No. 1, 1982 (Regular issue).....	14.00	19.00
Vol. 2, 1981 (Regular issue).....	14.00	19.00
Vol. 1, 1980 (Regular issue)	14.00	19.00

NAFO Scientific Council Reports - Available FREE online at www.nafo.int

This publication contains reports of Scientific Council Meetings held through each year since NAFO replaced ICNAF. (The comparable publication during ICNAF was entitled the *Redbook*).

2016 (issued Apr 2017)	Online only	
2015 (issued Apr 2016)	Online only	
2014 (issued Jan 2015)	Online only	
2013 (issued Jan 2014)	Online only	
2012 (issued Jan 2013)	36.00	41.00
2011 (issued May 2012)	36.00	41.00
2010 (issued April 2011)	36.00	41.00
2009 (issued May 2010)	36.00	41.00
2008 (issued May 2009)	36.00	41.00

2007 (issued May 2008)	36.00	41.00
2006 (issued May 2007)	36.00	41.00
2005 (issued May 2006)	36.00	41.00
2004 (issued January 2005)	36.00	41.00
2002/2003 Supplement (issued January 2004)	21.00	26.00
2002/2003 (issued August 2003)	31.00	36.00
2002 (issued January 2003)	31.00	36.00
2001 (issued January 2002)	31.00	36.00
2000 (issued January 2001)	29.00	34.00
1999 (issued January 2000)	29.00	34.00
1998 (issued January 1999)	26.00	31.00
1997 (issued January 1998)	23.00	28.00
1996 (issued January 1997)	23.00	28.00
1995 (issued January 1996)	23.00	28.00
1994 (issued January 1995)	21.00	26.00
1993 (issued January 1994)	21.00	26.00
1992 (issued December 1992)	18.00	23.00
1991 (issued December 1991)	16.00	21.00
1990 (issued December 1990)	14.00	19.00
1989 (issued December 1989)	14.00	19.00
1988 (issued December 1988)	12.00	17.00
1987 (issued December 1987)	12.00	17.00
1986 (issued December 1986)	12.00	17.00
1985 (issued December 1985)	12.00	17.00
1984 (issued December 1984)	12.00	17.00
1983 (issued December 1983)	12.00	17.00
1982 (issued December 1982)	12.00	17.00
1981 (issued December 1981)	12.00	17.00
1979–80 (issued December 1980)	12.00	17.00

NAFO Scientific Council Studies - Available FREE online at www.nafo.int

This publication includes papers of topical interest and importance to the current and future activities of Scientific Council.

No. 47 (2015) Kenchington, E., L. Beazley, F. J. Murillo, G. Tompkins MacDonald and E. Baker. 2015. Coral, Sponge, and Other Vulnerable Marine Ecosystem Indicator Identification Guide, NAFO Area	Online only	
No. 46, 2014 (Protocols of the EU bottom trawl survey of Flemish Cap)	Online only	
No. 45, 2013 (NAFO Secretariat. NAFO Research Vessel Stock-by-Stock Surveys Summary 2000–2010)	Online only	
No. 44, 2012 (Report of the Workshop on Implementation of Stock Reproductive Potential into Assessment and Management Advice for Harvested Marine Species)	Online only	
No. 43, 2010 (Sponge Identification Guide - NAFO Area)	Online only	
No. 42, 2009 (Coral Identification Guide - NAFO Area)	Online only	
No. 41, 2008 (Report of the Greenland Halibut (<i>Reinhardtius hippoglossoides</i>) Age Determination Workshop)	Online only	
No. 40, 2007 (Identification of Wolffish, Hake and Rockling in the Northwest Atlantic)	Online only	
No. 39, 2005 (Workshop on Mapping and Geostatistical Methods for Fisheries Stock Assessment)	31.00	36.00
No. 38, 2005 (Yellowtail Flounder Ageing Manual)	36.00	41.00
No. 37, 2003 (Working Group on Reproductive Potential)	41.00	46.00
No. 36, 2003 (Workshop on Assessment Methods)	41.00	46.00
No. 35, 2002 (Workshop: The Canada-United States Yellowtail Flounder Age Reading)	31.00	36.00
No. 34, 2001 (Regular issue)	31.00	36.00
No. 33, 2000 (Regular issue)	31.00	36.00
No. 32, 1999 (Regular issue)	31.00	36.00
No. 31, 1998 (Regular issue)	31.00	36.00
No. 30, 1997 (Miscellaneous Selected Papers)	26.00	31.00
No. 29, 1997 (Selected Studies Related to Assessment of Cod in NAFO Divisions 2J+3KL)	26.00	31.00
No. 28, 1996 (Assessment of Groundfish Stocks Based on Bottom Trawl Survey Results)	26.00	31.00
No. 27, 1996 (Miscellaneous Selected Papers)	26.00	31.00
No. 26, 1996 (Selected Papers on Harp and Hooded Seals)	26.00	31.00
No. 25, 1996 (Flemish Cap Selected Environmental and Other Papers)	26.00	31.00

No. 24, 1996 (Symposium on Impact of Anomalous Oceanographic Conditions at the Beginning of the 1990s in the Northwest Atlantic on the Distribution and Behaviour of Marine Life)	26.00	31.00
No. 23, 1995 (Miscellaneous Selected Papers).....	26.00	31.00
No. 22, 1995 (Miscellaneous Selected Papers).....	26.00	31.00
No. 21, 1994 (Collection of Papers Related to Northern Cod and Seals in NAFO Divisions 2J and 3KL)	26.00	31.00
No. 20, 1994 (Miscellaneous Selected Papers).....	26.00	31.00
No. 19, 1993 (Miscellaneous Selected Papers).....	21.00	26.00
No. 18, 1993 (Symposium on Changes in Abundance and Biology of Cod Stocks and Their Possible Causes)	21.00	26.00
No. 17, 1993 (Workbook: Introduction to Sequential Population Analysis)	18.00	23.00
No. 16, 1991 (Special Session on Management Under Uncertainties, 5–7 September 1990)	21.00	26.00
No. 15, 1991 (Miscellaneous Selected Papers)	18.00	23.00
No. 14, 1990 (Miscellaneous Selected Papers)	18.00	23.00
No. 13, 1989 (Miscellaneous Selected Papers)	18.00	23.00
No. 12, 1988 (Miscellaneous Selected Papers)	18.00	23.00
No. 11, 1987 (Miscellaneous Selected Papers)	18.00	23.00
No. 10, 1986 (Miscellaneous Selected Papers)	18.00	23.00
No. 9, 1985 (Special Session on Squids, September 1984)	21.00	26.00
No. 8, 1985 (Miscellaneous Selected Papers).....	16.00	21.00
No. 7, 1984 (Miscellaneous Selected Papers).....	14.00	19.00
No. 6, 1983 (Miscellaneous Selected Papers)	14.00	19.00
No. 5, 1982 (Symposium on Environmental Conditions in the Northwest Atlantic During 1970–79, September 1981)	14.00	19.00
No. 4, 1982 (Special Session on Remote Sensing September 1981).....	14.00	19.00
No. 3, 1982 (Miscellaneous Selected Papers).....	12.00	17.00
No. 2, 1982 (Manual on Groundfish Surveys in the Northwest Atlantic).....	11.00	16.00
No. 1, 1981 (Miscellaneous Selected Papers)	12.00	17.00

Information for Preparing Manuscripts for NAFO Scientific Publications

Journal of Northwest Atlantic Fishery Science

The Journal is for the primary publication of original practical and theoretical research that is unpublished and is not being submitted for publication elsewhere. While it is intended to be regional in scope, papers of general applicability and methodology may be considered. Space is also provided for notes, letters to the editor and notices. Each paper is assigned to an Associate Editor of the Journal's Editorial Board, and is normally reviewed by two referees regarding suitability as a primary publication.

NAFO Scientific Council Studies

The Studies publishes papers which are of topical interest and importance to the current and future activities of the Scientific Council, but which do not meet the high standards or general applicability required by the Journal. Such papers have usually been presented as research documents at Scientific Council meetings and nominated for publication by the Standing Committee on Publications. Studies papers are not peer reviewed.

Content of Paper

The paper should be in English. The sequence should be: Title, Abstract, Text, References, Tables and Figures.

Title

The paper should start with the title, followed by the name(s), address(es) and emails of the author(s) including professional affiliation, and any related footnotes.

Abstract

An informative concise abstract should be provided along with key words listed alphabetically.

Text

In general, the text should be organized into Introduction, Materials and Methods, Results, Discussion, and Acknowledgements. Authors should be guided by the organization of papers that have been published in the NAFO Journal or Studies.

Introduction should be limited to the purpose and rationale of the study.

Materials and Methods should describe in sufficient detail the materials and methods used, so as to enable other scientists to evaluate or replicate the work.

Results should answer the questions evolving from the purpose of the study in a comprehensive manner and in an orderly and coherent sequence, with supporting tables and figures.

Discussion should explain the main contributions from the study, with appropriate interpretation of the results focusing on the problem or hypothesis. Comparisons with other studies should be included here.

Acknowledgements should be limited to the names of individuals who provided significant scientific and technical support, including reviewers, during the preparation of the paper, and the names of agencies which provided financial support.

References

The references cited in the text should be listed alphabetically. References should be mainly restricted to significant published literature. Unpublished documents and data, papers in preparation, and papers awaiting acceptance to other journals, may be cited with full contact addresses as unpublished or personal communications.

Examples:

KING, M. 1995. Fisheries biology, assessment and management. Fishing News Books, UK, 341 p.

CROWDER, L. B., and S. A. MURAWSKI. 1998. Fisheries by-catch: implications for management. *Fisheries*, **23**: 8–16. doi:10.1577/1548-8446(1998)023<0008:FBIFM>2.0.CO;2

ÁVILA DE MELO, A. M., D. POWER, and R. ALPOIM. MS 2005. An assessment of the status of the redfish in NAFO Division 3LN, *NAFO SCR Doc.*, No. 52, Serial No. N5138, 19 p.

Text citations of the above would be (King, 1995; Crowder and Murawski, 1998; Ávila de Melo *et al.*, MS 2005). The surnames of two authors may be used in a citation, but *et al.* should be used for more than two authors. The citation of mimeographed reports and meeting documents should contain the abbreviation "MS". Abbreviations of periodicals can be found ftp://ftp.fao.org/fi/asfa/Monitoring_List/MASTER.txt. The Digital Object Identifier (doi) should be included if available. <http://www.crossref.org/freeTextQuery/> can be used to check this.

Tables and Figures

All Tables and Figures must be cited in the text. Tables and Figures must be numbered consecutively and correspond with the order of presentation in the text. Figure captions should be included as a separate page. Each table and figure should have a complete concise descriptive caption. Figures should always be submitted in black and white. Colour plots and photographs are acceptable only if colour is essential to the content.

All figures should be submitted as separate files in .eps or .ps format. Photographs, maps and contour plots can also be submitted in high quality .jpg format.

If using excel, open the files in R and save the graphs by right clicking and saving as metafiles or postscript files. If using SlideWrite copy the files as Metafiles (WMF). Do not save them as bitmap files. They are not editable.

Paper Submission

Papers should be submitted by email to General Editor, at journal@nafo.int

