NOT TO BE CITED WITHOUT PRIOR REFERENCE TO THE AUTHOR(S)

Northwest Atlantic



Fisheries Organization

Serial No. N252

NAFO SCR Doc. 80/XI/165

SPECIAL MEETING OF SCIENTIFIC COUNCIL - NOVEMBER 1980

Canadian observations on the shrimp fishery at West Greenland in 1980

Ьy

D.G. Parsons Northwest Atlantic Fisheries Center Department of Fisheries and Oceans St. John's, Newfoundland A1C 5X1

Introduction

Observers from the Newfoundland Region of the Department of Fisheries and Oceans spent a total of 260 sea days on board shrimp vessels fishing in Subareas O and 1. Information obtained from observers' written reports and data summaries include fishing patterns, shrimp discards and by-catch. All reports from Canadian observers are not currently available, therefore the following presentation can be considered preliminary.

The Canadian quota in 0+1 for 1980 was 2500 MT (500 MT in Subarea 0 and 2000 MT in Subarea 1). This amount was accounted for relatively early in the year and the area was closed on July 16.

Results

Fishing Pattern

During April 1980 most observed fishing occurred in Div 1B, 1C and 1D with some effort in western 1B when ice conditions were favourable. Heavy ice was reported in 1B for both April and May and vessels were often forced to move out of the western portions. By late June most fishing was reported from Div 1B.

Reports for the July period are lacking but boardings made during August and September indicate heavy fishing in the OA-1B area. Two reports stated that vessels fishing in the area were constantly moving back and forth over the boundary between the two divisions and another reported considerable effort just east of the line.

Shrimp Discards

The practice of discarding small, soft-shelled and/or broken shrimp occurs quite frequently in the West Greenland fishery and the pattern appears to be variable. Vessels with peelers on board have little or no discards and, presumably, vessels which land their product for peeling ashore would have little discarding as well. The market for the frozen, cooked product appears to dictate a preference for large animals and often results in discards in excess of 30%. (One report had discards as high as 36% for a two week period in June.)

Often the discarded animals consist of sizes which would otherwise be acceptable or, at least, one would interpret this from length frequencies obtained in 1980 (Fig. 1, 2 and 3) especially from sets made in deep water. It has been reported at times that when catch rates are considerably high (and where there is a preference for large animals) all medium and small shrimp are discarded due to processing limitations. This alternative is preferred rather than a reduction in fishing effort and, apparently, no records of these discarded shrimp are kept.

By-catch

Most vessels fishing shrimp on the 0+1 grounds are not equipped to handle by-catches of groundfish and in many instances all by-catch is discarded. It has been reported that some cod and halibut are, at times, kept in relatively small amounts. Major by-catch species include redfish, turbot, cod, halibut, American plaice, thorny skate and wolfish. All sizes of these species are susceptible because of the small mesh used (40-45 mm stretched observed in 1980). Small redfish (< 20 cm) are especially vulnerable and often occur in large quantities.

Tables 1 and 2 present a species breakdown of catches by month for each NAFO division. It is obvious that for most months in all Divisions redfish comprises the greatest amount of by-catch, the proportions in these examples varying from 2 to 33% of the total catch. Data are insufficient to detect any seasonal trend in redfish abundance but it appears that their occurrence is considerably less in Div 1C. No length distributions are available to supplement these figures but it is assumed that quite often the redfish are small.

By-catches of other species do not appear to be disproportionate. These, however, should be judged more objectively in relation to existing knowledge on stock sizes. The high proportion of halibut for Div 1D in April may be anomalous.

- 2 -

Discussion

The observer programme conducted in the shrimp grounds in Subareas O+1 reveals a number of problem areas relating to catch and effort, by-catch and the general state of the shrimp resource.

- 3 -

- In many instances processed weight is recorded rather than actual catch weight. This not only involves the discard problem but weight loss in processing as well (e.g. peeling).
- Discards (in relation to 1 above) are seldom, if ever, reported and often consist of marketable-sized shrimp.
- By-catches are not reported. This may be especially significant in the case of redfish.
- 4) When fishing is possible in Div OA and 1B later in the season, vessels often fish extremely close to the boundary and frequently cross back and forth in response to fishing success. Accurate catch and effort may be difficult to obtain for either area.

				4 –		
Calcm	® ~ ~ ○ ○ ○ ○ ○ ○ ○ ○ ○ ○ ○ ○ ○ ○ ○ ○ ○	0000000 0000000 0000000 00000000 000000	100.00 10 10 10 10 10 10 10 10 10 10 10 10 1	NOUUNNFOONOO FOUOOOOOOO BOOOOOOOOOOOOOOOOOOOOOOOOO	100°00	
LB) WEIGHT (MT	NOS & SOB N NOS & SOB N NON NOB N NON O 0 0 0 N NON O 0 0 N N N N N N N N N N N N N N N N N N	00000000000000000000000000000000000000	182.665 (OA and 1B) weight (mt	M M M M M M M M M M M M M M M M M M M	212.361	
SPECIES NAME	SMRIMP (PANDALUS) SMRIMP (P. MONTAGUI) AMERICAN PLAICE COD MALIBUT REDFISH (UNSPECIFIED) TURBOT	CONGENCRN SCULPIN SWATE (THORNY) WOLFFISH (BROADHEAD) WOLFFISH (STRIPED) WOLFFISH (SPOTTED) WOLFFISH (SPOTTED) GREENLAND SMARK CTMERLAND SMARK	TOTALS August Species name	SHRIMP (PANDALUS) SHRIMP (P. MONTAGUI) AMERICAN PLAICE COD HALIBUT (UNSPECIFIED) HALIBUT (UNSPECIFIED) WITCH CAPELIN CAPELIN SKATE (THORNY) WOLFFISH (BROADHEAD) WOLFFISH (UNSPECIFIED)	TOTALS (0A) (EIGHT (MT) % CATCH	N 00004000000 00000000 00000000 00000000
		- F O N - F O O O O O O O O O O O O O O O O O O	× Catch	00000000000000000000000000000000000000	CIES NAME September	IMP (PANDALUS) IMP (P. MONTAGUI) RICAN PLAICE TIC COD IBUT FISH (UNSPECIFIED) FFISH (STRIPED) FFISH (STRIPED) FFISH (UNSPECIFIED) FFISH (UNSPECIFIED) FFISH (UNSPECIFIED) FFISH (UNSPECIFIED) FFISH (UNSPECIFIED) FFISH (UNSPECIFIED) FFISH (UNSPECIFIED) FFISH (UNSPECIFIED)
April (Jb)	SPECIES NAME SHRIMP (PANDALUS) 169.893 SHRIMP (P. MONTAGUI) 0.360 AMERICAN PLAICE 5.219 COD 0.779 ARCTIC COU 0.078 HALIBUT COU 0.035 REDFISH (UNSPECIFIED) 22.047	TURBOT CAPELIN SKATE (THORNY) WOLFFISH (BROADHEAD) 6.000 WOLFFISH (STRIPED) 0.502 WOLFFISH (STRIPED) 0.5031 WOLFFISH (SPOTTED) 1.533 GREENLAND SHARK 0) 1.533 GREENLAND SHARK 0) 0.400	June (18) Species Name weight	SHRIMP (PANDALUS) SHRIMP (P. MONTAGUI) SHRIMP (P. MONTAGUI) COD COD HADDOCK HADDOCK HADDOCK CNSPECIFIE CUNSPECIFIED) SKATE (THORNY) CULFFISH (BROV) OTHER TOTALS TOTALS COO NOUTHER COO NOU NOUTHER COO NOUTHER COO NOU NOU NO	U B S S	NNAA HAFN SNAAHAFN SNAAHAFN SNAAHAFN SPITTT STORATTT

_
8
19
Ċ,
p
ar
َن
>
5
÷-
σ
ě
Ś
ŝ
å
0
t P
Ē
B
>
ثمَ
S
e.
ပ္ပ
be
S
\geq
<u>_</u>
5
÷
S
ğ
Б
Ó
Ę.
ц С
S
_
•
\sim
e
9
Ta
•

.....

	IC	· ·		
April			May	
SPECIES NAME	WEIGHT (MI) % CAICH	SPECIES NAME	*ELGHT (MI	X CATCH
SHRIMP (PANDALUS) SHRIMP (PANDALUS) SHRIMP (PANDALUS) AMERICAN PLAICE COD HADDOCK HADDOCK REDFISH (UNSPECIFIED) GRENADIERS (ROUGHHEAD) GRENADIERS (ROUGHHEAD) GRENADIERS (ROUGHHEAD) GRENADIERS (ROUGHHEAD) WOLFFISH (STRIPED) WOLFFISH (STRIPED) WOLFFISH (NNSPECIFIED) WOLFFISH (UNSPECIFIED) WOLFFISH (UNSPECIFIED) TOTALS	56 57 57 57	SHAIMP SHAIMP SHAIMP SANAJAP COMPACTION ACOMPACIAN COMPA	US) FAGUI) CIFIED) CIFIED) CIFIED) COSC CIFIED) COSC	Q Q Q Q Q Q Q Q Q Q Q Q Q Q
	JU SPECIES NAME	INE WEIGHT (MT) % CAT	I	
	SHRIMP (PANDALUS) SHRIMP (PANDALUS) AMERICAN PLAICE COD HALIBUT REDFISH (UNSPECIF TURBOT GRENADIERS (ROUGH SKATE (THORNY) WOLFFISH (BROADHE WOLFFISH (BROADHE WOLFFISH (BROADHE)	I I		
	TOTALS	104°187 100°0	0	
April	1D		May	
SPECIES NAME	WELGHT (MT) % CAICH	SPECLES NAME	AELGHT (A	I) % CATCH
SHRIMP (PANDALUS) SHRIMP (P. MONTAGUI) COD HALIBUT REDFISH (UNSPECIFIED WITCH SKATE (THORNY) SKATE (THORNY) UTHER	Contraction Contracti	SHRIMP (PANDA Shrimp (p. Mo Halibut Redfish (unsp Totals	LUS) NTAGUI) 0.000 ECIFLED) 0.010	40 50 50 50 50 50 50 50 50 50 5

- 5 -

TOTALS

100°001

2,495

% CATCH





