NOT TO BE CITED WITHOUT PRIOR REFERENCE TO THE SECRETARIAT



Fisheries Organization

Serial No. N783

NAFO SCS Doc. 84/VI/4

SCIENTIFIC COUNCIL MEETING - JUNE 1984

Canadian Request for Scientific Advice on Management

in 1985 of Certain Stocks in Subareas 0 to 4

by

L. S. Parsons

Assistant Deputy Minister for Atlantic Fisheries Department of Fisheries and Oceans, Ottawa, Canada

. Canada requests that the Scientific Council, at its meeting in advance of the 1984 Annual Meeting, provide advice on the scientific basis for the management of the following fish and invertebrate stocks in 1985:

Cod (Div. 2J, 3K and 3L; Div. 3N and 30, Div. 3Ps) Redfish (Div. 3L and 3N) American plaice (Div. 3L, 3N and 30) Witch flounder (Div. 3N and 30) Yellowtail flounder (Div. 3L, 3N and 30) Greenland halibut (Subarea 2 and Div. 3K and 3L) Roundnose grenadier (Subareas 2 and 3) Silver hake (Div. 4V, 4W and 4X) Capelin (Div. 3L, Div. 3N and 30) Squid (Subareas 3 and 4)

It is further suggested that, subject to the concurrence of the other coastal states concerned, the Scientific Council, prior to the 1984 Annual Meeting of NAFO, provide advice on the scientific basis for management in 1985 of the following stocks:

> Shrimp (Subareas 0 and 1) Greenland halibut (Subareas 0 and 1) Roundnose grenadier (Subareas 0 and 1)

- 2. Canada requests the Scientific Council to consider the following options in assessing and projecting future stock levels for those stocks listed above and for the Flemish Cap (Div. 3M) stocks:
 - a) For those stocks subject to analytical dynamic-pool type assessments, the status of the stock should be reviewed and management options evaluated in terms of their implications for fishable stock size in both the short and long term. In those cases where present spawning stock size is a matter of scientific concern in relation to the continuing productive potential of the stock, management options should be evaluated in relation to spawning stock size. As a general reference point, the implications of continuing to fish at F_{0.1} in 1985 and subsequent years should be evaluated. The present stock size should be described in relation to those observed historically and those expected at the F_{0.1} level. Opinions of the Scientific Council should be expressed in regard to stock sizes, catch rates, and TACs implied by these management strategies for 1985 and the long term.
 - b) For those stocks subject to general production-type assessments, the status of the stock should be reviewed and management options evaluated in the way described above to the extent possible. In this case, the general reference point should be the level of fishing effort (F) which is two thirds that calculated to be required to take the MSY catch in the long term.
 - c) For those resources on which only general biological and/or catch data are available, no standard criteria on which to base advice can be established. The evidence on stock status should, however, be weighed against a strategy of optimum yield management and maintenance of stock biomasss at levels of about two-thirds that of the virgin stock.

