



Serial No. N2269

NAFO SCR Doc. 93/84

SCIENTIFIC COUNCIL MEETING - JUNE 1993

Assessment of Shrimp in the Denmark Strait

by

U. Skúladóttir¹, M. Andersen², D. M. Carlsson³, P. Kannevorff³,
D. G. Parsons⁴ and H. Siegstad².

¹ Marine Research Institute, Skúlagata 4
P.O. Box 1390, 121 Reykjavik Iceland.

² Greenland Fisheries Research Institute, P.O. Box 570
DK-3900 Nuuk, Greenland.

³ Greenland Fisheries Research Institute, Tagensvej 135, 1
DK-2200 Copenhagen N., Denmark.

⁴ Science Branch, Department of Fisheries and Oceans
P.O. Box 5667, St. John's, Newfoundland, Canada, A1C 5x1.

1. INTRODUCTION

The fishery in Denmark Strait began in 1978 with a catch of about 360 tons and subsequently increased to around 12 500 tons in 1988. In 1989 and 1990 the nominal catch decreased to less than 11 000 tons and in 1991 and 1992 declined further to 8 600 and 7 200 tons, respectively. Vessels from Denmark, Faroe Islands, France, Greenland, Iceland and Norway were active in the fishery from 1980 to 1991, inclusive. No fishing was reported from France in 1992. Reported catches and TACs are given in Table 1.

The shrimp fishery in Denmark Strait occurs primarily in the area of Strede and Dohrn Banks as well as on the slopes of Storffjord Deep (Fig. 1). The available fishing grounds at a given time depend heavily on the ice conditions. The main fishing area extends from approximately 65°00'N to 67°30'N and between 26°W and 34°W.

2. INPUT DATA

2.1. COMMERCIAL FISHERY DATA (SCR Doc. 93/51, 93/60, 93/63)

2.1.1 Trends in catch and effort

Catch and effort data were available from logbooks from Greenland, Norway, Iceland and France since 1980. Some Greenlandic data from the early months of 1993 were also available. Catches and corresponding effort were compiled by month and by fleet. CPUEs were calculated by month and the mean weighted CPUE of two periods of the year (January to June and July to December) was then applied to the total catch of the period to estimate the total effort.

Total catches increased rapidly from 1978 to 1980, decreased in 1981 and remained stable to 1983. Catches increased steadily from 1983 to 1988 and then decreased to 1992 (Fig. 2). Total effort values show about the same pattern as catch. Between 1980 and 1989, effort increased from about 35 000 hours to more than 100 000 hours, declining thereafter to about 82 000 hours in 1992.

2.1.2 Standardization of the catch rates

The catch and effort data from Norway and Iceland from 1987 to 1992 were analyzed by country using SAS multiple regression procedures to account for the seasonality (months) of the fishery. The 1978 and 1979 catch rates corresponded to very low effort and, therefore were not comparable to the later years.

The catch rates of 27 Greenland trawlers from 1987 to 1992 were also standardized for the total catch.

2.1.3 Trends in catch rates

Abundance indices were calculated from both the unstandardized and standardized series using 1992 as the reference point (Table 2).

Unstandardized catch rates (Fig. 3) were without trend from 1983 to 1987 followed by a substantial decline to 1989. Values for 1990, 1991, and 1992 were similar to the low 1989 level. The 1989-92 level was about 50% of the level seen from the early to mid-1980s period.

Standardized catch rate series (Fig. 4) for Greenland, Norway, and Iceland showed that the Greenlandic rates declined steadily since 1987 whereas the Norwegian rates declined from 1987 to 1989, remaining stable thereafter. The Icelandic catch rate remained relatively stable between 1987 and 1989, increased substantially between 1989 and 1991, and stabilized again between 1991 and 1992.

2.2 COMMERCIAL BIOLOGICAL DATA (SCR Doc. 93/65)

2.2.1 Icelandic fishery data

The Icelandic samples (Fig. 5) taken in the fall of 1987 and 88 show that the catches east of the midline were comprised mainly of female shrimp with a distinct mode at 30 mm CL. The 1990 fall samples show the increased importance of the male component (about 50% compared to 32 and 26% in 1987 and 1988). The 1991, 1992 and 1993 samples taken in spring show that male shrimp dominated at this time in all three years.

The occurrence of a component of female shrimp with a mode at 26 mm in the 1990 samples suggested that sex change occurred earlier than normal. A study of female maturity ogives in Icelandic samples in the years 1985 to 1992 suggested a drop in the values of L_{50} for the year 1990 only.

2.2.2 Greenlandic fishery data

The samples from the Greenlandic fishery were not considered representative of the total fishery, due to poor coverage in time and area.

2.3 RESEARCH SURVEY DATA (SCR Doc. 93/63)

2.3.1 Biomass estimates

A stratified-random trawl survey was conducted by Greenland in Denmark Strait in October, 1992. The biomass estimate of 1044 tons was lower than the 1990 estimate (1 860 tons) for the same area. The Norwegian surveys from 1985 to 1989 produced much higher estimates between 25 000 and 50 000 tons. Investigations show that differences in survey design should cause the Greenlandic survey (stratified random) to provide biomass estimates of only 20-30% of the Norwegian (fixed station) results.

2.3.2 Demographic structure

Greenland survey samples from 1989, 1990 and 1992 (Fig. 6) showed an increase in the proportion of males over the period, which is consistent with a trend evident from the 1985 to 1989 Norwegian surveys. However, overall abundance declined, especially for females.

	Percent males							
	1985	1986	1987	1988	1989	1990	1991	1992
Norway	43.8	41.4	53.5	58.5	58.0			
Greenland					63.1	62.5	-	78.3

The data also showed that males in 1990 and 1992 did not attain the large sizes evident in 1989. Further, the proportion of small females was higher in 1992, suggesting the possibility of sex change occurring at an earlier age or migration of shrimp from Icelandic stocks.

The survey data showed that both males and females decreased by a factor of 10 in the stratum with highest abundance in 1990 while abundance increased in the area of the mid-line between Greenland and Iceland.

3. SUMMARY OF ALL INDICES

Trends in catches

Increase from 1978 to 1980.
Decrease in 1981 and stabilization in 82 and 83 at around 4 700 tons.
Steady increase from 1983 to 1988 to 12 500 tons.
Decrease from 1988 to 1992.

Trends in effort

General increase from 1979 to 1989.
Decrease from 1989 to 1992.

Trends in catch rates

Overall declining trend in the unstandardized catch rate from 1980 to 1992. Unstandardized rates without trend from 1983 to 1987. A decrease for the standardized catch rate from 1987 to 1992 for Greenland. Stability in standardized catch rate from 1989 to 1992 for Norway. Relative stability in standardized catch rate between 1987 to 1989, increasing substantially between 1989 and 1991 for Iceland.

Biomass estimates from research surveys

Biomass in 1992 lowest observed.

Demographic structure

After 1988, increasing proportions of male shrimp (<28 mm) in Icelandic samples. Indications of earlier sex change occurred between 1989 and 1990, and the condition has not reversed.

Increase in proportion of males since 1985 from Norwegian and Greenlandic surveys.
Differences in distribution of size groups over the survey area between years.

4. SUMMARY OF ADVICE FROM PREVIOUS YEARS

The interpretation of the effects of fishing on the stock in the Denmark Strait has changed since the first assessment was conducted in 1980. In 1981, it was thought that the decreases observed in the spring catch rates were due to heavy exploitation. Also, it was considered that the stock was at the northern limit of the species distribution range, and as such, could be more sensitive to exploitation. Therefore, a cautious approach for the exploitation was recommended, and the TAC was set at 4 200 tons (the estimated MSY). By 1985, it was interpreted that the stock was stable and the TAC of 5 000 tons (average catch 81-84) was advised.

No TAC advice was provided for 1986, 1987 or 1988 because the catch rates could not be interpreted as an index of stock abundance. In 1988, it was observed that increased catches over the previous several years had no apparent effect on the resource and catch levels at around 10 000 tons were recommended as an exploratory level for several years.

Catch rates declined in 1987 and 1988, however catch composition and biomass estimates from 1985 to 1989 suggested that the stock was stable and in 1990 it was recommended that the TAC remain at 10 000 tons. The 1989 Norwegian survey showed that the stock was dispersed and the sexes well-mixed. In 1991, the catch rate series was standardized to account for changes in seasonality and fleet composition and it was interpreted that the stock in 1989-90 was substantially lower than in the period of stabilized catch rates. Also more males appeared in the catches and there were indications of earlier sex change. These concerns resulted in an arbitrary reduction of the TAC from 10 000 to 8 000 tons. The depressed conditions were still evident in the 1991 data and, in 1992 a further reduction to 5 000 tons was advised for 1993 and several years thereafter in an attempt to protect the spawning biomass and rebuild the stock.

5. STATUS OF THE RESOURCE

As interpreted in recent years, the abundance of the resource on the fishing grounds is thought to be at a lower level than it was during the first half of the 1980s. The decrease of CPUE associated with an increasing proportion of male shrimp in the survey data reflects a decrease in the abundance of the females. The inferences for a decrease in the size at sex reversal might indicate a response to a decrease in population abundance or environmental influences.

Standardized catch rates showed different trends for the Icelandic, Norwegian and Greenlandic fleets over the same period. The Greenlandic data show a declining trend from 1987 to 1992, whereas the Norwegian data suggest stability since 1989 and the Icelandic an increase since 1988. Therefore, there is some uncertainty whether the stock has stabilized since 1989 at a low level, as indicated in the unstandardized catch rates and Norwegian data, or is still declining as suggested in the Greenlandic CPUE and in the estimates of biomass. At present, it is not possible to determine which is correct.

6. PROGNOSIS

Given the current low level of the shrimp stock abundance in Denmark Strait, the TAC for 1994 should be 5 000 tons and maintained at that level over the next several years in an attempt to increase the catch rates and prevent a possible over-exploitation of the apparently low spawning biomass.

7. REFERENCES

- Carlsson, D.M., and P. Kannevorff. 1993a. The commercial shrimp fishery in Denmark Strait in 1992 and early 1993. NAFO SCR Doc. 93/60 Serial No. N2243
- 1993b. Stratified-random trawl survey for shrimp (*Pandalus borealis*) in Denmark Strait in 1992.
- Skúladóttir, U. 1993a. The catch statistics of the shrimp fishery (*Pandalus borealis*) in the Denmark Strait in the years 1980-1992. NAFO SCR Doc. 93/63 Serial No. N2246.
- 1993b. The Icelandic shrimp fishery (*Pandalus borealis*) in the Denmark Strait in 1992 and early 1993. NAFO SCR Doc. 93/51 Serial No. N2232.
- 1993c. The sexual maturity of female shrimp (*Pandalus borealis*) in the Denmark Strait in the years 1985-1992 and a comparison to the nearest Icelandic shrimp stocks in 1992. NAFO SCR Doc. 93/65 Serial No. N2249.

Table 1. Nominal catch (tons) of shrimp in the Denmark Strait.

Country	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	*** 1993
Denmark	-	-	702	581	740	204	443	353	500	555	444	366	390	358	160	
Faroe Islands	-	-	4233	713	737	443	668	674	727	595	679	595	843	1007	1095	
France	-	-	50	353	414	291	500	642	780	1030	494	381	51	118	-	
Greenland	-	-	200	1004	1115	1467	2250	2596	5781	6627	7456	5976	6211	4202	2012	
Iceland	363	485	759	125	0	43	742	1794	1150	1330	1424	1326	281	465	1440	
Norway	-	800	2461	2016	1896	1727	2128	2051	2026	2041	2052	2098	2500	2504	2500	
Total catch	363	1285	8405	4792	4902	4175	6731	8110	10964	12178	12549	10742	10276	8654	7207	
Total catch eastern side	363	485	759	125	0	43	742	1794	1150	1330	1424	1326	281	465	1440	
Total catch western side	0	800	7646	4667	4902	4132	5989	6316	9814	10848	11125	9416	9995	8189	5767	
Advised TAC	-	-	-	-	4200	4200	4200	5000	-	-	-	-	10000*	10000*	8000	5000
Effective TAC western side	-	-	-	8000	4500	5725	5245	6090	7525**	7725**	8725**	9025**	14100	14500	13000	9563

* Advised for a few years as a precautionary measure.

** Not including Greenland fishery north of 66° 30' N.

*** Preliminary.

Table 2. Standardized catch rate indices by fleets (Greenland, Norway and Iceland), and unstandardized catch rate indices for all countries combined.

YEAR	GRL	NOR	ICE	ALL , unst.
1980				2.760
1981				2.780
1982				2.410
1983				2.000
1984				2.400
1985				1.780
1986				2.190
1987	3.260	2.060	0.738	1.890
1988	2.860	1.459	0.543	1.430
1989	1.900	1.061	0.563	1.140
1990	1.800	1.110	0.725	1.240
1991	1.400	0.999	1.050	1.020
1992	1.000	1.000	1.000	1.000




Fig. 1. The strata numbers in the Denmark Strait. The general fishery area is shaded and the area covered by the Norwegian observer is the black spot. A much wider area of likely distribution is indicated by a solid line.




Fig. 2. Catch and effort from the logbooks weighted by nominal catches




Fig. 3. The unstandardized catch rate indices of all countries combined.




Fig. 4. The standardized catch rate indices of the Greenland, Norway and Iceland fleets.

Per mille









Fig. 5 The Icelandic samples in the years 1987, 1990 to 1993 in the eastern part of the area.




Fig. 6 The Greenlandic survey samples in the year 1989, 1990 and 1992 in the Denmark Strait area. In pooling the samples were weighted by catch and stratum area.