

Northwest Atlantic



Fisheries Organization

Serial No. N2445

NAFO SCR Doc. 94/67

SCIENTIFIC COUNCIL MEETING – SEPTEMBER 1994

Stability of Water Masses – Impact on Cod Recruitment Off West Greenland?

by

M. Stein

Institut für Seefischerei, Bundesforschungsanstalt für Fischerei  
Palmaille 9, D-22767 Hamburg, Federal Republic of Germany

and

J. Llorc \*)

\*) Present affiliation: Institut für Seefischerei, Bundesforschungsanstalt für Fischerei  
Palmaille 9, D-22767 Hamburg, Federal Republic of Germany

**ABSTRACT**

Based upon the hypothesis of MEYER (1968) that periods of low stability of the water column in autumn should parallel good cod year classes in the following year, the paper analyses available time series of cod recruitment, subsurface oceanographic stability data and wind stress data off West Greenland. Correlation analysis yields no significant results between stability, wind stress and recruitment when considering the *entire length of available time series*. Instead, splitting of the autumn based oceanographic data set into a pre-seventies part (when warm climatic conditions and high recruitment prevailed), and a recent part (for the past twenty years of cooling climate) yields significant coherence of recruitment and environmental conditions of the cold climate period. For the warm climate period only occasional consistency is shown. The possible influence of the Great Salinity Anomaly on recruitment of cod is discussed. It is suggested that during the past twenty years a new correlation mechanism, an advective coupling instead of a low stability coupling explains the variability of recruitment from the environmental point of view.

**INTRODUCTION**

The fluctuations of several cod stock fisheries, e.g. Baltic, Arcto-Norwegian, Icelandic, West and East Greenland, and Canadian Atlantic cod stocks, are among the best documented around the world (HERMANN and HORSTED, 1976; BJORKE and SUNDBY, 1984; KOSIOR and NETZEL 1989; ROSE and LEGETT, 1989; HOVGARD and BUCH, 1990; SUNDBY and FOSSUM, 1990; JAKOBSSON, 1992). The intensity of the fluctuation has been different from one stock to another but they all have suffered great fluctuations over the decades. Considering the economical and social importance of cod fisheries, much investigation has been conducted into this problem, specially on the recruitment variability. As pointed out by KOSIOR and NETZEL (1989), "the abundance of a cod year class is only slightly affected by the abundance of its spawning stock and the quality of reproducers. What is important is the survival rate of eggs, larvae and juveniles". Among various physical processes affecting cod recruitment, larval and egg transport from the spawning grounds to their nurseries, temperature, salinity and turbulence are the most important. From the biological processes, predation and food availability are the dominant controlling

mechanisms for recruitment success. Climate appears as a key to link those biological and physical processes. At West Greenland cod occurs at one of its northern limits (HANSEN and HERMANN, 1965). This stock is sustained not only by his own reproduction in the waters off West Greenland and in local inshore fjords, but also by the eggs and larvae advected from cod spawning sites at Iceland (Icelandic cod stock). The main feature of that West Greenland cod stock related to the fisheries is that both abundance and spatial distribution have shown considerable changes in historical time. The large variations are reflected in the annual yield which have varied by a factor of  $10^2$  within the present century (BUCH and HANSEN, 1988). The climate at West Greenland has also undergone very marked changes (STEIN and MESSTORFF, 1990). These changes in the West Greenland climate are in part due to variations of the main current components of the West Greenland Current system, and due to air temperature variations. Periods of warm water temperatures correlate with air temperature as demonstrated by STEIN and BUCH (1991). These periods coincide with periods of high cod abundance. The general warming of the northern hemisphere around 1920 evidently lead to the establishment of a self sustaining and very abundant West Greenland cod stock which through the 1930s-60s produced good year classes at relatively short intervals. But since the late 1960s the West Greenland cod stock has not produced any good year classes and fisheries have dwindled to nearly 100%. At present, cod abundance is at a very low level, with the 1992 catches being the lowest experienced since 1925. The effect of the climate changes in West Greenland on cod stocks appears to be evident but the way those climate changes have affected and affect the size of the year-classes is poorly understood.

DICKSON and BRANDER (1993) suggested that the intensification of the Irminger current during warm periods increased the drift of cod larvae from the spawning grounds off south-west Iceland, thus leading to strong year-classes off West Greenland. They also pointed at the possibility that cod year classes which survive the drift to Greenland as larvae may later return as adults to Iceland. CUSHING (1988, 1990a,b) concludes that there is an impact of the Great Salinity Anomaly (GSA, DICKSON et al., 1988) on Northwest Atlantic cod stocks. Accordingly, poor recruitment was observed due to low zooplankton abundance or due to mismatch of cod larvae and zooplankton abundance peak. MERTZ and MYERS (1994) indicate that cod recruitment off West Greenland showed a reduction in the peak GSA years. MEYER (1968) speculated that the amount of food for cod larvae (which hatch in April) in the waters off West Greenland, might be closely related to the quantity of nutrients in the surface layer at the time of the plankton bloom, which in turn may be dependent on the effectiveness of the vertical convection process during the previous winter. This vertical convection depends on two factors:

- i) the stability (E) of the water column during autumn, and
- ii) the wind intensity during winter.

To examine the hypothesis of MEYER (1968) mean stability (E) at standard oceanographic depths down to 200m at 5 NAFO Standard Oceanographic Stations off West Greenland in autumn was calculated and correlated with recruitment data series, as well as with wind stress data.

## MATERIALS AND METHODS

The Standard Oceanographic Stations (STEIN, 1988) chosen to calculate the stability are given in Fig. 1: Cape Farewell 4 (CF4: 59°00'N 45°20'W), Cape Desolation 3 (CD3: 60°28'N 50°00'W), Frederikshab 3 (FR3: 61°47'N 51°09'W), Frederikshab 4 (FR4: 61°41'N 51°45'W), and Fylla Bank 4 (FY4: 63°53'N 53°22'W). The formula used to

calculate the stability was:

$$E = 1 / \rho * dp / dz$$

where  $\rho$  is the mean density of individual layers and  $dp / dz$  is the rate of change in density with depth. Temperature and salinity series to calculate the stability were available from own measurements of the Institut für Seefischerei, Hamburg and the Greenland Fisheries Institute, Copenhagen. Those series have been collected nearly every year in July and November or December since 1950 (July-data, FY4), 1963 (Autumn-data, FY4) and since 1982 (for the other stations). It has to be considered that the understanding of the dynamics of a fish population in relation to climate and fisheries, requires long time series such as those presented from Fylla Bank which cover low and high exploitation under different environmental conditions.

Year class size age 3 data were taken from literature and cover all the West Greenland area. The year class size age 3 data from 1953 to 1979 had been estimated by Virtual Population Analysis (VPA, HANSEN and BUCH, 1986), and the year class size age 3 data from 1982 to 1992 had been calculated proportionally using age compositions from East Greenland surveys (ANON, 1984, 1988). With the autumn oceanographic data, a correlation between the year class strength and the water mass stability of the year preceeding the year class, was calculated for standard oceanographic depths. To check the accuracy of the derived stability data, correlation analysis was done between the initial parameters, temperature anomaly, salinity anomaly, and year class strength, and between temperature anomaly, salinity anomaly and stability.

Other climatic data used in this paper were air temperatures from Nuuk (64°11'N, 51°44.5'W), and estimates of seasonal surface averages of the wind stress for the area (c.f. Fig. 1). From the air temperature data the annual mean from 1950 to 1993 relative to the climatological mean (1961-90) was calculated (Fig. 2). Wind data are available from literature and as ASCII-File (DRINKWATER and PETTIPAS, 1993). These data, U- and V-components of the wind stress vector, cover the period from 1946 to 1991. The stress direction was the wind direction after adjustment for frictional effects, plus 180° to convert it into oceanographic convention, i.e. the direction towards which the wind blows. Further details are given in (DRINKWATER and PETTIPAS, 1993). Seasonal averages were calculated taking December to February as winter, March to May as spring, June to August as summer, and September to November as autumn. These seasonal means were correlated with the stability data at stations FY4, FR3, FR4, CD3 and CF4, and with the year-class strength data. For the July-data (ICES data), the 1953-1979 stability of FY4 was computed at standard oceanographic depths and correlated with the year class of the same year, as well as with U, V wind stress components.

All correlations were calculated with the programme QUATTRO PRO for Windows, Version 5.0.

## RESULTS and DISCUSSION

### Climatic Background

The climatic conditions of the West Greenland area are given in Fig. 2, based upon the air temperature anomalies during the past 43 years. Accordingly, the climate of the region was characterized during the fifties and sixties by warmer than normal air temperatures, which exceeded the climatic mean (1961-90: -1.4°C) by 1 to 2K. The downward trend in mean air temperatures, as experienced from the late sixties onwards, was dominated by anomalous cold events during the beginning of the seventies, eighties and nineties. During

the mid seventies and eighties warmer than normal events were recorded. Since 1989, mean annual air temperature anomalies were well below normal.

Wind stress components, derived from the geostrophic wind field at selected sites in the Labrador Sea (DRINKWATER and PETTIPAS, 1993) reveal characteristic differences for the winter and spring season, and for the summer and autumn season (Fig. 3). The wind stress differs by about one order of magnitude from winter to summer, and is about constant during spring and autumn. The anomalous event during 1983 (upper panels of Fig. 3) did only emerge during the first two seasons. This event, which is unique throughout the 45 years of record, deviates by a factor of about five to six from the normal level, indicating stronger than normal winds into a southeasterly direction.

Stability of the water masses at Station 4 of the Fylla Bank during the past thirty years is displayed for the surface layer 0-50m (Fig. 4), and for the sub-surface layer 50-200m (Fig. 5). It would appear that there is an increase in surface layer stability from the mid eighties onwards, reflecting the presence of diluted water masses (see below).

Temperature and salinity anomalies were found to be correlated positive, indicating that cold, diluted water as of East Greenland Current origin alternates with more saline, warmer oceanic water (Fig. 6). Stability correlated negatively with the salinity anomaly at all stations. Comparing stability and temperature anomaly, only two stations (CF4 and FR3) indicated negative correlations; the rest of the stations (FY4, CD3 and FR4) did not give any correlation. Thus, salinity as advected by the oceanic Irminger component of the West Greenland Current system, has the dominating influence on the stability.

#### Impact on Cod Recruitment

Is there an impact of water mass stability on the recruitment of cod? Does wind stress influence recruitment? Correlation analysis performed for the available sub-surface ocean time series (CF4, CD3, FR3, F4, FY4), wind stress data and recruitment data off West Greenland reveal no significant dependence between stability and recruitment, as well as between wind stress components and recruitment. Only trends are detectable.

Fylla Bank Station 4 July stability data compared to recruitment data derived from VPA (HANSEN and BUCH, 1986) for the years 1953 to 1979, yield positive but not significant correlation ( $r = 0.45$ ,  $p > 0.01$ ). The results (Fig. 7) corroborate in part the theory that low stability favours good recruitment, and vice-versa (MEYER, 1968). However, despite low stability levels from 1974 onwards, no significant recruitment has been encountered thereafter.

A composite of two recruitment time series (from HANSEN and BUCH, 1986 and ANON., 1984) was used to compare recruitment and stability until the end of the eighties (Fig. 8). Low recruitment during the eighties, except for the strong 1984 year class and the good 1985 year class, characterizes the last decade.

From the late sixties onwards, the climatic conditions of West Greenland were dominated by negative air temperature anomalies (Fig. 2). Stability of the surface layers increased during the seventies with the trend still persisting (Fig. 4). The general coincidence of the pre-seventies recruitment curves with water mass stability (c.f. also MERTZ and MYERS, 1994 for the West Greenland cod stock) suggests a splitting of the time series. Before 1970, West Greenland cod stocks were self sustaining and with high spawning stock biomass based on several good recruitment events (1953, 1957, 1960, 1961 to indicate the strong year classes). These recruitment events happened during warm climatic conditions at lower than average water mass stability. During this period, the model assumption as anticipated by (MEYER, 1968) holds quite well. However, when climate started to be colder than normal, and the West Greenland cod stocks had undergone heavily exploitation, the recruitment/stability relation was not any longer valid. There has been a period of low recruitment from 1969 to 1972 (Fig. 7) which MERTZ and MYERS (1994) attribute to the passing of the Great Salinity Anomaly (GSA). Our

analysis reveals, however, that there is only one high stability event (1969 in Fig. 4) in the surface layer 0-50m which they referred to. In the deeper layer, 50-200m (Fig. 5), the years 1968 and 1969 emerge as high stability events. Since our data are the autumn data previous to the considered year classes, recruitment of 1969 and 1970 cod thus could have been influenced by the GSA. Fig. 9 displays year class strength and water mass stability of the previous autumn at Fylla Bank station 4. Visual inspection reveals high consistency of both curves. Correlation analysis yields significant results ( $r = 0.69$ ,  $p = 0.0018$ ). West Greenland cod recruitment was at a very low level during the past two decades. Considerable recruitment injections from Iceland/East Greenland were observed during 1973 and 1984. These recruitment peaks parallel high stability events (Fig. 9). As discussed above, salinity has the dominating influence on the stability of water masses. Thus, high stability should coincide with the advection of low saline water. The thermohaline properties of the 150-200m layer at station 4 of Fylla Bank are influenced by the Irminger component of the West Greenland Current system. The significant positive correlation of recruitment and stability of this layer - with no significant correlations observed in the upper layers - suggests a coupling of recruitment and changes in the inflow of Irminger (Atlantic) water to the West Greenland area. It would appear that during the past twenty years a new correlation mechanism, an advective coupling instead of a low stability coupling explains the variability of recruitment from the environmental point of view.

## References

- ANON., 1984. Report of the Working Group on Cod Stocks off East Greenland. ICES, Doc. C.M.1984/Assess:5.
- ANON., 1988. Report of the Working Group on Cod Stocks off East Greenland. ICES, Doc. C.M.1988/Assess:11.
- BJORKE, H. and S. SUNDBY 1984. Distribution and abundance of post larval Northeast Arctic cod and haddock (Reproduction and Recruitment of Arctic Cod). *Raspredelenie i chislennost' postlarval' noj severo-vostochnoj arkticheskoy treski i pikshi (Vosproizvodstvo i Popolnenie Treski)* S.A.Studenetskij (Ed.): 72-94.
- BUCH, E. and H.H. HANSEN 1988. Climate and cod fishery at West Greenland. pp. 345-364. In: Wyatt and Larraneta (Eds.), *Long-term Changes in Marine Fish Populations*, Proc. Vigo Symposium, November 1986.
- CUSHING, D.H. (1988) The northerly wind. In: *Toward a Theory of Biological-Physical Interactions in the World Ocean*. B.J. Rothschild (ed.). Kluwer, Dordrecht: NATO Advanced Research Workshop, 239, pp. 235-244.
- CUSHING, D.H. (1990a) Plankton production and year-class strength in fish populations: an update of the match/mismatch hypothesis. *Adv. Mar. Biol.* 26:249-293.
- CUSHING, D.H. (1990b) Recent studies on long-term changes in the sea. *Freshwater Biol.* 23:71-84.
- DICKSON, R.R. and K.M. BRANDER, 1993. Effects of changing windfield on cod stocks of the North Atlantic. ICES 1993/CCC Symposium/No. 16A.
- DICKSON, R.R., J. MEINCKE, S.-A. MALMBERG and A.J. LEE 1988. The "Great Salinity Anomaly" in the Northern North Atlantic 1968-1982. *Prog. Oceanog.* (20): 103-151.
- DRINKWATER, K.F. and E.G. PETTIPAS, 1993. Climate Data for the Northwest Atlantic: Surface Wind Stresses off Eastern Canada, 1946-1991. *Can. Data Rep. Hydrogr. Ocean Sci.* 123:iv + 139p.
- HANSEN, H. and E. BUCH, 1986. Prediction of Year-class Strength of Atlantic Cod (*Gadus morhua*) off West Greenland. *NAFO Sci. Coun. Studies*, 10:7-11.
- HANSEN, P.M. and F. HERMANN, 1965. Effect of Long-Term Temperature Trends on Occurrence of Cod at West Greenland. ICNAF Environmental Symposium, ICNAF Special Publication No. 6:817-819.

**HERMANN, F. and Sv.A. HORSTED 1976.** Fluctuations in the cod stock and the ocean climate at West Greenland. Book of Abstracts of papers presented at Joint Oceanographic Assembly, Edinburg UK, 13-24 September 1976; FAO, Rome (Italy), S.7, 52.

**HOVGARD, H. and E. BUCH 1990.** Fluctuation in the Cod Biomass of the West Greenland Sea Ecosystem in Relation to Climate. In: Large Marine Ecosystems (Ed. K. Sherman, L.M. Alexander and B.D. Gold) AAAS Selected Symposium.

**KOSIOR, M. and J. NETZEL 1989.** Eastern Baltic Cod stocks and environmental conditions. Rapp. P.-V. Réun.Cons.int.Explor. Mer 190: 159-162.

**JAKOBSSON, J. 1992.** Recent variability in the fisheries of the North Atlantic. In: ICES mar. Sci. Symp., 195: 291-315.

**MERTZ, G. and R.A.MYERS 1994.** The ecological impact of the Great Salinity Anomaly in the northern North-west Atlantic. Fish. Oceanogr. 3:1, 1-14.

**MEYER, A. 1968.** 1967 ein erfolgreiches Jahr für die deutsche Grönlandfischerei. Hansa., 16: 2-4.


**ROSE, G.A. and W.C. LEGETT 1989.** Interactive effects of geophysically-forced sea temperatures and prey abundance on mesoscale coastal distributions of a marine predator, Atlantic cod (*Gadus morhua*). Can.J.Fish.Aquat.Sci. 46(11):1904-1913.

**STEIN, M. and J. MESSTORFF 1990.** Relationship of Fluctuations in Cod Recruitments off West Greenland to Long-term Variations of the Physical Environment. NAFO Sci.Coun.Studies, 14: 39-44.

**STEIN, M. 1988.** Revision of list of NAFO standard oceanographic sections and stations. NAFO SCR Doc. 88/01: 1-9.

**STEIN, M. and E. BUCH 1991.** Are Subsurface Temperatures Predictable at Fylla Bank, West Greenland? NAFO Sci.Coun.Studies, 15: 25-30.

**SUNDBY, S. and P. FOSSUM 1990.** Feeding conditions of Arcto-norwegian cod larvae compared with the Rothschild-Osborn theory on small-scale turbulence and plankton contact rates. J.Plankton Res. 12(6): 1153-1162.



**Fig. 1** Location of sites used for correlation analysis of available time series




Fig. 2 Annual Mean Air Temperature Anomaly at Nuuk (rel. 1961-1990 climatic mean)




Fig. 3a Wind Stress components [pa] Site 03: Winter (1947-1991), Spring (1946-1991)









Fig. 3b Wind Stress components [pa] Site 03: Summer (1946-1991), Autumn (1946-1991)




Fig. 4 Stability of Surface layer 0-50m Fylla Bank Station 4 Autumn




Fig. 5 Stability of Deep layer 50-200m Fylla Bank Station 4 Autumn

# Correlations Between Temperature Anomaly and Salinity Anomaly CF4, CD3, FR3, FR4, FY4




Fig. 6 Correlation between temperature Anomaly and Salinity anomaly at oceanographic sites CF4, CD3, FR3, FR4, FY4




Fig. 7 Year class strength of cod (black boxes, 1953-1979) and Stability of the near-surface layer 10-20m at Station 4 of Fylla Bank during July




Fig. 8 Year class strength (lower curve, 1964-1989) and Stability of the deep layer 100-150m at Station 4 of Fylla Bank during autumn of the year before hatching




Fig. 9 Year class strength (lower curve, 1973-1989) and Stability of the deep layer 150-200m at Station 4 of Fylla Bank during autumn of the year before hatching