

Northwest Atlantic



Fisheries Organization

Serial No. N2541

NAFO SCR Doc. 95/32

SCIENTIFIC COUNCIL MEETING - JUNE 1995

Climatic Conditions Around Greenland - 1994

by

M. Stein

Institut für Seefischerei, Palmaille 9
D-22767 Hamburg, Federal Republic of Germany

Abstract

Based on air temperature data from three sites of West and East Greenland, the annual variability of climate is shown. Mean monthly air temperature data from Nuuk/West Greenland reveal the long-term interannual changes of air temperature anomalies. It is shown that on the seasonal basis, variation of mean amplitudes differs most when considering **winter** (JFM) and **summer** (JAS) data sub-sets. As a consequence of warm summer and autumn air temperatures, the **sea ice** cover is **less** expressed than in previous years of this decade. Subsurface observations on Fylla Bank/West Greenland, indicate that during the past 30 years **cold** and **diluted** events occurred **simultaneously**. This points at the **advective influence** of the **East Greenland Current** component on the water mass properties of the West Greenland Bank and slope area. The **vertical magnitude** of influence of this Polar Water amounts to **150m to 200m**. Warmer than normal subsurface conditions are attributed to a northerly extension of the North Atlantic Current.

Introduction

The climate at West Greenland has undergone very marked changes (**BUCH and STEIN, 1989; STEIN and MESSTORFF, 1990**). These changes in the West Greenland climate are in part due to variations of the main current components of the West Greenland Current system, and due to air temperature variations. Periods of warm water temperatures correlate with air temperature as demonstrated by **STEIN and BUCH (1991)**. The necessity of collecting oceanographic data at comparable standard stations has been demonstrated recently (**STEIN, 1990**). As an example of systematic sampling and reporting on the results in the Northwest Atlantic, the publications by **DRINKWATER and TRITES (1993)**, and **DRINKWATER et al. (1994)** might be taken. Similarly, a closer view at the climatic conditions of the Greenland area deemed necessary. Started as a project during 1993, this paper continues the series which provides an annual overview on environmental conditions around Greenland. The data used in the context of this paper, deal with data on air temperatures, the distribution of sea ice and subsurface observations. The latter data originate from oceanographic

observations performed at Standard Oceanographic Stations (STEIN, 1988) by the Federal Republic of Germany during the 1994 annual groundfish survey in the area off West and East Greenland. It has been shown that the availability of long-term time series enables a better view on the variability of the ocean, especially when correlating variability in the biota and the environment (STEIN and LLORET, 1995).

In this paper the available climatic data for 1994 off Greenland are presented. To enable a closer view on thermohaline conditions off West Greenland, isopleth diagrams from Fylla Bank are presented and discussed.

Data and Methods

Data on the atmospheric climate of Greenland were sampled by the Danish Meteorological Institute at Nuuk (64°11'N, 51°44.5'W), Egedesminde (68°42.5'N, 52°53'W) and Angmagssalik (65°36'N, 37°40'W). Whereas the first data set was mutually supplied by the Danish Meteorological Institute in Copenhagen and the Seewetteramt, Hamburg, the latter data sets are taken from ANON. (1994). The climatic mean which the air temperature anomaly charts are referenced to is 1961-1990. Ice charts were constructed from NOAA satellite ice charts. Anomalies of ice edge are referenced to sea ice normals as displayed by BUCH and STEIN (1989). The approximate location of the ice edge is given in the selected figures and in a computer slide show. The temperature anomaly maps for the Northwest Atlantic were also taken from ANON. (1994). Presently, only the ice charts are available from the author on request as computer slide shows. Subsurface ocean data are available from German measurements for the West Greenland area.

For graphical display of isopleth diagrams and sections the SURFER Software (Version 4) was used.

Results

Air Temperature and Sea Ice Anomaly during 1994

Northwest Atlantic

As in previous years (STEIN, 1995a, b), air temperatures over West Greenland during the first quarter of 1994 were colder than normal. Especially the month of March (Fig. 2a) showed the typical cold air mass centered over the town of Egedesminde. Anomalies at this site were less than 6K. Warmer than normal conditions were encountered in the month of July at Egedesminde and Angmagssalik, whereas Nuuk had summer temperatures below the norm (Fig. 2b).

Air Temperatures and Climatic Means

Mean monthly air temperatures at the three selected sites of the Greenlandic coast vary from about -15°C to +5°C at Egedesminde, -7.5°C to +7°C at Nuuk, and -7.5°C to +7.5°C at Angmagssalik (Figs. 3 to 5). Although Nuuk and Angmagssalik have about the same mean conditions, year-to-year changes, as well as seasonal changes differ significantly from West to East Greenland. Colder than normal conditions during January to March at the west coast, contrast with a colder than normal January, a warmer than normal February, and an about normal March at the east coast. From April onwards, Egedesminde air temperatures were above or at mean conditions until

December when winter cooling started. An outlook for 1995 indicates winter air temperatures even colder than the 1994 conditions. January to March revealed temperatures around -22°C . Nuuk experienced a colder than normal summer during 1994, a normal autumn, and winter cooling from December onwards. Similar to Egedesminde, Nuuk air temperatures during January to March 1995 were well below the 1994 conditions. At Angmagssalik about normal air temperatures were encountered from February to September. October and December were well below the norm, whereas November was about normal. During the first quarter of 1995, air temperatures were colder than during 1994.

Climatic Variability off West Greenland

After the cold years 1992 and 1993, 1994 indicated an increase of the annual mean air temperature. Climatic conditions at Nuuk are, however, still well below the norm (Figs. 6). As indicated by STEIN (1995a), the low frequency variation as given by the thick line in Fig. 6, points at longer than decadal variations of the mean air temperatures at Nuuk. Splitting of the air temperature anomalies data set in four monthly subsets, the means of January to March (JFM), April to June (AMJ), July to September (JAS), and October to December (OND), reveal different amplitudes of warming and cooling, as well as different low frequency variations (Fig. 7a-d). Largest variation in air temperatures is observed during the winter season, ranging from -2K to $+2\text{K}$ (Fig. 7a), variability during summer (JAS, Fig. 7c) is smallest. Steep downward trends after the warm period between the 20's and 60's are visible in the first two seasons (Fig. 7a, b), and a **surprisingly long warm period** which extends far into the 80's is encountered in the **autumn data** (OND, Fig. 7d). The warm periods in the seasonally splitted data sets were from 1918 to 1975 (JFM), from 1919 to 1964 (AMJ), 1918 to 1965 (JAS), and 1917 to 1985 (OND2). With the exception of the autumn season, the warm period data are in the range of previously published material (HOVGARD and BUCH, 1990; STEIN, 1995a,b).

Ice Conditions in the Northwest Atlantic

Two examples of sea ice conditions during 1994 are given in Figs. 8 and 9. The largest extension of the ice edge was encountered during February with anomalous presence of ice in the vicinity of Nuuk and in Julianehaab Bight/Southwest Greenland (Fig. 8). As visible from the individual ice maps which are plotted on 14 day's interval (see Data and Methods), the year 1994 was a warm year as concerns sea ice distribution around Greenland. Unusual cover of sea ice was observed during mid July at Cape Farewell, and during the first half of August around Baffin Island. Ice-free conditions were observed between mid of September and mid of October when ice started to form again off East Greenland and around Baffin Island.

Subsurface Observations off West Greenland

Thermohaline conditions along the Fylla Bank Section (STEIN, 1988) are given in Figs. 10a and 10b. The vertical field of temperature reveals warm conditions in the depth range between 100m and 200m with temperatures around 6°C . T,S-properties of the Fylla Bank Section are given for standard depth data in Fig. 11. Temperature and salinity anomalies for Fylla Bank Station 4 are given in Figs. 12 and 13 for the depth ranges 0-50m, 0-200m,

and 200-300m. Based on standard depth data, the temporal variation between 1963 and 1994 (autumn) of thermohaline changes of the upper 300m at Station 4 of the Fylla Bank Section are displayed in **Figs. 14 and 15**.

Discussion

Generally, near surface mean air temperatures were about 0.4K warmer than normal for the area 85°N to 5°N, 110°W to 70°E (ANON, 1995). From 1990 to 1994 these means amounted to +0.45K, +0.35K, -0.12K, +0.07K, and 0.38K. 1992 conditions were influenced by the Pinatubo-Effect (HENNING, 1994). Changes to positive anomalies occurred mainly over large areas of North America, Great Britain, the Iberian Peninsula/Morocco, Eastern Europe. Increasing warming tendencies which surmounted the 1993 anomalies, were encountered over Northern to Southern Europe, Northwest Africa. Except for the region off East Greenland (**Fig. 1**), air temperatures over the Northwest Atlantic were colder than normal. The negative anomalies over Baffin Bay/Davis Strait and Greenland, however, weakened as compared to previous years.

At Greenland, air temperature conditions during 1994 were similar to 1992 and 1993 as concerns the cold winter. From April onwards, however, temperatures increased to above normal conditions (**Figs. 3 to 5**), and led to mostly warmer than normal summer temperatures. This gave rise to fairly good sea ice conditions. From 27 April onwards, the west coast of Greenland was ice free, a month later Disko Bight was open. Anomalous ice conditions were encountered at Cape Farewell during June and July.

Comparison of monthly mean air temperature time series at Nuuk indicates different amplitudes of warming/cooling when considered on quarter year basis. Winter (JFM) amplitudes range from -2K to +2K, summer (JAS) amplitudes range from -0.5K to +1K. The analysis reveals a **surprisingly long warm period** which extends far into the 80's when considering the **autumn data** (OND, **Fig. 7d**).

Warm water off West Greenland might be derived by **two ways** (BUCH, 1994): Via the Irminger Current component of the West Greenland Current system; and from the northern part of the North Atlantic Current. The latter water mass originates from the area south to southwest of Cape Farewell, has temperatures above 4.5°C, and salinities between 34.5 and 34.9 PSU. It is observed during autumn off the shelf slope of the West Greenland banks at depths below 100m (**Fig. 10a**).

Thermohaline conditions during autumn 1994 indicate positive anomalies which are largest in the near surface layer (0-50m, **Figs. 12 and 13**). After the very cold year 1992, the years 1993 and 1994 might point at intermediate warming as anticipated recently by STEIN (1995). Isopleth diagrams of temperature and salinity (**Figs. 14 and 15**) reveal cooling and intermediate warming at station 4 of the Fylla Bank section throughout the last thirty years. The data show that cooling is accompanied by tremendous dilution of the water column, with the depth of the signal being the same for both parameters, temperature and salinity. Since these events are concurrent, the mode of formation must be of advective nature. In contrast to previous discussions, the coincidence of cold and diluted events points at changes due to the East Greenland Current component, which has a larger influence on the thermohaline properties at Fylla Bank during autumn than anticipated (e.g. BUCH and STEIN, 1989; STEIN and BUCH, 1991). Taking the autumn

T,S-characteristics of East Greenland Polar Water (3 to 5°C, < 34.4 PSU; **BUCH, 1994**), the 34.25 PSU isoline in **Fig. 15** roughly delineates the lower boundary of this water mass at Fylla Bank station 4 throughout the past thirty years. The vertical extension of this Polar Water varies between 150m and 200m.

References

- ANON., 1994.** Die Witterung in Übersee. Vol. 1-8.
- ANON., 1995.** Beilage zur Wetterkarte des Deutschen Wetterdienstes. 31/1995, 2p.
- BUCH, E., M. STEIN 1989.** Environmental Conditions off West Greenland, 1980-85. J.Northw.Atl.Fish.Sci., Vol. 9: 81-89.
- BUCH, E. 1994.** A Monograph on the Physical Oceanography of the Greenland Waters. Royal Danish Administration of Navigation and Hydrography, Copenhagen, Denmark, 405 p.
- DRINKWATER, K.F., and R.W. TRITES 1993.** Overview of Environmental Conditions in the Northwest Atlantic in 1990. NAFO Sci.Coun. Studies, 19: 15-29.
- DRINKWATER, K.F., B. PETRIE and S. NARAYANAN 1994.** Overview of Environmental Conditions in the Northwest Atlantic in 1991. NAFO Sci.Coun. Studies, 20: 19-46.
- HENNING, D. 1994.** Anomalien der Lufttemperatur im Raum Nordamerika, Nordatlantik, Europa, Nordafrika für das Jahr 1993. Der Wetterlotse 565: 9-10.
- HOVGARD, H. and E. BUCH 1990.** Fluctuation in the Cod Biomass of the West Greenland Sea Ecosystem in Relation to Climate. In: Large Marine Ecosystems (Ed. K. Sherman, L.M. Alexander and B.D. Gold) AAAS Selected Symposium.
- STEIN, M. 1988.** Revision of list of NAFO standard oceanographic sections and stations. NAFO SCR Doc., No. 1, Serial No. N1432, 9p.
- STEIN, M. 1990.** Some Remarks On Time-Series Sampled On Annual Intervals. NAFO SCR Doc. 90/16: 1-6.
- STEIN, M. and E. BUCH 1991.** Are Subsurface Temperatures Predictable at Fylla Bank, West Greenland?. NAFO Sci.Coun.Studies, 15: 25-30.
- STEIN, M. and J. MESSTORFF 1990.** Relationship of Fluctuations in Cod Recruitments off West Greenland to Long-term Variations of the Physical Environment. NAFO Sci.Coun.Studies, 14: 39-44.
- STEIN, M. 1995a.** Climatic Conditions around Greenland - 1992. NAFO Sci.Coun.Studies, 22: (in press).
- STEIN, M. 1995b.** Climatic Conditions around Greenland - 1993. NAFO Sci.Coun.Studies, 22: (in press).
- STEIN, M. and J. LLORET 1995.** Stability Of Water Masses - Impact On Cod Recruitment Off West Greenland? Fishery Oceanography 4, 2 (in press).




Fig. 1 Mean air temperature anomalies over the Northwest Atlantic during 1994 (E=Egedesminde, N=Nuuk, A=Angmagssalik)




Fig. 2a Mean air temperature anomalies over the Northwest Atlantic during March, 1994 (E=Egedesminde, N=Nuuk, A=Angmagssalik)




Fig. 2b Mean air temperature anomalies over the Northwest Atlantic during July, 1994 (E=Egedesminde, N=Nuuk, A=Angmagssalik)




Fig. 3 Monthly mean air temperature at Egedesminde during 1994 and climatic mean (1961-1990)




Fig. 4 Monthly mean air temperature at Nuuk during 1994 and climatic mean (1961-1990)




Fig. 5 Monthly mean air temperature at Angmagssalik during 1994 and climatic mean (1961-1990)




Fig. 6 Time series of annual mean air temperature anomalies at Nuuk (1880-1994, rel. 1876-1994) and 13 year running mean




Fig. 7 Time series of three monthly mean air temperature anomalies at Nuuk (1880-1994, rel. 1876-1994)
a) January, February, March (JFM)
b) April, May, June (AMJ)
c) July, August, September (JAS)
d) October, November, December (OND)




Fig. 8 Ice edge during March 23, 1994; dark shaded areas indicate anomalous extent of ice edge during the month of February





Fig. 9 Ice edge during September 28, 1994

FYLLA BANK T [C] 941101



FYLLA BANK Salinity [PSU] 941101




Fig. 10 Vertical distribution of temperature (a) and salinity (b) along the Fylla Bank Section during November 1, 1994




Fig. 11 TS-diagram of Fylla Bank Section (Standard depth data are used)




Fig. 12 Fylla Bank Station 4: Temperature Anomaly 0-50m, 0-200m, 200-300m




Fig. 13 Fylla Bank Station 4: Salinity Anomaly 0-50m, 0-200m, 200-300m

Fylla Bank Station 4 Temperature [C]




Fig. 14 Fylla Bank Station 4: Isopleth diagram of Temperature, Depth, Year

Fylla Bank Station 4 Salinity [PSU]




Fig. 15 Fylla Bank Station 4: Isopleth diagram of Salinity, Depth, Year