

Northwest Atlantic



Fisheries Organization

Serial No. N2816

NAFO SCS Doc. 96/18


SCIENTIFIC COUNCIL MEETING - NOVEMBER 1996

Report of Scientific Council

Special Meeting, 15-18 November 1996

CONTENTS

	<u>Page</u>
I. Plenary Sessions	1
II. Fishery Science	1
III. Formulation of Advice	1
Summary Sheets:	
- Shrimp in Subareas 0 and 1	2
- Shrimp in Denmark Strait	3
IV. Other Matters	4
V. Adoption of Reports	4
VI. Adjournment	4
 APPENDIX I. Report of Standing Committee on Fishery Science (STACFIS)	 5
I. Stock Assessments	5
1. Shrimp in Subareas 0 and 1	5
2. Shrimp in Denmark Strait	9
3. Other Business	12
 APPENDIX II. Agenda Scientific Council - 15-18 November 1996	 13
Annex 1. Canadian Request for Scientific Advice on Management in 1997 of Certain Stocks in Subareas 0 to 4	 14
Annex 2. Denmark (Greenland) Request for Scientific Advice on Management of Certain Stocks in 1997	 15
 APPENDIX III. List of Research and Summary Documents - November 1996	 17
 APPENDIX IV. List of Representatives, Advisers/Experts - November 1996	 19



Participants of Special Meeting, 15-18 November 1996



Standing: A. Nicolajsen, O. Folmer, D. M. Carlsson, L. Savard, H. Powles, D. G. Parsons, T. Amaratunga

Seated: W. B. Brodie, H. Siegstad, U. Skúladóttir, W. R. Bowering, C. Hvingel



REPORT OF SCIENTIFIC COUNCIL

Special Meeting, 15-18 November 1996

Chairman: W. R. Bowering

Rapporteur: T. Amaratunga

I. PLENARY SESSIONS

The Scientific Council met at NAFO Headquarters, Dartmouth, Nova Scotia, Canada, during 15-18 November 1996. Representatives attended from Canada, Denmark (in respect of Faroe Islands and Greenland) and Iceland. The Executive Secretary and Assistant Executive Secretary were in attendance.

The opening session was called to order on 15 November 1996 at 1100 hr.

The Chairman, W. R. Bowering (Canada), welcomed representatives to this Special Meeting of the Scientific Council to conduct assessments on shrimp in Subareas 0 and 1, and Denmark Strait. A special appreciation was extended to W. B. Brodie (Canada) who kindly undertook the task of chairing the STACFIS sessions of this meeting, after his tenure of STACFIS chairmanship had ended in September 1996. The Assistant Executive Secretary was appointed rapporteur. The Provisional Agenda was **adopted** (Appendix II).

The Council noted that STACFIS would undertake the assessments of the stocks, while the prognoses and the advice would be undertaken by the Council.

The session was adjourned at 1110 hr.

The concluding sessions were convened on 18 November 1996, noting that the shrimp assessment reports had been prepared by STACFIS. The Council then addressed the requests of the Coastal States and considering the results of the assessments provided advice and recommendations. The meeting was adjourned at 1708 hr.

Summary reports of the assessments and other matters considered by the Scientific Council are given below in Sections II-IV. The Agenda, List of Research (SCR) and Summary (SCS) Documents, and the List of Participants of this meeting are given in Appendix II, III and IV, respectively.

II. FISHERY SCIENCE (see STACFIS report, App. I)

The Council noted that matters referred to STACFIS relating to assessments of shrimp in Subareas 0 and 1 and Shrimp in Denmark Strait were addressed. The complete reports are given in Appendix I.

III. FORMULATION OF ADVICE

The Council reviewed the STACFIS assessments of shrimp in Subareas 0 and 1, and Denmark Strait and the agreed summaries are as follows:


Shrimp in Subareas 0 and 1

Background: A small scale inshore fishery began in SA 1 during the 1930s. Since 1969 an offshore fishery has developed and the shrimp fishery is now the largest in Davis Strait.



Fishery and catches: The fishery is conducted mainly by Greenland and Canada. Recent catches from the stock are as follows:

Year	('000 tons)	
	Inshore	Offshore
1990	13.6	55.7
1991	16.3	59.6
1992	20.6	66.2
1993 ¹	17.8	57.8
1994 ¹	18.1	58.5
1995 ¹	16.4	54.3
1996 ¹ (to Oct)	10.5	44.6

¹ Provisional.



Biomass: Survey biomass indices indicate a relatively stable stock size from 1988 to 1996.



Data: Catch, effort and biological sampling data were available from the offshore fishery, and catch and effort data from the inshore fleet. Time series of biomass indices and stock composition data were available from research surveys from both offshore and inshore areas.

Assessment: No analytical assessment is available and fishing mortality is unknown. Evaluation of the status of the stock is based on interpretation of commercial fishery data (catch, effort and standardized catch rates), time series of research biomass indices and stock composition data.

CPUE: A single combined index covering the whole area in the period 1976-96 indicates two levels of stock abundance. From 1977 to 1988 the indices fluctuated at a level higher than the 1989-96 period.

Recruitment: Survey length distributions indicate a relatively strong 1990 year-class and the presence of several year-classes of smaller shrimp. The 1993 year-class seems very abundant. The 1990 year-class will likely maintain the catch rates in 1997, as it recruits to the female component of the stock. If the 1993 year-class is as strong as indicated, the catches will contain high proportions of small shrimp in 1997.

State of the Stock: Stock seems to be relatively stable, but at a lower level than in the 1970s to late-1980s. The presence of several year-classes recruiting to the fishable stock further suggests that there is no concern for recruitment in the short or medium term.

Recommendations: TACs advised for both 1995 and 1996 were 60 000 tons. The current assessment does not show any significant change in the status of the stock, which could justify modifying the advice.

Special Comments: The Scientific Council noted in its November 1995 report that an increase in TAC (to 67 000 tons) based on an upward revision of the average inshore catch was not warranted. The previously advised TAC of 60 000 tons is lower than the recent catches and may allow the stock size to increase to the higher level observed in the 1980s. However, a catch of 67 000 tons is also lower than recent catches and may be sustainable given the relative stability of the stock for the 1990s. An increase in the TAC to 67 000 tons would likely decrease the probability that the stock will increase from the current lower level, but Scientific Council is unable to quantify this probability.

Sources of Information: SCR Doc. 96/106, 109, 110, 111, 112, 113, 114, 115.

Shrimp in Denmark Strait

Background: The fishery for shrimp in limited areas of the Denmark Strait began in 1978. The fishery started exploiting new areas after 1992.


Fishery and Catches: This soon became a multi-national fishery with recent catches and TACs as follows:

Year	Catch	('000 tons)	
		TAC Recommended	TAC ¹ Effective
1992	7.5	8	13.0
1993 ²	7.6	5	9.6
1994 ²	9.8	5	9.6
1995 ²	9.5	5	9.6
1996 ² (to Oct)	6.5	5	9.6

¹ On western side of midline.

² Provisional.

Effort has declined substantially since the late-1980s.




Data: Catch, effort and biological sampling data were available from the trawlers of several nations. Two time series of survey biomass indices were available, one from Norway for the years 1985 to 1989 and another from Greenland for the years 1989 to 1996, with associated biological samples.

Assessment: No analytical assessment is available and fishing mortality is unknown. Evaluation of the status of the stock is based on interpretation of commercial fishery data, the time series of survey biomass indices and biological data from both sources.

CPUE: Standardized CPUE indices in the traditional northern area have declined from peak values in 1987 to minimum values in 1992-93, subsequently increasing in 1994, remaining stable in 1995 and declining in 1996. However the unstandardized index for northern and southern areas combined increased from 1993 to 1994 and stabilized thereafter.

Recruitment: There are no immediate concerns for recruitment since the number of males in the surveys has increased substantially in recent years.

Abundance: The abundance index from the Greenlandic survey from the northern area declined from 1989 to 1992, and increased thereafter. The 1996 value is the highest in the series but is based on incomplete coverage.



State of the Stock: Although changes in fishing patterns make assessing stock status difficult, it seems that the stock has recently improved, but remains below the level of the early- to mid-1980s.

Recommendations: For 1997 there is no biological basis for advising any change to the TAC from the 1996 value of 5 000 tons.

Special Comment. Scientific Council noted that catch levels have substantially exceeded advised TACs in recent years. Although these recent catch levels have not resulted in stock decline, reducing catches to the advised TAC level would improve chances of stock rebuilding.

Sources of Information: SCR Doc. 96/107, 108, 116, 117, 118).

IV. OTHER MATTERS

The Council noted that the Working Group on Shrimp in Div. 3M will meet immediately after this meeting of the Scientific Council, with the Designated Expert, D. G. Parsons (Canada), as Chairman. The Working Group will present its report in the form of an SCS Document for consideration by the Council in 1997.

V. ADOPTION OF REPORTS

The Council met briefly at 1700 hr on 18 November 1996 and **adopted** the STACFIS Report. The report is given in Appendix I. The Council then **adopted** its own report.

VI. ADJOURNMENT

There being no further business, the Chairman thanked the participants, especially the Designated Experts, the Chairman of STACFIS, W. B. Brodie (Canada), who kindly offered to undertake the task of chairing this STACFIS meeting and the Secretariat for the able assistance in the conduct of the meeting.

APPENDIX I. REPORT OF STANDING COMMITTEE ON FISHERY SCIENCE (STACFIS)

Chairman: W. B. Brodie

Rapporteur: Various

The Committee met at NAFO Headquarters, Dartmouth, Nova Scotia, Canada, during 15-18 November 1996, to review the status of the shrimp stocks in Subareas 0 and 1, and Denmark Strait, as referred to it by the Scientific Council. Representatives attended from Canada, Denmark (in respect of Faroe Islands and Greenland) and Iceland.

I. STOCK ASSESSMENTS

1. Shrimp in Subareas 0 and 1 (SCR Doc. 96/106, 109, 110, 111, 112, 113, 114, 115)

a) Introduction

In accordance with the recommendation by STACFIS in November 1993, the entire shrimp stock in Div. 0A, and Subarea 1 both north and south of 71°N, as well as inshore, is assessed as a single population. Overall catches in the entire stock area increased until 1992, then decreased from 1993 to 1995 (Fig. 1). Catches in 1996 are projected to be slightly below the 1995 level, because measures have been taken by Greenland authorities to reduce the catch by 5% per year starting in 1995.

It has been recognized that shrimp catches include a small component of the species *P. montagui*. In 1995 and 1996, separate quotas were set by Greenland for catches of *P. montagui*. These catches amounted to 374 tons and 432 tons, respectively, and were not included in the nominal catches for *P. borealis*.

Recent nominal catches and TAC (tons) for shrimp in Div. 0A and Subarea 1 are as follows:

	1986	1987	1988	1989	1990	1991	1992	1993 ¹	1994 ¹	1995 ¹	1996 ^{1,2}
Div. 0A Total	2 995	6 095	5 881	7 235	6 177	6 788	7 493	5 491	4 766	2 361	2 100
SA 1 Offshore	52 634	50 720	44 159	45 198	49 478	52 834	58 664	52 280	53 693	51 900	42 466
SA 1 Inshore	7 500	6 921	10 233	13 224	13 630	16 258	20 594	17 843	18 118	16 429	10 533
SA 1 Total	60 134	57 641	54 392	58 422	63 184	69 092	79 258	70 123	71 811	68 329	52 999
SA 0+1 Total	63 129	63 736	60 273	65 657	69 361	75 880	86 751	75 614	76 577	70 690	55 099
0+1 offshore catch	55 629	56 815	50 040	52 433	55 731	59 662	66 157	57 771	58 459	54 261	44 566
0+1 advised TAC ³	36 000	36 000	36 000	44 000	50 000	50 000	50 000	50 000	50 000	60 000	60 000

¹ Provisional data.

² January-October.

³ Until 1994 the advised TAC was only for offshore south of 71°N. After 1994, the advised TAC includes offshore north of 71°N and inshore.

The nominal catch of shrimp in the offshore areas of Subarea 1 and the adjacent part of Subarea 0 (Div. 0A) increased from less than 1 000 tons before 1972 to almost 43 000 tons in 1976, fluctuated thereafter, stabilized around a level of 54 000 tons during 1985-88, then increased to 66 000 tons in 1992 and decreased thereafter to 54 000 tons in 1995. Preliminary statistics for the offshore area in 1996 (January to October, Subarea 1) show total catches of about 44 000 tons (compared to 43 000 tons in the same months in 1995). The offshore fishery has been regulated by TAC since 1977.

During the history of this fishery, the fishing grounds in Div. 1B have been the most important. The fishery started expanding southward in 1989 and this expansion continued until 1993.

The fishery in Div. 0A usually takes place from July to November. In Subarea 1 the fishery occurs in all months of the year, however, early in the year it is often confined to the southern Divisions due to ice coverage in Div. 1A and Div. 1B. The northwestern fishing grounds in Div. 1B and Div. 0A were fished less intensively in 1995 and 1996.




Fig. 1. Shrimp in Subareas 0 and 1: total catches and standardized effort.

The West Greenland inshore shrimp fishery was relatively stable from 1972 to 1987 with estimated catches of 7 000-8 000 tons annually (except for 10 000 tons in 1974). A revision of the inshore catch statistics showed that catches in recent years have increased to over 20 500 tons in 1992, but decreased to 16 000 tons in 1995. Preliminary data for 1996 (January-October) indicate that catches are at the same level as for the same period in 1995.

b) **Input Data**

i) **Commercial fishery data**

Fishing effort and CPUE. Catch and effort data from the shrimp fishery in 1996 were available from fishing records from Canadian vessels in Div. 0A (SCR Doc. 96/106) and from Greenland logbooks for Subarea 1 (SCR Doc. 96/109, 111).

Five time series of standardized CPUE indices were available, including both the inshore and offshore areas, as well as the small vessel component. The five indices were: 1) Seven trawler index in Div. 1B (offshore) of total shrimp catches from 1976 to 1990, 2) Div. 1B trawler index (offshore) of large shrimp (count 120 per kg or less) from 1987 to 1996, 3) Div. 1CD trawler index (offshore) of large shrimp (count 120 per kg or less) from 1988 to 1996, 4) The Greenland small vessel (<80 GRT) index (inshore and near shore) of total shrimp catches from 1988 to 1996, 5) the Canadian 0A trawler index (offshore) of total shrimp catches from 1981 to 1996. A single index was constructed by combining the five separate indices covering the period 1976-96 (SCR Doc. 96/109, 96/111). The last value of the index (1996) was calculated from the historic model.

During the 1976-88 period, the index fluctuated at a level substantially higher than the level during 1989-96. In the latter period the index fluctuated without trend (Fig. 2).

Up to 1986, the standardized effort showed a slight increasing trend. Effort more than doubled between 1987 and 1992, and then varied without a trend until 1995. The 1996 value (projected to the end of the year) is expected to be lower than the 1995 value (Fig. 1). Twin trawls introduced in 1995 on several Greenlandic trawlers have been accounted for in analyses of effort data.




Fig. 2. Shrimp in Subareas 0 and 1: standardized combined CPUE index.

Length and age composition. Length frequency distributions obtained by observers were available from the commercial fishery in Div. 0A from 1981 to 1996 (SCR Doc. 96/106) and in Subarea 1 from 1991 to 1996 (SCR Doc. 96/109).

Standardized catch rates for males, which included several year-classes, increased from 1993 to 1996 while for females they fluctuated between 1991 and 1996. There are indications that the 1993 year-class is abundant. Data from Div. 0A show that the 1993 year-class is very strong relative to other year-classes at age 3 in this fishery.

Shrimp discards. In Div. 0A in 1995 and 1996, discarding was lower than in previous years.

ii) **Research survey data**

Biomass and abundance estimates. Trawl surveys have been conducted from 1988 in offshore areas (Subarea 1 + Div. 0A) and from 1991 in inshore Subarea 1 (SCR Doc. 96/112, 96/114). Re-analyses of survey results showed that *P. montagui* occurred in small quantities (around 100 tons per year) from 1988 to 1991 and increased from 1992 to 1995 from 1 100 tons to about 15 000 tons. In 1996 the estimated biomass of *P. montagui* decreased to 2 800 tons (SCR Doc. 96/113).

The estimates of trawlable biomass for *P. borealis* are as follows:

Biomass ('000 tons)	1988	1989	1990	1991	1992	1993	1994	1995	1996
Offshore (Subarea 1+Div. 0A)	182	207	212	125	192	222	178	145	186
Inshore (Div. 1A)	-	-	-	49	45	32	41	47	55
Total	-	-	-	174	237	254	219	192	241

Offshore: In July-September 1996, a stratified-random trawl survey was carried out in the main area of shrimp distribution in Div. 1A to 1E and the adjacent part of Div. 0A. The survey was carried out with a two-phase design, applying more stations in strata with high shrimp densities (SCR Doc. 96/114).

Biomass estimates from the survey in the period 1988-96 were variable around a mean level of 180 000 tons (Fig. 3).




Fig. 3. Shrimp in Subareas 0+1: combined biomass estimates from inshore and offshore surveys.

Survey catches were dominated by males in 1996. Length distributions in 1996 indicated a relatively strong 1990 year-class and the 1991 and 1992 year-classes appeared to be about average. The 1993 year-class in 1996 is more abundant than previous year-classes observed at age 3 from 1993 to 1995.

Abundance-at-age (in billions) for shrimp from Greenland offshore research survey data is given in the following table:

Age	1988	1989	1990	1991	1992	1993	1994	1995	1996
1						0.5	0.3	0.5	0.7
2	0.4	0.4	0.8	0.2	0.7	2.2	1.3	0.5	1.9
3	0.9	4.5	1.1	0.6	2.5	3.4	2.4	1.1	8.0
4	3.4	16.0	3.2	1.7	3.2	7.2	6.5	3.6	7.0
5	7.1	7.0	11.7	2.2	5.7	10.2	7.0	7.6	6.0
6	6.3	3.9	5.2	7.5	8.9	8.4	7.5	4.7	9.3
7+	7.7	6.0	8.0	4.4	5.5	7.9	6.4	5.1	5.6
Total	25.8	37.9	29.9	16.6	26.4	39.7	31.4	23.1	38.5

Inshore: In August 1996, a stratified-random trawl survey also using a two-phased approach was conducted by Greenland in the inshore areas in Disko Bay and Vaigat (Div. 1A) (SCR Doc. 96/112). The biomass estimates from the survey series in 1991-96 were variable around 45 000 tons with an increasing trend from 1993 to 1996 (Fig. 3).

The overall size composition of shrimp from the inshore survey in 1996 was similar to that of the offshore in relation to the occurrence of modes. In the 1996 survey males were most abundant in southwestern and central Disko Bay. Females were most abundant in central Disko Bay and northern Vaigat.

c) **Assessment Results**

Indices from the commercial fishery show that the abundance of shrimp in 1989-96 fluctuated without trend, but at a lower level than in 1976 to 1988. The decrease from 1987 to 1989 was coincident with a substantial increase in effort. The survey indices from 1988 to 1996 also indicate that the abundance has fluctuated without trend.

The combined inputs to the assessment indicate a stable stock size. The fishery in 1997 will depend on the relatively strong 1990 year-class and as it recruits to the female component it should maintain catch rates. The presence of several recruiting year-classes, further suggests that there is no concern about recruitment in the short or medium term.

If the 1993 year-class is as strong as indicated, the catch will contain a high proportion of small shrimp in 1997.

d) **Research Recommendations**

STACFIS was pleased to note that the recommendations from the 1995 November Meeting of the Scientific Council were fulfilled.

For shrimp in Div. 0A and Subarea 1, STACFIS **recommended** that *for consideration at the November 1997 Meeting of the Scientific Council, sampling of the commercial fishery be improved to cover all components of the fishery by area and month.*

2. **Shrimp in Denmark Strait (SCR Doc. 96/107, 108, 116, 117, 118)**a) **Introduction**

The fishery in Denmark Strait started in 1978 and has taken place primarily in the area of Strede Bank and Dohrn Bank as well as on the slopes of Storfjord Deep. The available fishing grounds at any given time depend heavily on the ice conditions. The traditional area extends from approximately 65°N to 67°30'N and between 26°W and 34°W. In 1993, a fishery started in areas between 60°30'N and 65°N and west of 35°W. In 1996 the fishery was located mainly in the southern area, with some indication of activity in nearshore areas of East Greenland.

Catches in the northern (traditional) area increased rapidly to 1980, declined and remained stable from 1981 to 1983, increased gradually to 1988 (12 500 tons) and then decreased again to 1994, staying relatively constant from 1994 to 1996. Catches from the southern fishing area increased from 3 000 tons to 6 600 tons in 1994, then declined to 4 400 tons in 1996 (preliminary data). Catches for the whole area increased from 1993 to 1994-95 and then declined in 1996 (preliminary data) (Fig. 4).

Recent catches and TACs (tons) are as follows:

	1986	1987	1988	1989	1990	1991	1992	1993 ¹	1994 ¹	1995 ¹	1996 ^{1,2}
Catch north of 65°N											
eastern side	1 150	1 330	1 424	1 326	281	465	1 750	2 553	1 514	1 151	566
western side	9 814	10 848	11 125	9 416	9 994	8 192	5 764	3 950	3 358	4 052	1 582
Catch south of 65°N	-	-	-	-	-	-	-	2 995	6 641	5 461	4 380
Total	10 964	12 178	12 549	10 742	10 275	8 657	7 514	7 638	9 778	9 512	6 528
Advised TAC	-	-	-	10 000	10 000	10 000	8 000	5 000	5 000	5 000	5 000
Effective TAC western side	7 525 ³	7 725 ³	8 725 ³	9 025 ³	14 100	14 500	13 000	9 563	9 563	9 563	9 563

¹ Provisional catches.

² January-October.

³ Not including Greenland fishery north of 66°30'N.




Fig. 4. Shrimp in Denmark Strait: catches. The 1996 catch is to the end of October.

b) Input Data

i) Commercial fishery data

Fishing effort and CPUE. Catch and effort data from logbooks were available from Greenland, Norway, Iceland, Faroe Islands and EU-Denmark since 1980, and from EU-France for the years 1980 to 1991. Because of uncertainty regarding the area fished by Norway in 1993 to 1995, Norwegian data have been excluded from some catch-effort analyses.

In the northern area, between 1980 and 1989, total unstandardized effort increased from about 35 000 hours to more than 100 000 hours, declining thereafter to about 16 000-24 000 hours in 1994-95. The fishery in the July-December period became more important at the end of the 1980s, accounting for approximately 50% of the total annual effort, whereas in the 1990s the effort in spring has been the most important. In the southern area, effort was between 36 000-40 000 hours in 1993-95. For the whole area, effort has declined from 80 000 hours in 1993 to 63 000 hours in 1995.

In the northern area (excluding Norway) unstandardized catch rates (Fig. 5) declined from 1980 to 1983, fluctuated from 1983 to 1987 then declined again to 1989 (SCR Doc. 96/107, 96/108). Values for 1990-93 were similar to the low 1989 value at about 50% of the level seen in the early- to mid-1980s. In 1994 there was a considerable rise in the catch rate but this subsequently declined in 1995 and 1996.

For north and south areas combined, the unstandardized catch rate rose from 1993 to 1994 and then remained relatively constant.




Fig. 5. Shrimp in Denmark Strait: unstandardized catch rates (scaled to the 1996 value in the northern area).

Standardized catch-rate series for Greenland vessels for large shrimp and all shrimp in the northern area (Fig. 6) showed a continuous decline from 1987 to 1993 and a considerable increase in 1994. The 1995 value was approximately the same as that for 1994 but in 1996 the index declined to the lowest value in the series (SCR Doc. 96/117).




Fig. 6. Shrimp in Denmark Strait: standardized catch rate indices. Survey abundance index is shown as points. All indices are for the northern areas only and are relative to the 1996 value.

Biological data. Samples from the Icelandic and Greenlandic fisheries in the late-1980s were comprised mainly of females. Throughout the 1990s males have dominated the catches except in the Icelandic fishery of 1995 which may have been affected by sampling problems (SCR Doc. 96/118).

Commercial samples from both the Greenlandic and Icelandic fisheries indicate no concerns for recruitment for the next 1-3 years.

ii) **Research survey data**

A trawl survey was conducted by Greenland in the Denmark Strait in September 1996, based on a two-stage sampling method using a spline technique. Survey coverage was incomplete due to bad weather, and the results were not strictly comparable to earlier years.

The abundance index declined from 1989 to 1992, and increased from 1992 to 1996 (SCR Doc. 96/116). The 1996 value was about 50% higher than 1995 but this value should be treated with caution. Because variance estimates were not available, it was not known whether differences between years were statistically significant.

The Greenland survey showed an increase in the proportion of males from 1989 to 1992, which continued a trend from the Norwegian surveys of 1985 to 1989. In 1994 and 1995 the proportions of males were almost the same as in 1992, but a further increase was recorded in 1996:

Country	Percent males										
	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996
Norway	41.4	53.5	58.5	58.0							
Greenland				63.1	62.5	-	78.3	-	74.5	74.2	81.0

c) **Assessment Results**

The changes in fishing patterns (decline in catches in north area, increased in south area) make assessment of this stock difficult. The decline in catch rate in the north area is difficult to reconcile with the other indices. Catch rates are increasing in the southern area and are stable considering north and south areas combined. The 1996 survey indicates that abundance is being maintained in the northern area. Length frequencies indicate no immediate concern for recruitment. Despite the uncertainty of the present assessment, it seems that the stock has recently improved, however, it remains below the level of the early- to mid-1980s.

d) **Research Recommendations**

For shrimp in the Denmark Strait, STACFIS recommended that *the annual survey be expanded to cover the whole distribution of shrimp in this area.*

3. **Other Business**

There being no other business, the Chairman thanked the participants, and particularly the Designated Experts, for their work during the meeting. After thanking the Secretariat for their help, the meeting was adjourned.

APPENDIX II. AGENDA SCIENTIFIC COUNCIL - 15-18 NOVEMBER 1996

- I. Opening (Chairman: *W. R. Bowering*)
 1. Appointment of rapporteur
 2. Adoption of agenda
 3. Plan of work

- II. Fishery Science (STACFIS Chairman: *W. B. Brodie*)
 1. Stock assessments (see Annexes 1 and 2)
 - Northern shrimp (Subareas 0 and 1)
 - Northern shrimp (in Denmark Strait and off East Greenland)
 - [Note: For northern shrimp in Subareas 0 and 1, the assessment and TAC advice should include, if possible, the areas north of 71°N in Subarea 1 as well as the inshore region of Subarea 1.]
 2. Other business

- III. Formulation of Advice
 1. Northern shrimp (Subareas 0 and 1)
 2. Northern shrimp (Denmark Strait and off East Greenland)

- IV. Other Matters

- V. Adoption of Reports

- VI. Adjournment

**ANNEX 1. CANADIAN REQUEST FOR SCIENTIFIC ADVICE ON MANAGEMENT IN 1997
OF CERTAIN STOCKS IN SUBAREAS 0 TO 4**

1. Canada requests that the Scientific Council, at its meeting in advance of the 1996 Annual Meeting, provide advice on the scientific basis for the management of the following fish stocks in 1997:

Roundnose grenadier (Subareas 2 and 3)
Silver hake (Div. 4V, 4W and 4X)

It is also suggested that, subject to the concurrence of Denmark (Greenland), the Scientific Council, prior to the 1996 Annual Meeting of NAFO, provide advice on the scientific basis for management of the following fish or invertebrate stocks or groups of stocks in 1997:

Greenland halibut (Subareas 0 and 1)
Roundnose grenadier (Subareas 0 and 1)
Shrimp (Subareas 0 and 1)

The Scientific Council has noted previously there was no biological basis for making two separate assessments for the Greenland halibut throughout Subareas 0-3, but has advised that separate TACs be maintained for different areas of the distribution of Greenland halibut. The Council is asked therefore, subject to the concurrence of Denmark (Greenland) as regards Subarea 1, to provide an overall assessment of status and trends in the total stock throughout its range and comment on its management in Subareas 0+1 for 1997. In particular, the Council is asked to advise on appropriate TAC levels separately for SA 0+1, for SA 2 + Division 3K and for Divisions 3LMNO, and to make recommendations on the distribution of fishing effort within each of these three geographic areas. The Council is asked also to provide information on present harvest patterns in terms of yield per recruit and on distributional variation of the resource in recent years.

With respect to shrimp, it is recognized that the Council may, at its discretion, delay providing advice until later in the year, taking into account data availability, predictive capability, and the logistics of additional meetings.

2. Canada requests the Scientific Council to consider the following options in assessing and projecting future stock levels for those stocks listed above:
- a) For those stocks subject to analytical dynamic-pool type assessments, the status of the stock should be reviewed and implications of fishing at $F_{0.1}$ in 1997 and subsequent years should be evaluated. The present stock size should be described in relation to those observed historically and those to be expected at the $F_{0.1}$ level in both the short and long term. In those cases where present spawning stock size is a matter of scientific concern in relation to the continuing productive potential of the stock, management options should be considered to rebuild the spawning stock. All results should be expressed in terms of stock sizes, catch rates and TACs implied for 1997 and the long term.
 - b) For those stocks subject to general production-type assessments, the status of the stock should be reviewed and management options evaluated in the way described above to the extent possible. In this case, the general reference point should be the level of fishing effort (F) which is two-thirds that calculated to be required to take the MSY catch in the long term.
 - c) For those resources on which only general biological and/or catch data are available, no standard criteria on which to base advice can be established. The evidence on stock status should, however, be weighed against a strategy of optimum yield management and maintenance of stock biomass at levels of about two-thirds that of the virgin stock.
3. The Scientific Council is requested to review the status of the cod stock in Divisions 2J+3KL and to provide estimates of the current size of the total and spawning biomass, together with a description of recent trends. The Council is asked further to provide estimates of the immediate and long-term outlook for the abundance of this stock, including both total and spawning biomass.

William A. Rowat
Deputy Minister
Department of Fisheries and Oceans
Ottawa, Canada

**ANNEX 2. DENMARK (GREENLAND) REQUEST FOR SCIENTIFIC ADVICE ON
MANAGEMENT OF CERTAIN STOCKS IN 1997**

1. Denmark, on behalf of Greenland, requests the Scientific Council of NAFO in advance of the 1996 Annual Meeting, provide advice on the scientific basis for management of the following stocks in Subarea 1 in 1997 and as many years forward as data allow:
- i) Redfish (by species, if possible)
 - ii) Any other stock of invertebrates and finfish of commercial interest, for which data allow a status report

It is also suggested that, subject to the concurrence of Canada, advice be given for the following stocks overlapping Subareas 0 and 1:

- i) Greenland halibut
- ii) Roundnose grenadier

2. In the analyses on which management advice will be based, the following should be included:

In its 1993 report, the Scientific Council has noted that the offshore component of **Greenland halibut**, in Subareas 0 and 1, was distributed equally between these Subareas. Further in its 1995 report, the Scientific Council noted that the biomass of the inshore component in Subarea 1 was unknown. The Council is therefore asked to provide further information on following topics.

- a) Allocation of TACs to appropriate Subareas (Subareas 0 and 1).
- b) Allocation of TAC for Subarea 1 inshore areas.

3. Following bilateral consultations Canada and Denmark on behalf of Greenland, requests the Scientific Council of NAFO to provide advice on Greenland halibut in Subareas 0 and 1. Given that the assessment of the stock has been impeded by inconsistencies in age readings both within institutes and among institutes involved in the fishery, the Council is asked to:

establish an exchange program on Greenland halibut otoliths in order to calibrate the age reading methods between readers from the different countries involved in the fishery.


The Greenland halibut stock in Subareas 0 and 1 is at present being exploited by a number of different gears (trawl, long-line and gill net). The Council is asked to provide any new information on:

the impact on the stock composition of different exploitation patterns in terms of yield per recruit, long term sustainable yield and spawning stock biomass.

4. Denmark, on behalf of Greenland, further requests that the Scientific Council of NAFO before December 1996, provide advice on the scientific basis for management of the Northern shrimp (*Pandalus borealis*) in Subareas 0 and 1 in 1997 and as many year forward as data allow.

Further, in cooperation with ICES, the Scientific Council is requested to advise on the scientific basis for management of the Northern shrimp (*Pandalus borealis*) stock in the Denmark Strait.

Brent Buch, Director
On behalf of
Ministry for Fisheries, Hunting & Agriculture
Aslisarnermut, Piniarnermut, Nunalerinermullu Pisortaqaarfik
Direktoratet for Fangst, Fiskeri og Landbrug



APPENDIX III. LIST OF RESEARCH AND SUMMARY DOCUMENTS - NOVEMBER 1996

RESEARCH DOCUMENTS (SCR)

SCR #	Ser. #	Title
96/106	N2803	PARSONS, D. G., and P. J. VEITCH. The Canadian fishery for northern shrimp (<i>Pandalus borealis</i>) in NAFO Division 0A and Subarea 1, 1979-1996
96/107	N2804	SKÚLADÓTTIR, U. The catch statistics of the shrimp fishery (<i>Pandalus borealis</i>) in the Denmark Strait in the years 1980-1996.
96/108	N2805	SKÚLADÓTTIR, U. The Icelandic shrimp fishery (<i>Pandalus borealis</i> Kr.) in the Denmark Strait in 1995-1996 and some reflection on age groups in the years 1991-1996.
96/109	N2806	HVINGEL, C., H. SIEGSTAD, and O. FOLMER. The Greenland fishery for northern shrimp (<i>Pandalus borealis</i>) in Davis Strait in 1995 and January-October 1996.
96/110	N2807	HVINGEL, C. Geographical changes in the fishing pattern of Greenlandic shrimp fishery in the Davis Strait, 1987-1996.
96/111	N2808	HVINGEL, C., H. LASSEN, and D. G. PARSONS. A biomass index for northern shrimp (<i>Pandalus borealis</i>) in Davis Strait based on multiplicative modelling of commercial catch-per-unit-effort data (1976-1995).
96/112	N2809	FOLMER, O., D. M. CARLSSON, C. HVINGEL, and P. KANNEWORFF. Stratified random trawl survey for shrimp (<i>Pandalus borealis</i>) in Disko Bay and Vaigat, inshore West Greenland, 1996.
96/113	N2810	FOLMER, O. Occurrence of striped shrimp (<i>Pandalus montagui</i>) along the west coast of Greenland from 1988 to 1996.
96/114	N2811	FOLMER, O., D. M. CARLSSON, C. HVINGEL, and P. KANNEWORFF. Offshore trawl survey for shrimp (<i>Pandalus borealis</i>) in NAFO Subareas 0 and 1 in 1996.
96/115	N2812	SIEGSTAD, H. Preliminary assessment of shrimp (<i>Pandalus borealis</i>) in Davis Strait, 1996 (Subareas 0+1).
96/116	N2813	CARLSSON, D. M. Trawl survey for shrimp (<i>Pandalus borealis</i>) in Denmark Strait, 1996.
96/117	N2814	HVINGEL, C., H. SIEGSTAD, and O. FOLMER. The Greenland fishery for northern shrimp (<i>Pandalus borealis</i>) in Denmark Strait in 1995 and January-October 1996.
96/118	N2815	SKÚLADÓTTIR, U. Preliminary assessment of shrimp in the Denmark Strait in 1996.

SUMMARY DOCUMENTS (SCS)

SCS #	Ser. #	Title
96/18	N2816	NAFO. Report of Scientific Council, Special Meeting, 15-18 November 1996.

1880

1881

1882

1883

APPENDIX IV. LIST OF REPRESENTATIVES, ADVISERS/EXPERTS - NOVEMBER 1996**CANADA****Representatives:**

- Bowering, W. R. Northwest Atlantic Fisheries Centre, P. O. Box 5667, St. John's, Newfoundland A1C 5X1
Phone: +709-772-2054 - Fax: +709-772-4188 - E-mail: bowering@athena.nwafc.nf.ca
- Powles, H. Sr. Policy Advisor, Invert. & Pacific Marine Fish, 200 Kent St., Ottawa, Ontario K1A 0E6
Phone: +613-990-0285 - Fax: +613-954-0807
- E-mail: howard.powles@ncr.ottwpo.dfo-mpo.X400.gc.ca

Advisers/Experts:

- Brodie, W.B. Northwest Atlantic Fisheries Centre, P. O. Box 5667, St. John's, Newfoundland A1C 5X1
Phone: +709-772-3288 - Fax: +709-772-4188 - E-mail: brodie@athena.nwafc.nf.ca
- Parsons, D.G. Northwest Atlantic Fisheries Centre, P. O. Box 5667, St. John's, Newfoundland A1C 5X1
Phone: +709-772-2093 - Fax: +709-772-4105 - E-mail: parsons@athena.nwafc.nf.ca
- Savard, L. Dept. of Fish. & Oceans, Maurice Lamontagne Inst., P.O. Box 1000, Mont-Joli, Quebec G5H 3Z4
Phone: +418-775-0621 - Fax: +418-775-0740

DENMARK**GREENLAND****Representative:**

- Siegstad, H. Greenland Institute of Natural Resources, P. O. Box 570, DK-3900, Nuuk, Greenland
Phone: +299 21095 - Fax: +299 25957 - E-mail: helle@natur.centadm.gh.gl

Advisers/Experts:

- Carlsson, D.M. Greenland Institute of Natural Resources, P. O. Box 2151, Pilestræde 52,
DK-1016 Copenhagen K, Denmark
Phone: +45 33 14 5524 - Fax: +45 33 13 4250 - E-mail: grfidmc@inet.uni-c.dk
- Folmer, O. Greenland Institute of Natural Resources, P. O. Box 570, DK-3900, Nuuk, Greenland
Phone: +299 21095 - Fax: +299 25957 - E-mail: folmer@natur.centadm.gh.gl
- Hvingel, C. Greenland Institute of Natural Resources, P. O. Box 570, DK-3900, Nuuk, Greenland
Phone: +299 21095 - Fax: +299 25957 - E-mail: hvingel@natur.centadm.gh.gl

FAROE ISLANDS**Adviser/Expert:**

- Nicolajsen, Á Fiskorannsóknarstovan, Noatun, Postboks 3051, FR-110 Torshavn
Phone: +298 1 50 92 - Fax: +298 1 8264 - E-mail: aminic@frs.fo

ICELAND**Representative:**

- Skúladóttir, U. Marine Research Institute, Skulagata 4, P. O. Box 1390, 121-Reykjavik
Phone: +354 552 0240 - Fax: +354 562 3790 - E-mail: unnur@hafro.is