

Northwest Atlantic



Fisheries Organization

Serial No. N3005

NAFO SCR Doc. 98/21

SCIENTIFIC COUNCIL MEETING - JUNE 1998

Abundance, Biomass and Size Composition of Dominant Demersal Fish Stocks and
Trend in Near Bottom Temperature off West and East Greenland, 1982-97

by

Hans-Joachim Rätz

Institute for Sea Fisheries, Branch Office Bremerhaven
Fischkai 35, FRG 27572 Bremerhaven

Abstract

Detailed survey informations about structures and changes in the dominant demersal fish stocks off West Greenland (NAFO Divisions 1B-1F) and East Greenland (ICES Sub-area XIV) are presented together with temporal and geographical properties of the near bottom temperature regime.

After severe declines some stocks showed first and very slight indications of stock recovery due to increased numbers of recruits while their mature parts remained to be depleted, i.e. deep sea redfish, American plaice, Atlantic wolffish, spotted wolffish and starry skate. The recovery indications are more pronounced off East Greenland. All analysed stocks showed significantly lower mean individual weights off West Greenland throughout the survey period. This effect indicates continued recruitment failure off West Greenland which might be related to the shrimp directed fishing effort. No positive signals are derived from survey catches for Atlantic cod and golden redfish.

Almost all shallow strata showed the same trend in near bottom temperature close to the swept area of the fishing stations. This trend was characterized by a very cold event in 1982-84, warmer conditions in 1985-86, a decreasing trend in 1987-89 and warming since then. The 1996 measurements were the highest of the past 16 years and indicated a continued positive trend which was confirmed by the high near bottom temperature in 1997 (4.2 °C).

Introduction

The paper presents abundance and biomass indices as well as size structures of the dominant demersal fish stocks off West and East Greenland south of 67° and 66° northern latitude, respectively, from the 3 mile limit down to 400 m depth. Stock specific information as derived from German groundfish surveys (Rätz, 1997) is tabulated and illustrated for Atlantic cod (*Gadus morhua*), golden, deep sea and juvenile redfish (*Sebastes marinus*, *S. mentella*, and *S. spp*), American plaice (*Hippoglossoides platessoides*), Atlantic and spotted wolffish (*Anarhichas lupus* and *A. minor*) as well as starry skate (*Raja radiata*). These eight ecologically and economically important species/categories contributed more than 90 % to the overall survey catch in numbers and biomass. The trend in near bottom temperature is estimated from CTD-profiles close to the areas swept by the survey gear.

Materials and Methods

Abundance, biomass estimates and length compositions were derived from annual groundfish surveys covering shelf areas and the continental slope off West and East Greenland. Surveys commenced in 1982 and were primarily designed for the assessment of cod. Because of favourable weather and ice conditions and to avoid spawning concentrations, autumn was chosen for the time of the surveys. These were carried out by the research vessel (R/V) WALTHER HERWIG (II) throughout most of the time period. In 1984 R/V ANTON DOHRN was used and she was replaced by the new R/V WALTHER HERWIG III since 1994, respectively.

The fishing gear used was a standardized 140-feet bottom trawl, its net frame rigged with heavy ground gear because of the rough nature of the fishing grounds. A small mesh liner (10 mm) was used inside the cod end. The horizontal distance between wing-ends was 25 m at 300 m depth, the vertical net opening being 4 m. In 1994, smaller Polyvalent doors (4.5 m^2 , 1 500 kg) were used for the first time to reduce net damages due to overspread caused by bigger doors (6 m^2 , 1 700 kg), which have been used earlier. All calculations of abundance and biomass indices were based on the 'swept area' method using 22 m horizontal net opening as trawl parameter, i. e. the constructional width specified by the manufacturer. The towing time was normally 30 min. at a speed of 4.5 knots. Trawl parameters are listed in Table 1. Hauls which received net damage or became hangup after less than 15 minutes were rejected. Some hauls of the 1987 and 1988 surveys were also included although their towing time had been intentionally reduced to 10 minutes because of the expected large cod catches as observed from echo sounder traces.

The surveys were primarily designed for the assessment of cod. In order to reduce the error of abundance estimates, the subdivision of shelf areas and the continental slope into different geographic and depth strata was required due to a pronounced heterogeneity of cod distribution (Rätz, 1996 a and b). The survey area was thus split into seven geographic strata. Each stratum was itself subdivided into two depth strata covering the 0-200 m and 201-400 m zones. Figure 1 and Table 2 indicate the names of the 14 strata, their geographic boundaries, depth ranges and areas in nautical square miles (nm^2). All strata were limited at the 3 mile offshore line.

The applied strategy was to distribute the sampling effort according both to the stratum areas and to cod abundance. Consequently, fifty percent of the hauls were allocated proportionally to strata by stratum area while the other fifty percent were apportioned on the basis of a review of the historical mean cod abundance/ nm^2 , all hauls being randomly distributed within trawlable areas of the various strata. Non-trawlable areas were mainly located inshore. During 1982-96, 2,343 successful sets were carried out, the numbers of valid sets by year and stratum being listed in Table 3. Apart from stratum 7.2 (Dohrn Bank), East Greenland strata were not covered adequately in 1984, 1992 and 1994 due to technical problems. In 1995, the survey area off West Greenland was incompletely covered for the first time again due to technical problems. Only 50 % of the strata of West Greenland were covered, namely the southern strata 3.1, 3.2, 4.1, and 4.2. Stratum 7.1 has a very low area and therefore never been covered. In 1996 and 1997, the entire survey area was covered. Figure 1 shows the positions of hauls conducted during the most recent survey.

Fish were identified to species or lowest taxonomic level and the catch in number and weight was recorded. Redfish (≥ 17 cm) were separated to *Sebastes marinus* L. or deep sea *Sebastes mentella* Travin, whereas juvenile redfish (< 17 cm) were classified as *Sebastes spp.* due to time-consuming and difficult species identification. Total fish lengths were measured to cm below. Stratified abundance estimates were calculated from catch-per-tow data using the stratum areas as weighting factor (Cochran, 1953; Saville, 1977). Strata with less than five valid sets were rejected from the calculation. The coefficient

of catchability was set arbitrarily at 1.0, implying that estimates are merely indices of abundance and biomass. Respective confidence intervals (CI) were set at the 95% level of significance of the stratified mean.

As a standard procedure, near bottom temperatures were measured directly before or after trawling in the vicinity of the swept area by a CTD-sonde with a precision of a hundredth °C. Table 20 lists the available numbers of temperature values by stratum and year.

Results

Tables 4 and 5 list abundance and biomass estimates for **Atlantic cod**. The values are shown in Figures 2 and 3 appended by the trend in mean individual weight and recent length frequencies in Figures 4 and 5, respectively. The increase in stock abundance and biomass during 1984-87 to 830 million individuals and 690 000 tons was due to the recruiting process of the year classes 1984 and 1985. Until 1992, the stock abundance and biomass collapsed by nearly 100% and remained at that low level. The 1997 indices amounted to 4 million specimens and 14 000 tons. Since the stock collapse in the early 90s, the majority of the stock was distributed off East Greenland in contrast to earlier observations. The recent length structure was of the stock is hardly derivable from the scarce length data (Fig. 5). However, current recruitment is estimated to be very poor and does not indicate stock recovery in the near future. The fish appeared to be smaller off West Greenland (Fig. 4) at all times surveyed.

In comparison with the 80s, the most recent abundance and biomass estimates of **golden redfish** (≥ 17 cm) displayed a significant decrease by more than 90 % (Tab. 6 and 7, Fig. 6 and 7). The great majority of golden redfish was always found to be distributed off East Greenland. Until the end of the decade of the 80s, the abundance of golden redfish (≥ 17 cm) off West Greenland decreased to non-recognizable values and did not show any signs of recovery since then. Figure 8 illustrates that the calculated size reductions of the specimens were more significant off West Greenland than off East Greenland where bigger fish were found throughout the time series. The length compositions in 1995-97 were dominated by immature fish of 20-22 cm and 25-27 cm and around 30 cm indicating an annual growth increment of about 3-5 cm (Fig. 9).

As observed for the golden redfish (≥ 17 cm) the stock of **deep sea redfish** (≥ 17 cm) was found to be distributed almost exclusively off East Greenland (Tab. 8 and 9, Fig. 10 and 11). During the 80s, abundance and biomass estimates for deep sea redfish (≥ 17 cm) varied without any distinct trend. Since 1991, both estimates indicate enormous increases in abundance and biomass due to good recruitment. Compared with the development off East Greenland where 99 % of the fish were recently distributed, the highest abundance value of the time series off West Greenland in 1997 is considered insignificant but does indicate a close relation between the deep sea redfish off West and East Greenland. Deep sea redfish were estimated to be significantly smaller off West Greenland for most of the survey years. Furthermore, Figure 12 displays an increase in mean weight since 3 years off East Greenland after significant size reductions which continued off West Greenland where the body weight of the fish was reduced by 50 %. Recent length compositions are dominated by juvenile fish and indicate an annual growth rate of about 2-3 cm (Fig. 13).

Juvenile and unspecified redfish (< 17 cm) dominated the abundance of finfish by far. Since 1993, this category was very abundant and concentrated mainly off East Greenland (Tab. 10 and 11, Fig. 14 and 15). Driven by the recruitment, the mean individual weight varied between 7 and 64 g (Fig. 16). Juvenile redfish (< 17 cm) off East Greenland were generally bigger as compared with fish off West Greenland. Annual growth increments of 4 cm were indicated by pronounced peaks in the length compositions at 9-11 and 13-15 cm (Fig. 17).

The abundance and biomass indices of **American plaice** off West and East Greenland are listed in Tables 12 and 13 and shown in Figures 18 and 19, respectively. The most recent records indicate a slight increase in abundance and biomass from the lowest values of the time series in the early and mid 90s (Lloret, 1997). The fish off West Greenland were generally smaller as compared to East Greenland (Fig. 20). The first indications of a slight stock recovery are due to the increased presence of recruits < 20 cm (Fig. 21).

Recently, **Atlantic wolffish** increased in abundance and biomass off East Greenland to the highest values of the time series. Off West Greenland, the abundance and biomass estimates indicate fewer fish and a slight recovery during the most recent years from the lowest values in the early and mid 90s (Tab. 14 and 15, Fig. 22 and 23). The relatively high abundance with low biomass indices result in small individual sizes. The mean individual weight decreased significantly and almost

continuously to 0.3 kg in 1997 (Fig. 24). In 1996 and 1997, the size composition displayed higher abundance of recruits <30 cm (Fig. 25).

Spotted wolffish were caught rarely during the whole survey period, but abundance and biomass estimates decreased significantly off West Greenland while East Greenland abundance estimates were high since 1994. The biomass off East Greenland increased since the early 90s (Tab. 16 and 17, Fig. 26 and 27). During the 90s, the majority of the fish were distributed off East Greenland. During the previous decade, the stock was found to be mainly distributed off East Greenland. The mean individual weight decreased more pronounced off West Greenland (Fig. 28). The length compositions in 1994, 1995 and 1996 were scattered due to small catches (Fig. 29).

During the decade of the 80s, the majority of starry skates were distributed off West Greenland, only 2 % of abundance and 8 % of biomass were found off East Greenland (Tab. 18 and 19, Fig. 30 and 31). Both abundance and biomass indicate a slight increase from the lowest level due to recruitment. There was a significant negative trend in body weight since 1982. Again individuals of West Greenland were smaller throughout the survey period (Fig. 32). The higher abundance of small recruits <15 cm is illustrated in Figure 33.

Table 21 lists mean near bottom temperatures by stratum and weighted means by stratum area, 1982-97. Respective values are illustrated in Figure 34. The northern shallow strata (0-200 m) off West Greenland 1.1 and 2.1 on the left hand panel in Figure 34 showed lower temperatures as compared with the weighted mean and similar trends during the survey period, the weighted mean temperature being illustrated as bold line. A very cold event around 2 °C was identifiable for the period 1982-84 followed by a warming of 2 °C to an overall mean of more than 4 °C. During 1987-89, a less pronounced cooling was observed. Subsequently, there was again an increasing trend exceeding 4 °C in the most recent years. The estimated near bottom temperature for 1996 indicated the warmest conditions of the entire survey period. The 1997 estimate was slightly lower and amounted to 4.2 °C. In contrast to this trend, the temperature regime of the deep strata (201-400 m) appeared more constant and generally warmer than the mean, the only exception being stratum 7.2 off East Greenland.

Discussion

During the period 1982-97, survey results indicated fundamental shifts in abundance, biomass and size composition of the stocks inhabiting the shelf and continental slope off West and East Greenland down to 400 m in depth. After severe declines some stocks showed first and very slight indications of stock recovery due to increased numbers of recruits while their mature parts remained to be depleted, i.e. deep sea redfish, American plaice, Atlantic wolffish, spotted wolffish and starry skate. It is important to note, that the recovery indications are more pronounced off East Greenland. No positive signals are derived from survey catches for Atlantic cod and golden redfish.

The possibly adverse impact of the shrimp fishery at traditional fishing grounds off West Greenland on the recruitment and recovery process of the fish stocks remains unproven due to unknown by-catch rates. However, the geographical differences between individual size off West and East Greenland were pronounced. All analysed stocks showed significantly lower mean individual weights off West Greenland throughout the survey period. This effect indicates continued recruitment failure off West Greenland and is consistent with the effort distribution of the Greenlandic shrimp fleet which expends more than 150 000 hours trawling per year off West and only 15 000 hours off East Greenland (Hvingel et al., 1997 a and b) without any selective device.

Almost all shallow strata showed the same trend in near bottom temperature close to the swept area of the fishing stations. This trend was characterized by a very cold event in 1982-84, warmer conditions in 1985-86, a decreasing trend in 1987-89 and warming since then. The 1996 measurements were the highest of the past 16 years and indicated a continued positive trend which was confirmed by the high near bottom temperature in 1997 (4.2 °C). These findings are in agreement with the description of the climatic conditions around Greenland as derived from air temperatures and oceanographic standard sections (Stein, 1997). The average or warmer near bottom temperatures during the 90s did not indicate any unfavorable environmental conditions for fish growth or reproduction.

References

- Cochran, W. G. 1953. Sampling techniques. John Wiley & Sons Inc., New York: 1-330
- Hvingel, C., O. Folmer and H. Siegstad 1997 a. The Greenlandic fishery for northern shrimp (*Pandalus borealis*) in Denmark Strait 1996 and Januar-October 1997. NAFO SCR Doc. 97/97, Ser. No. N2954:1-19
- Hvingel, C., O. Folmer and H. Siegstad 1997 b. The Greenland fishery for northern shrimp (*Pandalus borealis*) off West Greenland, 1970-1997. NAFO SCR Doc. 97/98, Ser. No. N2955:1-24
- Lloret, Josep 1997. Population dynamics of American plaice (*Hippoglossoides platessoides*) off West Greenland (NAFO Divisions 1B-1F), 1982-94. NAFO Sci. Coun. Studies, 30:89-107
- Rätz, H.-J. 1996 a. Relevance of some Environmental Parameters to Distribution Patterns of Groundfish and Implications for Reasonable Survey Design: Case Study Atlantic Cod off Greenland. NAFO Sci. Coun. Studies, 28:73-78
- Rätz, H.-J. 1996 b. Efficiency of Geographical and Depth Stratification in Error Reduction of Groundfish Survey Results: Case Study Atlantic Cod off Greenland. NAFO Sci. Coun. Studies, 28:65-71
- Rätz, H.-J. 1997. Structures and Changes of the Demersal Fish Assamblage off Greenland and Trends in Near Bottom Temperature, 1982-96. NAFO SCR Dod. 97/5, Ser. No. N2830:1-32
- Saville, A. 1977. Survey methods of apprising fishery resources. FAO Fish. Tech. Pap. 171: 1-76
- Stein, M. 1997. Climatic conditions around Greenland - 1996. NAFO SCR Doc. 97/4, Ser. No. N2829:1-12

Table 1 Trawl parameters of the survey.

Gear	140-feet bottom trawl
Horizontal net opening	22 m
Standard trawling speed	4.5 kn
Towing time	30 minutes
Coefficient of catchability	1.0

Table 2 Specification of strata.

Stratum	geographic boundaries			depth (m)	area (nm ²)	
	south	north	east	west		
1.1	64°15'N	67°00'N	50°00'W	57°00'W	1-200	6805
1.2	64°15'N	67°00'N	50°00'W	57°00'W	201-400	1881
2.1	62°30'N	64°15'N	50°00'W	55°00'W	1-200	2350
2.2	62°30'N	64°15'N	50°00'W	55°00'W	201-400	1018
3.1	60°45'N	62°30'N	48°00'W	53°00'W	1-200	1938
3.2	60°45'N	62°30'N	48°00'W	53°00'W	201-400	742
4.1	59°00'N	60°45'N	44°00'W	50°00'W	1-200	2568
4.2	59°00'N	60°45'N	44°00'W	50°00'W	201-400	971
5.1	59°00'N	63°00'N	40°00'W	44°00'W	1-200	2468
5.2	59°00'N	63°00'N	40°00'W	44°00'W	201-400	3126
6.1	63°00'N	66°00'N	35°00'W	41°00'W	1-200	1120
6.2	63°00'N	66°00'N	35°00'W	41°00'W	201-400	7795
7.1	64°45'N	67°00'N	29°00'W	35°00'W	1-200	92
7.2	64°45'N	67°00'N	29°00'W	35°00'W	201-400	4589
Σ					37463	

Table 3 Numbers of valid hauls by stratum, West, East Greenland and total, 1982-97.

Year	1.1	1.2	2.1	2.2	3.1	3.2	4.1	4.2	5.1	5.2	6.1	6.2	7.1	7.2	Σ West	Σ East	Σ
1982	20	11	16	7	9	6	13	2	1	10	3	12	1	25	84	52	136
1983	26	11	25	11	17	5	18	4	3	19	10	36	0	18	117	86	203
1984	25	13	26	8	18	6	21	4	5	4	2	8	0	5	121	24	145
1985	10	8	26	10	17	5	21	4	5	21	14	50	0	28	101	118	219
1986	27	9	21	9	16	7	18	3	3	15	14	37	1	34	110	104	214
1987	25	11	21	4	18	3	21	3	19	16	13	40	0	18	106	106	212
1988	34	21	28	5	18	5	18	2	21	8	13	39	0	26	131	107	238
1989	26	14	30	9	8	3	25	3	17	18	12	29	0	11	118	87	205
1990	19	7	23	8	16	3	21	6	18	19	6	15	0	13	103	71	174
1991	19	11	23	7	12	6	14	5	8	11	10	28	0	16	97	73	170
1992	6	6	6	5	6	6	7	5	0	0	0	0	0	6	47	6	53
1993	9	6	9	6	10	8	7	0	9	6	6	18	0	14	55	53	108
1994	16	13	13	8	10	6	7	5	0	0	0	0	0	6	78	6	84
1995	0	0	3	0	10	7	10	5	8	6	6	17	0	12	35	49	84
1996	5	5	8	5	12	5	10	5	7	9	5	13	0	9	55	43	98
1997	5	6	5	5	6	5	8	5	5	5	4	8	0	8	45	30	75

Table 4 Abundance indices (1 000) by year, stratum, West and East Greenland and total for Atlantic cod (*Gadus morhua*), 1982-97. Confidence intervals (CI) are given at the 95% level of significance in per cent of the stratified mean. Incomplete survey coverage off East Greenland in 1984, 1992 and 1994!

Year	1.1	1.2	2.1	2.2	3.1	3.2	4.1	4.2	5.1	5.2	6.1	6.2	7.1	7.2	Σ West	Σ East	Σ	CI
1982	5092	729	47957	1888	15114	3706	17790		468	6173	1449	92276	8090	100366	28			
1983	431	467	16013	5170	14881	2326	10916		2228	1274	2276	2213	50204	7991	58195	25		
1984	377	179	4714	171	5201	689	5353		4063		1750	790	16684	6603	23287	32		
1985	19630	2428	13222	4395	10531	1638	7499		3564	373	3978	3348	1141	59343	12404	71747	33	
1986	32438	1236	50908	229	37446	1321	22104			780	6950	6676	828	145682	15234	160916	32	
1987	330944	1651	248002		154681		51114		18317	9832	6527	6081	878	786392	41635	828027	59	
1988	92024	2423	338740	84935	47336	.89	60946		7985	8085	2060	4375	1083	626493	23588	650081	48	
1989	2497	920	27930	673	281502		65203		30906	38407	11600	9383	1436	358725	91732	450457	59	
1990	965	513	4155	362	6014		10303	12213	4956	2524	4533	9041	4200	34525	25254	59779	43	
1991	268	205	180	152	1027	611	1839	523	2343	1786	779	1958	3541	4805	10407	15212	29	
1992	552	622	117	137	121	74	151	269					658	2043	658	2701	50	
1993	566	457	178	127	80	31	0		1252	98	922	502	527	1437	3301	4738	36	
1994	206	103	33	33	72	23	82	22					801	574	801	1375	36	
1995					138	67	58	15	265	78	2933	3654	257	278	7187	7465	93	
1996	152	126	76	38	121	0	298	0	290	0	260	382	515	811	1447	2258	38	
1997	0	47	35	0	120	5	108	0	74	0	624		3456	315	4153	4469	75	

Table 5 Biomass indices (t) by year, stratum, West and East Greenland and total for Atlantic cod (*Gadus morhua*), 1982-97. Confidence intervals (CI) are given at the 95% level of significance in per cent of the stratified mean. Incomplete survey coverage off East Greenland in 1984, 1992 and 1994!

Year	1.1	1.2	2.1	2.2	3.1	3.2	4.1	4.2	5.1	5.2	6.1	6.2	7.1	7.2	Σ West	Σ East	Σ	CI
1982	2378	307	63684	2632	20319	8745	30426		1927	14563	7127	128491	23617	152108	25			
1983	353	205	20215	7827	22806	9594	21374		6147	3512	11344	13154	82374	34157	116531	25		
1984	824	234	7508	234	7218	1055	8493		10387		4110		5237	25566	19744	45310	34	
1985	2528	251	12869	2351	10731	990	5952		7073	1356	9955	9437	5744	35672	33565	69237	39	
1986	10641	484	26098	80	28510	1423	19483			2645	18631	16543	3366	86719	41185	127904	26	
1987	283591	545	200632		116610		37210		10315	9054	9291	17616	5316	638588	51592	690180	63	
1988	94175	1367	333848	77967	44593	93	55945		8750	18204	8162	16258	3572	607988	52946	660934	46	
1989	727	228	25829	441	231239		75388		40614	127865	34957	31324	4786	333850	239546	573396	46	
1990	224	114	3552	190	5778		13185	11388	9229	6813	12954	24408	12560	34431	65964	100395	34	
1991	91	72	73	45	1208	589	2621	451	4236	5779	1263	7467	14006	6150	32751	37901	36	
1992	135	195	23	36	21	14	81	102					1216	607	1216	1823	69	
1993	135	88	49	33	44	10	0		862	60	1742	1078	1860	359	5600	5959	41	
1994	27	33	6	23	23	11	4	13					2792	140	2792	2932	68	
1995					28	13	11	7	93	185	1115	13750	382	57	15525	15582	155	
1996	23	64	23	20	51	0	192	0	187	0	755	1004	1673	373	3599	3972	56	
1997	0	40	24	0	107	4	110	0	57	0	1193		12473	284	13722	14007	90	

Table 6 Abundance indices (1 000) by year, stratum, West and East Greenland and total for golden redfish ≥ 17 cm (*Sebastes marinus*), 1982-97. Confidence intervals (CI) are given at the 95% level of significance in per cent of the stratified mean. Incomplete survey coverage off East Greenland in 1984, 1992 and 1994!

Year	1.1	1.2	2.1	2.2	3.1	3.2	4.1	4.2	5.1	5.2	6.1	6.2	7.1	7.2	Σ West	Σ East	Σ	CI
1982	7015	6340	88792	5512	5736	14876	4087		195798		312132	38899	132358	546829	679187	55		
1983	4025	3186	3355	6523	4043	5885	1697		140766	453	264813	14365	28714	420397	449111	53		
1984	1324	3438	460	1209	10671	2776	4214		6888		47974	9890	24092	64752	88844	65		
1985	4658	10451	6158	1569	3220	14441	4973		78118	32397	1787	141500	25944	45470	279746	325216	52	
1986	6327	4324	2077	3483	21503	2883	2717		124613	470	298708		22234	43314	446023	489337	53	
1987	906	653	1327		9612		659		50961	9422	245	507387	27920	13157	595935	609092	39	
1988	831	2239	342	2255	5938	1954	731		3012	5015	148	132458	34352	14290	174985	189275	54	
1989	421	422	776	690	6489		361		4003	33320	625	110663	76934	9158	225545	234704	60	
1990	120	433	279	709	1038		146	2271	14974	72316	391	653009	37483	4996	778173	783169	75	
1991	227	256	96	691	236	527	21	1671	1385	13237	172	64692	28201	3725	107687	111412	51	
1992	126	106	73	190	183	477	192	835					32622	2192	32622	34814	151	
1993	169	481	59	267	80	132	0		175	6043	77	54424	4170	1188	64889	66077	93	
1994	111	325	156	167	65	46	151	247					3348	1268	3348	4616	41	
1995					51	67	38	146	348	1521	153	38892	2060	302	42972	43274	97	
1996	152	267	22	244	381	383	29	298	647	3145	494	21110	2366	1776	27762	29538	47	
1997	252	609	18	175	120	311	36	552	721	913	21257		1611	2072	24501	26573	40	

Table 7 Biomass indices (t) by year, stratum, West and East Greenland and total for golden redfish ≥ 17 cm (*Sebastes marinus*), 1982-97. Confidence intervals (CI) are given at the 95% level of significance in per cent of the stratified mean. Incomplete survey coverage off East Greenland in 1984, 1992 and 1994!

Year	1.1	1.2	2.1	2.2	3.1	3.2	4.1	4.2	5.1	5.2	6.1	6.2	7.1	7.2	Σ West	Σ East	Σ	CI
1982	1798	1354	34440	2558	3206	9794	2532		155971	194379	30115	55682	380465	436147	54			
1983	846	945	1572	3042	1873	4815	1084		161687	269	229541	15607	14177	407104	421281	61		
1984	308	894	196	519	4935	2284	2089		3601		21281	12052	11225	36934	48159	55		
1985	1020	1819	2968	472	1427	9209	2718		8613	22453	1317	65299	23762	19633	121444	141077	35	
1986	1282	1215	752	1229	10122	1705	1762			43119	382	213268	24368	18067	281137	299204	38	
1987	255	247	660		4954		438		9539	5346	106	230844	19327	6554	265162	271716	38	
1988	146	404	118	942	2570	1342	382		1092	4930	68	98131	48262	5904	152483	158387	60	
1989	182	137	272	249	2619		209		970	14920	442	54589	34360	3668	105281	108949	47	
1990	39	149	75	275	479		79	1343	6761	27245	154	130530	14723	2439	179413	181852	45	
1991	44	83	24	226	120	273	3	1007	725	10631	120	34265	62979	1780	108720	110500	98	
1992	18	35	20	61	53	241	70	447					12076	945	12076	13021	130	
1993	46	112	19	114	39	55	0		75	1377	30	20179	2899	385	24560	24945	68	
1994	34	146	48	64	26	35	40	80					1540	473	1540	2013	38	
1995					19	19	20	43	114	712	51	8896	1141	101	10914	11015	38	
1996	64	102	4	60	128	118	8	132	139	1714	196	10855	1408	618	14312	14928	40	
1997	41	261	5	61	35	188	10	246	163	447		15411	1225	847	17246	18092	58	

Table 8 Abundance indices (1 000) by year, stratum, West and East Greenland and total for deep sea redfish ≥ 17 cm (*Sebastes mentella*), 1982-97. Confidence intervals (CI) are given at the 95% level of significance in per cent of the stratified mean. Incomplete survey coverage off East Greenland in 1984, 1992 and 1994!

Year	1.1	1.2	2.1	2.2	3.1	3.2	4.1	4.2	5.1	5.2	6.1	6.2	7.1	7.2	Σ West	Σ East	Σ	CI
1982	0	390	17	348	0	2360	0			9275		19370	58822	3115	87467	90582	65	
1983	40	1011	70	2528	0	5236	0			15820	0	42393	28378	8885	86591	95476	42	
1984	41	2967	7	1276	0	1115	0			18			34633	76541	5406	111192	116598	93
1985	0	369	31	27	55	328	0		34904	16909	105	38689	81487	810	172094	172904	47	
1986	2141	414	38	292	5	444	0			6932	27	76655	67172	3334	150786	154120	36	
1987	987	13679	42		56		0			0	18340	64	7182	62458	14764	88044	102808	45
1988	150	3187	25	777	60	4619	0		22025	28158	74	176839	25344	8818	252240	261058	58	
1989	0	186	9	102	0	0	8			847	3067	0	72046	222281	305	298241	298546	60
1990	0	10	4	705	50		0	3881	329	12453	2354	13513	16046	4650	44695	49345	43	
1991	0	0	0	0	0	652	0	1773	0	10707	46	724504	234748	2425	970005	972430	81	
1992	0	35	0	15	0	106	0						60064	156	60064	60220	165	
1993	0	24	0	159	7	0	0	0	62	3528	140	1258378	121927	190	1384033	1384223	86	
1994	0	271	20	95	94	162	0	36					77891	678	77891	78569	168	
1995					29	234	96	1468	265	24463	1173	2394064	83314	1827	2503279	2505106	55	
1996	1527	619	0	236	0	1921	29	7135	398	176448	1215	4246101	75011	11467	4499171	4510638	64	
1997	252	1759	0	381	37	3204	144	30742	165	22270		6257093	628353	36518	6907882	6944399	62	

Table 9 Biomass indices (t) by year, stratum, West and East Greenland and total for deep sea redfish ≥ 17 cm (*Sebastes mentella*), 1982-97. Confidence intervals (CI) are given at the 95% level of significance in per cent of the stratified mean. Incomplete survey coverage off East Greenland in 1984, 1992 and 1994!

Year	1.1	1.2	2.1	2.2	3.1	3.2	4.1	4.2	5.1	5.2	6.1	6.2	7.1	7.2	Σ West	Σ East	Σ	CI
1982	0	96	6	114	0	893	0			5178		4843	22795	1109	32816	33925	68	
1983	16	213	26	1158	0	2857	0			8701	0	21047	12747	4270	42495	48765	47	
1984	6	798	4	490	0	472	0			2		12786	35202	1770	47990	49760	97	
1985	0	96	15	11	27	110	0		2960	7169	40	17011	38533	259	65713	65972	35	
1986	223	39	20	110	3	179	0			3943	15	29277	31333	574	64568	65142	36	
1987	84	1184	9		31	0	0	0	4891	17	2328	23264	1308	30500	31808	46		
1988	20	425	21	159	45	1878	0		3542	10166	9	55838	11607	2548	81162	83710	56	
1989	0	23	7	15	0	1			90	655	0	21151	45452	46	67348	67394	63	
1990	0	5	2	87	7		0	542	62	2741	329	1961	3275	643	8368	9011	44	
1991	0	0	0	0	153	0	445	0	2959	30	211468	69454	598	283911	284509	80		
1992	0	3	0	2	0	28	0					19856	33	19856	19889	160		
1993	0	5	0	23	2	0	0	0	34	483	19	194675	34102	30	229323	229353	61	
1994	0	31	3	10	12	25	0	3				7122	84	7122	7206	128		
1995					5	25	10	159	29	2859	207	355946	16505	199	375546	375745	52	
1996	5	55	0	19	0	235	4	689	13	24445	124	837222	14503	1007	876307	877314	59	
1997	20	141	0	38	2	320	18	2973	20	3445		1323985	162744	3512	1490174	1493686	59	

Table 10 Abundance indices (1 000) by year, stratum, West and East Greenland and total for juvenile redfish <17 cm (*Sebastes spp.*), 1982-97. Confidence intervals (CI) are given at the 95% level of significance in per cent of the stratified mean. Incomplete survey coverage off East Greenland in 1984, 1992 and 1994!

Year	1.1	1.2	2.1	2.2	3.1	3.2	4.1	4.2	5.1	5.2	6.1	6.2	7.1	7.2	Σ West	Σ East	Σ	CI
1982	1057	358	121	27	8	42	22			152		607		1553	1835	2312	3947	44
1983	3956	505	14	138	9	17	21			92	8	1709		859	4660	2668	7328	56
1984	5021	3714	20	219	141	28	14			129		693		206	9157	1028	10185	67
1985	4889	9615	54	2712	47	87	55		817414	149899	210	5068	98	17439	972689	990128	164	
1986	10740	237636	113	1811	54	218	38			2651	69	12312		5757	250610	20789	271399	168
1987	12455	113990	4		20		18		2343	2580	132	8961		123715	126487	137731	264218	87
1988	19679	42481	0	107	20	139	0		1579	2983	896	13064		18457	62426	36979	99405	41
1989	7717	13160	3071	5370	18	0	69		1331	3171	150	4274		2155	29405	11081	40486	36
1990	11256	35932	15417	1538	73	0	6199	848	2267	3183	482	13708		4358	71263	23998	95261	52
1991	51939	59845	34871	22668	13692	2508	892	1541	45453	3051	209	1708		622	187956	51043	238999	38
1992	25715	19084	12691	17277	17463	13973	41	13718						1373	119962	1373	121335	54
1993	5460	39035	664	11331	355	2773	14		3401243	2403634	244	810639		6009	59632	6621769	6681401	111
1994	3405	12002	9827	4013	1189	1731	10843	9867						57889	52877	57889	110766	95
1995					399	10236	855	34694	274128	2671933	4072	188899		3061	46184	3142093	3188277	106
1996	457	14357	5210	9377	26961	11571	2488	107237	405272	223348	1373189	2423		3071	177658	2007303	2184961	98
1997	6519	47117	0	15852	43421	20194	444	68931	225859	89354		374542		1372	202479	691127	893605	62

Table 11 Biomass indices (t) by year, stratum, West and East Greenland and total for juvenile redfish <17 cm (*Sebastes spp.*), 1982-97. Confidence intervals (CI) are given at the 95% level of significance in per cent of the stratified mean. Incomplete survey coverage off East Greenland in 1984, 1992 and 1994!

Year	1.1	1.2	2.1	2.2	3.1	3.2	4.1	4.2	5.1	5.2	6.1	6.2	7.1	7.2	Σ West	Σ East	Σ	CI
1982	37	13	6	1	0	2	1			11		36		72	60	119	179	41
1983	103	21	1	6	0	1	1			5	0	73		17	133	95	228	51
1984	91	104	1	5	5	1	1			4		19		9	208	32	240	71
1985	82	367	2	58	2	3	1		15335	7129	6	200		5	515	22675	23190	142
1986	454	6645	3	77	2	6	1			123	3	218		73	7188	417	7605	168
1987	265	5021	0		1	0			147	137	4	288		6502	5287	7078	12365	93
1988	218	1491	0	4	1	5	0		67	144	42	618		1414	1719	2285	4004	56
1989	111	270	22	49	0	0	1		81	167	7	317		135	453	707	1160	42
1990	99	369	63	20	0	0	9	2	67	118	20	833		268	562	1306	1868	58
1991	198	797	73	242	29	24	2	15	563	94	4	63		34	1380	758	2138	46
1992	152	385	49	111	74	220	1	65						18	1057	18	1075	54
1993	72	512	17	265	6	77	1		51857	75876	12	48523		260	950	176328	177278	90
1994	26	216	55	57	30	64	141	277						2704	866	2704	3570	132
1995			6	330	10	347	3834	40792	46	9749		190		693	54611	55304	97	
1996	3	285	13	117	91	297	19	3301	5840	10853	26882	135		171	4126	43881	48007	96
1997	61	344	0	214	163	544	15	2437	5017	2141		16112		73	3779	23344	27123	81

Table 12 Abundance indices (1 000) by year, stratum, West and East Greenland and total for American plaice (*Hippoglossoides platessoides*), 1982-97. Confidence intervals (CI) are given at the 95% level of significance in per cent of the stratified mean. Incomplete survey coverage off East Greenland in 1984, 1992 and 1994!

Year	1.1	1.2	2.1	2.2	3.1	3.2	4.1	4.2	5.1	5.2	6.1	6.2	7.1	7.2	Σ West	Σ East	Σ	CI	
1982	31584	5093	29597	5735	2844	2133	1042			912	1094		2939	78028	4945	82973	30		
1983	46602	6481	55494	2870	2725	461	811			1095	210	2787		7268	115444	11360	126804	49	
1984	18250	6258	53765	4366	2929	2244	1793			1311		2809		3812	89605	7932	97537	43	
1985	21387	5973	22820	6186	2632	239	3162			1884	2823	422	7029		5966	62399	18124	80523	23
1986	22038	11393	58741	9556	2937	2388	4462			1817	186	10365		5105	111515	17473	128988	39	
1987	23322	3314	26226		2356		1030			2415	5636	293	6758		2700	56248	17802	74050	26
1988	10963	3475	8026	5698	3565	799	1036			417	1638	180	8629		2807	33562	13671	47233	19
1989	9371	4454	11363	3775	8764	0	1445			722	637	304	4605		3852	39172	9920	49092	28
1990	8617	6464	8227	2614	1083	0	1492	605	995	1994	440	6417		5813	29102	15459	44561	25	
1991	7826	4536	5168	1899	1517	639	1249	952	1327	3913	1240	8895		6964	23786	22339	46125	18	
1992	8529	4997	3019	2704	1233	1707	1743	174						6696	24106	6896	30802	28	
1993	5856	3284	1202	1213	631	694	398			1734	3996	1111	17920		4991	13278	29752	43030	17
1994	2212	3524	1488	1514	624	282	1661	189						2490	11494	2490	13984	21	
1995														4249	3886	32915	36801	18	
1996	3718	1337	956	1424	1946	772	1566	472	3549	4366	5842	25777		6773	12191	46307	58498	17	
1997	8656	3262	2585	3543	2973	1288	2427	109	1920	2995				23845	4548	24842	33307	58149	21

Table 13 Biomass indices (t) by year, stratum, West and East Greenland and total for American plaice (*Hippoglossoides platessoides*), 1982-97. Confidence intervals (CI) are given at the 95% level of significance in per cent of the stratified mean. Incomplete survey coverage off East Greenland in 1984, 1992 and 1994!

Year	1.1	1.2	2.1	2.2	3.1	3.2	4.1	4.2	5.1	5.2	6.1	6.2	7.1	7.2	Σ West	Σ East	Σ	CI	
1982	6048	947	7797	1151	919	376	156		265		211		682	17394	1158	18552	32		
1983	7450	1154	11772	607	1008	88	166		340	53	747		2191	22245	3331	25576	41		
1984	1704	761	8663	807	607	387	365		240		571		941	13294	1752	15046	45		
1985	1940	601	3862	1062	520	49	321		329	654	68	1034		1596	8355	3681	12036	22	
1986	2149	1147	8429	1385	703	452	460		844	45	2621		1867	14725	5377	20102	30		
1987	3129	338	5470		645	228			992	1103	62	1245		799	9810	4201	14011	30	
1988	919	293	1699	808	814	137	236		79	436	28	1586		752	4906	2881	7787	20	
1989	520	296	1477	371	2120	0	288		110	115	54	847		737	5072	1863	6935	40	
1990	393	397	1220	314	213	0	287		221	180	423	75	1073		918	3045	2669	5714	22
1991	349	399	487	280	265	125	188		172	213	1423	262	1748		1571	2245	5217	7462	18
1992	582	419	229	183	151	250	152	25							1063	1991	1063	3054	26
1993	324	222	83	102	67	71	26		174	418	159	2386		850	895	3987	4882	17	
1994	145	416	134	143	64	34	109	28							631	1073	631	1704	25
1995					70	154	123		58	134	524	246	3275		635	405	5014	5419	21
1996	214	100	67	164	158	78	149	38	316	521	1195	3987		1163	968	7182	8150	22	
1997	490	265	209	343	353	168	185	7	193	308	3874			1019	2020	5392	7412	27	

Table 14 Abundance indices (1 000) by year, stratum, West and East Greenland and total for Atlantic wolffish (*Anarhichas lupus*), 1982-97. Confidence intervals (CI) are given at the 95% level of significance in per cent of the stratified mean. Incomplete survey coverage off East Greenland in 1984, 1992 and 1994!

Year	1.1	1.2	2.1	2.2	3.1	3.2	4.1	4.2	5.1	5.2	6.1	6.2	7.1	7.2	Σ West	Σ East	Σ	CI	
1982	11113	2956	3457	2318	1821	458	946		830		486		604	23069	1920	24989	23		
1983	7570	3186	1720	485	1472	211	785		449	172	915		1831	15429	3367	18796	24		
1984	5774	1278	1542	186	1382	111	750		1237		2807		2747	11023	6791	17814	27		
1985	5372	2718	1420	369	955	999	906		1995	2645	386	3476		4249	12739	12751	25490	19	
1986	4960	1705	1968	635	1500	512	811			1645	347	2444		6843	12091	11279	23370	19	
1987	5326	1644	888		1023		686		4481	3648	638	4265		6925	9567	19957	29524	15	
1988	4927	1833	892	449	1136	555	705		1825	4956	271	2797		4207	10497	14056	24553	21	
1989	3672	674	1668	516	2900		1130		1249	2014	471	1639		3684	10560	9057	19617	21	
1990	3511	1076	1980	295	1899		1048		606	2114	1182	622	2042		5336	10415	11296	21711	17
1991	2841	1011	967	756	2592	639	563		494	3232	2562	448	1625		2275	9863	10142	20005	21
1992	3013	375	1509	937	3360	916	947	2107							5437	13164	5437	18601	26
1993	4724	1713	703	425	573	326	384		1231	5380	112	5379		4452	8848	16554	25402	28	
1994	1543	1197	1725	567	1225	153	4516	1046							5523	11972	5523	17495	48
1995				1349	563	528	705	2043	3976	601	5901		5051	3145	17572	20717	26		
1996	713	943	253	381	2574	622	797	1076	4815	4210	1299	13100		8814	7359	32238	39587	21	
1997	3001	1279	1126	335	3771	1149	1538	2863	4413	5124	11264		7489	15062	28289	43351	30		

Table 15 Biomass indices (t) by year, stratum, West and East Greenland and total for Atlantic wolffish (*Anarhichas lupus*); 1982-97. Confidence intervals (CI) are given at the 95% level of significance in per cent of the stratified mean. Incomplete survey coverage off East Greenland in 1984, 1992 and 1994!

Year	1.1	1.2	2.1	2.2	3.1	3.2	4.1	4.2	5.1	5.2	6.1	6.2	7.1	7.2	Σ West	Σ East	Σ	CI		
1982	9910	2974	5202	3926	2274	475	1241			854	202		207	26002	1263	27265	31			
1983	3663	3441	2085	471	1770	271	1087			368	65	519		921	12788	1873	14661	31		
1984	3089	673	1284	190	820	72	870			600		1211		754	6998	2565	9563	24		
1985	1836	1134	1019	199	581	557	633			698	1329	65	1108		2155	5959	5355	11314	17	
1986	1784	912	1440	434	973	458	768				794	97	988		3730	6769	5609	12378	16	
1987	2192	520	574		1049		615			1625	1705	83	1688		4509	4950	9610	14560	16	
1988	1104	384	798	299	881	340	697			848	1804	39	791		2559	4503	6041	10544	16	
1989	689	222	620	247	1749		1037			436	900	134	800		1689	4564	3959	8523	19	
1990	710	177	497	111	655		660			321	799	649	135	550		1897	3131	4030	7161	16
1991	457	165	160	181	675	149	249			214	1208	679	159	621		956	2230	3623	5853	20
1992	437	79	321	237	832	201	231	631							2221	2969	2221	5190	27	
1993	647	315	101	80	130	67	108			403	1236	22	1125		1446	1448	4232	5680	22	
1994	214	209	376	97	285	26	865	171							2291	2243	2291	4534	43	
1995					248	69	131	113		592	793	101	2610		1809	561	5905	6466	25	
1996	64	262	43	68	487	114	169	238		1231	972	412	4611		2452	1445	9678	11223	19	
1997	306	239	89	77	665	164	352	372		859	731	2705		2304	2264	6599	8863	24		

Table 16 Abundance indices (1 000) by year, stratum, West and East Greenland and total for spotted wolffish (*Anarhichas minor*), 1982-97. Confidence intervals (CI) are given at the 95% level of significance in per cent of the stratified mean. Incomplete survey coverage off East Greenland in 1984, 1992 and 1994!

Year	1.1	1.2	2.1	2.2	3.1	3.2	4.1	4.2	5.1	5.2	6.1	6.2	7.1	7.2	Σ West	Σ East	Σ	CI
1982	382	154	352	152	129	14	325		105	49	124	1508	278	1786	29			
1983	186	38	148	4	90	39	368		49	38	81	153	873	321	1194	32		
1984	183	60	176	14	89	23	243		55	292	172	788	519	1307	30			
1985	204	114	105	4	30	61	110		19	50	57	76	313	628	515	1143	33	
1986	481	203	109	38	68	16	117		70	12	55	409	1032	546	1578	22		
1987	306	211	63		129	238			107	102	32	131	248	947	620	1567	28	
1988	232	151	44	15	145	33	315		220	44	26	112	172	935	574	1509	25	
1989	245	131	50	25	281		112		92	65	7	10	297	844	471	1315	34	
1990	107	201	38	19	100		151	6	118	142	56	175	211	622	702	1324	27	
1991	335	141	27	33	54	5	76	51	173	43	13	115	129	722	473	1195	27	
1992	42	47	15	107	36	23	14	29					200	313	200	513	63	
1993	85	223	49	51	36	17	69		51	0	0	146	405	530	602	1132	35	
1994	63	108	61	62	22	9	27	7					1259	359	1259	1618	79	
1995													787	37	1009	1048	72	
1996	0	56	55	38	18	11	0	7	68	117	25	179	916	185	1303	1488	45	
1997	102	58	16	38	60	16	49	73	17	188	257		578	413	1040	1453	46	

Table 17 Biomass indices (t) by year, stratum, West and East Greenland and total for spotted wolffish (*Anarhichas minor*), 1982-97. Confidence intervals (CI) are given at the 95% level of significance in per cent of the stratified mean. Incomplete survey coverage off East Greenland in 1984, 1992 and 1994!

Year	1.1	1.2	2.1	2.2	3.1	3.2	4.1	4.2	5.1	5.2	6.1	6.2	7.1	7.2	Σ West	Σ East	Σ	CI
1982	2195	470	1897	826	532	42	1989		409	90	179	7951	678	8629	43			
1983	1674	256	656	6	562	183	2357		200	201	417	1007	5694	1825	7519	37		
1984	848	196	1036	16	447	55	1358		316	1539	529	3956	2384	6340	34			
1985	13	118	570	0	134	307	681		779	172	98	369	702	1823	2120	3943	44	
1986	1155	307	567	63	371	35	1004		239	48	360	1312	3502	1959	5461	27		
1987	654	125	334		1029	2035			1090	512	264	529	954	4177	3349	7526	29	
1988	137	85	194	87	1141	101	3011		2007	177	151	756	674	4756	3765	8521	38	
1989	373	32	168	40	1382		847		519	485	70	27	855	2842	1956	4798	36	
1990	83	82	199	7	666		1216	3	859	576	128	1153	616	2256	3332	5588	31	
1991	28	30	2	9	251	5	723	179	1547	239	55	663	195	1227	2699	3926	41	
1992	7	6	0	7	30	4	36	35					1109	125	1109	1234	104	
1993	65	40	16	33	34	17	210		283	0	0	194	843	415	1320	1735	43	
1994	29	25	74	26	10	2	142	3					1772	311	1772	2083	76	
1995					67	40	219	0	432	1	48	350	1550	326	2381	2707	64	
1996	0	137	33	42	38	8	0	10	468	150	89	276	2480	268	3443	3711	68	
1997	75	9	26	38	37	2	23	57	49	122	1161		1303	267	2636	2903	69	

Table 18 Abundance indices (1 000) by year, stratum, West and East Greenland and total for starry skate (*Raja radiata*), 1982-97. Confidence intervals (CI) are given at the 95% level of significance in per cent of the stratified mean. Incomplete survey coverage off East Greenland in 1984, 1992 and 1994!

Year	1.1	1.2	2.1	2.2	3.1	3.2	4.1	4.2	5.1	5.2	6.1	6.2	7.1	7.2	Σ West	Σ East	Σ	CI
1982	5385	1625	1412	473	556	83	163		0	0	172	9697	172	9869	38			
1983	4798	589	818	360	350	28	59		0	0	0	0	7000	0	7000	87		
1984	2740	1673	653	505	149	481	114		37		146	309	6315	492	6807	42		
1985	2241	2393	1848	689	397	56	256		0	0	0	18	166	7878	184	8062	44	
1986	2178	2807	766	326	295	131	203		0	0	39	177	6706	216	6922	46		
1987	1792	537	653		290		64		10	29	3	22	181	3336	245	3581	30	
1988	3879	1046	995	769	334	39	85		4	44	0	37	73	7147	158	7305	39	
1989	11966	2141	3860	694	607		150		11	0	0	30	187	19418	228	19646	38	
1990	7142	1981	2489	548	272	549	345	10	49	7	0	489	13326	555	13881	51		
1991	1970	480	1220	261	810	130	96	65	12	53	25	73	97	4832	260	5092	26	
1992	4456	598	2843	1531	496	523	206	58					200	10711	200	10911	50	
1993	2263	352	684	279	189	264	96		0	78	14	97	196	4127	385	4512	39	
1994	2529	379	872	271	232	79	398	15					1059	4775	1059	5834	43	
1995					181	301	115	15	0	0	7	34	315	612	356	968	59	
1996	1273	126	429	76	115	111	58	29	53	26	25	179	324	2217	607	2824	29	
1997	4886	493	879	46	120	122	180	0	17	141	218		707	6726	1083	7809	41	

Table 19 Biomass indices (t) by year, stratum, West and East Greenland and total for starry skate (*Raja radiata*), 1982-97. Confidence intervals (CI) are given at the 95% level of significance in per cent of the stratified mean. Incomplete survey coverage off East Greenland in 1984, 1992 and 1994!

Year	1.1	1.2	2.1	2.2	3.1	3.2	4.1	4.2	5.1	5.2	6.1	6.2	7.1	7.2	Σ West	Σ East	Σ	CI
1982	2992	810	1328	340	409	58	154		0	0	0	0	182	6091	182	6273	36	
1983	969	193	703	132	332	27	58		0	0	0	0	0	2414	0	2414	34	
1984	731	334	403	96	136	126	94		19		153		307	1920	479	2399	31	
1985	494	427	803	181	159	46	57		0	0	0	23	217	2187	240	2407	23	
1986	517	526	421	84	122	64	40		0	0	52		242	1774	294	2088	28	
1987	415	149	306	185		12		20	45	1	11		221	1067	298	1365	29	
1988	654	123	502	238	175	19	34		0	66	0	14		88	1745	168	1913	28
1989	2077	429	979	107	314		90		1	0	0	6	256	3996	263	4259	31	
1990	977	263	528	56	91	113	201		0	24	0	0	610	2229	634	2863	45	
1991	278	80	182	36	245	42	11	33	1	3	1	88	94	907	187	1094	28	
1992	328	93	138	135	222	89	23	27					290	1055	290	1345	49	
1993	343	88	82	31	29	24	3		0	2	0	61	177	600	240	840	28	
1994	229	72	142	30	91	14	55	11					1315	644	1315	1959	62	
1995	95	24	38	23	21	16	9	13	18	6	1	113	194	239	332	571	44	
1997	354	96	181	6	16	29	33	0	0	13	55		197	715	264	979	35	

Table 20 Number of temperature measurements by stratum and total, 1982-97.

Year	1.1	1.2	2.1	2.2	3.1	3.2	4.1	4.2	5.1	5.2	6.1	6.2	7.1	7.2	Σ	
1982	20	11	16	7	9	6	13	2	0	0	0	0	3	0	5	92
1983	26	11	25	11	17	5	18	4	1	6	3	14	0	7	148	
1984	23	10	18	7	13	4	16	4	4	4	2	7	0	3	115	
1985	10	8	26	10	17	5	21	4	5	21	14	49	0	28	218	
1986	27	9	16	7	16	7	18	3	3	15	14	37	1	34	207	
1987	25	11	21	4	18	3	21	3	4	6	3	16	0	7	142	
1988	34	20	28	5	18	5	16	2	17	8	5	36	0	24	218	
1989	23	10	23	8	3	2	22	3	13	11	9	23	0	11	161	
1990	15	6	23	7	14	3	17	4	15	15	6	15	0	11	151	
1991	16	9	21	5	12	4	13	1	0	0	0	0	0	0	81	
1992	6	6	6	5	6	5	6	4	0	0	0	0	0	5	49	
1993	9	6	8	6	10	8	7	0	8	5	6	18	0	14	105	
1994	16	13	12	8	10	6	7	4	0	0	0	0	0	0	76	
1995	0	0	3	0	10	7	10	5	8	6	6	17	0	12	84	
1996	5	5	8	5	12	5	10	5	6	9	5	13	0	9	97	
1997	5	6	5	5	6	5	8	5	5	5	4	8	0	8	75	

Table 21 Mean near bottom temperature ($^{\circ}$ C) by stratum and weighted total (by stratum area), 1982-97. Incomplete coverage of the survey area in 1982, 1984, 1991, 1992, and 1994!

Year	1.1	1.2	2.1	2.2	3.1	3.2	4.1	4.2	5.1	5.2	6.1	6.2	7.1	7.2	Total	
1982	2.540	3.627	1.953	3.100	3.256	3.633	2.623								4.600	3.139
1983	2.028	3.713	1.420	3.819	2.139	4.808	2.160								2.943	3.012
1984	1.365	2.790	1.617	3.886	2.462		2.519								4.129	2.698
1985	4.190	5.154	3.116	4.612	2.614	4.336	4.444		5.040	5.190	4.421	4.300			3.300	4.181
1986	3.669	4.393	4.014	5.073	4.203	5.066	4.102			4.796	4.042	4.516			3.347	4.136
1987	3.086	4.890	3.393		3.504		3.530			4.467		4.400			3.300	3.783
1988	2.548	4.328	3.034	4.956	4.228	5.234	4.332		4.479	4.559	4.298	4.578			3.792	3.959
1989	2.323	3.953	2.718	4.525			2.579		3.392	3.743	3.648	4.064			3.146	3.295
1990	2.497	3.922	3.000	4.809	3.421		2.516		4.395	4.570	3.252	4.019			3.025	3.461
1991	3.533	4.726	3.477	4.204	3.016		2.997								3.558	
1992	3.900	4.418	2.911	4.457	2.985	4.691	1.938								3.472	3.489
1993	3.007	4.003	2.360	3.360	4.711	4.959	2.773		3.771	4.056	4.327	4.394			2.820	3.597
1994	2.914	4.436	3.747	4.641	3.847	5.109	3.773								3.620	
1995					4.229	4.614	3.469	4.242	2.601	3.623	3.683	4.318			3.834	3.862
1996	4.614	5.506	4.414	5.688	5.610	5.700	5.057	5.732	4.505	5.129	5.320	4.903			2.848	4.709
1997	3.304	4.938	4.022	5.180	4.570	5.478	4.606	5.540	4.578	4.742		4.266			3.464	4.189




Fig. 1 Stratification scheme of the survey area as specified in Table 2 and positions of survey hauls in 1997.




Fig. 2 Abundance indices off West, East Greenland and total for Atlantic cod (*Gadus morhua*) as listed in Table 4, 1982-97.




Fig. 4 Mean individual weight off West, East Greenland and total for Atlantic cod (*Gadus morhua*) as derived from Tables 4 and 5, 1982-97.




Fig. 3 Biomass indices off West, East Greenland and total for Atlantic cod (*Gadus morhua*) as listed in Table 5, 1982-97.




Fig. 5 Length composition for Atlantic cod (*Gadus morhua*), 1995-97.




Fig. 6 Abundance indices off West, East Greenland and total for golden redfish ≥ 17 cm (*Sebastes marinus*) as listed in Table 6, 1982-97.




Fig. 8 Mean individual weight off West, East Greenland and total for golden redfish ≥ 17 cm (*Sebastes marinus*) as derived from Tables 6 and 7, 1982-97.




Fig. 7 Biomass indices off West, East Greenland and total for golden redfish ≥ 17 cm (*Sebastes marinus*) as listed in Table 7, 1982-97.




Fig. 9 Length composition for golden redfish ≥ 17 cm (*Sebastes marinus*), 1995-97.




Fig. 10 Abundance indices off West, East Greenland and total for deep sea redfish ≥ 17 cm (*Sebastes mentella*) as listed in Table 8, 1982-97.




Fig. 12 Mean individual weight off West, East Greenland and total for deep sea redfish ≥ 17 cm (*Sebastes mentella*) as derived from Tables 8 and 9, 1982-97.




Fig. 11 Biomass indices off West, East Greenland and total for deep sea redfish ≥ 17 cm (*Sebastes mentella*) as listed in Table 9, 1982-97.




Fig. 13 Length composition for deep sea redfish ≥ 17 cm (*Sebastes mentella*), 1995-97.




Fig. 14 Abundance indices off West, East Greenland and total for juvenile redfish <17 cm (*Sebastes spp.*) as listed in Table 10, 1982-97.




Fig. 16 Mean individual weight off West, East Greenland and total for juvenile redfish <17 cm (*Sebastes spp.*) as derived from Tables 10 and 11, 1982-97.




Fig. 15 Biomass indices off West, East Greenland and total for juvenile redfish <17 cm (*Sebastes spp.*) as listed in Table 11, 1982-97.




Fig. 17 Length composition for juvenile redfish <17 cm (*Sebastes spp.*), 1995-97.




Fig. 18 Abundance indices of West, East Greenland and total for American plaice (*Hippoglossoides platessoides*) as listed in Table 12, 1982-97.




Fig. 20 Mean individual weight off West, East Greenland and total for American plaice (*Hippoglossoides platessoides*) as derived from Tables 12 and 13, 1982-97.




Fig. 19 Biomass indices off West, East Greenland and total for American plaice (*Hippoglossoides platessoides*) as listed in Table 13, 1982-97.




Fig. 21 Length composition for American plaice (*Hippoglossoides platessoides*), 1995-97.




Fig. 22 Abundance indices off West, East Greenland and total for Atlantic wolffish (*Anarhichas lupus*) as listed in Table 14, 1982-97.




Fig. 24 Mean individual weight off West, East Greenland and total for Atlantic wolffish (*Anarhichas lupus*) as derived from Tables 14 and 15, 1982-97.




Fig. 23 Biomass indices off West, East Greenland and total for Atlantic wolffish (*Anarhichas lupus*) as listed in Table 15, 1982-97.




Fig. 25 Length composition for Atlantic wolffish (*Anarhichas lupus*), 1995-97.




Fig. 26 Abundance indices off West, East Greenland and total for spotted wolffish (*Anarhichas minor*) as listed in Table 16, 1982-97.




Fig. 28 Mean individual weight off West, East Greenland and total for spotted wolffish (*Anarhichas minor*) as derived from Tables 16 and 17, 1982-97.




Fig. 27 Biomass indices off West, East Greenland and total for spotted wolffish (*Anarhichas minor*) as listed in Table 17, 1982-97.




Fig. 29 Length composition for spotted wolffish (*Anarhichas minor*), 1995-97.




Fig. 30 Abundance indices off West, East Greenland and total for starry skate (*Raja radiata*) as listed in Table 18, 1982-97.




Fig. 32 Mean individual weight off West, East Greenland and total for starry skate (*Raja radiata*) as derived from Tables 18 and 19, 1982-97.



Fig. 31 Biomass indices off West, East Greenland and total for starry skate (*Raja radiata*) as listed in Table 19, 1982-97.




Fig. 33 Length composition for starry skate (*Raja radiata*), 1995-97.




Fig. 34 Mean near bottom temperature by stratum and total off Greenland as listed in Table 21, 1982-97. Weighted (by stratum area) mean temperature off Greenland is illustrated as a bold line, respectively.