



**SCIENTIFIC COUNCIL MEETING – JUNE 1999**

**A review on Roughhead Grenadier (*Macrourus berglax*) Population Structure  
on Flemish Cap (NAFO Div. 3M) 1991-1998**

by

**H. Murua<sup>1</sup>, A. Sarasua<sup>1</sup>, E. Rodríguez-Marín<sup>2</sup>**

<sup>1</sup> AZTI, Tecnological Institute for Fisheries and Food, Satrustegi 8, 20008 Donostia, Spain

<sup>2</sup> Instituto Español de Oceanografía. Apto. 240, 39080 Santander, Spain

**ABSTRACT**

The European Union has conducted since 1988 an annual bottom trawl survey in Flemish Cap (NAFO Div. 3M) in the 200-720 m depth range. Roughhead grenadier (*Macrourus berglax*) is considered a target species within this project and its population structure is being studied.

The information on roughhead grenadier recorded during the last 8 EU surveys (1991-1998) in Flemish Cap is presented in this paper. Age and length composition of the catches showed clear differences between the two sexes. The importance of males in the capture declines in larger fish, disappearing from the capture in largest length classes. Capture is dominated by the 6-8 age-classes.

The 1984-1986 cohorts dominated the catches during the first years. The importance of these annual classes have declined sharply during last 4 years and the captures now are dominated by the abundant 1990 cohort.

The oldest male found was 14 years old, while oldest female was 18 years old. Results show that *M. berglax* has a prolonged life cycle and multiaged population structure with differences in growth and mortality between males and females.

**INTRODUCTION**

The roughhead grenadier (*Macrourus berglax* Lacépède, 1802) is an abundant and widespread fish species in the north Atlantic and is usually found both on the shelf and on the continental slope (Scott and Scott, 1988; Savvatimsky, 1994). It is predominant in depths ranging from 400 to 1200 m, although they may inhabit depths between 200-2000 m (Snelgrove and Haedrich, 1985; Cardenas *et al.* 1996), and it has been found in depths up to 2700 m (Wheeler, 1969).

Roughhead grenadier is becoming an important commercial fish in NAFO Regulatory Area and reliable information is needed for its assesment. The fishery for *M. berglax* is unregulated as it has been mainly taken as by-catch in traditional fisheries. The impact of the fishing pressure is unknown at present (Atkinson, 1996).

Spain has been developing a deep water trawl fishery in the NAFO Regulatory Area since 1990, with Greenland halibut as target species. The fleet started operating in the northern area of Flemish Pass, but has gradually extended the area of its activities and now fishes in practically all the continental slope of the NAFO Regulatory Area (Cardenas, 1996). At the beggining of the fishery, grenadiers (*Coryphaenoides rupestris* and *Macrourus berglax*) were usually discarded. Since then, its commercial importance increased and it has become the main component in the by-catch of the Spanish and Portuguese Greenland halibut fishery (Paz and Iglesias, 1994; Alpoim *et al.*, 1997).

Since 1988 EU has conducted an annual random-stratified bottom trawl survey in Flemish Cap (NAFO Div. 3M) in the 200-720 m depth range. The objective of this scientific survey is to obtain abundance indices and to study the population structure and biological parameters of the main species in the area. Since 1994 *M. berglax* is considered a target species within this project and its population structure is being studied (Casas, 1994; Sainza, 1996; Alpoim, 1997; Sarasua *et al.*, 1998).

Limited information on age structure and growth rate of *M. berglax* is available in scientific literature. Savvatimsky (1971, 1984, 1989, 1994) and Jorgensen (1996) have carried out studies on this species in the NW Atlantic (NAFO Divisions 0,2GHJ,3K and 1ABCD respectively), basing findings on age readings from scales. The age structure and growth parameters of roughhead grenadier have been estimated by Casas (1994), Sainza (1996), Alpoim (1997), Sarasua *et al.*, (1998) and Rodríguez-Marín *et al.*, (1998) from otolith readings of specimens captured in NAFO Divisions 3LM. Eliassen (1983) also performed age estimation by otolith reading from roughhead caught in the continental slope of Norway.

The population structure of roughhead grenadier on Flemish Cap in the period 1991-1998 is presented in this paper. The results are presented taking into account that the survey only covers the shallowest distribution area of *M. berglax*.

## MATERIAL AND METHODS

Data were collected on the annual random-stratified bottom trawl surveys carried out by the European Union on the area. Methodology is described by Vázquez (1998). Data on *M.berglax* of the 1991-1998 period will be used in this paper. Data on age structure and sex-ratio are only available for the 1993-1998 period. Otolith sampling began in 1994, and since then a total of 2177 otoliths have been read. Annual length-age keys have been applied for each year. For years when otoliths were not sampled the mean 96-97 age-length key (the most consistent readings) were applied.

Otoliths were broken through the nucleus and read by transmitted light (Casas, 1994). Many difficulties in reading Macrouridae age from otoliths and scales have been reported previously (Savvatimsky, 1984). Age reading in larger fish (more than 9 years old) is even more complicated, because many rings are present and they lie close to each other. Nevertheless, intercalibration of readings between three readers has been done and 80% of agreement has been reached. Differences were  $\pm 1$  year in otoliths between 2-10 years and 1, 2 years in older than 10 years (Rodríguez-Marín *et al.*, 1998).

Individuals were measured from tip of snout to base of first anal-fin ray, in 0.5 cm intervals, as adopted by NAFO in June 1980 (Atkinson, 1991) as a standard measurement for roundnose and roughhead grenadiers. Data are given in 1 cm intervals. Length is presented as anal-fin length (AFL). Total weight was recorded accurate to the nearest 10 g.

## RESULTS AND DISCUSSION

Table 1 shows length distributions of roughhead grenadier for the 1991-1998 period. Captures are dominated by the 14-20 cm length classes, 54.1 % of the total catch. The average AFL for both sexes is 15,47 cm. This value is smaller than the values found by Savvatimsky (1994) and Cardenas *et al.* (1996). The former gives an average AFL of 19.8 cm (51.25 cm total length) for Div. 3K, and the latter gives a mean AFL of 21,12 cm (54,39 cm TL) for specimens caught in a long-line survey in Div. 3LMN. These differences are related to the fishing gear employed and depth where fish were caught, because length has a tendency to increase with depth, from the shallowest stratum to the deepest (Cárdenas *et al.*, 1996).

Annual length frequencies by sex are presented in Table 2. The importance of males in the capture declines in larger fish and they disappear from the capture in largest length classes. Largest male found in the scientific surveys was 25 cm while females are larger reaching 32 cm long. Average AFL for females is also greater than for males. Female's mean AFL for the 1993-1998 period in Flemish Cap was 16.69 cm, while mean AFL for males was 15.41 cm. This sexual difference is consistent with data found in the literature. Savvatimsky (1989) gives an average AFL of 18 cm (47 cm total length) and 21.2 cm (54.6 cm total length) for males and females respectively in 3KLN.

Those differences have also been seen in the commercial fleet in 3LN, where females are larger than males (Godinho *et al.*, 1996).

The mean AFL-age key for 1994-1998, as well as mean length at age and standard deviation are given by sex in Table 3. Mean length at age is similar for males and females for ages under 9 years, but males grow slower from this length onwards. Mean lengths at age are higher than those obtained by Savvatimsky (1994) for NAFO Divs. 0B, 2GH and 3K. Savvatimsky (1994) and Jorgensen (1996) described similar growing pattern using scales for aging fish, they found that the differences between sexes in size at age come about from 10 years onwards. This fact could be explained due to the different aging method used in their studies and in ours, or due to different latitude of the sampling areas where specimens were obtained, because temperature differences would cause slower growth, and a delay in reaching sexual maturity (Rodríguez-Marín *et al.*, 1998).

Table 4 shows age composition by sexes for *Macrourus berglax* on Flemish Cap in 1993-1998. Females are more abundant in almost all year-classes, but differences increase in largest groups (>12 year-class). Only females appear in year-classes older than 14. The oldest male found in the study period was 14 years old, while oldest female was 18 years old.

Sexual differences in the growing pattern seem to be related with reaching maturity (Cárdenas, 1996; Murua and Motos, 1997; Scott and Scott, 1988). Scott and Scott (1988) affirm that females grow faster than do males from first maturity in the following species: Atlantic halibut (*Hippoglossus hipoglossus*) and yellowtail flounder (*Limanda ferruginea*). Similar growth differences between sexes have been also found in megrim (*Lepidorhombus whiffiagonis*) (Landa *et al.*, 1996) and anglerfish (*Lophius piscatorius*) (Alfonso-Dias *et al.*, 1996). We also have to bear in mind that a greater longevity of females is not an uncommon phenomenon in many fish species, especially Pleuronectiforms (Scott & Scott, 1988). Mean age for females in Flemish Cap in the 1993-1998 period was 7.9 years, while mean age for males was 7.3 years. Savvatimsky (1994) for Div. 0B, 2GH and 3K found similar differences.

Interannual differences in length and age are shown in Figure 1. The 1984-1986 cohorts dominated the catches during the first years. The importance of these annual classes have declined sharply during last 4 years and the abundant 1990 cohort now dominates captures. The strength of this year-class has been seen in previous surveys (Alpoim, 1997; Sarasua *et al.*, 1998) and it has been confirmed by the 1998 survey (Sarasua *et al.*, 1999).

Average frequency of length and age composition for the whole study period are presented in Figure 2 and Figure 3 respectively. The length and age range found during these 8 surveys are very similar to those cited before in the Northwestern Atlantic (3KLN). Chrzan (1969) gives values of 11.2-34.1 cm AFL and 2-17 years; Savvatimsky (1971) found values of 9.1-26 cm and 3-16 years; and Savvatimsky (1984), ranges of 15-34.9 cm and 6-23 years. Differences can be attributed to differences in depth and sampling gear.

Female-ratio in the whole study period is 51%. This value is lower than the one found by Cardenas *et al.* (1996) in 3LMN, where females made up 71,4% of the catch. However, this difference could be easily explained due to the different area covered by both surveys. As length increases with depth in many species (Cárdenas *et al.*, 1996; Junquera *et al.*, 1992), female ratio might also increase in deeper areas.

Figures 4 and 5 present sex-ratio by age and by length respectively, for the whole study period. In the sex-ratio by age, female proportion fluctuates around 50%-60% the first 12 years. It increases from age 13 upwards. Female-proportion reaches 73% in year-group 13 and 80% in year-group 14. Females are 100% of the captures after that. Similar sex-ratio, with males being more abundant in the central part of the population, is described by Savvatimsky (1994) for Northwestern Atlantic.

Five different areas can be described in the sex-ratio by length figure. In the first length-classes (5-10 cm AFL), females are more abundant than males. This might be due to data scarcity or due to the great amount of specimens with indetermined sex for those length-classes. Between 10-13 cm, sex-ratio fluctuates around 50%. After that, female-ratio decreases to around 40%. This fact is a result of the different growth rate between sexes. Male's growth rate declines when reaching 18 cm long, around 9 years old, while females do not decline growing until reaching 34-35 cm, around 20 years old (Rodríguez-Marín *et al.*, 1998). Therefore, male proportion increases in those central lengths. Data show that after this period, females proportion increases very fast and reaches 100% at

length-classes over 25 cm AFL. The increment in the female-ratio can be due to three different reasons: sexual differences in growth rate, in mortality or a combination of both. In this case, there are certainly sexual differences in growth, which are reflected in the mean length at age (Rodríguez-Marín *et al.*, 1998). And it seems that there are also some differences in mortality, since males disappear from the capture in larger length-classes.

Total mortality ( $Z$ ) by sex was calculated from catch curves, fitting regression lines by sex to ages fully recruited to the fishery. The catch curve (Figure 6). was elaborated from data for five years (1994-1998) (Table 4). Both sexes are fully recruited at age 7, and in fact a different mortality is obtained: 0.28 for females and 0.47 for males. The same fully recruited age and mortality differences between sexes have been found in the Spanish commercial bottom trawl fishery for this species, with practically identical values of  $Z$  (Rodríguez-Marín, pers. com.)

Data available show that *M. berglax* has a prolonged life cycle and multiaged population structure with differences in growth and mortality between males and females. The complex multi-mode length structure and a slow growth is characteristic of deepwater fishes, including grenadiers (Hureau *et al.*, 1979; Casas, 1994; Savvatimsky, 1994). All this results must be taken with care due to the small proportion of the roughhead grenadier distribution area covered by the survey.

Acknowledgements. This study was supported by the European Commission (DG XIV, Study 96-030), Basque Government, IEO, CSIC, and IPIMAR.

## REFERENCES

- Alfonso-Dias, I. P. And J.R.G. Hislop. 1996. The reproduction of anglerfish *Lophius piscatorius* Linnaeus from the north-west coast of Scotland. *Journal of Fish Biology* **49**: (Supplement A), 18-39.
- Alpoim, R. 1997. Age Structure of Roughhead Grenadier (*Macrourus berglax*) on Flemish Cap, 1996. *NAFO SCR Doc.* 97/58. Serial No. N2892. 6p.
- Alpoim, R. M.L. Godinho, A. Avila do Melo and E. Santos. 1997. Portuguese research report for 1996. *NAFO SCS Doc.* 97/9 Ser. No N2857, 43p.
- Atkinson, D.B. 1991. Relationships Between Pre-anal fin Length and Total Length of Roughhead Grenadier (*Macrourus berglax* Lacépède) in The Northwest Atlantic. *J. Northw. Atl. Fish. Sci.*, **11**: 7-9.
- Atkinson, D.B. 1996. Roundnose Grenadier (*Coryphaenoides rupestris*) and Roughhead Grenadier (*Macrourus berglax*) in NAFO Subareas 2+3. *NAFO SCR Doc.* 96/69. Ser. No N2745.
- Casas, J.M. 1994. Age Structure of Roughhead Grenadier (*Macrourus berglax*) on Flemish Cap, 1994. *NAFO SCR Doc.* 94/80. Serial No. N2459. 5p.
- Chrzan, F. 1969. Polish research report. 1968. *ICNAF Redbook*, Part II:63-75.
- De Cardenas, E. 1996. The Female Ratio by Length as an Indicator of Sexual Differences in Mortality of Greenland Halibut (*Reinhardtius hippoglossoides*) at Ages 8+. *NAFO SCR Doc.* 96/35 Ser. No N2710.
- De Cardenas, E., J.M. Casas, R. Alpoim, and H. Murua. 1996. Preliminary Results of the european Long-line Suuvey in the NAFO Regulatory Area. *NAFO SCR Doc.* 96/34 Ser. No N2709, 6p.
- Eliassen, J.E. 1983. Occurrence, gonad development and age distribution of the roughhead grenadier (*Macrourus berglax* Lacepede) (Gadiformes, Macrouridae) along the Continental slope of northern Norway. *ICES, Demersal Fish Committee*, C.M. 1983 /G:41.
- Godinho, M.L., R. Alpoim, A.M. Avila do Melo, and E. Santos. 1996. Portuguese Research Report for 1995. *NAFO SCS Doc.* 96/12. Ser. No N2703. 61 pp.

- Hureau, J.C., P. Geistdorfer, and M. Rannou. 1979. The cology of deep-sea benthic fish. *Sarsia*, **64**(1-2):103-108.
- Jorgensen, O.A. 1996. Distribution and Biology of Grenadiers (Macrouridae) in West Greenland Waters. *J. Northw. Atl. Fish. Sci.* **18**: 7-29
- Junquera, S., S. Iglesias, and E. de Cardenas. 1992. Spanish fishery of Greenland halibut (*Reinhardtius hippoglossoides*) in 1990-1991. *NAFO SCS Doc.* 92/28 Ser. No N2075, 14 pp.
- Landa, J., C. Pineiro, and N. Pérez. 1996. Megrim (*Lepidorhombus whiffiagonis*) growth patterns in the northeast Atlantic. *Fisheries Research* **26**: 279-294.
- Murua, H. and L. Motos. 1997. Length at first maturity of roughhead grenadier, *Macrourus berglax*, in NAFO Divisions, 3MNL. *NAFO SCR Doc.* 97/19. Serial No. N2849.
- Paz, J. And S. Iglesias. 1994. Grenadiers in the spanish Fishery of Greenland Halibut, NAFO Divisions 3LM and N, 1991-1993. *NAFO SCR Doc.* 94/23. Serial No. N2389.
- Rodríguez-Marín, E., M. Ruiz, and A. Sarasua. 1998. Validation of roughhead grenadier (*Macrourus berglax*) otolith reading. *NAFO SCR Doc.* 98/33 Ser. No 3020.
- Sainza, C. 1996. Age Structure of Roughhead Grenadier (*Macrourus berglax*) on Flemish Cap, 1995. *NAFO SCR Doc.* 96/58. Serial No. N2734. 5p.
- Sarasua, A., M. Ruiz, and E. Rodríguez-Marín. 1998. A review on Roughhead Grenadier (*Macrourus berglax*) Population Structure on Flemish Cap (NAFO Div. 3M) 1991-1997. *NAFO SCR Doc.* 98/43. Ser. No 3034. 14pp.
- Sarasua, A., H. Murua, and E. Rodríguez-Marín. 1999. Roughhead Grenadier (*Macrourus Berglax*) Population Structure on Flemish Cap, 1998.. *NAFO SCR Doc.* 99/ Ser. No
- Savvatimsky, P.I. 1971. On age readings in grenadiers (Macrouriformes). *Vopr. Ikhtiol.*, **11**(3): 495-501.
- Savvatimsky, P.I. 1984. Biological Aspects of Roughhead Grenadier (*Macrourus berglax*) from Longline Catches in the Eastern Grand Bank Area, 1982. *NAFO Sci. Coun. Studies*, **7**: 45-51.
- Savvatimsky, P.I. 1989. Investigations of Roughhead Grenadier (*Macrourus berglax* L) in the Northwest Atlantic, 1967-83. *NAFO Sci. Coun. Studies*, **13**: 59-75.
- Savvatimsky P.I. 1994. Age Structure of of Roughhead Grenadier (*Macrourus berglax*) in the Northwest Atlantic, 1985. *NAFO Sci. Coun. Studies*, **20**: 53-64.
- Scott, W. B., and M. G. Scott. 1988.. Atlantic fishes of Canada. *Can Bull. Fish. Aquat. Sci.* 219, 731pp.
- Snelgrove, P.V.R. and R.L. Haedrich. 1985. Structure of the deep demersal fish-fauna off Newfoundland. *Mar. Ecol. Prog. Ser.* **27**: 99-107
- Vazquez, A. 1998. Results from the bottom trawl survey of Flemish Cap in July 1997. *NAFO SCR Doc.* 98/
- Wheeler, A. 1969. *The fishes of the British Isles and Northwest Europe: Anacanthini* (p. 255.259). MacMillan and Co. Ltd., London England, 613 p.

Table 1. Roughhead grenadier length distribution and mean AFL (,000) for each year of the 1991-1998 period.

	91	92	93	94	95	96	97	98	Proportion (%)
3	7	29	0	0	0	7	7	113	0,50
4	14	104	104	0	37	59	35	33	1,20
5	24	321	379	47	45	215	16	95	3,54
6	33	82	120	31	46	63	52	65	1,52
7	77	120	515	65	60	31	111	57	3,21
8	25	57	226	169	72	56	98	66	2,38
9	41	37	112	98	139	46	76	208	2,35
10	40	36	182	231	342	113	94	177	3,76
11	97	49	156	196	295	116	31	121	3,29
12	215	53	200	117	527	160	173	155	4,96
13	253	98	177	100	271	255	195	217	4,85
14	275	259	307	255	131	308	395	496	7,52
15	208	298	560	236	185	212	317	577	8,03
16	221	256	890	213	275	308	208	489	8,86
17	271	187	715	426	332	244	90	345	8,09
18	315	197	613	469	412	244	127	197	7,97
19	266	155	505	440	433	308	163	215	7,70
20	226	210	364	248	280	270	148	159	5,90
21	86	112	282	166	151	114	107	144	3,60
22	81	81	193	101	73	101	113	157	2,79
23	45	74	244	71	16	31	83	50	1,90
24	31	35	95	73	26	46	82	88	1,47
25	0	35	50	72	26	21	45	68	0,98
26	22	62	78	43	19	29	52	38	1,06
27	16	29	31	24	28	21	53	23	0,70
28	25	29	58	8	13	8	22	7	0,53
29	16	21	58	46	0	0	15	23	0,55
30	16	44	17	19	9	8	0	17	0,40
31	0	10	0	31	6	7	7	0	0,19
32	0	14	0	11	7	14	8	0	0,17
33	0	0	0	0	0	0	0	0	0,00
34	0	0	0	0	0	0	0	7	0,02
<b>Total</b>	<b>2946</b>	<b>3094</b>	<b>7231</b>	<b>4006</b>	<b>4256</b>	<b>3415</b>	<b>2923</b>	<b>4407</b>	<b>100,00</b>
<b>Mean AFL (cm)</b>	<b>16,03</b>	<b>15,04</b>	<b>15,31</b>	<b>16,45</b>	<b>15,04</b>	<b>15,24</b>	<b>15,81</b>	<b>15,13</b>	<b>15,47</b>

Table 2. Length distribution (,000) and mean AFL by sex, Flemish Cap 1993-1998.

Length	1993		1994		1995		1996		1997		1998		Total	
	Males	Females	Males	Females	Males	Females	Males	Females	Males	Females	Males	Females	Males	Females
3									7		26	13	33	13
4							23		7	14	13	20	43	34
5				9		6	7	24			28	54	35	99
6		17		8	6	16	31	16	15	15	20	38	78	126
7	45	142	17	15	9	26	0	31	29	82	14	43	123	365
8	70	104	17	32	22	31	16	32	39	59	22	44	208	333
9	60	44	17	32	57	76	30	16	16	60	112	96	349	400
10	67	115	92	65	128	171	70	37	40	54	79	98	604	711
11	100	39	54	42	175	89	38	62		31	79	42	621	394
12	89	111	16	60	217	267	76	71	107	59	104	51	826	886
13	91	86	19	32	89	169	149	100	100	95	98	119	635	770
14	184	123	81	54	73	34	205	103	171	224	305	184	1092	756
15	435	109	55	24	79	88	127	79	139	170	327	250	1241	808
16	520	362	111	17	130	145	183	125	106	102	273	216	1453	1112
17	374	341	155	95	237	70	152	92	45	45	154	191	1354	904
18	256	332	162	69	281	112	191	45	69	58	142	55	1382	783
19	301	196	111	113	270	156	215	87	96	67	145	70	1408	845
20	202	162	84	85	112	156	185	85	88	60	68	91	851	795
21	92	190	33	70	32	119	73	41	70	37	63	81	395	657
22	34	159	17	62	9	58	37	64	14	99	20	137	140	637
23	26	218	14	19		16		31	7	76		50	47	426
24		95		73		26		46		82	7	81	7	429
25	9	41		63		26		21		45		68	9	290
26		78		32		19		29		52		38	0	267
27		31		24		28		21		53		23	0	208
28		58		8		13		8		22		7	0	129
29		58		46				0		15		23	0	142
30		17		19		9		8				17	0	79
31				20		6		7		7			0	46
32				11		7		14		8			0	47
33													0	0
34												7	0	7
<b>Total</b>	<b>2955</b>	<b>3228</b>	<b>1055</b>	<b>1199</b>	<b>1926</b>	<b>1939</b>	<b>1808</b>	<b>1295</b>	<b>1165</b>	<b>1691</b>	<b>2099</b>	<b>2207</b>	<b>12934</b>	<b>13498</b>
<b>Mean AFL (cm)</b>	<b>15,93</b>	<b>17,57</b>	<b>15,95</b>	<b>18,57</b>	<b>15,25</b>	<b>15,70</b>	<b>15,78</b>	<b>16,57</b>	<b>15,01</b>	<b>16,71</b>	<b>14,59</b>	<b>16,15</b>	<b>15,41</b>	<b>16,69</b>

Table 3. Mean Age-Length key (1994-1998).

FEMALES	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	Total
3	2																		2
4		5																	5
5		10																	10
6		7	7																14
7		6	18																24
8		1	19	7															27
9			13	19	1														33
10			6	25	8														39
11				15	16	1													32
12				8	15	15	1	1											40
13				1	17	20	8	1	1										48
14					12	19	10	6											47
15					4	19	14	11	4	1									53
16					2	11	18	10	3	2									46
17						6	19	14	8	1									48
18							12	14	10	3									39
19							3	16	21	8	2	2							52
20								8	16	15	7	5							51
21								4	11	14	9	3	2						43
22									4	13	15	11	5	2					50
23									2	7	5	6			2				24
24									1	5	7	7	11	3	1				35
25										1	6	3	8	7	2				27
26										1	1	10	3	1	5	2			23
27										1	1	1	5	6	4		1		19
28													3	3	1	1			8
29													1	1	3	2			8
30															3				6
31											1				1	1	1	1	3
32													1		1	1			4
34																1			1
<b>Total</b>	<b>2</b>	<b>29</b>	<b>63</b>	<b>75</b>	<b>75</b>	<b>91</b>	<b>85</b>	<b>85</b>	<b>81</b>	<b>67</b>	<b>56</b>	<b>49</b>	<b>45</b>	<b>24</b>	<b>23</b>	<b>8</b>	<b>2</b>	<b>1</b>	<b>861</b>
<b>Mean AFL</b>	<b>3,0</b>	<b>5,6</b>	<b>7,9</b>	<b>10,0</b>	<b>12,3</b>	<b>14,1</b>	<b>15,9</b>	<b>17,3</b>	<b>19,1</b>	<b>20,8</b>	<b>22,5</b>	<b>23,4</b>	<b>24,7</b>	<b>26,0</b>	<b>27,2</b>	<b>29,4</b>	<b>28,5</b>	<b>31,0</b>	
<b>St. Dv.</b>		<b>1,12</b>	<b>1,15</b>	<b>1,18</b>	<b>1,56</b>	<b>1,51</b>	<b>1,67</b>	<b>2,05</b>	<b>2,00</b>	<b>2,19</b>	<b>2,07</b>	<b>2,63</b>	<b>2,12</b>	<b>2,23</b>	<b>2,47</b>	<b>2,83</b>	<b>2,12</b>		

MALES	1	2	3	4	5	6	7	8	9	10	11	12	13	14	Total
3	3	1													4
4	1	5													6
5		5													5
6		6	5												11
7		1	9												10
8		2	9	5											16
9		1	11	13											25
10			6	22	6										34
11			2	13	14	2									31
12			1	2	19	19	4								45
13				2	14	22	9								47
14					12	24	12	6	1						55
15					2	17	17	8	2						46
16						2	14	24	16	9	2	1			68
17							9	16	13	10	9	1			58
18							1	10	20	14	12	10	2	1	70
19								6	12	16	16	13	7	3	74
20									3	12	12	8	15	5	56
21									1	9	4	7	10	5	37
22											2	2	3	1	9
23														1	1
24															1
<b>Total</b>	<b>4</b>	<b>21</b>	<b>43</b>	<b>57</b>	<b>69</b>	<b>108</b>	<b>98</b>	<b>79</b>	<b>73</b>	<b>57</b>	<b>42</b>	<b>37</b>	<b>16</b>	<b>5</b>	<b>709</b>
<b>Mean AFL</b>	<b>3,3</b>	<b>5,5</b>	<b>8,3</b>	<b>10,0</b>	<b>12,4</b>	<b>14,1</b>	<b>15,7</b>	<b>17,1</b>	<b>18,4</b>	<b>18,8</b>	<b>19,3</b>	<b>20,1</b>	<b>20,3</b>	<b>21,2</b>	
<b>St. Dv.</b>	<b>0,50</b>	<b>1,54</b>	<b>1,49</b>	<b>1,13</b>	<b>1,44</b>	<b>1,63</b>	<b>1,79</b>	<b>1,65</b>	<b>1,72</b>	<b>1,40</b>	<b>1,33</b>	<b>1,00</b>	<b>1,25</b>	<b>1,72</b>	



Table 4. Roughhead grenadier age composition (,000), Flemish Cap 1993-1998.

Age	1993			1994			1995			1996		
	Males	Females	Total	Males	Females	Total	Males	Females	Total	Males	Females	Total
2		8	8		9	9	20	38	58	46	24	70
3	112	248	359	17	36	53	92	104	196	35	79	114
4	173	162	335	143	121	264	190	247	437	116	84	200
5	120	151	271	108	139	247	360	443	803	149	172	321
6	445	207	652	187	101	288	324	259	583	365	194	559
7	621	570	1190	224	104	328	385	180	565	191	157	348
8	437	413	850	169	134	303	343	240	583	232	127	359
9	309	353	663	124	160	284	162	192	354	167	85	252
10	278	257	535	42	133	175	41	111	152	199	127	326
11	248	327	575		113	113		39	39	226	97	323
12	128	205	334	28	50	78		33	33	68	52	120
13	69	149	218		47	47		27	27	15	34	49
14	6	62	67					7	7		44	44
15		85	85		22	22		22	22		5	5
16		31	31								14	14
17												
18												
<b>Total</b>	<b>2946</b>	<b>3228</b>	<b>6174</b>	<b>1042</b>	<b>1169</b>	<b>2211</b>	<b>1917</b>	<b>1942</b>	<b>3859</b>	<b>1809</b>	<b>1295</b>	<b>3104</b>
<b>Mean Age</b>	<b>7,79</b>	<b>8,39</b>	<b>8,10</b>	<b>6,79</b>	<b>7,93</b>	<b>7,39</b>	<b>6,33</b>	<b>6,57</b>	<b>6,45</b>	<b>7,62</b>	<b>7,60</b>	<b>7,61</b>
<b>S.D.</b>	<b>2,37</b>	<b>3,10</b>	<b>2,79</b>	<b>1,93</b>	<b>2,88</b>	<b>2,54</b>	<b>1,76</b>	<b>2,56</b>	<b>2,20</b>	<b>2,53</b>	<b>3,09</b>	<b>2,78</b>

Age	1997			1998			Total		
	Males	Females	Total	Males	Females	Total	Males	Females	Total
1				26	13	39	26	13	39
2	14	21	35	63	130	193	143	230	373
3	83	149	232	119	135	254	458	751	1208
4	53	106	159	125	121	246	800	841	1641
5	73	89	162	90	49	139	900	1043	1943
6	214	230	444	141	201	342	1676	1192	2868
7	284	309	593	411	322	733	2116	1642	3757
8	59	110	169	549	410	959	1789	1434	3223
9	71	111	182	125	155	280	958	1056	2015
10	78	63	141	156	120	276	794	811	1605
11	74	130	204	101	73	174	649	779	1428
12	103	127	230	99	149	248	426	616	1043
13	44	110	154	60	147	207	188	514	702
14	15	55	70	34	68	102	55	236	290
15		56	56		61	61		251	251
16		18	18		31	31		94	94
17					15	15		15	15
18		18	18					18	18
<b>Total</b>	<b>1165</b>	<b>1702</b>	<b>2867</b>	<b>2099</b>	<b>2200</b>	<b>4299</b>	<b>10978</b>	<b>11536</b>	<b>22514</b>
<b>Mean Age</b>	<b>7,60</b>	<b>8,24</b>	<b>7,98</b>	<b>7,53</b>	<b>8,12</b>	<b>7,83</b>	<b>7,34</b>	<b>7,87</b>	<b>7,61</b>
<b>S.D.</b>	<b>2,84</b>	<b>3,60</b>	<b>3,33</b>	<b>2,74</b>	<b>3,58</b>	<b>3,21</b>	<b>2,46</b>	<b>3,23</b>	<b>2,89</b>











Figure 1. Annual length distribution and age composition, Flemish Cap 1991-1998.











Figure 1 (continued)




Figure 2. AFL average frequency (%) for the 1993-1998 surveys.




Figure 3. Distribution of age average frequency (%) in 1994-1998 surveys. (a) Both sexes together; (b) sex distribution.




Figure 4. Female ratio by age, Flemish Cap 1993-1998




Figure 5. Female ratio by length, Flemish Cap 1993-1998.




Figure 6. Catch curves for roughhead grenadier, females and males.