



Northwest Atlantic

Fisheries Organization

Serial No. N4330

NAFO SCR Doc. 00/ 73

SCIENTIFIC COUNCIL MEETING – NOVEMBER 2000

**Oceanographic conditions on the Flemish Cap during the summer of 2000 with comparisons to the previous year and the 1961-1990 average**

by

E. Colbourne

Science Branch, Department of Fisheries and Oceans  
P.O. Box 5667 St. John's, Newfoundland A1C 5X1

**ABSTRACT**

Oceanographic data from the summer of 2000 on the Flemish Cap are examined and compared to the long-term (1961-1990) average and to conditions during the summer of 1999. The cold near-surface temperatures (0.5 to 2°C below normal) experienced over the Cap from 1993 - 1996 had warmed to 0.5 - 1.5°C above normal by July of 1997, which increased to 2°C above normal by the summer of 1998 and 1999. Surface temperatures during the summer of 2000 decreased somewhat but remained above normal in some areas over the Cap. Bottom temperatures over the Cap were slightly below normal during 1997, up to 0.5°C - 1°C above normal during 1998 to 1999 and near normal during the summer of 2000. Salinities during 1999 ranged from near normal to slightly above normal and during the summer of 2000 they ranged from near normal to slightly below normal over most areas. In general the colder than normal temperatures experienced over the continental shelf and on the Flemish Cap from the late 1980s up to 1995 moderated by the summer of 1996 and continued above normal during 1999. During the summer of 2000 the measurements shows evidence of a reversal in the recent warm trend in some areas of the water column. As in previous years, summer chlorophyll levels in the upper 100-m of the water column over the Cap were higher compared to the adjacent Grand Bank and dissolved oxygen levels were about normal for the region. Both the measured currents and the geostrophic estimates, while showing considerable differences and variability between years, indicate a general anticyclonic circulation around the Flemish Cap.

**1. Introduction**

The Flemish Cap is an isolated bank located east of the Grand Banks of Newfoundland centered at about 47° N, 45° W with minimum water depths of 126-m. To the west, the Flemish Pass with maximum water depths of about 1100 m separates the Cap from the Grand Bank. The water mass over the Flemish Cap is a mixture of Labrador Current and North Atlantic Current water, the general circulation of which is shown in Fig. 1a. Since the early 1970s the water masses over the Atlantic continental shelves have been dominated by three anomalous periods: early 1970s, mid 1980s and the early 1990s (Colbourne et al. 1994). During these periods strong positive winter North Atlantic Oscillation (NAO) index anomalies were mainly responsible for colder than normal air temperatures over the Northwest Atlantic resulting in increased ice cover and eventually colder and fresher than normal oceanographic conditions over the continental shelf of Atlantic Canada (Drinkwater 1996). An examination of the time series of historical temperature data from the Flemish Cap indicated similar conditions

existed on the Flemish Cap, however recent data indicates that conditions have returned to more normal conditions (Colbourne 1999).

The monthly mean temperature and salinity along the standard Flemish Cap Section have been published by Keeley (1981) and includes observations from 1910 to 1980. Drinkwater and Trites (1986) published spatially averaged temperature and salinity from all available bottle data from 1910 to 1982 over the Flemish Cap area. Numerous reviews and studies of the physical oceanography around the Flemish Cap were conducted during the Flemish Cap Project of the late 1970s and early 1980s (Hays et al. 1978, Bailey 1982, Akenhead 1981). More recent reviews of oceanographic conditions in the region compared 1993 and 1995-1998 observations with the long-term mean and also presented time series of temperature and salinity anomalies at various depths around the Cap (Colbourne 1993, 1995, 1996, 1997, 1998, 1999). Stein (1996), summarized the most recent studies of oceanographic conditions on the Flemish Cap. Additionally, Cervio and Prego (1996, 2000) presented hydrographic conditions on the Flemish Cap in July of 1996 and 1999 from fisheries research surveys conducted by the European Union. This manuscript presents an update to these studies by including data up to the summer of 2000 and represents the eighth such review of summer oceanographic conditions on the Flemish Cap in support of the annual Shrimp assessment for NAFO Division 3M.

The report describes oceanographic conditions on the Flemish Cap during the summer of 2000 with a comparison to 1999 conditions and to the long-term mean, based on all available historical data. The normal has been defined as the 30-year period from 1961-1990 in accordance with the convention of the World Meteorological Organization and recommendations of the NAFO Scientific Council. The 1999 and 2000 observations were made by the Canada's Department of Fisheries and Oceans on oceanographic surveys in mid-July of each year. During these surveys oceanographic observations were made along the standard NAFO Flemish Cap transect at 47° N latitude (Fig. 1b). Physical oceanographic measurements included vertical profiles of currents, temperature, salinity, and chlorophyll and dissolved oxygen.

## **2. Average Temperature and Salinity**

The vertical distributions (depth versus horizontal distance from the Avalon Peninsula of Newfoundland) of the average temperature and salinity over the Flemish Cap along 47° N for the period July 3 to August 2, based on all available historical data from 1961-1990 are shown in Fig. 2. These dates were chosen to span a one-month time period centered on the 1999 observations. No other attempts were made to adjust the mean for possible temporal biasing arising from variations in the number of observations within the time interval.

The average temperature for this time period (Fig. 2, upper panel) ranges from 4 to 5°C at 50-m depth to above 10°C near the surface. In deeper water (50-m to the bottom) temperatures range from 2 to 4°C in the Flemish Pass area, in the offshore branch of the Labrador Current, and from 3.5 to 6°C offshore of the Cap, where the influence of the Gulf Stream is evident. The corresponding average surface salinities (Fig. 2, bottom panel) range from about 33 psu in the Flemish Pass to about 33.5 psu over the Cap. Near bottom over the Cap in water depths of 150 to 300-m salinities range from 34.5 to 34.75 psu. In water depths greater than 300 m, salinities range from 34.75 to 34.85 psu.

## **3. 1999 and 2000 Temperature**

The vertical temperature distribution in July of 2000 (Fig. 3, bottom panel) over the

Flemish Cap show temperatures ranging from about 4 to 6°C at 50-m depth to above 11°C near the surface, somewhat colder than the 1999 values (Fig. 3, upper panel). In the depth range of 50-100 m temperatures ranged from 3 to 4°C, except to the west of the Cap where they were between 1 to 3°C. In the depth range of 100-m to the bottom over the Cap temperatures were about 4°C, somewhat warmer than the 1997 and 1998 values but slightly colder than 1999 values.

The vertical distribution of temperature anomalies for 1999 and 2000 over the Flemish Cap referenced to the 1961-1990 normal is shown in Fig. 4. A temporal adjustment of the 1999 calculations was made by restricting the historical data in the time period of July 3 to August 2. The 2000 observations were made near the middle of the month hence no adjustment of the data range was made. An examination of the historical data distribution for that time period during 1999 shows that only about 17% of the profiles were collected before July 18, the mid-point of the 1999 observations, with a median date of July 27. The annual temperature cycle over the Flemish Cap indicates that the temperature normally changes by approximately 0.5°C during the time period from July 18 to 27 in the near surface layers (0 to 20 m) and about 0.1°C at 50-m depth. This indicates that in near surface zone, where the annual cycle is the strongest, the temperature anomalies may be biased low, assuming normal atmospheric heat flux.

During July 2000, surface layer temperature anomalies over the Cap (Fig. 4 bottom panel) ranged from 1 to 1.5°C above normal, similar to the 1999 values (Fig. 4 top panel). Below the surface layer however, except directly on top of the Cap temperatures were 0 to 0.5°C below normal, colder than the 1999 values. Bottom temperatures on the cap were near normal. The positive surface anomalies over the Flemish Cap during 1997 to 1999 and to a lesser extent during 2000 are in contrast to the generally negative anomalies experienced during July of 1993, 1995 and 1996.

### ***Long-term Trends***

Similar to the Newfoundland Shelf the monthly temperature anomalies on the Flemish Cap (Fig. 5) are characterised by 3 major cold periods: most of the 1970s, mid 1980s and the late 1980s to the mid-1990s. The cold period beginning around 1971 continued until about 1977. From 1978 to 1984 temperature anomalies showed a high degree of variability in the upper water column with a tendency towards positive anomalies. By 1985 in the top 100-m of the water column negative temperature anomalies had returned. This cold period moderated briefly in 1987 but returned again by 1988 and continued until the mid-1990s. From 1995 to 1998 temperatures have moderated and remained above normal during the summer of 1999 in the depth range from 0 to 100-m. During 2000 temperatures decreased over 1999 values but remained near normal over most of the water column.

### **4. 1999 and 2000 Salinity**

The vertical distribution of salinities for July of 2000 (Fig. 6, bottom panel) show values ranging from less than 33 near the surface in the Flemish Pass, where the influence of the Newfoundland Shelf Water is evident. Across the Cap in the upper 50-m of the water column salinities were between 33.25 to 33.75, somewhat lower than the 1999 values except over the Flemish Pass where they were higher than the 1999 values (Fig. 6, upper panel). In the depth range of 50-m to the bottom salinities generally ranged from 34.5 to 34.75 in 1999 and from 33.73 to 34.75 during 2000.

The corresponding salinity anomalies during 1999 were slightly saltier than normal (by

0.2 to 0.4) over most of the Cap and over the Pass in the surface layer (Fig. 7). On the eastern side of the Cap during 1999 salinities were slightly below normal indicating the possible influence of the offshore branch of the Labrador Current, this is also evident in the temperature anomaly plot of Figure 4. In the deeper water (below 300-m) of the Flemish Pass and on the continental slope to the east of the Cap, values were near normal. During July of 2000 salinities were fresher-than-normal over the Cap and to the east of the Cap, slightly saltier than normal near the surface and about normal near bottom.

### ***Long-Term Trends***

The time series of salinity anomalies (Fig. 8) shows fresher than normal conditions from 1971 to 1976 and from 1983 to 1986 in the upper 100-m of the water column with peak amplitudes reaching 0.9 psu below normal. The trend in salinity values during the early 1990s range from slightly above normal at the surface to below normal at 20 to 100-m depth. During the summer of 1999 salinities were near the long-term average on the Flemish Cap but fell to slightly below normal by the summer of 2000. In general, the temperature and salinity (except for the 1990s) anomalies are very similar to those at Station 27 and elsewhere on the continental shelf over similar depth ranges (Colbourne 1998b).

### **5. 1998 and 1999 Chlorophyll and Dissolved Oxygen**

The vertical distributions of chlorophyll and dissolved oxygen saturation for July of 1999 and 2000 along the standard NAFO transect across the Grand Bank and the Flemish Cap are shown in Figs. 9 and 10. These data were collected in conjunction with the temperature and salinity data using a YSI type polarographic element dissolved oxygen sensor and a fluorometer interfaced to a Seabird-9 CTD system. The oxygen sensor was factory calibrated at zero and air-saturated water oxygen levels and also field calibrated by taking water samples at standard depths. The oxygen levels of the samples were determined by semi-automated analytical chemistry using a modified Winkler titration technique. The sensor readings were then corrected by using a least-squares fit of the titration measurements to the electronic sensor measurements. The 2000 standard depth chlorophyll values were determined using a laboratory fluorometer. The electronic fluorometer measurements were then corrected by using a least-squares fit of the laboratory measurements and the electronic measurements. No field calibrations were applied to the 1999 chlorophyll values presented here.

The chlorophyll concentrations show relatively high values over the Flemish Cap compared to over the adjacent Grand Bank during both 1999 and 2000. Maximum values were confined to a sub-surface layer of about 50-m thick (Fig. 9). The higher chlorophyll values over the Flemish Cap during mid summer appear to be a common occurrence and may indicate a delayed or extended offshore plankton bloom relative to the Newfoundland Shelf areas.

Over the Flemish Cap dissolved oxygen saturation levels during both 1999 and 2000 ranged from 95 to 105% from the surface to about 50-m depth (Fig. 10). Below 50-m depth values ranged from 85 to 95%, except near bottom where values ranged from 75 to 80% in 1999. Overall, oxygen saturation values were very similar during both 1999 and 2000 except near bottom over both the Grand Bank and the Cap. The super-saturated values in the top 50-m of the water column correspond to the high chlorophyll concentrations encountered over the Flemish Cap. These saturation levels are similar to that observed during 1995 to 1998 and are typical for this region.

## 6. Circulation

The general circulation in the Flemish Cap region consists of the offshore branch of the Labrador Current which flows through the Flemish Pass more or less trapped to the Grand Bank side of the continental slope area and a jet that flows to the east, north of the Cap. To the south the Gulf Stream flows to the Northeast merging with the Labrador Current to form the North Atlantic Current (Fig. 1a). In the absence of strong wind forcing the circulation over the Flemish Cap is dominated by a topographically induced anticyclonic gyre over the central portion of the bank (Kudlo et al. 1984, Ross 1981). The stability of this circulation pattern may influence the retention of ichthyoplankton on the bank and is probably a factor in determining the year-class strength of various fish species, such as cod, redfish and shrimp (Kudlo and Borovkov 1977; Kudlo and Boytsov 1979). This hypothesis however, has never been investigated in detail.

From 1993 to 2000 (except 1994) currents on the Flemish Cap during mid-July were measured with hull-mounted 150 kHz RDI acoustic Doppler current profilers (ADCPs) at a spatial resolution of 4-m vertically by approximately 1.5 km horizontally. Measurements were restricted to water depths less than 500 where bottom referencing was possible. The useful range of the 150 kHz ADCP for current measurements in this area is about 10 to 300-m depth.

The circulation around the Flemish Cap is predominately anticyclonic in all surveys since 1993 with typical re-circulation times ranging from 50 to 70 days. Figure 11 shows vertical cross-sections of the north-south currents over the Flemish Cap during July of 1999 and 2000 along 47° N latitude. The 1999 measurements show a northward flow in the Flemish Pass area and over the western side of the Cap and the southward flowing surface water on the eastern side of the Cap. The July 2000 measurements show a northward component ranging from 5 to 20 cm/s over the shoreward portion of the Cap in the Flemish Pass area and over the Cap in water depths below 50-m depth. In the surface layer and east of the Cap currents were generally southward with speeds ranging from 5 to 20 cm/s very similar in magnitude to those reported for July of 1998.

The anticyclonic motion of the water mass around the Flemish Cap was first described by Kudlo and Burmakin (1972), Kudlo and Borovkov (1975) and Kudlo et al. (1984) using geostrophic currents estimated from density measurements. The geostrophic currents perpendicular to the 47° N transect estimated from the density data collected during 1999 and 2000 are shown in Fig. 12. These calculations which are referenced to 300-m, or the bottom, in water depths less than 300-m, show some of the well-known features of the circulation. The strong baroclinic component of the offshore branch of the Labrador Current shoreward of the Cap and the northward flowing water of the North Atlantic Current east of the Cap are evident. The estimates over the Cap itself show strong evidence of anticyclonic circulation during both 1999 and 2000. The results however differ significantly from the direct current measurements made with the ADCPs, thus showing the potential importance of wind driven and tidal currents on the Flemish Cap. Finally, both the measured currents and the geostrophic estimates while showing considerable variability between years, nevertheless show similar features in the overall circulation pattern.

## ACKNOWLEDGEMENTS

I thank C. Fitzpatrick, P. Stead, W. Bailey, C. Bromley, J. Craig and S. Kennedy of the oceanography section at the Canadian Northwest Atlantic Fisheries Center for data collection and processing and for computer software support. I would also like to thank the captain and crew of the CCGS Teleost for field support and to the many scientists whom over the years have contributed to the national database.

## REFERENCES

- Akenhead, S. 1981. Local sea-surface temperature and salinity on the Flemish Cap. NAFO SCR Doc. 87/66, Ser. No. N426 20 p.
- Bailey, W.B., 1982. A time-series of sea-surface temperature on the Flemish Cap, 1962-81. NAFO SCR Doc. 82/4, Ser. No. N489 7 p.
- Cervifio, S. and R. Prego. 1996. Hydrographic conditions on the Flemish Cap in July 1996. NAFO SCR Doc. 97/17, Ser. No. N2847 13 p.
- Colbourne, E.B. S. Narayanan, and S. Prinsenberg. 1994. Climate change and environment conditions in the Northwest Atlantic during the period 1970-1993. ICES mar. Sci. Symp., 198:311-322.
- Colbourne, E. 1993. Oceanographic conditions on the Flemish Cap during the summer of 1993, with comparisons to the long-term average. NAFO SCR Doc. 93/107, Ser. No. N2300 36p.
- Colbourne, E. 1995. Oceanographic conditions on the Flemish Cap during the summer 1995, with comparisons to the 1961-1990 average. NAFO SCR Doc. 95/102, Ser. No. N2625 12p.
- Colbourne, E. 1996. Oceanographic conditions on the Flemish Cap during the summer 1996, with comparisons to the previous year and the 1961-1990 average. NAFO SCR Doc. 96/87, Ser. No. N2770 12p.
- Colbourne, E. 1997. Oceanographic conditions on the Flemish Cap during the summer 1997, with comparisons to the previous year and the 1961-1990 average. NAFO SCR Doc. 97/84, Ser. No. N2930 15p.
- Colbourne, E. 1998a. Oceanographic conditions on the Flemish Cap during the summer 1998, with comparisons to the previous year and the 1961-1990 Average. NAFO SCR Doc. 98/86, Ser. No. N3087 16p.
- Colbourne, E. and K. D. Foote. 1998b. Oceanographic variability on the Flemish Cap. NAFO SCR Doc. 98/35, Ser. No. N3022 56p.
- Colbourne, E. 1999. Oceanographic conditions on the Flemish Cap during the summer 1999, with comparisons to the previous year and the 1961-1990 Average. NAFO SCR Doc. 99/101, Ser. No. N4180 18p.
- Drinkwater, K. F., and R. W. Trites. 1986. Monthly means of temperature and salinity in the Grand Banks region. Can. Tech. Rep. Fish. Aquat. Sci. 1450:iv+111 p.
- Drinkwater, K. F. 1996. Climate and oceanographic variability in the Northwest Atlantic during the 1980s and early-1990s. J. Northw. Atl. Fish. Sci. 18: 77-97.
- Keeley, J.R., 1981. Mean conditions of potential temperature and salinity along the standard Flemish Cap Section. Mar. Environ. Data Serv. Tech. Rep. No. 9: 148 pp.

- Kudlo, B.P., V.A. Borovkov and N.G. Sapronevskaya. 1984. Water circulation patterns on the Flemish Cap from observations in 1977-82. NAFO Sci. Coun. Studies, 7:27-37.
- Kudlo, B.P., and V.V. Burmakin. 1972. Water circulation in the South Labrador and Newfoundland areas in 1970-71. ICNAF Redbook 1972 (iii):22-33.
- Kudlo, B.P., and V. A. Borovkov. 1975. Circulation of waters in the ICNAF area in 1973-1974. ICNAF Res. Doc. No. 79, Serial No. 3506, 12 p.
- Kudlo, B.P., and V.A. Borovkov. 1977. Hydrological regime of the Labrador Current and reproduction of commercial fishes. Syezd Sovetskikh Okeanologov, Tezisy Dokladov, Moskva vyp. ii:133-134.
- Kudlo, B.P., and V. D. Boytsov. 1979. The effect of water dynamics on year-class strength of cod on Flemish Cap. ICNAF Sel. Papers, 5: 7-9.
- Ross, C.K., 1981. Drift of satellite-tracked buoys on the Flemish Cap, 1970-80. NAFO Sci. Coun. Studies, No. 1: p47-50.
- Stein, M. 1996. Flemish Cap - A review on Research Activities with Focus on Oceanographic Conditions. NAFO Sci. Coun. Studies, 25:1-24.




Fig. 1. (a) The major circulation features around the Flemish Cap area (Adapted from Anderson, 1984) and (b) the location of stations occupied during the summer of 1999 and 2000. Bathymetry lines are 2000, 1000, 500, 400, 300, 200 and 150 m.






Fig. 2. The vertical distribution of the average temperature and salinity over the Flemish Cap based on all available historical data from 1961-1990 for the time period of July 3 to August 2.




Fig. 3. The vertical distribution of temperature over the Flemish Cap (along  $47^{\circ}\text{N}$ ) for July of 1999 and 2000.




Fig. 4. The vertical distribution of temperature anomalies over the Flemish Cap (along  $47^{\circ}$  N) for July of 1999 and 2000. Negative anomalies are shaded.




Fig. 5. Time series of monthly temperature anomalies at standard depths of 0, 20, 50 and 100 m on the Flemish Cap in NAFO Division 3M. The solid line represents the smoothed temperature anomalies.




Fig. 6. The vertical distribution of salinity over the Flemish Cap (along  $47^{\circ}\text{N}$ ) for July of 1999 and 2000.




Fig. 7. The vertical distribution of salinity anomalies over the Flemish Cap (along  $47^{\circ}$  N) for July of 1999 and 2000. Negative anomalies are shaded.




Fig. 8. Time series of monthly salinity anomalies at standard depths of 0, 20, 50 and 100 m on the Flemish Cap in NAFO Division 3M. The solid line represents the smoothed salinity anomalies.




Fig. 9. The vertical distribution of chlorophyll concentrations (mg/l) along the 47°N transect for July of 1999 and 2000.






Fig. 10. The vertical distribution of dissolved oxygen percent saturation along  $47^{\circ}\text{N}$  for July of 1999 and 2000.




Fig. 11. A vertical cross-section of the N-S current field in cm/s over the Flemish Cap (along  $47^{\circ}$  N) during July of 1999 and 2000 measured with a 150 kHz ADCP. Negative (shaded) currents are southward and positive are northward.




Fig. 12. The vertical distribution of the N-S geostrophic current field in cm/s over the Flemish Cap during July of 1999 and 2000 calculated from the density data. Negative (shaded) currents are southward and positive are northward.