

Northwest Atlantic



Fisheries Organization

Serial No. N4331

NAFO SCR Doc. 00/74

SCIENTIFIC COUNCIL MEETING – NOVEMBER 2000

The Icelandic Shrimp Fishery (*Pandalus borealis* Kr.) at Flemish Cap in 1993-2000

by

U. Skúladóttir
Marine Research Institute, Skúlagata 4,
P.O. Box 1390, 121 Reykjavík, Iceland

Abstract

Some 7 Icelandic vessels have been fishing for shrimp in the waters at Flemish Cap in 2000 compared to 10 in 1999. In this paper there are logbook information on the Icelandic fishery for the years 1993 through 1999. The unstandardized catch rate has recently increased considerably or from 172 kg/hour in January-July 1997 to 282 in 1998 and was 270 and 276 kg in 1999 and 2000, respectively.

The observer samples have been pooled by months and depth

Introduction

The Spanish investigators (EU) have been measuring the biomass index of northern shrimp at the Flemish Cap since 1988 in their annual bottom trawl survey at Flemish cap. In 1993 the fishery was initiated by Canada, followed closely by Faroe Islands and Iceland.

The fishery was some 24-33 thousand tons in the years 1993-1995 to increase in 1996 to 48 thousand tons. Since then the fishery decreased to some 25 thousand tons in 1997. The total catch of all countries has since increased to about 42 thousand tons in 1999.

In this paper all the information from the Icelandic investigators is gathered. From the logbooks comes effort, catch and from this CPUE is calculated. From the biological samples taken by Icelandic observers comes various information on length and sex distribution of shrimp. From these the age assessments can be carried out.

Materials and Methods

Icelandic observers sampled shrimp onboard all Icelandic vessels in the years 1996 through 2000 at Flemish Cap. The shrimp was measured fresh to the nearest 0.5 mm using Vernier callipers. Observers then sorted each length class into males and females using the method of Rasmussen (1953) and the females further into primiparous and multiparous using the sternal spine criterion of McCrary (1971).

The deviation method (Sund, 1930; Skuladottir, 1981) was applied to length frequency distributions (lfd) of several months. The lfd of all samples within a month was combined and turned into a promille distribution. Then as an example all the lfd of June of the years 1993-2000 were summed to calculate an overall promille lfd. Then the lfd of June in each year was subtracted from the overall lfd. From this positive anomalies could be detected as indicators of a year class stronger than average.

The logbook data include catch and effort. Not all skippers send in the logbooks, but information on landings can be obtained from the Fisheries Directorate in Iceland. Thus effort was raised by dividing the nominal catch of each month with the calculated CPUE from the logbooks in the years 1993-1996. In 1997 and the effort is first raised to the nominal effort by every half year. The overall CPUE of the January-July was then obtained by summing nominal catch of all months and corresponding effort. Nominal catch for the whole period was then divided by "nominal effort" to get the CPUE for the period January-July. When twin trawls were used the effort was always multiplied by 1.9 for those but the catch was kept the same.

Catch and Effort Data

In 2000 the fishery was carried out since January. The catch in 2000 so far is 7 437 tons (Table 2). Iceland increased the total allowable catch (TAC) for Icelandic vessels from 6 800 tons in 1998 to 9 300 tons for the year 1999 and 10 100 tons for year 2000.

The distribution of effort is shown by months and years in Fig. 1-7. Note the difference between the years 1998 and 1999 for the lack of tows in the southeast area in 1998 and an increase in 1999. In 2000 the pattern of tow stations was similar to that of year 1999. Looking at distribution of tows by months, the months of March and April of 1999 are different from other months in that there are quite many tows at shallow depths in the northwest and southwest areas. The same pattern occurs again in February and March in year 2000.

It was decided in 1999 to close the area of shallow water during the summer in order to protect the small shrimp. This corresponds approximately to depth less than 140 fathoms. In Table 3 is shown how the mean size of shrimp increases with depth in years 1999 and 2000. The biggest shrimp is caught at depths greater than 300 fm. In Table 4. The percentages of catch by depth are shown. Most of the shrimp is caught between 141 and 300 fathoms.

The mean CPUE for the year 1997 was the lowest ever for Iceland or 172 kg per trawling hour for the period January through July (Table 1). In 1998 the mean CPUE for the same period was much higher or 282 kg and rather similar in 1999, namely 270 kg. The average size of gear used was about 3 000 meshes in most years, but increased to 3 520 meshes in year 2000. At the same time the use of twin trawls has increased in 1998 from a little less than 60% in 1995-1997 to about 81% in 2000. Although the CPUE unstandardized does not seem to be falling recently, The CPUE when calculated for a standard size of 3 000 meshes trawl (effort *1.9 for double trawls) is declining since 1998.

Length Frequencies and Age Groups

The length frequency distributions of Icelandic samples from 1999 and 2000 are shown in Fig. 8-20. In 1999 and 2000, 3 year-old males are the most prominent peaks of about 18 mm CL. Two year olds are seen in March and April 1999 about 12 mm CL. The assumed 4 year olds started changing sex in late 1998 and continued to change sex as 5 year olds in early-1999. The primiparous peak appears to be sometimes bimodal and broad, but later in the year it appears to be unimodal (see Fig. 9).

The differing height of peaks can be studied further in relation to depth and month. On the whole the 2 group seems to have a tendency to occur at less depth than other groups. The older animals have generally a tendency to be more numerous at greater depths. (Fig. 10-20).

It is rather difficult to figure out the proper age so the deviation method (Sund, 1939; Skúladóttir 1981) was applied to the monthly length frequency distributions. The 1993 year-class, which appeared first in March 1995 was about 14 mm and is assumed to be two years old then. This same year-class can be followed until it is 5 years old in 1998 about 25.7 mm CL on Fig. 21. This same year class can be traced in several months or on Fig. 22 through Fig. 27. In October 1998 this year-class is assessed to be 26.5 mm CL.

By-catch

The by-catch was about 1% in the years 1999 and 2000 as compared to 0.8% of the shrimp catch in 1998, 1.8% in 1997 and 3 % in 1996. Most of this was redfish or 0.7% in both 1999 and 2000. Other species were wolffish, Greenland halibut and American plaice. Cod was seen for the first time in April 1999, but has not been seen since then (Skúladóttir, 1997; 1998).

References

- McCrary, J.A. 1971. Sternal spines as a characteristic for differentiating between females of some Pandalidae. *J. Fish. Res. Board Can.*, **28**: 98-100.
- Rasmussen, B. 1953. On the geographical variation in growth and sexual development of the deep sea prawn (*Pandalus borealis* Kr.). *Norweg. Fish. and Mar. Invest. Rep.*, **10** (3): 1-160.
- Skúladóttir, U. 1981. The deviation method. A simple method for detecting year-classes of a population of *Pandalus borealis* from length distributions. T. Frady (Ed.), Proc., Int. Pandalid Shrimp Symp., Kodiak Alaska, Feb. 13-15. pp. 283-306. *Univ. Alaska, Sea Grant Rep.* 81-3.
- Skúladóttir, U. 1999. The Icelandic shrimp fishery (*Pandalus borealis* Kr.) at the Flemish Cap in 1993-1999. *NAFO SCR Doc.*, No. 116, Serial No. N4197, 27 p.
- Skúladóttir, U. 1997. The by-catch in the shrimp fishery of Iceland at Flemish Cap in 1997 and 1998. *NAFO SCR Doc.*, No. 80, Serial No. N 2918, 4 p.
- Skúladóttir, U. 1998. The by-catch in the shrimp fishery of Iceland at Flemish Cap in 1997 and 1998. *NAFO SCR Doc.*, No. 29, Serial No. N3016, 5 p.
- Sund, O. 1930. The renewal of fish population studied by means of measurement of commercial catches. Example: the arcto Norwegian cod stock. *Rapp. P.-v. Réun. Cons. Int. Explor. Mer.*, **65**: 10-17.

Table 1. Catch (tons) effort (trawling hours *1.9 when double trawl) and CPUE (kg/hr) of Icelandic vessels at Flemish Cap.

Year	January - July				August - December			
	Month	CPUE	Effort	Catch	Month	CPUE	Effort	Catch
1993					Aug	320.4	1334	427.4
					Sep	349.8	1034	361.7
					Oct	231.7	334	77.4
	Jun	380.2	1767	671.8	Nov	306.8	588	180.4
	Jul	342.4	1097	375.6	Dec	236.5	537	127.0
	Subtotal	365.7	2864	1047.4	Subtotal	306.7	3827	1173.9
	Total	365.7	2918	1067.0	Total	306.7	3834	1176.0
1994	Jan	228.5	144	32.9	Aug	175.3	1657	290.4
	Feb	371.8	510	189.6	Sep	126.9	476	60.4
	Mar	295.5	531	156.9	Oct	125.4	492	61.7
	Jun	256.4	1297	332.5	Nov	115.5	181	20.9
	Jul	212.9	2653	564.8	Dec	75.0	8	0.6
	Subtotal	248.6	5135	1276.7	Subtotal	154.2	2814	434
	Total	248.6	6693	1664.0	Total	154.2	4123.7	636
1995	Feb	280.0	65	18.2	Aug	178.0	4869	866.9
	Mar	246.8	711	175.5	Sep	134.1	2928	392.5
	Apr	149.9	1487	222.9	Oct	166.3	2088	347.2
	May	260.1	2617	680.7	Nov	144.4	1074	155.1
	June	248.9	3733	929.2	Dec	174.5	740	129.1
	Jul	249.5	6625	1653.0				
	Subtotal	241.5	15238	3679.5	Subtotal	161.6	11699	1890.8
	Total	241.5	16932	4088.5	Total	161.6	21868	3534.4
1996	Jan	207.2	1755	363.7				
	Feb	251.7	1326	333.7				
	Mar	261.8	4604	1205.1	Aug	165.4	8156	1349.4
	Apr	211.2	10754	2271.2	Sep	167.1	8089	1351.7
	May	189.1	12749	2410.2	Oct	129.7	5482	711.2
	Jun	202.5	13933	2821.5	Nov	137.9	1456	200.8
	Jul	235.9	11963	2821.5	Dec	158.1	253	40.0
	Subtotal	214.2	57084	12226.9	Subtotal	155.9	23436	3653.1
	Total	214.2	64760	13871.0	Total	155.9	43689	6810.0
1997	Jan	175.8	413	72.6	Aug	206.7	4252	879.0
	Feb	214.7	621	133.3	Sep	202.4	3476	703.6
	Apr	135.0	514	69.4	Oct	222.0	2519	559.1
	May	141.4	3736	528.2	Nov	192.5	1039	200.0
	Jun	167.7	5386	903.2	Dec	176.9	429	75.9
	Jul	209.2	5802	1213.7				
	Subtotal	177.3	16472	2920.4	Subtotal	206.4	11715	2417.6
	Total	177.3	19478	3453.3	Total	206.4	14681	3029.6
1998 *	Feb	217.2	297	64.5	Aug	256.4	3184	816.3
	Mar	206.8	812	167.9	Sep	184.5	5028	927.5
	Apr	229.5	880	202.0	Oct	196.3	3612	708.9
	May	261.4	2820	737.2	Nov	204.6	1761	360.3
	Jun	330.7	3537	1169.7	Dec	222.5	644	143.3
	Jul	285.3	4117	1174.7				
	Subtotal	282.1	12463	3516.0	Subtotal	207.8	14229	2956.3
	Total	282.1	12657	3570.8	Total	207.8	14447	3001.5
1999 *	Feb	350.5	382	133.9	Aug	250.8	3642	913.4
	Mar	289.4	1851	535.7	Sep	235.5	1371	322.9
	Apr	253.0	3483	881.2	Oct	255.6	2150	549.6
	May	249.5	5941	1482.3	Nov	256.2	2173	556.8
	Jun	285.8	5993	1712.7	Dec	230.6	989	228.1
	Jul	280.4	5224	1464.6				
	Subtotal	271.5	22874	6210.4	Subtotal	249.0	10325	2570.8
	Total	271.5	24009	6518.6	Total	249.0	10837	2698.4
2000 *	Jan	263.8	1050	277.0	Aug	246.0	2353	578.9
	Feb	280.5	2206	618.8	Sep	258.6	2165	559.9
	Mar	306.7	3292	1009.8				
	Apr	281.0	4398	1235.8				
	May	231.9	4947	1147.3				
	Jun	304.5	3680	1120.7				
	Jul	251.0	3056	767.0				
	Subtotal	272.9	22629	6176.4	Subtotal	252.1	4518	1138.8
	Total	272.9	22629	6176.4	Total	252.1	4518	1138.8

Table 2. Landings and some averages calculated from the Icelandic logbooks.
CPUE is only from the period January - July. The effort of twin trawls * 1.9.

Year	Nominal Catch Tons	Twin trawls % of catch	Trawl size No. of meshes	Unstandardized CPUE	CPUE at size 3000 trawl
1993	2 243	46.0	3086	366	363
1994	2 300	56.2	2975	249	240
1995	7623	57.6	2688	242	283
1996	20681	57.8	2839	214	217
1997	6483	54.6	2932	177	192
1998	6572	74.7	2939	282	294
1999	9217	73.5	3339	270	252
2000	7437	81.1	3520	276	245

Table 3. Mean lengths (CL in mm) by depth strata at Flemish Cap 1999-2000.

1999	Depth fm 1-100	Depth fm 101-140	Depth fm 141-200	Depth fm 201-300	Depth fm >301
Month	Mean Cl	Mean Cl	Mean Cl	Mean Cl	Mean Cl
1					
2			21.2	21.6	24.3
3		19.8	20.6	22.0	22.3
4		19.1	18.9	22.0	
5			19.8	22.2	24.0
6			21.0	23.0	25.8
7			20.8	21.4	23.7
8		21.6	20.4	20.9	21.9
9		18.9	19.9	21.5	22.4
10		20.4	20.0	22.3	
11		19.8	20.4	21.3	
12				21.8	23.0

2000	Depth fm 1-100	Depth fm 101-140	Depth fm 141-200	Depth fm 201-300	Depth fm >301
Month	Mean Cl	Mean Cl	Mean Cl	Mean Cl	Mean Cl
1		19.1	20.2	20.8	22.8
2		19.0	20.1	23.5	
3		19.4	19.2	20.1	
4		19.2	18.8	21.0	22.1
5		18.0	19.7	22.8	24.0
6			21.3	23.6	20.1
7		23.0	20.8	22.1	24.8
8		19.7	20.7	21.1	24.7
9		21.2	20.3	21.6	
10					
11					
12					

Table 4. Catch of shrimp (kgs) from log books by depth strata on the Flemish Cap 1994-2000.

1994		Depth fm 1-100		Depth fm 101-140		Depth fm 141-200		Depth fm 201-300		Depth fm >301		Total	Total
Month		Catch kg	%	Catch kg	%	Catch kg	%	Catch kg	%	Catch kg	%	Catch kg	%
1						30577	97.4	815	2.6			31392	100.0
2				349	0.2	8082	4.4	166290	90.9	8201	4.5	182922	100.0
3						40734	27.4	29402	19.8	78634	52.9	148770	100.0
4													
5													
6						228336	72.3	87396	27.7	125	0.04	315857	100.0
7						259372	48.0	281127	52.0			540499	100.0
8						67250	23.7	213102	75.2	3093	1.1	283445	100.0
9						31448	59.5	21391	40.5			52839	100.0
10						46415	75.3	14950	24.2	300	0.5	61665	100.0
11						18017	93.0	1356	7.0			19373	100.0
12						601	100.0					601	100.0
Total 1994		0	0.0	349	0.02	730832	44.6	815829	49.8	90353	5.5	1637363	100.0

1995		Depth fm 1-100		Depth fm 101-140		Depth fm 141-200		Depth fm 201-300		Depth fm >301		Total	Total
Month		Catch kg	%	Catch kg	%	Catch kg	%	Catch kg	%	Catch kg	%	Catch kg	%
1													
2						16750	92.3	1400	7.7			18150	100.0
3				47550	27.4	124750	71.8	1400	0.8			173700	100.0
4				37050	17.6	153929	73.1	19500	9.3			210479	100.0
5				1500	0.2	539106	81.0	124788	18.8			665394	100.0
6						259647	28.7	635954	70.4	8107	0.9	903708	100.0
7						823551	51.3	772532	48.1	10104	0.6	1606187	100.0
8				3117	0.4	284436	33.4	564065	66.2			851618	100.0
9				2600	0.7	299596	78.7	78253	20.6			380449	100.0
10				800	0.2	256380	74.3	87650	25.4			344830	100.0
11				1700	1.1	81373	52.8	69723	45.3	1200	0.8	153996	100.0
12				26260	20.3	90288	69.9	12512	9.7	50	0.04	129110	100.0
Total 1995		0	0.0	120577	2.2	2929806	53.9	2367777	43.5	19461	0.4	5437621	100.0

1996		Depth fm 1-100		Depth fm 101-140		Depth fm 141-200		Depth fm 201-300		Depth fm >301		Total	Total
Month		Catch kg	%	Catch kg	%	Catch kg	%	Catch kg	%	Catch kg	%	Catch kg	%
1				1940	0.5	242356	68.5	109339	30.9			353635	100.0
2				8500	2.5	263209	78.9	61986	18.6			333695	100.0
3				246715	20.5	896472	74.4	61437	5.1	500	0.04	1205124	100.0
4				488378	21.5	1084700	47.8	453478	20.0	244672	10.8	2271228	100.0
5				9931	0.4	1009597	42.2	1131708	47.3	243318	10.2	2394554	100.0
6				10102	0.4	977909	34.7	1773075	62.9	55910	2.0	2816996	100.0
7				2049	0.1	709740	33.6	1388454	65.8	10439	0.5	2110682	100.0
8						712341	52.8	612807	45.4	24276	1.8	1349424	100.0
9				33433	2.5	963094	71.3	353343	26.2			1349870	100.0
10				18957	2.7	478687	67.3	212991	29.9	581	0.1	711216	100.0
11				295	0.1	39133	19.5	161323	80.4			200751	100.0
12						33014	82.5	6986	17.5			40000	100.0
Total 1996		0	0.0	820300	5.4	7410252	49.0	6326927	41.8	579696	3.8	15137175	100.0

Table 4 (continued)

1997	Depth fm 1-100		Depth fm 101-140		Depth fm 141-200		Depth fm 201-300		Depth fm >301		Total	Total
Month	Catch kg	%	Catch kg	%	Catch kg	%	Catch kg	%	Catch kg	%	Catch kg	%
1					64021	88.2	8567	11.8			72588	100.0
2					49140	36.9	84141	63.1			133281	100.0
3												
4	1686	2.4			43871	63.2	23850	34.4			69407	100.0
5	1112	0.2	5187	1.0	275838	52.2	196892	37.3	49140	9.3	528169	100.0
6	1530	0.2			153081	16.9	571396	63.3	177155	19.6	903162	100.0
7	3300	0.3	509	0.04	697428	57.5	510075	42.0	2376	0.2	1213688	100.0
8					331232	37.7	547082	62.3	200	0.02	878514	100.0
9			2666	0.4	369438	52.5	330459	47.0	1056	0.2	703619	100.0
10	2590	0.5	1134	0.2	250855	45.0	301366	54.1	1226	0.2	557171	100.0
11					5504	2.8	187136	96.3	1755	0.9	194395	100.0
12							72112	95.0	3767	5.0	75879	100.0

Total 1997	10218	0.19	9496	0.18	2240408	42.03	2833076	53.15	236675	4.44	5329873	100.0
-------------------	-------	-------------	------	-------------	---------	--------------	---------	--------------	--------	-------------	---------	-------

1998	Depth fm 1-100		Depth fm 101-140		Depth fm 141-200		Depth fm 201-300		Depth fm >301		Total	Total
Month	Catch kg	%	Catch kg	%	Catch kg	%	Catch kg	%	Catch kg	%	Catch kg	%
1												
2					44656	69.2	19863	30.8			64519	100.0
3			1875	1.1	116085	70.1	47701	28.8			165661	100.0
4	5364	2.7	194233	96.1	2439	1.2					202036	100.0
5	3386	0.5			320321	43.5	274883	37.3	138580	18.8	737170	100.0
6	7051	0.6	24469	2.1	490260	41.9	611887	52.3	36004	3.1	1169671	100.0
7					299439	25.5	779266	66.3	96030	8.2	1174735	100.0
8					262978	32.4	544690	67.0	5100	0.6	812768	100.0
9			500	0.1	239746	25.8	647931	69.9	39288	4.2	927465	100.0
10			824	0.1	183710	25.9	519944	73.3	4448	0.6	708926	100.0
11			935	0.3	2615	0.7	269960	74.9	86742	24.1	360252	100.0
12					60952	42.5	30405	21.2	51975	36.3	143332	100.0

Total 1998	15801	0.24	222836	3.45	2023201	31.29	3746530	57.94	458167	7.09	6466535	100.0
-------------------	-------	-------------	--------	-------------	---------	--------------	---------	--------------	--------	-------------	---------	-------

1999	Depth fm 1-100		Depth fm 101-140		Depth fm 141-200		Depth fm 201-300		Depth fm >301		Total	Total
Month	Catch kg	%	Catch kg	%	Catch kg	%	Catch kg	%	Catch kg	%	Catch kg	%
1					64021	88.2	8567	11.8			72588	100.0
2			2600	1.9	54567	40.7	75842	56.6	900	0.7	133909	100.0
3			244274	45.6	184826	34.5	106410	19.9	161	0.0	535671	100.0
4			291696	33.1	556243	63.1	31077	3.5	2164	0.2	881180	100.0
5	3215	0.2			539847	36.4	593786	40.1	345465	23.3	1482313	100.0
6	7786	0.5			126598	7.4	1562477	91.5	10775	0.6	1707636	100.0
7					237537	16.2	1214893	83.0	12176	0.8	1464606	100.0
8			11350	1.2	212033	23.2	685934	75.1	4106	0.4	913423	100.0
9			57158	8.2	336417	48.2	302496	43.4	1459	0.2	697530	100.0
10			26290	4.8	290693	52.9	232628	42.3			549611	100.0
11			52929	9.5	397581	71.4	106334	19.1			556844	100.0
12							216711	95.0	11367	5.0	228078	100.0

Total 1999	11001	0.12	686297	7.44	3000363	32.53	5137155	55.70	388573	4.21	9223389	100.0
-------------------	-------	-------------	--------	-------------	---------	--------------	---------	--------------	--------	-------------	---------	-------

2000	Depth fm 1-100		Depth fm 101-140		Depth fm 141-200		Depth fm 201-300		Depth fm >301		Total	Total
Month	Catch kg	%	Catch kg	%	Catch kg	%	Catch kg	%	Catch kg	%	Catch kg	%
1			21689	7.8	144741	52.2	110607	39.9	0.0		277037	100.0
2			291663	47.1	277874	44.9	49223	8.0	0.0		618760	100.0
3			510041	50.5	406143	40.2	93611	9.3	0.0		1009795	100.0
4			211098	17.1	721950	58.4	283807	23.0	18947	1.5	1235802	100.0
5			134999	11.8	492183	42.9	371748	32.4	148414	12.9	1147344	100.0
6	300	0.0			934559	83.4	185810	16.6			1120669	100.0
7			2792	0.4	333740	43.6	429528	56.1			766060	100.0
8	9019	1.6	18283	3.2	258630	44.7	291174	50.3	1798	0.3	578904	100.0
9					207304	37.0	351734	62.8	861	0.2	559899	100.0
10												
11												
12												

Total 2000	9319	0.13	1190565	16.28	3777124	51.64	2167242	29.63	170020	2.32	7314270	100.0
-------------------	------	-------------	---------	--------------	---------	--------------	---------	--------------	--------	-------------	---------	-------




Fig. 1. Towing positions in the Icelandic fleet on Flemish Cap in 1996-1999.




Fig. 2. Towing positions in the Icelandic fleet on Flemish Cap in year 2000 and by months in 1999.




Fig. 3. Towing positions in the Icelandic fleet on Flemish Cap year 1999.




Fig. 4. Towing positions in the Icelandic fleet on Flemish Cap year 1999.




Fig. 5. Towing positions in the Icelandic fleet on Flemish Cap year 2000.




Fig. 6. Towing positions in the Icelandic fleet on Flemish Cap year 2000.




Fig. 7. Towing positions in the Icelandic fleet on Flemish Cap year 2000.




Fig. 8. The length frequency distribution of northern shrimp by Flemish Cap by months in 1999.




Fig. 9. The length frequency distribution of northern shrimp at Flemish Cap by months in 2000.




Fig. 10 The length frequency distribution of northern shrimp at Flemish Cap in February by depth in 1999.




Fig. 11 The length frequency distribution of northern shrimp at Flemish Cap in March and April by depth in 1999.




Fig. 12 The length frequency distribution of northern shrimp at Flemish Cap in May and June by depth in 1999.




Fig. 13 The length frequency distribution of northern shrimp at Flemish Cap in July and August by depth in 1999.




Fig. 14 The length frequency distribution of northern shrimp at Flemish Cap in September and October by depth in 1999.




Fig. 15 The length frequency distribution of northern shrimp at Flemish Cap in November and December by depth in 1999.




Fig. 16 The length frequency distribution of northern shrimp at Flemish Cap in January and February by depth in 2000.




Fig. 17 The length frequency distribution of northern shrimp at Flemish Cap in March and April by depth in 2000.




Fig. 18 The length frequency distribution of northern shrimp at Flemish Cap in May and June by depth in 2000.




Fig. 19 The length frequency distribution of northern shrimp at Flemish Cap in July and August by depth in 2000.




Fig. 20 The length frequency distribution of northern shrimp at Flemish Cap in September by depth in 2000.




Fig. 21. The deviations of length frequencies of northern shrimp by years in March on Flemish Cap from the mean length frequency distribution of the years 1994-2000 in the same month. 1994 and 1995 are data of Canada and other countries. Since 1996 data are solely from Iceland.




Fig 22. The deviations of length frequencies of northern shrimp by years in April on the Flemish Cap from the mean length frequency of the years 1993-2000 in the same month. 1993 through 1995 are data of Canada and other countries. Since 1996, data are solely from Iceland.




Fig 23. The deviations of length frequencies of northern shrimp by years in May on the Flemish Cap from the mean length frequency of the years 1993-2000 in the same month. 1993 through 1995 are data of Canada and other countries. Since 1996, data are solely from Iceland.




Fig 24. The deviations of length frequencies of northern shrimp by years in June on the Flemish Cap from the mean length frequency of the years 1993-2000 in the same month. 1993 through 1995 are data of Canada and other countries. Since 1996, data are solely from Iceland.




Fig 25. The deviations of length frequencies of northern shrimp by years in July on the Flemish Cap from the mean length frequency of the years 1993-2000 in the same month. 1993 through 1995 are data of Canada and other countries. Since 1996, data are solely from Iceland.




Fig 26. The deviations of length frequencies of northern shrimp by years in September on the Flemish Cap from the mean length frequency of the years 1993-2000 in the same month. 1993 through 1995 are data of Canada and other countries. Since 1996, data are solely from Iceland.




Fig 27. The deviations of length frequencies of northern shrimp by years in October on the Flemish Cap from the mean length frequency of the years 1993-2000 in the same month. 1993 through 1995 are data of Canada and other countries. Since 1996, data are solely from Iceland