

Northwest Atlantic



Fisheries Organization

Serial No. N4441

NAFO SCR Doc. 01/63

SCIENTIFIC COUNCIL MEETING – JUNE 2001

Stock Status Update of Witch Flounder in NAFO Divisions 3NO

by

W. R. Bowering

Northwest Atlantic Fisheries Center, P.O. Box 5667
St. John's, Newfoundland, Canada, A1C 5X1

Abstract

Biomass and abundance indices from Canadian spring surveys in Div. 3N have been at very low levels throughout the period since 1984. In most years the biomass index was estimated to be less than 1,000 tons or 2 million fish. For Div. 3O where most of the stock resides, estimates of stock size exhibited considerable annual fluctuations on average between 8,000 and 24,000 tons or 6-44 million fish particularly in the late 1980's. The data for Div. 3NO combined indicate an overall-declining trend in stock size with the estimates for the spring 1998 survey at the lowest level observed since 1984. Values have been somewhat higher since then.

Indices from Canadian fall surveys for Div. 3N are similar to the spring in both stock size estimates, which are very low but lack trend. Indices are highly variable for Div. 3O in the fall surveys and also lack any overall trend. Nonetheless, the estimates for each seasonal series are generally within the same numeric range.

Fisheries and Management

Catches in the 1960's peaked at 11,000-12,000 tons in 1967-68 and remained relatively high during the next several years (Table 1; Fig. 1). During the period 1971-84 catches ranged from a low of about 2,400 tons in 1980 and 1981 to as high as 15,000 tons in 1971 which is the highest recorded catch in the history of the fishery, however, from 1975-84 annual catches rarely exceeded 6,000 tons. Species specific catch statistics for flatfish prior to 1973 were largely developed from breakdowns of unspecified flounders and therefore should be quoted with caution.

As a result of an increase in fishing effort in the NRA during 1985 and 1986, especially by EU-Spain and EU-Portugal, catches rose rapidly to levels of 8,800 and 9,100 tons, respectively. This increased effort was primarily concentrated on the "tail" of the Grand Bank in the NAFO Regulatory area of Div. 3N. Non-Contracting parties such as South Korea, USA, Cayman Islands and Panama also contributed to increased catch levels during this period. Catches remained relatively high in 1987 and 1988 at 7,600 and 7,300 tons, respectively. During 1990-93 estimated catches were in the range of 4,200-5,000 tons. The estimated catch for 1994 was still in the order of 1,100 tons despite a moratorium being introduced on fishing this stock (Table 1; Fig. 1). The catch dropped to 300 tons in 1995 likely as a result of a substantial reduction in fishing effort for Greenland halibut where witch flounder comprises a by-catch. Since then catches have increased steadily and by 1999 was about 800 tons although declined again to under 600 tons in 2000 (Table 1; Fig. 1).

Historically, mostly Canada and the former Soviet Union conducted the fishery. Canadian catches fluctuated from between 1,200 and 3,000 tons from 1985-91 but increased to about 4,300 tons in 1992 and 4,200 in 1993 (Table 1). The increase in 1992 and 1993 was essentially the result of a quota transfer to Canada by the Russian

Federation. Canada has taken very little catch since then due to the moratorium. Catches by the USSR/Russian vessels declined from between 1,000 and 2,000 tons in the period 1982-88 to less than 100 tons in 1989-90 and little or no catch since then until 1999-2000 when Russia caught 86 and 50 tons, respectively (Table 1).

The first total allowable catch (TAC) for this resource was introduced by ICNAF in 1974 at a level of 10,000 tons largely based on average historical catches (Fig. 1). This level remained in effect until 1979 when it was reduced to 7,000 tons in consideration of declining commercial catch rates. It was further reduced to 5,000 tons in 1981 and remained at that level to 1993. The Scientific Council advised that for 1994 catches from this stock should not exceed 3,000 tons. A TAC of 3,000 tons was agreed to by the NAFO Fisheries Commission, however, it was also agreed that no directed fishery would be conducted for witch flounder in 1994 due to the poor state of the stock and to allow for rebuilding. The NAFO Fisheries Commission introduced a complete moratorium for directed fishing in 1995, which has continued through 2001.

Canadian Research Vessel Surveys

Stratified-random research vessel surveys have been carried out by Canada on the Grand Bank (including Div. 3NO) during spring since 1971 although during the early period coverage was limited and, in fact, for most years only surveyed to 366 meters. Since 1990, on the other hand, depth coverage was extended to 720 meters, which should be more representative of the stock distribution. Nevertheless, this still may not cover the entire range of depth distribution of witch flounder when compared to its distribution observed in other stock areas during recent years. In addition to spring surveys, a time series of fall surveys was begun in 1990 to investigate seasonal variation in stock distribution and abundance of various groundfish species. In fall 1998 the survey depth range was further extended to 1500 meters.

Beginning with the 1995 fall survey the survey gear was changed from an *Engel 145* groundfish trawl with steel bobbin footgear to a *Campelen 1800* shrimp trawl with rockhopper footgear. The data from these surveys have now been converted from Engel trawl catches to Campelen 1800 trawl catch equivalents. Only the converted survey data are presented here.

Survey Biomass and Abundance Indices

Biomass estimates by stratum are presented for the spring surveys in NAFO Div. 3N and 3O, respectively in Tables 2 and 3. Similar data are presented for abundance estimates from spring surveys in Tables 4 and 5, respectively. Fall survey results are shown in the same order as above for spring survey in Tables 6-9, inclusive. Graphical plots to better illustrate the comparative trends in stock biomass and abundance by season are presented by NAFO Div. 3N and 3O separately and combined in Fig. 2-4, respectively.

Estimated biomass and abundance from spring surveys (which are the longer time series) in Div. 3N have been at very low levels throughout the period since 1984. In most years trawlable stock size was estimated to be less than 1,000 tons or 2 million fish (Fig. 2; Tables 2 and 4) although some increase was observed in 2000 compared to previous years. For Div. 3O where most of the stock resides, estimates of stock size exhibited considerable annual fluctuations on average between 8,000 and 24,000 tons or 6-44 million fish particularly in the late 1980's (Fig. 2; Tables 3 and 5). The several high spikes in the time series appear related to distribution shifts between the deeper smaller strata and the more shallow large strata (see Fig. 6-9 for illustration). This would have the effect of giving lower estimates when fish are distributed deeper and higher estimates when fish are distributed more in over the bank. Nevertheless, the data indicate an overall-declining trend in stock size (Fig. 2 and 4) in both the lower estimates and the spikes with the estimates for the spring 1998 and 1999 surveys at the lowest level observed since 1984 in both trends, respectively. The 2000 estimate was about half the 1999 estimate but about double the 1998 estimate.

Results of the fall surveys for Div. 3N are similar to the spring in both stock size estimates, which are very low and lack trend (Fig. 3; Tables 6 and 8). The data trends for Div. 3O in the fall surveys are quite different than in the spring series (Fig. 3; Tables 7 and 9). There is an increasing trend for 1991-96, however, when the higher value for 1990 and the lower values for 1997 and 1998 are included the trend is removed (Fig. 3; Tables 7 and 9). Nonetheless, the estimates for each seasonal series are generally within the same numeric range. With Div. 3NO combined, the most recent biomass and abundance estimates from the spring surveys are among the lowest observed

although variable and illustrate a declining trend since the beginning of the data series in 1984. The fall survey series for Div. 3NO combined is less clear with no real trend in biomass. The high variability in annual estimates may also be related to distribution shifts similar to the spring series (see Fig. 6-7 for illustration). It should be emphasized as well that the more recent lower estimates are also based on more detailed survey coverage than in the earlier years (annual percentage contributions to the estimates are shown in Tables 2-9). Consequently, in reality the declining trends are stronger than illustrated in the figures.

Resource Status

Based on the 1998 and 1999 spring survey estimates it now appears that the resource remains at a lower level than during the 1980's. The general trend in this longer (spring) survey series would in fact suggest that the stock shows little or no improvement despite a commercial fishing moratorium being in effect for several years. No aging data have been available since 1994 and are not expected to be available in the foreseeable future. Therefore, it is difficult to comment on any recruitment prospects for the resource. Population abundance at length from true *Campelen 1800* surveys in the fall of 1995-2000 indicated a higher proportion of smaller fish in recent years especially in the 1998-2000 surveys (Fig. 5). However, it is quite variable from year to year which makes it difficult to track recruitment.

Table 1 - Catches and TACs (t) of Witch Flounder in Div. 3NO from 1960-2001.

Year	USSR			Total	TAC
	Canada	(Russia)	Other		
1960	-	-	-	5798	-
1961	-	-	-	4627	-
1962	-	-	-	1228	-
1963	895	485	893	2163	-
1964	1055	-	11	1066	-
1965	1324	849	4	2177	-
1966	3644	3828	50	7522	-
1967	2863	8565	75	11503	-
1968	1503	9078	18	10599	-
1969	479	4215	6	4700	-
1970	723	6039	1	6763	-
1971	178	14774	13	14965	-
1972	3419	5738	20	9177	-
1973	4943	1714	34	6691	-
1974	2807	5235	3	8045	10000
1975	1137	5019	12	6168	10000
1976	3044	2991	-	6035	10000
1977	3013	2742	4	5759	10000
1978	1165	2275	33	3473	10000
1979	1193	1868	16	3077	7000
1980	425	1994	1	2420	7000
1981	381	2044	-	2425	5000
1982	1760	1989	3	3752	5000
1983	1674	1942	-	3616	5000
1984	834	1955	13	2802	5000
1985	2748	1908	4117	8771	5000
1986	2937	1724	4470	9131	5000
1987	2829	1425	3342	7596	5000
1988	1927	1037	4381	7345	5000
1989	1241	81	2366	3688	5000
1990	2654	9	1516	4179	5000
1991	2624	-	2223	4847	5000
1992	4328	-	632	4960	5000
1993	4337	3	2506	4414	5000
1994 ^a	2	-	1117 ^b	1119	3000
1995 ^a	-	-	300 ^b	300	0
1996 ^a	64	-	294 ^b	358	0
1997 ^a	19	-	483 ^b	512	0
1998 ^a	2	5	605	612	0
1999 ^a	6	85	671	762	0
2000 ^a	12	50	483	545	0
2001 ^a	-	-	-	-	0

*Note: Although a TAC of 3000 tons was agreed by the PC, it was also agreed that no directed fishing be conducted in 1994 due to the poor state of the stock.

a = Provisional Data b = Estimated

Table 2. Biomass (tons) of which flounder from surveys in Div. 3N during spring 1984-2000. (Eggel data converted to Cartesian units for 1984-95).

Year	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	
Depth Range (meters)																		
Old Stratum Area (sq. n. mi.)																		
New Stratum Area (sq. n. mi.)																		
Stratum																		
<=56	1993	179	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<=56	1499	376	0	0	19	0	0	0	0	0	0	0	0	0	0	0	0	0
57 - 92	2992	1715	89	629	461	1519	175	0	0	29	165	0	0	115	33	120	266	0
57 - 92	1853	119	0	39	50	0	20	0	0	0	0	0	39	0	0	0	242	0
57 - 92	2520	0	82	23	18	147	0	0	0	0	0	0	0	0	0	0	0	0
57 - 92	2520	0	0	43	0	0	0	0	0	0	0	0	0	0	0	0	0	0
57 - 92	931	0	0	0	0	0	0	0	0	18	34	0	0	0	0	0	0	0
57 - 92	674	0	57	0	37	0	0	0	0	0	0	0	0	0	0	0	0	0
93 - 183	421	231	47	99	43	306	121	0	0	19	0	0	0	0	0	0	0	0
93 - 183	600	8	0	0	72	3	32	0	0	0	0	0	0	0	0	0	0	0
93 - 183	647	8	0	0	12	0	0	0	0	0	0	0	0	0	0	0	0	0
184 - 374	225	338	40	308	42	137	20	29	57	0	44	132	106	7	51	49	134	7
184 - 374	139	378	22	19	32	135	31	42	0	29	0	0	0	0	3	0	0	0
184 - 374	182	181	21	7	32	101	69	6	28	0	0	0	0	0	0	0	0	0
275 - 366	164	164	8	87	154	4	60	21	0	31	49	81	20	36	12	159	21	75
275 - 366	106	379	36	12	23	173	44	20	35	3	18	0	4	0	0	9	2	26
275 - 366	116	116	6	23	0	134	24	7	4	0	0	0	0	0	0	0	0	0
367 - 549	155	155	723					90	102	79	36	51	16	23	53	33	36	
367 - 549	105	105	725					62		40	44	0	5	28	4	20	32	
367 - 549	160	160	727					0	5	38	17	0	0	3	9	13	12	
550 - 731	124	124	734					327	181	218	51	36	28	157	53	105	106	
550 - 731	72	72	726					81	25	22	24	3	12	42	96	39	63	
550 - 731	156	156	728					92	19	82	22	152	21					
732 - 914			134															
732 - 914			154															
732 - 914			106															
915 - 1097			154															
915 - 1097			138															
915 - 1097			102															
1098 - 1280			171															
1098 - 1280			180															
1098 - 1280			99															
1281 - 1463			212															
1281 - 1463			385															
1281 - 1463			127															
1281 - 1463			261															
Biomass (>366 m)								682	333	480	284	242	84	255	230	362	290	
Percent >366 m								99.5	68.8	55.7	55.7	78.6	49.2	57.6	40.6	49.9	38.4	
Biomass (all strata)		2209	761	1078	1401	2217	485	164	655	484	862	310	308	170	443	566	325	1042

Table 4 Abundance (000's) of Witch flounder from surveys in Div. 3N during spring of 1984-2000 by the Waifred Templeman (Eggle data converted to Campelen units for 1984-95)

Year	Old Stratum (sq. n. mi.) Area (sq. n. mi.)	New Stratum (sq. n. mi.) Area (sq. n. mi.)	Stratum	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
<56	1499	1593	375	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
57 - 92	1499	1499	376	0	0	0	26	0	0	0	0	0	0	0	0	0	0	0	34	34
57 - 92	2992	2992	360	2234	129	728	741	2641	220	0	0	59	225	0	0	0	132	65	244	613
57 - 92	1853	1853	361	155	0	32	36	0	28	0	0	0	0	0	36	0	0	0	0	212
57 - 92	2420	2420	362	0	95	25	27	173	0	0	0	0	0	0	0	0	0	0	0	0
57 - 92	2520	2520	373	0	0	50	0	0	0	0	0	0	0	0	0	0	0	0	0	0
57 - 92	931	931	374	0	0	0	0	0	0	0	0	0	0	43	43	0	0	0	0	0
57 - 92	674	674	383	0	62	0	31	0	0	0	0	0	0	0	0	0	0	0	0	0
93 - 183	421	421	359	405	58	232	58	985	203	0	0	0	29	0	0	0	0	0	0	203
93 - 183	100	100	377	14	0	0	186	7	83	0	0	0	0	0	0	0	0	0	0	0
93 - 183	647	647	382	0	0	0	30	0	0	0	0	0	0	0	0	0	0	0	0	0
184 - 274	225	225	358	77	557	93	279	31	46	93	0	93	294	232	31	77	83	261	15	41
184 - 274	139	139	378	48	29	48	354	86	115	0	0	96	0	0	0	0	0	0	0	0
184 - 274	182	182	381	25	13	42	163	75	0	25	0	0	0	0	0	0	0	0	13	0
275 - 366	164	164	357	23	180	53	11	237	56	0	90	124	102	23	40	30	373	259	293	0
275 - 366	106	106	379	66	36	68	423	102	44	109	7	44	0	22	0	0	18	6	102	28
275 - 366	116	116	380	8	88	0	247	32	8	8	0	0	0	0	0	0	0	0	8	0
367 - 549	155	155	723	0	0	0	0	0	0	288	341	256	53	181	45	51	149	96	171	0
367 - 549	105	105	725	0	0	0	0	0	0	166	101	87	0	13	235	26	51	72	0	0
367 - 549	160	160	727	0	0	0	0	0	0	0	0	11	55	22	0	0	11	33	33	21
550 - 731	124	124	724	0	0	0	0	0	0	1134	580	597	188	119	128	432	144	550	509	0
550 - 731	72	72	726	0	0	0	0	0	0	213	59	30	114	5	33	183	322	213	198	0
550 - 731	156	156	728	0	0	0	0	0	0	182	21	139	29	172	134	0	64	158	145	0
732 - 914	174	174	753	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
732 - 914	106	106	756	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
732 - 914	154	154	760	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
915 - 1097	138	138	753	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
915 - 1097	102	102	757	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
915 - 1097	173	173	761	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
1098 - 1280	180	180	754	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
1098 - 1280	99	99	758	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
1098 - 1280	212	212	762	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
1281 - 1463	185	185	755	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
1281 - 1463	127	127	759	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
1281 - 1463	261	261	763	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Abundance >= 366 m (000's)				1984	1013	1178	712	477	353	913	738	1101	1107							
Percent >= 366 m				99.6	72.7	62.3	64.1	84.1	75.0	77.1	49.5	56.5	41.0							
Total abundance (000's)				3053	1246	1837	2595	4180	955	320	1901	1394	1892	1110	567	470	1184	1493	1947	2701

Table 5. Abundance (000's) of Witch flounder from surveys in Div. 30 during spring 1984-2000 by the Alfred Neelzer and Wilfrid Templeman. (Engel data converted to Ciempoles units for 1984-95)

Year	Old Stratum Area (sq. n. mi.)	New Stratum Area (sq. n. mi.)	Stratum	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
57 - 92	2089	330		0	0	0	0	32	0	0	0	0	0	0	0	0	73	36	210
57 - 92	456	331	456	3555	376	94	31	1004	0	0	0	0	0	0	0	63	0	0	1104
57 - 92	1898	338	1898	209	11804	1505	1944	5418	2480	587	0	131	479	0	305	1417	0	671	1973
57 - 92	1716	340	1716	59	210	0	26	0	0	52	0	142	0	0	0	0	0	0	0
57 - 92	2520	351	2520	924	231	495	267	1317	240	116	0	0	0	0	0	0	0	0	39
57 - 92	2580	352	2580	101	1807	431	2048	1839	928	1775	51	80	51	44	71	79	197	35	1814
57 - 92	1282	353	1282	9347	1235	1713	2146	13050	3880	2910	0	265	333	0	35	35	265	439	5035
93 - 183	1721	329		0	0	0	0	1454	53	34	763	0	0	12263	521	0	35	68	623
93 - 183	1047	332	1047	11018	16592	6529	7230	16023	2832	10372	4513	5761	504	432	3925	2927	5665	1085	5045
93 - 183	948	337	948	130	9181	2634	3543	2641	2556	2608	3182	815	2087	87	1239	826	469	848	3709
93 - 183	585	339	585	443	0	80	268	134	0	0	0	0	0	0	0	161	36	80	36
93 - 183	474	354	474	1174	239	3282	450	619	106	359	261	261	1663	0	0	98	33	363	3208
184 - 274	151	187	151	21	156	35	0	145	52	332	1361	187	301	13447	423	30	277	140	267
184 - 274	121	336	121	25	17	174	67	208	0	158	1365	3287	266	3029	125	432	682	150	173
184 - 274	103	355	103	92	418	128	135	0	383	310	340	28	99	340	90	168	193	157	38
275 - 366	92	334	92	95	165	63	95	44	51	38	272	63	2238	40	462	880	7	161	
275 - 366	98	335	98	0	203	40	8	148	68	331	169	2340	223	215	108	192	243	12	169
275 - 366	61	356	61	17	214	38	55	109	80	126	92	348	319	189	126	88	40	90	54
367 - 549	93	717	93	166	717						32	371	166	5960	228	1363	11506	710	237
367 - 549	76	719	76	76	719						288	2335	267	37	42	364	1161	150	112
367 - 549	76	721	76	76	721						235	209	94	193	42	42	63	214	152
550 - 731	111	718	111	134	718						282	122	512	1161	535	518	507	517	324
550 - 731	105	720	105	105	720						361	376	1026	498	43	101	518	186	104
550 - 731	93	722	93	93	722						45	166	512	518	601	274	819	177	364
732 - 914		764		103	764									217					
732 - 914		768		99	768									501					
732 - 914		772		135	772														
915 - 1097		765		124	765														
915 - 1097		769		138	769														
915 - 1097		773		128	773														
1098 - 1280		766		144	766														
1098 - 1280		770		128	770														
1098 - 1280		774		135	774														
1281 - 1463		767		158	767														
1281 - 1463		771		175	771														
1281 - 1463		775		155	775														
Abundance >366 m (000's)											1243	3770	2376	9086	1491	2662	14635	1054	1293
Percent >366 m											9.3	21.3	28.7	22.0	17.5	27.6	61.7	30.3	5.2
Total abundance (000's)				27114	62867	17347	18286	44236	13811	20521	13317	17795	8983	41371	8568	9639	21725	6449	24969

Table 6 Biomass (tons) of Witch flounder from surveys in Div. 3N during fall 1990-2000 by the Wilfred Templeman, Alfred Needler and Tekeot (Engel data converted to Campelen units for 1990-94).														
Year				1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
Depth Range (meters)	Old Stratum Area (sq. n. mi.)	New Stratum Area (sq. n. mi.)	Stratum											
<=56	1593	1593	375	0	73	.	0	0	0	0	0	0	0	0
<=56	1499	1499	376	0	0	0	0	0	14	0	22	0	0	0
57 - 92	2992	2992	360	265	171	1297	173	75	888	23	427	431	177	535
57 - 92	1853	1853	361	28	467	463	0	32	0	0	14	0	268	28
57 - 92	2520	2520	362	400	221	87	0	0	0	0	0	0	32	0
57 - 92	2520	2520	373	0	0	0	0	0	0	0	0	0	0	0
57 - 92	931	931	374	0	0	.	0	0	0	0	0	0	0	0
57 - 92	674	674	383	0	0	.	0	0	0	0	0	0	0	0
93 - 183	421	421	359	0	0	278	0	0	22	0	0	1213	1	0
93 - 183	100	100	377	0	.	0	0	8	0	0	0	0	0	0
93 - 183	647	647	382	0	0	0	0	0	0	0	0	0	0	0
184 - 274	225	225	358	0	20	66	24	0	74	0	11	30	19	40
184 - 274	139	139	378	0	41	15	0	0	0	0	1	0	0	0
184 - 274	182	182	381	.	0	.	0	0	0	0	1	0	0	0
275 - 366	164	164	357	0	234	9	187	43	85	0	27	0	.	52
275 - 366	106	106	379	4	.	4	0	0	0	1	7	0	0	2
275 - 366	116	116	380	.	0	.	0	0	0	0	0	1	2	5
367 - 549	155	155	723	.	41	.	163	180	57	15	28	74	27	28
367 - 549	105	105	725	.	.	15	376	46	19	0	135	10	33	19
367 - 549	160	160	727	.	.	.	0	38	0	0	29	7	4	0
550 - 731	124	124	734	.	172	.	414	180	104	60	197	72	181	87
550 - 731	72	72	726	.	.	.	310	54	48	40	21	38	34	16
550 - 731	156	156	728	153	35	21	76	78	106	153
732 - 914	.	134	752	120	.	23
732 - 914	.	106	756	124	.	51
732 - 914	.	154	760	88	.	41
915 -1097	.	138	753	0	.	0
915 -1097	.	102	757	0	.	0
915 -1097	.	171	761	46	.	147
1098 -1280	.	180	754	0	.	0
1098 -1280	.	99	758	0	.	0
1098 -1280	.	212	762	0	.	0
1281 -1463	.	385	755	0	.	0
1281 -1463	.	127	759	0	.	0
1281 -1463	.	261	763	0	.	19
Biomass (>731 m)												379		
Percent >731 m												16.2		
Biomass (all strata)				696	1441	2235	1647	808	1346	160	993	2333	884	1244

Table 7. Biomass (tons) of Witch flounder from surveys in Div. 30 during fall 1990-2000 by the Wilfred Templeman, Alfred Needler and Teleost (Engel data converted to Campelen units for 1990-94).

Year				1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
Depth Range	Old Stratum	New Stratum	Stratum											
(meters)	Area (sq. n. mi.)	Area (sq. n. mi.)												
57 - 92	2089	2089	330	122	67	79	0	0	247	0	72	168	208	48
57 - 92	456	456	331	22	315	134	0	0	108	0	0	256	946	243
57 - 92	1898	1898	338	2226	438	837	3966	2193	4685	503	1329	483	2736	375
57 - 92	1716	1716	340	173	280	63	0	0	204	0	22	0	415	104
57 - 92	2520	2520	351	1690	284	72	0	0	0	0	0	37	205	0
57 - 92	2580	2580	352	1415	896	1352	946	228	379	273	573	374	1491	920
57 - 92	1282	1282	353	2405	343	477	0	732	538	789	168	1066	2996	2379
93 - 183	1721	1721	329	99	85	0	18	0	417	0	173	305	0	0
93 - 183	1047	1047	332	2102	155	1724	813	321	1114	4569	190	245	1664	544
93 - 183	948	948	337	1333	188	954	563	2132	421	492	322	479	978	344
93 - 183	585	585	339	1132	224	651	119	742	1911	0	481	261		344
93 - 183	474	474	354	1291	23	316	75	210	191	4647	215	201	103	766
184 - 274	151	147	333	221	11	22	30	92	26		4	6	33	4
184 - 274	121	121	336	82	151	76	298	13	35	32	19	19	67	31
184 - 274	103	103	355		497	93	120	25	16	343	6	14	110	35
275 - 366	92	96	334	24	16	0	9	17	4		5	1	7	5
275 - 366	58	58	335	194	25	25	30	18	1	23	0	1	23	8
275 - 366	61	61	356		11	7	430	98	7	60	3	4	32	22
367 - 549	93	166	717	30			0	32	37		12	42	260	0
367 - 549	76	76	719	110	2		65	6	1	226	19	9	10	14
367 - 549	76	76	721		18		169	67	21	54	6	14	67	17
550 - 731	111	134	718				22	68	8		68	47	53	34
550 - 731	105	105	720				73	0	13	68		2	17	4
550 - 731	93	93	722		9		81	21	14	39	12	12	26	8
732 - 914		105	764									75		12
732 - 914		99	768									18		7
732 - 914		135	772											
915 - 1097		124	765									173		62
915 - 1097		138	769									24		4
915 - 1097		128	773									17		5
1098 - 1280		144	766									4		13
1098 - 1280		128	770											24
1098 - 1280		135	774											4
1281 - 1463		158	767											4
1281 - 1463		175	771											15
1281 - 1463		155	775											0
														0
Biomass (>731 m)												311		148
Percent > 731 m												7.15		2.31
Biomass (all strata)				14671	4036	6884	7827	7013	10397	12117	3698	4356	12446	6396

Table 8 Abundance (000s) of Witch flounder from surveys in Div. 3N during fall 1990-2000 by the Wilfred Templeman, Alfred Needler and Teleost (Engel data converted to Campelen units for 1990-94).														
Year				1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
Depth Range	Old Stratum	New Stratum	Stratum											
(meters)	Area (sq. n. mi.)	Area (sq. n. mi.)												
<=56	1593	1593	375	0	55	.	0	0	0	0	0	0	0	0
<=56	1499	1499	376	0	0	0	0	0	23	0	19	0	0	0
57 - 92	2992	2992	360	382	206	1646	320	103	1232	41	672	755	360	926
57 - 92	1853	1853	361	32	425	701	0	42	0	0	23	0	306	51
57 - 92	2520	2520	362	441	277	116	0	0	0	0	0	0	50	0
57 - 92	2520	2520	373	0	0	0	0	0	0	0	0	0	0	0
57 - 92	931	931	374	0	0	.	0	0	0	0	0	0	0	0
57 - 92	674	674	383	0	0	.	0	0	0	0	0	0	0	0
93 - 183	421	421	359	0	0	608	0	0	87	0	0	2722	29	0
93 - 183	100	100	377	0	.	0	0	7	0	0	0	0	0	0
93 - 183	647	647	382	0	0	0	0	0	0	0	0	0	0	0
184 - 274	225	225	358	0	46	108	31	0	234	0	31	93	46	69
184 - 274	139	139	378	0	105	19	0	0	0	0	9	10	0	0
184 - 274	182	182	381	.	0	.	0	0	0	0	7	13	0	0
275 - 366	164	164	357	0	384	23	338	135	180	0	60	0	.	124
275 - 366	106	106	379	7	.	15	0	0	0	19	22	0	0	6
275 - 366	116	116	380	.	0	.	0	0	0	0	0	8	8	24
367 - 549	155	155	723	.	53	.	330	394	117	21	88	313	85	104
367 - 549	105	105	725	.	.	36	701	173	49	0	237	29	101	71
367 - 549	160	160	727	.	.	.	0	44	11	0	55	11	11	0
550 - 731	124	124	724	.	444	.	1126	512	223	178	571	326	640	337
550 - 731	72	72	726	.	.	.	669	114	119	99	40	92	125	40
550 - 731	156	156	728	268	195	129	212	215	311	417
732 - 914	.	134	752	165	.	28
732 - 914	.	106	756	255	.	149
732 - 914	.	154	760	244	.	229
915 -1097	.	138	753	0	.	9
915 -1097	.	102	757	0	.	0
915 -1097	.	171	761	106	.	578
1098 -1280	.	180	754	0	.	0
1098 -1280	.	99	758	0	.	0
1098 -1280	.	212	762	0	.	0
1281 -1463	.	385	755	0	.	0
1281 -1463	.	127	759	0
1281 -1463	.	261	763	72
Abundance >731 m (000's)												770		1065
Percent >731 m												14.4		32.9
Total abundance (000's)				863	1995	3272	3515	1793	2470	488	2046	5355	2073	3232

Year	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000			
Depth Range (meter)	Old Stratum Area (sq. n. mi.)	New Stratum Area (sq. n. mi.)	Stratum											
57 - 92	2089	2089	330	131	144	72	0	0	517	0	96	335	383	192
57 - 92	456	456	331	42	302	125	0	0	408	0	0	596	4799	533
57 - 92	1898	1898	338	3264	627	1436	6893	4700	8459	522	2872	1723	7572	609
57 - 92	1716	1716	340	262	330	118	0	0	295	0	47	0	1652	189
57 - 92	2520	2520	351	1837	347	58	0	0	0	0	0	50	347	0
57 - 92	2580	2580	352	1597	1242	2011	1115	355	371	355	1141	754	1825	1668
57 - 92	1282	1282	353	2822	485	941	0	1176	999	882	573	5467	5996	6172
93 - 183	1721	1721	329	132	101	0	47	0	663	0	616	852	0	0
93 - 183	1047	1047	352	3625	396	5281	2064	960	5233	11954	1248	2544	7393	3249
93 - 183	948	948	337	2347	424	2347	1043	5216	1435	717	1130	1613	3738	1623
93 - 183	585	585	339	1556	241	724	121	966	2776	0	1086	356	0	3943
93 - 183	474	474	354	1891	33	685	359	424	489	8955	489	782	391	2478
184 - 274	151	147	333	582	52	83	62	312	187	0	192	147	152	27
184 - 274	121	121	336	222	466	216	633	42	549	208	100	215	300	141
184 - 274	103	103	355	0	1459	298	425	85	63	768	28	170	411	85
275 - 366	92	96	334	76	70	0	21	57	56	0	33	20	58	18
275 - 366	58	58	335	371	100	112	68	52	64	64	4	40	48	37
275 - 366	61	61	356	0	25	8	1255	252	40	113	13	34	75	55
367 - 549	93	166	717	122	0	0	96	703	0	46	833	2166	0	0
367 - 549	76	76	719	209	42	0	277	10	52	612	183	178	99	75
367 - 549	76	76	721	0	47	0	444	183	102	131	17	125	311	98
550 - 731	111	134	718	0	0	0	107	428	164	0	535	618	581	396
550 - 731	105	105	720	0	0	0	339	0	105	316	0	29	202	39
550 - 731	93	93	722	0	26	0	243	58	64	134	51	103	122	70
732 - 914	0	105	764	0	0	0	0	0	0	0	0	357	0	72
732 - 914	0	99	768	0	0	0	0	0	0	0	0	217	0	24
732 - 914	0	135	772	0	0	0	0	0	0	0	0	1514	0	669
915 - 1097	0	124	765	0	0	0	0	0	0	0	0	165	0	31
915 - 1097	0	138	769	0	0	0	0	0	0	0	0	180	0	38
915 - 1097	0	128	773	0	0	0	0	0	0	0	0	35	0	136
1098 - 1280	0	144	766	0	0	0	0	0	0	0	0	0	0	113
1098 - 1280	0	128	770	0	0	0	0	0	0	0	0	0	0	36
1098 - 1280	0	135	774	0	0	0	0	0	0	0	0	0	0	28
1281 - 1463	0	158	767	0	0	0	0	0	0	0	0	0	0	65
1281 - 1463	0	175	771	0	0	0	0	0	0	0	0	0	0	0
1281 - 1463	0	155	775	0	0	0	0	0	0	0	0	0	0	0
Abundance >731 m (000's)												2468		1213
Percent >731 m												12.3		5.3
Total abundance (000's)				21086	7158	14515	15517	15369	23795	25731	10499	20054	38620	22907




Fig. 1 Commercial catches of witch flounder in Div. 3NO from 1960-2000 and TAC's from 1974-2001. Catches in recent years include estimates of those not reported.

*Note: Although a TAC of 3000 tons was agreed by the Fisheries Commission, it was also agreed that no directed fishing on witch flounder in Div. 3NO take place during 1994 due to the poor state of the stock.




Fig. 2 Biomass (tons) and abundance estimates (000s) of witch flounder from Canadian spring surveys in Div. 3N and 3O during 1984-2000




Fig. 3 Biomass (tons) and abundance estimates (000s) of witch flounder from Canadian fall surveys in Div. 3N and 3O during 1990-2000.




Fig. 4 Comparison of biomass (tons) and abundance estimates (000s) of witch flounder for converted data from Canadian spring (1984-2000) and fall (1990-2000) surveys in Div. 3NO combined.




Fig. 5 Length frequency distributions of witch flounder from both spring and fall surveys in 1995-2000 using the Campelen1800 shrimp trawl. Estimates represent abundance at length (cm) for the surveyed area.




Fig. 6 Number per standard set of Witch flounder from Canadian spring and fall surveys during 1998-2000.




Fig. 7 Weight (kg) per standard set of Witch flounder from Canadian spring and fall surveys during 1998-2000.