

Northwest Atlantic



Fisheries Organization

Serial No. N4442

NAFO SCR Doc. 01/64

## SCIENTIFIC COUNCIL MEETING – JUNE 2001

An Update of Distribution and Population Trends of Witch Flounder in Divisions 2J, 3K and 3L

by

W.R. Bowering

Dept. of Fisheries and Oceans, Science, Oceans & Environment Branch  
P.O. Box 5667 St. John's, NF, Canada A1C 5X1

### Abstract

Canadian fall survey distribution data from the late-1970s and early-1980s indicated that witch flounder were widely distributed throughout the shelf area in deeper channels around the fishing banks primarily in Div. 3K. By the mid-1980s, however, they were rapidly disappearing and by the early-1990s had virtually disappeared from the area entirely except for some very small catches along the slope and more to the southern area. They now appear to be located only along the deep continental slope area, especially in Div. 3L both inside and outside the Canadian 200-mile fishery zone. The results from the fall 1998-2000 surveys confirm that this distribution pattern remains. For the three divisions combined, there has been a very steady and rather systematic decline in the biomass index from about 65,000 tons in 1984 to less than 1,000 tons in 1995, by far the lowest in the time series. A small increase was observed between 1995 and 1996 and was almost exclusively a result of inclusion of the deeper strata surveyed in Div. 3L. The estimates have essentially remained the same since then. The stock size remains extremely low compared to the early-1980s.

### Fisheries and Management

The fishery for witch in this area began in the early-1960s and increased steadily from about 1,000 tons in 1963 to a peak of over 24,000 tons in 1973 (Table 1; Fig. 1). Catches declined rapidly to 2,800 tons by 1980 and subsequently fluctuated between 3,000 and 4,500 tons to 1991. The catch in 1992 declined to about 2,700 tons, the lowest since 1964, and further declined to around 400 tons by 1993 (Table 1). Until the late-1980s, the fishery was conducted by Poland, USSR and Canada (Table 1) mainly in Div. 3K (Table 1; Fig. 1). More recently, the regulated fishery has been mainly Canadian although EU (Portugal and Spain) has taken increased catches in the NAFO Regulatory area of Div. 3L since the mid-1980s. Although only 12 tons were reported for 1994, a catch of 491 tons was indicated for Spain in the Spanish Research Report (SCS Doc. 95/15) for the Regulatory Area of Div. 3L. In 1995 and 1996 total catches were estimated to be about 780 and 1370 tons, respectively. However, it is believed that these catches could be overestimated by 15-20% because of misreported Greenland halibut. The catches in 1997 and 1998 were estimated to be about 850 and 1,100 tons, respectively most of which was reported from the NAFO Regulatory Area of Div. 3L. The 1999 and 2000 catches were estimated to be about 300 and 700 tons, respectively.

During 1988-92, the Canadian fishery was particularly successful by fishing on prespawning concentrations in the deep slopes of Div. 3K, especially in depths beyond 700 m. Between 1988 and 1993, however, the area fished had become increasingly smaller and substantially deeper as the resource became depleted. The fishery during the winter of 1993 was very poor with the best catch rates occurring in depths greater than 1400 m. No directed fishing by Canada has been permitted since 1994 due to the poor state of the stock.

The stock has been regulated by TAC since 1974 (first introduced by ICNAF) and managed by Canada within its zone since the introduction of the 200 mile national limit and has been under moratorium from 1995 to the

present (Fig. 1). Because of the poor state of the stock, the NAFO Fisheries Commission agreed to extend the moratorium to the NAFO Regulatory Area in 1998 and has continued to 2001.

### **Canadian Research Vessel Surveys**

#### *Distribution*

Changes in spatial distribution patterns of witch flounder over the 20-year history of the surveys from 1978-97 were presented in the previous assessment as graphical distribution maps (ACON plots) (SCR Doc. 98/64). Survey distribution data from the late-1970s and early-1980s indicated that witch flounder were widely distributed throughout the shelf area in deeper channels around the fishing banks primarily in Div. 3K. By the mid-1980s, however, they were rapidly disappearing and by the early-1990s had virtually disappeared from the area entirely except for some very small catches along the slope and more to the southern area. They now appear to be located only along the deep continental slope area, especially in Division 3L both inside and outside the Canadian 200-mile fishery zone. The results from the fall 1998 (SCR Doc. 99/35), 1999 (SCR Doc. 00/13) and 2000 (Fig. 2) surveys confirm that this distribution remains.

#### *Biomass and Abundance Indices*

Stratified-random research vessel surveys have been conducted in the fall in Div. 2J, 3K and 3L since 1977, 1978 and 1981 respectively. As indicated above, up until 1994, the surveys were conducted using an *Engel 145'* high-rise groundfish trawl whereas the 1995-2000 surveys were carried out with a much more efficient *Campelen 1800* shrimp trawl. All data presented here are now in *Campelen 1800* trawl catch equivalents for 1977-94 with the actual data for 1995-2000.

For Div. 2J, biomass estimates ranged from as high as 5,900 tons in 1986 to a low of less than 300 tons in 1995 with only marginal increases since then with the 2000 estimate still only 500 tons (Table 2; Fig. 3).

In Div. 3K, during 1979-85, there was a period of relative stability where most annual biomass estimates were near 50,000 tons (Table 3; Fig. 3). Since that time estimates have declined considerably to less than 200 tons in 1995, the lowest in the time series. Estimates increased slightly after 1995 ranging from 900-1,300 tons between 1996-2000 (Table 3; Fig. 3).

For Div. 3L, biomass estimates varied generally between 7,000 and 10,000 tons from 1983 to 1990 but declined rapidly since then to a low of less than 400 tons in 1995 (Table 4; Fig. 3). The 1996 estimate increased to nearly 1,800 tons, however, more than half this estimate was based on the inclusion of deep-water strata (at depths of 732-1097 m) that weren't surveyed previously (Table 4). The 1997 estimate then declined to 1,100 tons although there was equal coverage to that of 1996 with 70% of the estimate attributed to the deeper strata. The 1998 estimate was similar to 1996 with more than half being attributed also to the inclusion of the new deeper strata. The 1999 estimate of about 800 tons is the lowest since the extension of the survey coverage to deeper water in 1996 with about 30% of the estimate accounted for by the new deep strata (Table 4; Fig. 3). Little change occurred in the 2000 survey, however, the biomass and abundance in the deeper strata appear to have been declining since 1996 (Table 4).

The abundance indices followed similar trends as biomass and are shown in Tables 5-7 for Div. 2J, 3K and 3L, respectively, and illustrated in Fig. 3 by Division and Fig. 4; Table 9 for the Divisions combined.

For the three Divisions combined, there has been a very steady and rather systematic decline in the biomass index from about 65,000 tons in 1984 to less than 1,000 tons in 1995, by far the lowest in the time series (Fig. 4; Table 8). Although there was a small increase between 1995 and 1996 there has been little change since then. The current level of stock size is still extremely low compared to the early-1980s.

### **Current Status**

The stock remains at an extremely low level with current indices of stock size based on survey trends at about 5% of the average of the early-1980s when the stock was considered at a reasonably healthy level.



Table 2 Estimated biomass (tons) of Witch Flounder (M+F) in each stratum from surveys in Div. 21 during fall of 1977-2000.  
(Engel 145 data converted to Campelen Units for 1977-94).

Year	Old Stratum Area (sq. n. mi.)	New Stratum Area (sq. n. mi.)	Stratum	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
101 - 200	1427	633	201	0	0	0	0	0	0	0	61	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	1823	1594	205	0	0	0	0	0	11	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	2582	1870	206	114	0	0	0	0	37	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	2246	2264	207	0	0	0	0	0	21	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		733	237																								
		778	238																								
201 - 300	440	621	202	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	1608	680	209	103	14	48	122	0	83	123	19	152	0	0	0	0	14	0	0	0	0	0	0	0	0	0	0
	774	1035	210	133	45	121	338	24	129	0	286	0	0	38	0	22	0	0	0	0	0	0	0	0	0	0	0
	1725	1583	213	265	249	160	298	280	371	197	118	102	130	98	21	56	0	0	0	0	0	0	0	0	0	0	0
	1171	1341	214	193	54	0	58	65	122	74	21	106	71	0	16	14	19	0	0	0	0	0	0	0	0	0	0
	1270	1302	215	193	33	11	0	82	67	0	45	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	1428	2196	228	508	134	301	543	183	678	264	467	79	728	93	123	151	76	0	44	35	0	0	0	0	0	0	0
301 - 400	508	530	234	0	35	36	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	480	487	203	0	0	0	0	0	54	112	0	0	19	0	0	20	0	0	0	0	0	0	0	0	0	0	0
	448	588	208	178	36	75	367	91	638	80	95	608	91	0	0	27	0	0	0	0	0	0	0	0	0	0	0
	330	251	211	447	198	100	289	70	242	12	99	72	27	38	0	34	0	0	0	0	0	0	0	0	0	0	0
	384	360	216	0	0	27	42	56	63	85	0	54	13	10	16	0	0	0	0	0	0	0	0	0	0	0	0
	441	450	222	197	99	29	103	155	285	69	26	46	0	0	173	46	0	10	0	0	0	0	0	0	0	0	0
	567	536	229	183	177	118	215	127	139	155	103	52	837	70	145	596	32	31	28	15	13	0	0	0	0	0	0
401 - 500	354	288	204	57	0	38	0	85	125	13	91	0	71	14	42	58	14	0	0	0	0	0	0	0	0	0	0
	268	241	217	0	0	15	0	0	0	0	0	0	54	64	44	72	6	0	0	0	13	0	0	0	0	0	0
	180	158	223	13	0	0	0	37	0	0	31	0	139	116	59	64	18	18	8	8	14	0	0	0	0	0	0
	686	598	227	161	123	44	482	180	358	211	85	147	329	411	203	228	1837	207	125	132	0	0	0	0	0	0	0
	420	414	235	813	0	456	430	502	371	908	517	399	121	168	0	62	149	37	20	0	41	0	16	3	22	0	0
	133	133	240																								
501 - 750	664	557	212	1564	106	640	193	630	1116	1390	822	1253	3139	834	392	588	639	111	272	44	52	71	96	90	184	261	125
	420	362	218	0	0	0	0	0	0	0	15	0	0	44	114	79	58	13	0	8	19	0	17	10	12	9	4
	270	228	224	0	0	0	0	0	0	0	0	0	32	48	120	125	17	49	33	0	23	0	18	23	16	1	15
	237	185	230	0	0	0	15	0	0	0	17	0	57	15	101	396	771	1711	346	85	105	69	126	176	26	161	36
		120	239																								
751 - 1000	213	283	219																								
	182	186	231	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	15	8	0	13	9	14	0	3
	122	193	236	0	0	0	0	0	0	0	0	0	0	0	0	0	0	457	176	197	118	115	6	36	228	143	283
1001 - 1250	324	303	220																								
	177	195	225	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	236	228	232	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
1251 - 1500	286	330	221																								
	180	201	226																								
	180	237	233																								
Biomass (t)				5123	1302	2218	3494	2582	4909	3693	2903	3030	5920	2063	1571	2653	3672	2669	1102	627	462	255	370	465	649	752	497

[illegible]

Table 4 Estimated biomass (tons) of Witch Flounder (M+F) per stratum from surveys in Div. 3L during fall of 1983-2000 (Engel 145 data converted to Campelen Units for 1983-94).																						
Year	Depth Range	Old Stratum Area (sq. n. mi.)	New Stratum Area (sq. n. mi.)	Stratum	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
				800																3		
	30 - 56		268	784														0	0	0	1	
	57 - 92	2071	2071	350	0	136	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
		1780	1780	363	0	85	0	50	0	0	0	264	33	41	0	0	0	0	0	0	0	
		1121	1121	371	0	46	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
		2460	2460	372	0	144	0	0	0	16	0	38	8	0	0	0	27	0	0	0	0	
		1120	1120	384	120	98	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
			465	785														0	0	0	0	
	93 - 183	1519	1519	328		45	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
		1574	1574	341	0	230	0	0	34	34	0	0	0	0	0	0	0	0	0	1	0	
		585	585	342	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
		525	525	343	0	84	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
		2120	2120	348	26	334	0	0	44	0	0	0	0	0	0	0	0	0	0	0	1	
		2114	2114	349	0	306	0	155	0	36	0	145	0	0	0	0	0	0	0	0	2	
		2817	2817	364	50	202	0	143	0	39	0	27	0	0	0	0	0	0	0	0	0	
		1041	1041	365	0	100	0	68	29	18	0	0	36	0	0	0	0	0	0	0	0	
		1320	1320	370	0	190	0	0	34	0	0	0	0	0	0	0	0	0	0	0	0	
		2356	2356	385	0	340	0	79	58	27	0	0	0	0	0	0	0	0	0	0	0	
		1481	1481	390	0	159	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
			84	786														1	0	0	0	
			613	787														0	0	0	0	
			261	788														0	0	0	0	
			89	790														0	0	1	0	
			72	793														0	0	0	0	
			216	794														0	0	0	0	
			98	797														0	0	0	0	
			72	799														0	0	0	0	
	184 - 274	1494	1582	344	159	159	37	29	127	0	0	0	0	0	0	0	0	0	0	0	1	
		983	983	347	41	467	0	42	0	154	66	0	0	0	0	0	0	0	0	0	0	
		1394	1394	366	0	186	355	307	171	110	187	27	0	7	0	0	0	0	0	0	0	
		961	961	369	181	374	570	706	320	1061	429	473	162	0	0	0	0	0	0	0	0	
		983	983	386		168	519	1082	1518	1750	442	218	307	875	0	0	0	0	0	0	0	
		821	821	389		196	133	760	250	138	21	79	0	27	0	0	38	0	0	0	11	
		282	282	391	0	0	32	0	9	0	0	0	70	22	0	0	36	0	25	0	0	
			164	795														0	0	0	0	
			72	789														0	0	0	0	
			227	791														6	0	0	0	
			100	798														0	2	21	3	
	275 - 366	1432	1432	345	5808	4484	1227	617	3693	2099	2358	750	0	61	73	0	10	3	5	35	3	
		865	865	346	2134	1423	2240	3321	1201	1823	1287	1863	203	40	14	0	0	12	3	1	20	
		334	334	368		47	29	386	23	64	144	106	39	14	0	0	22	0	0	0	6	
		718	718	387		169	404	276	572	1775	1546	3668	159	52	32	12	63	8	2	0	5	
		361	361	388		1229	48		589	92	126	0	125	173	0	14	0	0	0	12	5	
		145	145	392	17	55	13	20	50	13	0	0	0	0	0	4	0	0	0	0	0	
			175	796														0	1	2	0	
	367 - 549	186	186	729		146	127	280				48	274	246	42	131	2	151	24	0	1	
		216	216	731		498	248					465	178	356	38	79	19		0	7	19	
		468	468	733		328	1164					1618	2110	610	183	60	24	12	0	41	54	
		272	272	735		367	34	1714				222	216	40	12	3	20	23	18	12	3	
			50	792														55	37	11	29	
	550 - 731	170	170	730		104	16					130	6	140	88	83	0	21	11	10	59	
		231	231	732		282	235					29	207	283	41	194	16	147	121	440	252	
		228	228	734		30	184					168	100	11	106	49	37	127	15	149	95	
		175	175	736	546		268	709				355	913	90	70	20	10	261	41	135	93	
	732 - 914		227	737														130	104	435	151	
			223	741														115	164	313	7	
			348	745														154	212	123	65	
			159	748														87	0	0	40	
	915 - 1097		221	738														331	127	24	0	
			206	742														31	3	9	0	
			392	746														120	126	0	0	
			126	749														33	29	0	0	
	1098 - 1280		254	739														0	0	0	0	
			211	743														0	0	0	0	
			724	747														0	0	107	0	
			556	750														0	0	0	0	
	1281 - 1463		264	740														0	0	0	0	
			280	744														0	0	0	0	
			229	751														0	0	0	0	
Biomass >731 m																		1002	765	1011	262	
Percent >731 m																		55.4	70.3	53.1	31.8	
Biomass (tons)					9082	13210	7881	10743	8679	9294	6606	10341	5274	3131	778	663	390	1806	1087	1903	826	

Table 5 Abundance (000s) per stratum of Witch flounder (M+F) from research vessel surveys in Div. 2J during fall 1977-2000 (Engel data converted to Campelen Units for 1977-94).

[illegible]





Table 7 Abundance (000s) per stratum of Witch flounder (M+F) from research vessels in Div. 3L during the fall 1984-2000 (Engel data converted to Campelen Units for 1984-94).																				
Year				1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
Depth Range	Old Stratum	New Stratum	Stratum																	
(meters)	Area (sq. n. mi.)	Area (sq. n. mi.)																		
30 - 56			800															178		
		268	784													0	0	0		74
57 - 92	2071	2071	350	166	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	1780	1780	363	92	0	35	0	0	0	306	43	39	0	0	0	0	0	0	0	0
	1121	1121	371	44	0	0	0	0	0	0	0	0	0	0	0	0	44	0	0	0
	2460	2460	372	182	0	0	0	26	0	34	13	0	0	0	34	0	0	0	0	0
	1120	1120	384	128	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		465	785													0	0	0		
93 - 183	1519	1519	328	52	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	1574	1574	341	217	0	0	24	27	0	0	0	0	0	0	0	0	0	43	0	0
	585	585	342	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	40	0
	525	525	343	90	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	2120	2120	348	292	0	0	0	58	0	0	0	0	0	0	0	0	0	0	0	0
	2114	2114	349	291	0	162	0	32	0	166	0	0	0	0	0	0	42	0	0	42
	2817	2817	364	271	0	155	0	55	0	32	0	0	0	0	0	0	0	43	43	0
	1041	1041	365	143	0	57	48	29	0	0	48	0	0	0	0	0	0	0	0	0
	1320	1320	370	233	0	0	30	0	0	0	0	0	0	0	0	0	0	0	0	0
	2356	2356	385	324	0	122	36	25	0	0	0	0	0	0	0	0	0	0	0	0
	1481	1481	390	136	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		84	786													90	36	23		40
		613	787													0	0	0		0
		261	788													0	0	18		18
		89	790													6	18	55		0
		72	793													0	0	0		0
		216	794													0	0	0		0
		98	797													7	0	0		0
		72	799													0	0	0		0
184 - 274	1494	1582	344	206	46	117	154	0	0	0	0	0	0	0	0	0	0	0	0	64
	983	983	347	586	0	34	0	135	108	0	0	0	0	0	0	0	0	0	0	0
	1394	1394	366	157	362	431	219	110	164	32	0	8	0	0	0	0	38	0	38	0
	961	961	369	359	507	661	330	1348	529	463	162	0	0	0	39	0	0	0	0	0
	983	983	386	186	568	1082	1792	1974	352	237	270	1262	0	0	0	0	0	0	0	0
	821	821	389	169	158	875	226	169	28	75	0	38	0	0	33	0	0	0	0	301
	282	282	391	0	39	0	19	0	0	0	91	26	0	0	34	0	19	0	0	0
		164	795													0	0	0		56
		72	789													0	5	5		22
		227	791													42	62	0		0
		100	798													7	7	172		135
275 - 366	1432	1432	345	6895	1488	739	4531	2589	3180	2088	0	345	394	0	113	70	223	439	149	117
	865	865	346	2380	3498	3927	1487	2427	1606	2340	389	170	76	0	0	35	317	178	282	119
	334	334	368	46	46	459	23	69	207	115	69	14	0	0	23	0	23	0	20	23
	718	718	387	165	444	247	691	2025	1679	4971	198	66	33	77	99	49	44	0	44	593
	361	361	388	1440	50		819	149	149	0	116	199	0	14	0	0	0	149	0	124
	145	145	392	80	20	20	70	20	0	0	0	0	0	7	0	0	0	0	0	0
		175	796													0	107	24		21
367 - 549	186	186	729	217	192	409				64	341	422	51	290	34	375	115	0	0	34
	216	216	731	877	371					520	248	604	99	200	45		0	74	56	59
	468	468	733	338	1610					2221	2983	665	258	136	32	19	0	114	129	170
	272	272	735	661	37	2320					349	249	37	14	75	58	75	168	50	17
		50	792													901	423	279		915
550 - 731	170	170	730	105	23						117	12	195	171	108	0	47	19	21	58
	231	231	732	365	302					32	270	397	48	339	78	280	413	969	508	524
	228	228	734	21	267					251	110	16	141	146	44	467	70	380	173	125
	175	175	736		373	987				506	1613	217	241	34	75	782	277	1037	433	1432
732 - 914		227	737													468	297	1109	390	16
		223	741													291	460	892	14	0
		348	745													311	479	168	202	24
		159	748													186	0	0	50	0
915 - 1097		221	738													532	347	56	0	0
		206	742													43	14	14	13	0
		392	746													216	168	0	0	27
		126	749													61	43	0	0	0
1098 - 1280		254	739													0	0	0	0	0
		211	743													0	0	0	0	0
		724	747													0	0	100	0	0
		556	750													0	0	0	0	0
1281 - 1463		264	740													0	0	0	0	0
		280	744													0	0	0	0	0
		229	751													0	14	0		0
Abundance (000's)				17914	10401	12839	10500	11269	8002	14453	7428	4748	1572	1428	865	5297	4227	6754	2655	5361

Table 8 Estimates of biomass (tons) of witch flounder from Canadian fall surveys in Div. 2J, 3K and 3L during 1977-2000.

YEAR	DIV. 2J	DIV. 3K	DIV. 3L	TOTAL
1977	5123			
1978	1302	30353		
1979	2218	49789		
1980	3494	44962		
1981	2582	43405		
1982	4909	32429		
1983	3693	49250		
1984	2903	49038	13210	65151
1985	3030	35694	7881	46605
1986	5920	21359	10743	38022
1987	2063	21746	8679	32488
1988	1571	18110	9294	28975
1989	2653	8976	6606	18234
1990	3672	17088	10341	31101
1991	2669	4272	5274	12215
1992	1102	1863	3131	6095
1993	627	1327	778	2733
1994	462	846	663	1971
1995	255	184	390	828
1996	370	855	1806	3031
1997	465	1116	1087	2669
1998	649	1255	1906	3810
1999	752	881	828	2459
2000	497	1200	468	2165

Table 9 Estimates of abundance (000s) of witch flounder from Canadian fall surveys in Div. 2J, 3K and 3L during 1977-2000.

YEAR	DIV. 2J	DIV. 3K	DIV. 3L	TOTAL
1977	7106			
1978	1962	59729		
1979	3016	84954		
1980	4503	72871		
1981	3190	70058		
1982	6486	52145		
1983	4963	75267		
1984	3840	79554	17914	101307
1985	4089	70384	10401	84874
1986	9432	40917	12839	63188
1987	3337	37279	10500	51117
1988	2746	35486	11269	49501
1989	5377	22734	8002	36113
1990	8110	29338	14453	51901
1991	6941	10045	7428	24414
1992	2463	6377	4748	13588
1993	2588	8918	1572	13078
1994	2369	4815	1428	8612
1995	1696	2191	865	4753
1996	1724	5081	5297	12102
1997	1890	5716	4227	11833
1998	2505	7955	6755	17215
1999	2548	5441	2655	10644
2000	1964	7952	5361	15277




Fig. 1 Commercial catches and TAC's of witch flounder in Divisions 2J, 3K and 3L during 1963-2001. Catches in Division 3M are included for 1998-2000.




Fig. 2 Weight (kg) per set of Witch flounder from Canadian surveys in NAFO Divisions 2J, 3K and 3L during fall 2000 .




Fig. 3 Biomass (tons) and abundance (000s) of witch flounder by division from Canadian surveys in Div. 2J, 3K and 3L during 1977-2000. Data based on Campelen trawl catch equivalents.




Fig. 4 Biomass (tons) and abundance (000s) of Divisions 2J, 3K and 3L combined, of witch flounder from Canadian fall surveys based on Campelen trawl catch equivalents during 1984-2000.