



Serial No. N4532

NAFO SCR Doc. 01/138

SCIENTIFIC COUNCIL MEETING – SEPTEMBER 2001
(Deep-sea Fisheries Symposium – Poster)

Testicular Activity and Sperm Glycoproteins in Giant Red Shrimp *Aristaeomorpha foliacea*

by

S. Desantis, M. Labate, F. Cirillo, G. M. Labate, P. Maiorano*, and G. D'Onghia*

Laboratory of Comparative Anatomy and Cytology, * Laboratory of Ecology - Department of Zoology University of Bari, Via E. Orabona 4, 70125 Bari, Italy
e-mail: s.desantis@biologia.uniba.it

Abstract

By histological and histochemistry methods we found that in male *Aristaeomorpha foliacea*, collected from the late winter to the summer in the north-western Ionian Sea (Mediterranean Sea), the spermiogenesis and the glycoprotein pattern undergo seasonal changes and that the sperm glycoproteins mature as gametes transit from the testis to the terminal ampulla. In serial sections stained with hematoxylin and eosin the testicular activity appeared discontinuous: in late winter, testes did not show meiotic and seminiferous epithelium consisted of interkinetic spermatogonia and spermatozoa; in spring, the high spermiogenetic activity occurred and the seminiferous epithelium was mainly constituted of spermatocytes and spermatozoa; in summer, the testicular activity seemed locked again since testes containing both spermatocytes and spermatozoa were lacking. The use of twelve different lectins evidenced that the intratesticular spermatozoa from late winter to summer contain surface binding sites for SNA, MAA, Con A and KOH-sialidase (si)-WGA; in March and July they exhibited also nuclear and cytoplasmic reactivity for SNA and Con A. In the hemispermaphore the spermatozoa displayed a more complex lectin binding pattern because they also reacted with PNA, DBA, HPA, GSA II. The staining with DBA, KOH-si-DBA, and GSA II evidenced differences between the spermatozoa from late winter-spring hemispermaphore and summer hemispermaphore ones: the former showed a nuclear affinity whereas the latter displayed surface and/or cytoplasm staining. No reaction was observed with SBA, GSA I-B₄, UEA I, and LTA.

Introduction

The giant red shrimp *Aristaeomorpha foliacea* (Risso, 1827), generally distributed at depths between 300-700 m, is a crustacean widespread in the eastern and western Atlantic, Indian Ocean and western Pacific, in the waters of Japan, Australia, New Zealand and in the Mediterranean Sea (Holtius, 1980). This species plays an important role in the overall biomass of the muddy bottoms of the Mediterranean Sea and represents an important commercial resource among crustaceans since it is much appreciated by the consumer.

In *Aristaeomorpha foliacea*, studies have been carried on the general biology, distribution and population dynamics (D'Onghia *et al.*, 1998 for references) and, as in many crustacean species, the male reproductive tract has received little attention. Only histological observations of spermatophore formation (Tunesi, 1987) and, more recently, the ultrastructural aspects of spermatozoa (Medina, 1995) have been described, whereas studies on both the testicular activity and the glycoconjugate composition of spermatozoa are lacking.

Glycoconjugates are a fundamental component of eukaryotic cells, and although their biological or physiological functions are known in many cases, the biological roles of their oligosaccharides remain mostly undefined (Lis and Sharon, 1993; Varki, 1993). Lectins are useful probes for intracellular localization of sugar residues and characterization of distinct cellular populations as well as cell-to-cell interactions and variation of biological activity (Spicer and Schulte, 1992; Danguy *et al.*, 1994). Since *Aristaeomorpha foliacea* spermatozoa lack acrosome (Medina, 1995), the sperm surface as well as the cytoplasmic content seems to be of critical importance in the process of fertilization.

The aim of this study is to investigate the testicular activity as well as the glycoconjugate pattern in testicular and hemispermatozoa spermatozoa from late winter to summer using a series of lectins with the attainment of sperm physiological maturation in understanding the reproductive dynamics of this commercially important species.

Materials and Methods

A total of 15 mature males of *Aristaeomorpha foliacea* (carapace length >32 mm), considered sexually mature for the presence of the hemispermatozoa in the terminal ampullae and joined petasma (Sardà and Demestre, 1989), were collected in late winter (March) (n= 2), spring (April) (n= 8) and summer (July) (n=5) by commercial bottom-trawl gear in the north-western Ionian Sea (Mediterranean Sea) (Fig. 1). The testes and the vas deferentes were removed immediately after capture, fixed in Bouin's solution, dehydrated in an ascending ethanol series, and embedded in paraffin wax. Sections 5 µm thick were cut and, after dewaxing, were stained with Mayer's hematoxylin and eosin for histological observations or processed for lectin histochemical studies.

Lectin histochemistry

The lectins used are listed in Table 1. Dewaxed and rehydrated tissue sections were immersed in 3% H₂O₂ for 10 min to suppress the endogenous peroxidase activity, rinsed in 0.05 M Tris-HCl buffered saline (TBS) pH 7.4 and incubated in lectin solution at appropriate dilutions (10-25 µg/ml) for 1 h at room temperature (RT). After 3 rinsing in TBS, peroxidase activity was visualized by incubation in a solution containing 0.05% 3,3'-diaminobenzidine (DAB) and 0.003% H₂O₂ in 0.05 M TBS (pH 7.6) for 10 min at RT before dehydration and mounting. Tissue sections incubated in biotinylated lectins (SNA,MAA and GSA I-B₄) were rinsed 3 times with 0.05 M phosphate-buffered saline (PBS) and were incubated in streptavidin/peroxidase complex (Vector Lab. Inc., Burlingame, CA) for 30 min at RT. After washing in PBS, peroxidase was developed in a DAB- H₂O₂ solution.

Controls for lectin staining included: (1) substitution of the substrate medium with buffer without lectin; (2) incubation with each lectin in the presence of its haptan sugar (0.2 M).

Sialidase digestion

Before staining with SNA,MAA,PNA,DBA,WGA, some sections were incubated at 37 °C for 16 h in 0.1 Uml⁻¹ of sialidase (Type V, from *Clostridium perfringens*, from Sigma, St. Louis, MO) dissolved in 0.1 M sodium acetate buffer, pH 5.5, containing 10 mM CaCl₂. Prior to the neuraminidase treatment, a saponification technique was performed to render the enzyme digestion effective, with 0.5% KOH in 70% ethanol for 15 min at RT (Reid *et al.*, 1978). As controls of the enzyme digestion procedure, certain sections were incubated in the enzyme-free buffer solution under the same experimental conditions.

Results

Histology

The testes of *Aristaeomorpha foliacea* are a pair of convoluted tubules containing germinal cells, involved in the production of sperm, and somatic (accessory) cells. Spermatogonia are found along one margin, whereas the bulk is occupied by developing germ cells and/or spermatozoa (Fig. 2a,c,d,e,f).

Testes showed a different spermatogenetic activity from March to July. In late winter (March) the testes display interkinetic spermatogonia (Ø 12,08 ± 0,56 µm) and spermatozoa (Ø 3,14 ± 0.17 µm) (Fig. 2c). The spermatozoa were slightly elliptic in shape and were characterized by a strongly hematoxylinophil nucleus surrounded by an

unstained peripheral band of cytoplasm. In spring (April) the testes contained less spermatogonia, while they were packed with developing spermatocytes and spermatozoa (Fig. 2d). In summer (July) the testes were found in the following three conditions: 1) spermatogonia and primary spermatocytes (Fig. 2e), 2) spermatogonia and secondary spermatocytes ($\bar{O} 5,34 \pm 0,24$) (Fig. 2f), 3) spermatogonia and spermatozoa.

The hemispermatozoa are hardened structures constituted of four layers, the first of them surrounds the spermatozoa which are enmeshed in an extracellular matrix (Fig. 2b).

Lectin histochemistry

Since the cytoplasm of spermatozoa is constituted of a thin perinuclear band it was very difficult to clearly distinguish the cell surface from the cytoplasm, therefore the material stained outside the nuclear region we sometimes indicate as cytoplasm.

The lectin binding pattern of testicular (T) and hemispermatozoa (H) spermatozoa are summarized in Table 2.

SNA weakly bound the T spermatozoa surface in spring, the whole spermic cell in March and a few in July (Fig. 3a); this lectin moderately marked the surface and the cytoplasm in H spermatozoa (Fig. 3b).

MAA moderately marked the surface in T spermatozoa (Fig. 3c) and also the cytoplasm in H ones (Fig. 3d).

PNA did not find binding sites in T spermatozoa but weakly reacted with the cytoplasm of H spermatozoa. KOH-sialidase (si)-treatments revealed cryptic binding sites in the nucleus of H spermatozoa.

DBA did not show binding sites in spermatozoa of T, whereas the lectin gave a faintly visible reaction in the nuclear region of rare spermatozoa in the spring H, and a weak staining in the cytoplasm of spermatozoa contained in summer H. DBA affinity was increased by KOH-si-sialidase treatments only in cytoplasm of H sperm mass.

HPA did not marked T spermatozoa, whereas showed weak reactions in the cytoplasm of H spermatozoa.

Con A showed staining on the surface and in the cytoplasm of T and H spermatozoa (Fig. 3e); the H spermatozoa displayed binding sites also in the nucleus region (Fig. 3f).

The KOH-si-WGA procedure intensely marked the surface of T spermatozoa (Fig. 3g), and gave a faintly visible reaction in the nucleus and the cytoplasm of H spermatozoa (Fig. 3h).

GSA II did not react with T spermatozoa, whereas in H showed binding sites both for the cytoplasm and the nucleus in spring spermatozoa, and it stained only the cytoplasm in summer ones.

SBA, GSA I-B₄, UEA I, and LTA did not show binding patterns.

Discussion

The present study was performed on testes and hemispermatozoa from mature males of *Aristaeomorpha foliacea* collected in the Mediterranean Sea during the major mating and reproductive periods. In the Mediterranean Sea, the highest percentage of mature males has been found between January and July (Mura *et al.*, 1992; Ragonese and Bianchini, 1995; D'Onghia *et al.*, 1998). Coupling occurs a few months before ovulation because the highest percentage of females with spermatozoa in the thelycum has been observed during March and May (D'Onghia *et al.*, 1998), whereas female with mature ovaries (Levi and Vacchi, 1988; Desantis *et al.*, 2001) occur from May to September with the highest percentage during August.

As in other Decapoda (Kaestner, 1970), the testes of *Aristaeomorpha foliacea* are a pair of tubes in which the male reproductive cells are produced. In this shrimp the testicular activity is appeared discontinuous. In late winter, testes activity is locked since they did not show meiotic activity and seminiferous epithelium consisted of interkinetic spermatogonia and spermatozoa. These spermatozoa could represent the remains of a previous spermiogenesis period. The high spermiogenetic activity occurs in spring when the seminiferous epithelium was mainly constituted of primary spermatocytes and spermatozoa. In summer the testicular activity seems locked

again: testes containing both spermatocytes and spermatozoa are lacking. Seasonal changes of testicular activity have been reported in other crustaceans (King, 1948; Meusy, 1963; Du *et al.*, 1988; Sreekumar and Adiyodi, 1983; Pochon-Masson, 1994) as well as in the companion species *Aristeus antennatus* collected in the same basin (Desantis *et al.*, 1999). These changes can be related to seasonal changes in the androgenic gland (Taketomi, 1986; Legrand and Juchault, 1994), which is under the influence of the protocerebral neurohormones (Tourir, 1977; Payen and Amato, 1978; Martin, 1994).

The glycoproteins fall into two main categories according to the attachment of the oligosaccharide to the peptide (Kornfeld and Kornfeld, 1985). The two types include those in which a reducing terminal GalNAc is linked O-glycosidically to the hydroxyl of serine or threonine and those in which a reducing terminal of GlcNAc is bound N-glycosidically to the epsilon amine of asparagine. Among the lectins used in this study PNA, DBA, or HPA specifically identify the many O-linked oligosaccharides (Spicer and Schulte, 1992), whereas Con A visualizes specifically the glycoproteins containing N-linked oligosaccharides (Bernhard and Avrameas, 1971). The surface of intratesticular spermatozoa reacted with SNA, MAA, Con A and KOH-si-WGA showing the presence of N-linked oligosaccharides terminating or not with sialic acid, besides in March and July intratesticular spermatozoa showed nuclear and cytoplasmic binding sites for SNA and Con A, respectively. This similar lectin reactivity strengthens the assumption that late winter and summer testicular spermatozoa could represent the remains of a previous spermiogenesis period, whereas spring spermatozoa are a recent spermiogenic product. In hemispermaphore the spermatozoa displayed a more complex glycoprotein pattern than in testis because they also reacted with PNA, DBA, HPA (specific for O-linked oligosaccharides), and GSA II. In addition, the staining with DBA, KOH-si-DBA, and GSA II evidenced differences between the hemispermaphore late winter-spring spermatozoa and the summer hemispermaphore ones. During the late winter-spring the spermatozoa showed a nuclear affinity for DBA, increased after KOH-sialidase procedure, and Con A. They were wholly stained by GSA II. In summer the hemispermaphore spermatozoa displayed surface and/or cytoplasm affinity for DBA, KOH-si-DBA, GSA II. These findings suggest that the glycoprotein sperm pattern is seasonally different and that the spermatozoa mature as they transit from the vas deferens to the terminal ampulla. In other species, such as *Inachus falangium* (Diesel, 1989), it was found that mature spermatozoa are formed in the medial portion of the vas deferens. The function of the terminal ampulla is connected to maturation of spermatozoa in *Penaeus setiferus* and *P. vannamei* (Chow *et al.*, 1991). The referred cytoplasmic lectin binding pattern can hide surface glycoproteins or oligosaccharides contained in vesicles. The former could be involved in the interaction with oocyte and the latter may represent an acrosome-like structure since no acrosomal structure is recognizable in *Aristaeomorpha foliacea* (Medina, 1995). To our knowledge lectin histochemical studies on spermatozoa from crustaceans are lacking, but in vertebrates it is well known that spermatozoa undergo changes in their surface and acrosome glycoconjugates as they transit through extratesticular ductus (Arya and Vanha-Perttula, 1984; Burkett *et al.*, 1987; Eddy, 1988; Vreeburg *et al.*, 1992; Labate *et al.*, 1997; Ueda *et al.*, 1997). These modifications are of critical importance since the glycoconjugates are involved in inter- and intracellular processes of the fertilization.

In conclusion, this study shows that in mature male of *Aristaeomorpha foliacea* 1) seasonal changes of testicular activity occur, 2) sperm glycoprotein pattern of active spermiogenesis period (spring) is different from locked periods (late winter and summer), 3) spermatozoa undergo maturative changes in glycoconjugates during their transit from testis to hemispermaphore.

References

- ARYA, M., and T. VANHA-PERTTULA. 1984. Distribution of lectin binding in rat testis and epididymis. *Andrologia*, **16**: 495-508.
- BERNHARD, W., and S. AVRAMEAS. 1971. Ultrastructural visualization of cellular carbohydrate components by means of concanavalin A. *Exp. Cell Res.*, **64**: 232.
- BURKETT, B. N., B. A. SCHULTE, and S. S. SPICER. 1987. Histochemical evaluation of glycoconjugates in the male reproductive tract with lectin-horseradish peroxidase conjugates: II. Staining of ciliated cells, basal cells, flask cells, and clear cells in the mouse. *Am J Anat.*, **178**: 23-29.
- CHOW, S., M.M. DOUGHERTY, W.J. DOUGHERTY, and P.A. SANDIFER. 1991. Spermaphore formation in the white shrimps *Penaeus setiferus* and *P. vannamei*. *J. Crust. Biol.*, **11**: 201-216.

- D'ONGHIA, G., P. MAIORANO, A. MATARRESE, and A. TURSI. 1998. Distribution, biology, and population dynamics of *Aristaeomorpha foliacea* (Risso, 1827) (Decapoda, Natantia, Aristeidae) in the North-western Ionian Sea (Mediterranean Sea). *Crustaceana*, **71**: 518-544.
- DANGUY, A., F. AFIK, B. PAJAK, and H.J. GABIUS. 1994. Contribution of carbohydrate histochemistry to glycobiology. *Histol. and Histopathol.*, **9**: 155-171.
- DE SANTIS, S., M. LABATE, A. TURSI, G. D'ONGHIA, and P. MAIORANO. 1999. Testicular activity in the shrimp *Aristeus antennatus* (Risso, 1816). pp: 903-914. F. R. Schram and J. C. von Vaupel Klein (Eds.). *Crustaceans and the Biodiversity Crisis. Vol. I*
- DEMESTRE, M. and J. M. FORTUÑO. 1992. Reproduction of the deep-water shrimp *Aristeus antennatus* (Decapoda: Dendrobranchiata). *Mar. Ecol. Prog. Ser.*, **84**: 41-51.
- DIESEL, R. 1989. Structure and function of the reproductive system of the symbiotic spider crab *Inachus falganum* (Decapoda: Majidae): observation on sperma transfer, sperm storage, and spawning. *J. Crust. Biol.*, **9**: 266-277.
- DOUGHERTY, W. J., M. M., DOUGHERTY, and S. J. Harris. 1986. Ultrastructural and histochemical observation on electroejaculated spermatophores of the palaemonid shrimp, *Macrobrachium rosenbergii*. *Tissue Cell*, **18**: 709-724.
- DU, N., L. XUE, and W. LAI. 1988. Histology of the male reproductive system in *Eriocheir sinensis* (Decapoda, Crustacea). *Acta zool. Sinica*, **34**: 329-333.
- EDDY, E. M. 1988. The spermatoon. pp: 27-68. E. Kobil and J. Neill (Eds.). *The Physiology of Reproduction*.
- HOLTHUIS, L. B. 1980. FAO species catalogue. Shrimps and prawns of the world: an annotated catalogue of species of interest to fisheries. *FAO Fish. Synop.*, **125** (1): 1-261.
- KAESTNER, A. 1970. Crustacea. *Invertebrate zoology*, **3**: 266-366.
- KING, J. E. 1948. A study of the reproductive organs of the common marine shrimp, *Penaeus setiferus* (L.). *Biol. Bull., Woods Hole*, **94**: 244-262.
- KORNFELD, R., and S. KORNFELD. 1985. Assembly of asparagina-linked oligosaccharides. *Ann. Rev. Biochem.*, **54**: 631-664.
- LABATE, M., S. DESANTIS, and A. CORRIERO. 1997. Glycoconjugates during the annual spermatogenetic cycle in lizard epididymal ductuli efferentes: a histochemical study. *Eur. J. Histochem.*, **41**: 47-56.
- LEGRAND, J. J., and P. JUCHAULT. 1994. Ontogenèse du sexe et physiologie sexuelle. pp : 631-713. J. Forest (Ed.). *Traité de Zoologie, Crustacés. Vol. 7*.
- LEVI, D., and M. VACCHI. 1988. Macroscopic scale for simple and rapid determination of sexual maturity in *Aristaeomorpha foliacea* (Risso, 1826) (Decapoda, Penaeidae). *J. Crust. Biol.*, **8**: 532-538.
- LIS, H., and N. SHARON. 1993. Protein glycosylation. Structural and functional aspects. *Eur. J. Biochem.*, **218**: 1-27.
- MARTIN, G. 1994. Système nerveux. II. Neurohormones. pp : 301-344. J. Forest (Ed.). *Traité de Zoologie, Crustacés. Vol. 7*.

- MEDINA, A. 1995. The atypical sperm morphologies of *Aristeus antennatus* and *Aristaeomorpha foliacea* (Crustacea, Dendrobranchiata, Aristeidae) and their phylogenetic significance. *Mém. Mus. natn. Hist. nat., Paris*, **166**: 243-250.
- MEUSY, J. J. 1963. La gamétogénèse d'*Orchestia gammarella* Pallas, Crustacé Amphipode. *Bull. Soc. zool. France*, **88**: 197-220.
- MURA, M., S. CAMPISI, and A. CAU. 1992. Osservazioni sulla biologia riproduttiva negli aristeidi demersali del Mediterraneo centro-occidentale. *Oebalia*, **17** (suppl. 2): 61-66.
- PAYEN, G., and G. D. AMATO. 1978. Données actuelles sur le contrôle de la spermatogénèse chez les Crustacés Décapodes Reptantia. *Arch. Zool. exp. gén.*, **119**: 447-463.
- POCHON-MASSON, J. 1994. Les gamétogénèses. pp : 727-784. J. Forest (Ed.). *Traité de Zoologie, Crustacés. Vol. 7*.
- RAGONESE, S., and M. L. BIANCHINI. 1995. Size at sexual maturity in red shrimp females, *Aristaeomorpha foliacea*, from the Sicilian Channel (Mediterranean Sea). *Crustaceana*, **68**: 73-82.
- REID, P. F., C. F. CULLING, and W. L. DUNN . 1978. A histochemical method for the identification of 9-O-acetyl sialic acids. An investigation of bovine submaxillary gland and intestinal mucins. *J. Histochem. Cytochem.*, **26**: 187-192.
- SARDÀ, F., and M. DEMESTRE. 1989. Shortening of the rostrum and rostral variability in *Aristeus antennatus* (Risso, 1816) (Decapoda: Aristeidae). *J. Crust. Biol.*, **9**: 570-577.
- SPICER, S. S., and D. A. SCHULTE. 1992. Diversity of cell glycoconjugates shown histochemically: a perspective. *J. Histochem. Cytochem.*, **40**: 1-38.
- SREEKUMAR, S., and R. G. ADIYODI. 1983. Role of eyestalk principle(s) in spermatogenesis of the freshwater shrimp, *Macrobrachium idella*. *Proc. First int. biennal Conf. worm Water Aquacult; Hawaii*: 15-16.
- TAKETOMI, Y. 1986. Ultrastructure of androgenic gland of the crayfish, *Procambarus clarkii*. *Cell. Biol. int. Rep.*, **10**: 131-136.
- TOUIR, A. 1977. Données nouvelles concernant l'endocrinologie sexuelle des Crustacés Décapodes Natantia hermaphrodites et gonochoriques. II. Maintien des gonies et évolution des gamétogénèses in vivo et in vitro. *C. R. Acad. Sci., Paris, (D)* **284**: 2515-2518.
- TUNESI, L. 1987. La formation du spermatophore chez *Aristaeomorpha foliacea* (Risso, 1827). *Inv. Pesq.*, **51** (Supl. 1): 471-476.
- UEDA, T., O. FUJIMORI, A., and K. YAMADA. 1997. Histochemical analysis of sialic acids in the epididymis of the rat. *Histochem. Cell. Biol.*, **109**: 399-407.
- VARKI, A. 1993. Biological roles of oligosaccharides: all theories are correct. *Glycobiology*, **3**: 97-130.
- VREEBURG, J. T. M., M. K. HOLLAND, and M. C. ORGEVIN-CRIST. 1992. Binding of epididymal proteins to rat spermatozoa in vivo. *Biol. Reprod.*, **47**: 588-597.
- YUFENG, W., D. NANSHAN, and L. WEI. 1997. A histochemical study on the male reproductive system of *Macrobrachium rosenbergii*. *J. Fish. Sci. China*, **4**: 13-16.

Table 1. Lectin used, their sugar specificities and inhibitory sugar used in control experiments

Lectin abbreviation	Source of lectin	Sugar specificity	Inhibitory sugar (0.2 M)*
SNA	<i>Sambucus nigra</i>	NeuNAc ₂ Gal/GalNAc	NeuNAc
MAA	<i>Maackia amurensis</i>	NeuNAc ₂ ,3-Gal/ β 1,4-GlcNAc	NeuNAc
PNA	<i>Arachis hypogea</i>	Terminal β -D-Gal(1-3)-GalNAc	Galactose
DBA	<i>Dolichos biflorus</i>	Terminal FP>GalNAc _{1,3} GalNAc	GalNAc
SBA	<i>Glycine max</i>	Terminal α / β -D-GalNAc	GalNAc
HPA	<i>Helix pomatia</i>	Terminal α -GalNAc	GalNAc
Con A	<i>Canavalia ensiformis</i>	Terminal and internal α -D-Man> α -D-Glc	Mannose
GSA I-B4	<i>Bandeiraea simplicifolia</i>	Terminal α -D-Gal	Galactose
WGA	<i>Triticum vulgare</i>	Terminal and internal β -D-GlcNAc>> NeuNAc	GlcNAc
GSA II	<i>Bandeiraea simplicifolia</i>	Terminal D-GlcNAc	GlcNAc
UEA I	<i>Ulex europaeus</i>	Terminal α -L-Fuc	Fucose
LTA	<i>Lotus tetragonolobus</i>	Terminal α -L-Fuc	Fucose

FP, Forssman pentasaccharide GalNAc_{1,3}GalNAc_{1,3}Gal β 1,4Gal β 1,4GlcNAc; Fuc, Fucose; Gal, galactose; GalNAc, N-acetylgalactosamine; Glc, glucose; GlcNAc, N-acetylglucosamine; Man, mannose; NeuNAc, N-acetyl neuraminic acid; *, concentration of inhibiting sugars.

Table 2. Summary of lectin binding to the sperm mass in testis and hemispermatochore of *Aristaeomorpha flitacea*

Lectin	LATE WINTER-SPRING		SUMMER	
	T	H	T	H
SNA	+s/+wMr	++	+s/++w*	++
MAA	++s	++	++s	++
PNA	-	+	-	+
KOH-si-PNA	-	+ w	-	+w
DBA	-	-/ \pm n*	-	+
KOH-si-DBA	-	\pm / \pm n*	-	++
SBA	-	-	-	-
HPA	-	+	-	+
Con A	++s/+Mr	++/+n	+++s/+	+/+n
GSA I-B4	-	-	-	-
KOH-si-WGA	++s	++s/ \pm w	++s	++s/ \pm w
GSA II	-	++w	-	++
UEA I	-	-	-	-
LTA	-	-	-	-

H, hemispermatochore; Mr, March; n, nucleus; s, surface; si, sialidase (neuraminidase); T, testis; w, whole cell; *, rare positive reaction; -, negative reaction; \pm , faintly visible reaction; +, ++, +++, weak, moderate, intense positive reactions. Where not specified, the reactions concern both the surface and the cytoplasm.




Fig.1. Map of Ionian Sea (Mediterranean Sea). ? , investigated area.




Fig. 2. Hematoxylin-eosin staining. Seminiferous tubules of *Aristaeomorpha foliacea* collected in March (c), April (d), July (a,e,f); (b), cross section of hemispermatophore. sg, spermatogonia; spl, primary spermatocytes; spII, secondary spermatocytes; sz, spermatozoa; w, wall of hemispermatophore; ? , somatic cell.




Fig. 3. Lectin histochemistry. (a) summer testis and (b) hemispermatochore stained with SNA; (c) spring testis and (d) summer hemispermatochore incubated with MAA; (e) late winter testis and (f) spring hemispermatochore stained with Con A; (g) spring testis and (h) hemispermatochore incubated with KOH-si-WGA procedure. ? , wholly stained spermatozoa.