



**SCIENTIFIC COUNCIL MEETING – SEPTEMBER 2004**

The Impact of Condition on Reproduction in Flemish Cap Cod

M. Joanne Morgan and George R. Lilly

Dept. of Fisheries and Oceans, PO Box 5667, St. John's  
NL A1C 5X1, Canada

**Abstract**

The impacts of fish condition on reproduction have not been examined previously for Flemish Cap cod. It is known that a substantial portion of female cod skipped spawning during the late-1970s and 1980s (Walsh *et al.*, MS 1986) and it was speculated that this might be as a result of poor condition. The purpose of this paper was to determine if there is a relationship between condition and maturity in Flemish Cap cod, both male and female, and to determine if there was a relationship between condition and the probability of spawning in female cod. There was no significant effect of relative body condition ( $K_r$ ) on the probability of being an adult for either male or female cod. For males there was also no significant effect of relative liver condition ( $LK_r$ ) on the probability of being an adult. For female cod there was a significant effect of  $LK_r$  on the probability of being an adult, after accounting for the effects of age and length. However,  $LK_r$  accounted for only a small portion of the deviance (1%) compared to the deviance accounted for by the combined effects of age and length (69%). There was no significant effect of  $K_r$  on the probability of spawning in adult females, however there was a significant impact of  $LK_r$  with females with higher  $LK_r$  having a greater probability of spawning.

Key words: Condition, adult, skip spawning, cod

**Introduction**

The probability of a fish being an adult generally increases with age and size (Korsbrekke, 1999; Morgan and Colbourne, 1999). In addition there can be effects of growth rate with fish maturing younger and/or smaller at increased growth rate (Jorgensen, 1990; Rjinsdorp, 1993; Morgan and Colbourne, 1999). Increased temperature also has been shown to lead to earlier maturation through increased growth (Alm, 1957; Sandstrom *et al.*, 1995). There can also be an additional relationship with condition so that fish in better condition are more likely to be adults, although few studies have directly examined this aspect. Marteinsdottir and Begg (2002) found an increased proportion of adult cod (*Gadus morhua*) at a given size or age among those that were in better condition. Bromley *et al.* (2000) found that adolescent turbot (*Scophthalmus maximus*) failed to mature under low food rations. Morgan (2004) found that female American plaice (*Hippoglossoides platessoides*) with low condition were less likely to be adults.

Condition can continue to exert an influence on reproduction once fish have become adults. Fish in poor condition can have lower fecundity and produce larvae that are smaller and less likely to survive (Kjesbu *et al.*, 1991; Marteinsdottir and Steinarsson, 1998; Marshall *et al.*, 1998). Fish in poor condition may also suffer higher post spawning mortality (Lambert and Dutil, 2000). For some fish, poor condition can mean that they will forego spawning altogether and skip one or more spawning seasons (Burton and Idler, 1987; Rideout *et al.*, 2000).

The impacts of fish condition on reproduction have not been examined previously for Flemish Cap cod. It is known that a substantial portion of female cod skipped spawning during the late-1970s and 1980s (Walsh *et al.*, MS 1986) and it was speculated that this might be as a result of poor condition. The purpose of this paper was to

determine if there is a relationship between condition and maturity in Flemish Cap cod, both male and female, and to determine if female cod that were in lower condition were more likely to skip a spawning season.

### Materials and Methods

Data for these analyses came from Canadian research vessel surveys of the Flemish Cap from 1978-1985. These surveys were conducted in January/February using a stratified random survey design. The *Gadus Atlantica* conducted the surveys using an Engel 145 bottom trawl with a small mesh liner in the cod end. All cod were measured (3 cm groupings) and most cod were examined for sex and maturity. Fish were classed as adult or juvenile based on macroscopic examination of their gonads using the maturity classification of Tempelman *et al.* (1978). Otoliths were extracted from a length-stratified sample (up to 25-30 specimens per 3 cm grouping) in 1978-1982 and from virtually all fish caught in 1983-1985. A length-stratified sample of cod was frozen round at sea. In 1978-1982 up to 5-6 cod per 3 cm length-group were chosen haphazardly from the survey area as a whole, and in 1983-1985 up to 2 cod per 3 cm length-group were chosen from each of 4 subareas. These frozen cod were thawed in the laboratory on land for detailed observations, among which were sex, maturity, weights of whole fish, both round and gutted, and weights of various organs. Otoliths were removed for ageing.

Indices of condition were used to examine the effect of both liver and body weight. Only fish greater than 20 cm were used in the calculations because of variability in the weight measurements below this size. 1,309 fish were used in the analyses. For gutted body weight the commonly used index of Fulton's  $K$  ( $K = W/L^3$ , where  $W$  is gutted body weight and  $L$  is length) showed an increasing trend with body length, so an alternative index, relative  $K$  ( $K_r$ ) was used.  $K_r = W/\hat{W}$ , where  $\hat{W}$  is the predicted body weight from a length/weight relationship. The form of the length/weight relationship was  $\log(W) = \text{intercept} + \log(L)$ . Separate regressions were carried out for each sex. Data for all years combined were used in the regression. To calculate  $K_r$  for an individual fish, its gutted body weight was divided by the gutted body weight predicted by the length/weight regression for a fish of that length and sex. The use of this type of condition index removes the problems associated with systematic change in the index across length. However, the indices produced are not comparable between sexes.

For liver weight only fish greater than or equal to 30 cm in length were used because data below that length were highly variable. There were 1,184 fish with liver weight data that met this criterion. There was a pattern in the residuals of the linear regression of log liver weight against log length, especially for males. A model of the form  $\log(LW) = \text{intercept} + \log(L) + (\log(L))^2$  was fit to the data, where  $LW$  is liver weight. This model resulted in a significant fit for both males and females and minimized patterns in the residuals. The results of these models were used to produce an index of liver condition,  $LK_r = LW/L\hat{W}$ , where  $L\hat{W}$  is the predicted liver weight from the length/liver weight relationship.

To determine if there was a significant effect of  $LK_r$  or  $K_r$  on the probability of being an adult, generalized linear models were used (McCullagh and Nelder, 1983). In these models fish were classed as either adult (1) or juvenile (0). All models had a logit link function and a binomial error structure. All parameters were modelled as continuous variables. Since age and length are already known to affect the probability of being an adult in this species (Korsbrekke, 1999) the significance of  $LK_r$  or  $K_r$  was tested by determining if there was a significant decrease in the deviance with the addition of the index to a model already containing age and length effects.

Females which were skipping a spawning season (non-spawners) and which had weight information, occurred over a length range of 60-95 cm, therefore only fish in this length range were included in the analyses. Linear length/weight regressions, as described above, for gutted body weight were used to produce  $K_r$  and  $LK_r$  for spawning and non spawning females. There were only 24 non-spawning females with weight information in the data set. The effect of condition was tested by using generalized linear models with binomial distribution and logit link to determine whether there was a significant effect of condition on the probability of spawning. The effects of length and age were not removed since analyses showed that they did not affect the probability of spawning over this size range.

## Results

$K_r$  for males ranged from 0.7 to 1.3 and for females from 0.8 to 1.3. There was no significant effect of  $K_r$  on the probability of being an adult for either male ( $\chi^2 = 0.5$ ,  $df = 1$ , NS) or female ( $\chi^2 = 3.4$ ,  $df = 1$ , NS) cod.  $LK_r$  for males ranged from 0.2 to 2.7 and for females the range was from 0.2 to 3.4. For males there was also no significant effect of  $LK_r$  on the probability of being an adult ( $\chi^2 = 2.1$ ,  $df = 1$ , NS). For female cod there was a significant effect of  $LK_r$  on the probability of being an adult, after accounting for the effects of age and length ( $\chi^2 = 8.9$ ,  $df = 1$ ,  $p < 0.005$ ). However,  $LK_r$  accounted for only a small portion of the deviance (1%) compared to the deviance accounted for by the combined effects of age and length (69%). Parameter estimates from the fitted model and mean length-at-age for fish in the data set were used to estimate proportion mature at age for females with  $LK_r$  of 0.5, 1.0 and 1.5 (Fig. 1). This range of  $LK_r$  results in about a 0.5 year difference in age at 50% maturity from 5.5 years for fish with an  $LK_r$  of 0.5 to 5.9 years for fish with an  $LK_r$  of 1.5.

There was no significant effect of  $K_r$  on the probability of spawning ( $\chi^2 = 0.6$ ,  $df = 1$ , NS). However, there was a significant effect of  $LK_r$  on the probability of spawning ( $\chi^2 = 17.7$ ,  $df = 1$ ,  $p < 0.0001$ ). Model estimates of the probability of spawning increased from less than 0.5 to 1.0 over the range of  $LK_r$  observed in the data (Fig. 2).

## Discussion

There was a significant positive effect of liver condition on the probability of female cod on Flemish Cap being adult. This was in addition to the positive effects of age and size. Since many fish species store energy in their liver, indices of liver condition should be good indicators of overall fish condition (Lambert and Dutil, 1997). Previous studies have also shown a positive effect of condition on the probability of being adult (Ajiad *et al.*, 1999; Bromley *et al.*, 2000; Marteinsdottir and Begg, 2002; Morgan, 2004). Fish make tradeoffs in energy allocation between growth and reproduction (Calow, 1985) and those in better condition may have more surplus energy to devote to reproduction and be able to mature at a smaller size and younger age.

The effect of liver condition on the probability of being adult was small compared to the combined effect of length and age. Marteinsdottir and Begg (2002) for cod and Morgan (2004) for American plaice (*Hippoglossoides platessoides*) also found that condition explained much less of the variability in the probability of being adult than length and/or age. It would seem that the main determinants of maturation are size and age with condition playing a lesser role.

In this study there was no effect of relative gutted body weight on the probability of being adult. This is in contrast to previous studies on cod and American plaice (Marteinsdottir and Begg, 2002; Morgan, 2004). However, those other studies used whole body weight in their calculations of indices of body condition. Also, both studies indicated geographic variation in the effect of body condition on the probability of being adult. Marteinsdottir and Begg (2002) attributed this to differing environmental conditions in the two areas they studied, speculating that the influence of condition might be more pronounced in areas where the environment is less conducive to growth.

There was no significant impact of either gutted body condition or liver condition on the probability of being adult in males. As with many fish species (Rijnsdorp and Ibelings, 1989; Morgan and Colbourne, 1999), male cod mature at a younger age and smaller size than females (Olsen *et al.*, 2004). Related to this there can be differences in energy allocation between males and females when they are adults (Rijnsdorp and Ibelings, 1989; Bromley *et al.*, 2000). The differences in the effect of condition on maturation seen here may be a result of sex specific differences in energy allocation. However, male Icelandic cod did exhibit a significant impact of condition on the probability of being adult (Marteinsdottir and Begg, 2002). It is not known whether this is the result of differences in methodology between the two studies or different responses in the two populations.

Liver condition also affected the probability that adult females would spawn. Adult females were less likely to spawn when they had lower relative liver condition. Lower condition in skip spawning females was also observed for cod in the inshore of Newfoundland (Rideout *et al.*, 2000). Winter flounder (*Pseudopleuronectes americanus*) skipped spawning when kept under low ration levels during a critical period (Burton, 1994). The condition of the fish in this study was measured close to the spawning period and it is not known what their condition level was at any other point. The assumption is made that the fish in poor condition and failing to spawn were also in poor condition during some critical period for gametogenesis. There were fish in similar condition to those that skipped

spawning, that did not fail to spawn. It may be that individual fish make different 'decisions' regarding spawning or perhaps their condition during gametogenesis was higher than those that skipped spawning. It is probable that those cod that skipped a spawning season would spawn in subsequent years if their condition improved (Burton, 1991).

For female cod on Flemish Cap it would appear that condition can have a substantial impact on reproduction. It plays a major role in determining if adult females will spawn in a given spawning season. It also has an influence on the probability of females being adults. Together these two effects could have important implications for the reproductive potential of the stock.

### Acknowledgements

We thank all the scientific staff and ship's crew involved in the collection of these data. Eugene Murphy and Dave Downton helped with the compilation of data.

### References

- AJIAD, A., T. JAKOBSEN, and O. NAKKEN. 1999. Sexual difference in maturation of northeast Arctic cod. *J. Northw. Atl. Fish. Sci.*, **25**: 1-15.
- ALM, G. 1957. Connections between maturity, size and age in fishes. *Reports from the Institute of Freshwater Research, Drottningholm*, **40**: 5-145.
- BROMLEY, P.J., C. RAVIER, and P.R. WHITTHAMES. 2000. The influence of feeding regime on sexual maturation, fecundity and atresia in first-time spawning turbot. *J. Fish Biol.*, **56**: 264-278.
- BURTON, M.P.M. 1991. Induction and reversal of the non-reproductive state in winter flounder, *Pseudopleuronectes americanus* Walbaum, by manipulating food availability. *J. Fish Biol.*, **39**: 909-910.
- BURTON, M.P.M. 1994. A critical period for nutritional control of early gametogenesis in female winter flounder, *Pleuronectes americanus* (Pisces: Teleostei). *J. Zool. Lond.*, **233**: 405-415.
- BURTON, M.P., and D.R. IDLER. 1987. An experimental investigation of the non-reproductive, post-mature state in winter flounder. *J. Fish Biol.*, **30**: 643-650.
- CALOW, P. 1985. Adaptive aspects of energy allocation. IN *Fish Energetics: New Perspectives*. pp. 13-31. Ed. P. Tytler and P. Calow. The Johns Hopkins University Press, Baltimore.
- JORGENSEN, T. 1990. Long-term changes in age at sexual maturity of northeast Arctic cod (*Gadus morhua* L.). *J. du Conseil Perm. Inter. pour l'Exploration de la Mer*, **46**: 235-248.
- KJESBU, O.S., J. KLUNGSOYR, H. KRYVI, P.R. WITTHAMES, and M. GREER WALKER. 1991. Fecundity, atresia, and egg size of captive Atlantic cod (*Gadus morhua*) in relation to proximate body composition. *Can. J. Fish. Aquat. Sci.*, **48**: 2333-2343.
- KORSBREKKE, K. 1999. Variations in maturity of haddock in the Barents Sea in relation to year-class strength, age, size, sex and area. *J. North. Atl. Fish. Sci.*, **25**: 37-45.
- LAMBERT, Y., and J-D., DUTIL. 2000. Energetic consequences of reproduction in Atlantic cod (*Gadus morhua*) in relation to spawning level of somatic energy reserves. *Can. J. Fish. Aquat. Sci.*, **57**: 815-825.
- MARSHALL, C.T., O.S., KJESBU, N.A. YARAGINA, P. SOLEMDAL, and O. ULLTANG. 1998. Is spawner biomass a sensitive measure of the reproductive and recruitment potential of northeast Arctic cod? *Can. J. Fish. Aquat. Sci.*, **55**: 1766-1783.

- MARTEINSDOTTIR, G., and G.A. BEGG. 2002. Essential relationships incorporating the influence of age, size and condition on variables required for estimation of reproductive potential in Atlantic cod *Gadus morhua*. *Mar. Ecol. Prog. Ser.*, **235**: 235-256.
- MARTEINSDOTTIR, G., and A. STEINARSSON. 1998. Maternal influence on the size and viability of Iceland cod *Gadus morhua* eggs and larvae. *J. Fish Biol.*, **52**: 1241-1258.
- MCCULLAGH, P., and J.A. NELDER. 1983. Generalized linear models. Chapman and Hall, London. 261pp.
- MORGAN, M.J. 2004. The relationship between fish condition and the probability of being mature in American plaice (*Hippoglossoides platessoides*). *ICES J. Mar. Sci.*, **61**: 64-70.
- MORGAN, M.J., and E.B. COLBOURNE. 1999. Variation in maturity-at-age and size in three populations of American plaice. *ICES J. Mar. Sci.*, **56**: 673-688.
- OLSEN, E.M., M. HEINO, G.R. LILLY, M.J. MORGAN, J. BRATTEY, B. ERNANDE, and U. DIECKMAN. 2004. Maturation trends indicative of rapid evolution preceded the collapse of northern cod. *Nature*, **428**: 932-935.
- RIDEOUT, R.M., M.P.M. BURTON, and G.A. ROSE. 2000. Observations on mass atresia and skipped spawning in northern Atlantic cod, from Smith Sound, Newfoundland. *J. Fish Biol.*, **57**: 1429-1440.
- RIJNSDORP, A.D. 1993. Fisheries as a large-scale experiment on life-history evolution: disentangling phenotypic and genetic effects in changes in maturation and reproduction of North Sea plaice, *Pleuronectes platessa* L. *Oecologia*, **96**: 391-401.
- RIJNSDORP, A.D. and B. IBELINGS. 1989. Sexual dimorphism in the energetics of reproduction and growth of North Sea plaice, *Pleuronectes platessa* L. *J. Fish. Biol.*, **35**: 401-415.
- SANDSTROM, O., E. NEUMAN, and G. THORESSON. 1995. Effects of temperature on life history variables in perch. *J. Fish Biol.*, **47**: 652-670.
- TEMPLEMAN, W., V.M. HODDER, and R. WELLS. 1978. Sexual maturity and spawning in haddock, *Melanogrammus aeglefinus*, of the southern Grand Bank. *ICNAF Res. Bull.*, **13**: 53-65.
- WALSH, S.J., R. WELLS, and S. BRENNAN. MS 1986. Histological and visual observations on oogenesis and sexual maturity of Flemish Cap female cod. *NAFO SCR Doc. No. 11*, Ser. No. N1238, 11p.




Fig. 1. Model estimates of proportion mature at age for female cod from Flemish Cap with different relative liver condition. Lengths-at-age used in the calculation were the means for females in the data set.




Fig. 2. Model estimates of the probability of spawning for female cod on Flemish Cap at various levels of relative liver condition.