

Northwest Atlantic



Fisheries Organization

Serial No. N5054

NAFO SCR Doc. 04/ 84

SCIENTIFIC COUNCIL MEETING – OCTOBER/NOVEMBER 2004

An Update of the Icelandic Shrimp Fishery (*Pandalus borealis* Kr.) at Flemish Cap in 1993-2004

by

U. Skúladóttir
Marine Research Institute, Skúlagata 4,
P.O. Box 1390, 121 Reykjavík, Iceland

Abstract

One Icelandic vessel went fishing for shrimp in the waters at Flemish Cap in 2004 as compared to three in 2003. In this paper there is logbook information on the Icelandic fishery for the years 1993 through 2003. The catch rate of Icelandic vessels in January-September which was very high in the years 2001-2003 has now decreased, or from about 290 kg/hour to 250 kg/hr in 2004.

The biological samples show that the 1999 year-class, five year olds in 2004 are still prominent. The 2001 year-class is above average in all months. A new year-class the 2002 year-class appears to be quite strong as one year old in 2003 and two year old in 2004. So recruitment appears to be good.

Introduction

The Spanish investigators (EU) have been measuring the biomass index of northern shrimp at the Flemish Cap since 1988 in their annual bottom trawl survey at Flemish Cap. In 1993 the fishery was initiated by Canada, followed closely by Faroe Islands and Iceland.

The fishery was some 24-33 thousand tons in the years 1993-1995 to increase in 1996 to 48 thousand tons. Since then the fishery decreased to some 25 thousand tons in 1997. The total catch of all countries has since increased to about 62 thousand tons in 2003. Iceland has been catching a fair deal of the catch in some previous years. In later years however the catch has decreased substantially due to low prizes in shrimp.

In this paper all the information from the Icelandic side is gathered. From the logbooks come effort, catch and size of trawl. From this CPUE is calculated. From the biological samples taken by Icelandic observers comes various information on length and sex distribution of shrimp.

Materials and Methods

The logbook data include catch and effort. Sometimes information on landings as obtained from the Fisheries Directorate in Iceland exceeds the logbook information. The effort is then raised by dividing the nominal catch of each month/half year with the calculated CPUE from the logbooks. The overall CPUE of the January-July was then obtained by summing nominal catch of all months and corresponding effort. Nominal catch for the whole period was then divided by "nominal effort" to get the CPUE for the period January-July. When twin trawls were used the effort was always multiplied by 1.9 for those but the catch was kept the same. The same method was applied to the period January-September.

For calculation of standardized CPUE to the standard size of trawl of 3000 meshes, the catch and effort of a period like January to July was calculated in the manner described above. At the same time the average size of

trawl (no. of standard meshes (40 mm) in circumference of the belly), be it single or double was calculated. The CPUE for trawl size 3000 meshes was then considered to be proportional to the mean size of trawl in the same period.

Icelandic observers have sampled shrimp onboard Icelandic vessels since 1996 at Flemish Cap. The shrimp was measured fresh to the nearest 0.5 mm using Vernier calipers. Observers then sorted each length class into males and females using the method of Rasmussen (1953) and the females further into primiparous and multiparous using the sternal spine criterion of McCrary (1971).

The deviations from an overall mean length frequency distribution (lfd) are calculated using data from Canada, Faroe Islands and Iceland for the years 1993-1995. From 1996-2003 there are only Icelandic data. The basic unit is a promille length frequency distribution for each month where all the samples of that month are compiled. Then a mean overall promille lfd for say April months for all the years 1993-2003 is calculated. From this the overall mean lfd of April is subtracted from the lfd of April in 1993 and so on, every year (Fig. 8). What is unusual about each year appears as a deviation high or low. The positive modes are representative of stronger than usual year-classes. As each year-class is supposed to grow from one year to another, the positive bump one year moves to the right in the following year and so on.

Catch and Effort Data

In 2003 the fishery was carried out since January (Table 1). The catch in 2004 so far is 2 480 tons (Table 2). Iceland increased the total allowable catch (TAC) for Icelandic vessels from 6 800 tons in 1999, to about 10 000 tons for years 2000 to 2002 and to 13 500 for year 2003. In spite of this high TAC the total catch was only 5 300 tons in year 2001, 5 700 tons in 2002 and 4 700 in 2003.

The distribution of effort around the Cap is shown by years in Fig. 1-3 for the years 1993 through 2004. There appears to be a lack of tows in the south east of Flemish Cap area in the last two years but in other years the distribution of tows is traditional.

The mean CPUE for the year 1997 was the lowest ever for Iceland or 203 kg per trawling hour for the period January through September (Table 2). In 1998 the mean CPUE for the same period was much higher or 266 kg and decreased slightly in 1999 and 2000 to increase in 2001 and 2002 to 294 kg/hour. In 2003 the CPUE was 291 kg/hour. In 2004 the CPUE is again lower and has dropped to the level of the years 1998 to 2000.

The average size of gear used was about 3 000 meshes in most years, but increased to about 3 500 meshes in the years 1999 to 2001 and 4 460 meshes in 2004. The trawl size in year 2004 is by far the largest so the unstandardized CPUE (no correction for size of trawl) of 2004 gives an impression of the shrimp stock being quite large. Therefore it makes more sense to look at CPUE at a standard trawl size. At the same time the use of twin trawls has increased from little less than 60% in 1995-1997 to about 93%- 99% in the years 2003-2004.

Length Frequencies and Age Groups

The length frequency distributions of Icelandic samples from 2003 through 2004 are shown by months in Fig. 4 and 5. Two year olds are seen in May year 2001 about 15-16 mm CL and get more prominent in the latter part of the year, namely September to December (Skuladottir, 2002). This 1999 year-class is also very prominent in years 2002 and 2003 (Fig. 4 and 5).

The deviation method (Sund, 1939; Skuladottir, 1981) is very useful in detecting year-classes and can be of great aid in assessing age when it comes to applying the modal analysis. The major drawback of the modal analysis is the fact that it does not tell you how many components there should be in an lfd and sometimes there is e.g. no difference in fitting 3 components when there should indeed be 4. From the Fig. 6 to 16 it is possible to study the deviations as positive peaks and occasionally as a peak that is at the mean line like in Fig. 16, the 3 year olds of December 1998.

The afore-mentioned 1999 year-class is first seen as a positive peak in August 2001 as a 2 year old (Fig. 12). It can be seen in September as well and then again in November and December. In year 2002 the most prominent peak is that of the 1999 year-class as three year olds in all months from March and onwards (no samples in February). In 2003 this same year-class can be seen all year, although sometimes like in April and June 2003 it appears to be

mixed with the 1997 and 1998 year-classes. In 2004 one can still see the positive deviation of the 1999 year-class at around 22.4 mm in February. In June the 5 year-old can be guessed at the length of 24 mm. A drawback of the method as in all length based age assessments is when growth slows down there is a fusion with the adjacent year-classes which then form a single peak that may be broader than it should be. The 2000 year-class is weaker than the 1999 year-class and appears to fuse together with it in 2003. This is also shown by the deviation method. When later using the modal analysis, this information has to be used as there are 4 components in the male lfd where most people would think there are only 3 (Fig. 5), (Skúladóttir, 2003).

From the deviations in Fig. 6-16 it has been attempted to follow the various year-classes and the mean length of each is assessed by eye from the deviations. The results are listed in Table 3 where the 8 year-classes, namely 1993, 1994, 1995, 1996, 1997, 1999, 2001 and 2002 are followed. In Table 3, the year stated is the one where the appropriate deviation is assessed. So e.g. the 1993 year-class which was quite strong can be seen first as 2 year old in March 1995 at the CL 14 mm. In the last two years one year olds are detected for the first time as deviations. These are the 2001 and 2002 year-classes at 12.5 to 13.8 mm in the months September through December (Fig. 13 to 16, Table 3). Although the mean lengths at age by the deviation method appear to be some guesswork these can be used as inputs in the modal analysis.

By-catch

The by-catch was about 0.5 % in the years 2003 and 2004. The percent of by-catch has decreased from the high 2.1% and 1.8% in 1996 and 1997 to 0.8-1.0% in the years 1998-2001 (Skúladóttir, 2003c). By-catch was only 0.3% in 2002. Most of this was redfish or 0.6-0.8 % in the years 1999 to 2001. Other species were wolffish, Greenland halibut and American plaice. Cod was seen for the first time in April 1999, but has not been seen since.

References

- McCrary, J.A. 1971. Sternal spines as a characteristic for differentiating between females of some Pandalidae. *J. Fish. Res. Board Can.*, **28**: 98-100.
- Rasmussen, B. 1953. On the geographical variation in growth and sexual development of the deep sea prawn (*Pandalus borealis* Kr.). *Norweg. Fish. and Mar. Invest. Rep.*, **10** (3): 1-160.
- Skúladóttir, U. 1981. The deviation method. A simple method for detecting year-classes of a population of *Pandalus borealis* from length distributions. T. Frady (Ed.), Proc., Int. Pandalid Shrimp Symp., Kodiak Alaska, Feb. 13-15. pp. 283-306. *Univ. Alaska, Sea Grant Rep.*, 81-3.
- Skúladóttir, U. 1998. The Bycatch in the shrimp fishery of Iceland at Flemish Cap in 1997 and 1998. *NAFO SCR Doc.*, No. 29, Serial No. N3016, 5 p.
- Skúladóttir, U. 2003. An update of the Icelandic shrimp fishery (*Pandalus borealis* Kr.) at the Flemish Cap in 1993-2002. *NAFO SCR Doc.*, No. 83, Serial No. N4925, 20 p.
- Skúladóttir, U. 2003b. An update of the Assessment of the International Fishery for shrimp fishery (*Pandalus borealis*) in Division 3M (Flemish Cap) in 1993-2002. *NAFO SCR Doc.*, No. 91, Serial No. N4933, 14 p.
- Skúladóttir, U. 2003c. The Bycatch in the shrimp fishery of Iceland at Flemish Cap in 1993-2003. *NAFO SCR Doc.*, No. 85, Serial No. N4926, 6 p.
- Sund, O. 1939. The renewal of fish population studied by means of measurement of commercial catches. Example: the arcto Norwegian cod stock. *Rapp. P. - v. Réun. Cons. Int. Explor. Mer.*, **65**: 10-17.

Table 1. Catch (tons) effort (trawling hours *1.9 when double trawl) and unstandardized CPUE (kg/hr) of Icelandic vessels at Flemish Cap.

Year	January - July				August - December				Year	January - July				August - December			
	Month	CPUE	Effort	Catch	Month	CPUE	Effort	Catch		Month	CPUE	Effort	Catch	Month	CPUE	Effort	Catch
1993					Aug	320.4	1334	427.4	2001 *	Jan	285.9	538	153.7	Aug	292.6	2094	612.9
					Sep	349.8	1034	361.7		Feb	299.9	1593	477.6	Sep	277.3	1160	321.6
					Oct	231.7	334	77.4		Mar	303.6	2174	660.0	Oct	267.5	1563	418.1
	Jun	380.2	1767	671.8	Nov	306.8	588	180.4		Apr	239.6	45	10.8	Nov	253.4	1210	306.6
	Jul	342.4	1097	375.6	Dec	236.5	537	127.0		May	271.1	917	248.7	Dec	500.8	404	202.5
	Subtotal	365.7	2864	1047.4	Subtotal	306.7	3827	1173.9	Jun	282.9	2777	785.6					
	Total	365.7	2918	1067.0	Total	306.7	3834	1176.0	Jul	296.5	2992	887.2					
									Subtotal	292.1	11036	3223.6	Subtotal	289.5	6431	1861.7	
									Total	292.1	11036	3223.6	Total	289.5	7178	2077.8	
1994	Jan	228.5	144	32.9	Aug	175.3	1657	290.4	2002 *	Jan	292.6	372	108.9	Aug	311.7	1739	542.0
	Feb	371.8	510	189.6	Sep	126.9	476	60.4		Feb	343.4	705	242.0	Sep	313.2	1054	330.0
	Mar	295.5	531	156.9	Oct	125.4	492	61.7		Mar	264.6	1786	472.4	Oct	234.7	923	216.7
	Jun	256.4	1297	332.5	Nov	115.5	181	20.9		Apr	305.7	2056	628.4	Nov	312.9	559	174.9
	Jul	212.9	2653	564.8	Dec	75.0	8	0.6		May	330.8	2439	806.6	Dec	359.9	437	157.1
	Subtotal	248.6	5135	1276.7	Subtotal	154.2	2814	434	Jun	346.0	2113	731.1					
	Total	248.6	6693	1664.0	Total	154.2	4123.7	636	Jul	444.6	1241	551.7					
									Subtotal	330.6	10710	3541.1	Subtotal	301.6	4711	1420.7	
									Total	330.6	10711	3541.1	Total	301.6	7296	2200.3	
1995	Feb	280.0	65	18.2	Aug	178.0	4869	866.9	2003 *	Jan	384.5	162	62.1	Aug	391.3	943	369.0
	Mar	246.8	711	175.5	Sep	134.1	2928	392.5		Feb	422.1	715	301.8	Sep	293.5	1610	472.4
	Apr	149.9	1487	222.9	Oct	166.3	2088	347.2		Mar	506.6	1323	670.1	Oct	352.2	941	331.6
	May	260.1	2617	680.7	Nov	144.4	1074	155.1		Apr	349.5	2028	708.9	Nov	333.4	727	242.4
	June	248.9	3733	929.2	Dec	174.5	740	129.1		May	293.5	1827	536.2	Dec	659.6	310	204.7
	Subtotal	241.5	15238	3679.5	Subtotal	161.6	11699	1890.8	Jun	317.1	1211	383.9					
	Total	241.5	16932	4088.5	Total	161.6	21868	3534.4	Jul	371.1	1016	377.1					
									Subtotal	367.1	8282	3040.2	Subtotal	357.5	4532	1620.1	
									Total	367.1	8282	3040.2	Total	357.5	4629	1655.0	
1996	Jan	207.2	1755	363.7	Aug	165.4	8156	1349.4	2004 *	Jan	251.4	403	101.2	Aug	417.0	763	318.2
	Feb	251.7	1326	333.7	Sep	167.1	8089	1351.7		Feb	293.3	892	261.5	Sep	379.1	282	106.9
	Mar	261.8	4604	1205.1	Oct	129.7	5482	711.2		Mar	268.0	974	261.0	Oct			
	Apr	211.2	10754	2271.2	Nov	137.9	1456	200.8		Apr	280.3	1044	292.6	Nov			
	May	189.1	12749	2410.2	Dec	158.1	253	40.0		May	315.1	1089	343.0	Dec			
	Subtotal	214.2	57084	12226.9	Subtotal	155.9	23436	3653.1	Jun	403.5	1015	409.5					
	Total	214.2	64760	13871.0	Total	155.9	43689	6810.0	Jul	386.9	967	374.3					
									Subtotal	320.1	6383	2043.1	Subtotal	406.8	1045	425.1	
									Total	320.1	6383	2043.1	Total	406.8	1075	437.2	
1997	Jan	175.8	413	72.6	Aug	206.7	4252	879.0	1998 *	Feb	217.2	297	64.5	Aug	256.4	3184	816.3
	Feb	214.7	621	133.3	Sep	202.4	3476	703.6		Mar	206.8	812	167.9	Sep	184.5	5028	927.5
	Apr	135.0	514	69.4	Oct	222.0	2519	559.1		Apr	229.5	880	202.0	Oct	196.3	3612	708.9
	May	141.4	3736	528.2	Nov	192.5	1039	200.0		May	261.4	2820	737.2	Nov	204.6	1761	360.3
	Jun	167.7	5386	903.2	Dec	176.9	429	75.9		Jun	330.7	3537	1169.7	Dec	222.5	644	143.3
	Subtotal	177.3	16472	2920.4	Subtotal	206.4	11715	2417.6	Jul	285.3	4117	1174.7					
	Total	177.3	19478	3453.3	Total	206.4	14681	3029.6	Subtotal	282.1	12463	3516.0					
									Total	282.1	12657	3570.8					
1999 *	Feb	350.5	382	133.9	Aug	250.8	3642	913.4	1999 *	Feb	289.4	1851	535.7	Aug	250.8	3642	913.4
	Mar	289.4	1851	535.7	Sep	235.5	1371	322.9		Mar	253.0	3483	881.2	Sep	255.6	2150	549.6
	Apr	253.0	3483	881.2	Oct	255.6	2150	549.6		Apr	249.5	5941	1482.3	Nov	256.2	2173	556.8
	May	249.5	5941	1482.3	Nov	256.2	2173	556.8		Jun	285.8	5993	1712.7	Dec	230.6	989	228.1
	Jun	285.8	5993	1712.7	Dec	230.6	989	228.1		Jul	280.4	5224	1464.6				
	Subtotal	271.5	22874	6210.4	Subtotal	249.0	10325	2570.8	Subtotal	271.5	22874	6210.4	Subtotal	249.0	10325	2570.8	
	Total	271.5	24009	6518.6	Total	249.0	10837	2698.4	Total	271.5	24009	6518.6	Total	249.0	10837	2698.4	
2000 *	Jan	263.8	1050	277.0	Aug	244.9	2357	577.1	2000 *	Jan	263.8	1050	277.0	Aug	244.9	2357	577.1
	Feb	280.5	2206	618.8	Sep	239.0	2134	510.2		Feb	280.5	2206	618.8	Sep	239.0	2134	510.2
	Mar	306.3	3297	1009.8	Oct	274.8	1787	491.1		Mar	306.3	3297	1009.8	Oct	274.8	1787	491.1
	Apr	280.7	4378	1229.0	Nov	256.1	2984	764.3		Apr	280.7	4378	1229.0	Nov	256.1	2984	764.3
	May	231.9	4943	1146.6	Dec	267.5	798	213.5		May	231.9	4943	1146.6	Dec	267.5	798	213.5
	Subtotal	272.7	22618	6167.2	Subtotal	254.1	10060	2556.2	Subtotal	272.7	22618	6167.2	Subtotal	254.1	10060	2556.2	
	Total	272.7	22618	6167.2	Total	254.1	11051	2807.8	Total	272.7	22618	6167.2	Total	254.1	11051	2807.8	

Table 2. Nominal catch for the whole year and some averages calculated from the Icelandic logbooks to show trends in CPUEs and size of trawl. In calculations of CPUE the effort of twin trawls is multiplied by 1.9.

Year	Nominal Catch Tons	Twin trawls % of catch	Mean trawl size No. of meshes January-July	Unstandardized CPUE January-July	CPUE at size 3000 trawl January-July	Mean trawl size No. of meshes January-Sept	Unstandardized CPUE January-Sept	CPUE at size 3000 trawl January-Sept.
1993	2 243	43.4	3063	373	363	3102	356	344
1994	2 300	54.4	2994	238	240	2951	216	219
1995	7623	38.2	2779	254	283	2733	228	251
1996	20681	42.9	2803	206	218	2813	198	211
1997	6483	53.4	2780	188	192	2921	198	203
1998	6572	74.8	3016	288	294	2974	264	266
1999	9217	70.6	3441	280	252	3402	276	243
2000	8978	81.4	3528	287	245	3528	282	240
2001	5301	63.0	3571	328	290	3518	325	289
2002	5741	73.6	3713	370	305	3713	363	294
2003	4695	92.6	3949	367	302	4004	358	291
2004	2480	98.9	4460	320	227	4460	332	250
Mean 93-2004	8366	63	3240	289	271	3242	279	259

Table 3. Length at age as assessed by eye from the deviations in length frequency distributions each month in figures 6 to 16.

Months	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	Mean
1 years											
Sept.								12.5	12.2		12.35
Oct.									12.5		12.5
Nov.								13.8			13.8
Dec.								13.0	13		13.0
2 years											
Jan.											
Feb.									13.5	13.5	13.5
Mar.	14.0						14.0			13.5	13.8
Apr.	14.5			13.0	12.7				14.6	12.5	13.5
May	15.5										15.5
Jun.	15.7										15.7
Jul.	16.2			14.8					17.2	16.0	16.1
Aug.		15.5	15.5				16.5		16.7	16.0	16.0
Sept.	16.5						16.5	16.0		15.5	16.1
Oct.	17.5	16.0	16.0	17.5	15.5				15.8		16.4
Nov.	18.3		16.5	16.0	15.5			17.2			16.7
Dec.	18.2			16.0			16.2		16.7		16.8
3 years											
Feb.			18.5	17.5		15.7				18.0	17.4
Mar.		19.0		17.8	16.0	16.7		16.7		17.7	17.3
Apr.		20.0	18.5	18.3	16.2	16.8		17.5		17.5	17.8
May		20.0	18.7	18.5	17.2	17.6		18.5			18.4
Jun.		20.5	19.4	19.0	18.0	18.5		18.3		19.6	19.0
Jul.		20.7	19.0	19.2	18.3	18.7				20.0	19.3
Aug.		21.0	19.2	19.7	18.0	18.5		18.5		19.8	19.2
Sept.		20.8	19.5	19.5	18.5						19.6
Oct.		21.2	19.7	19.8	18.6			19.7			19.8
Nov.		22.0	20.3	20.5				20.3	20.5		20.7
Dec.		22.2	20.7	20.7	19.2	19.5		19.8			20.4
4 years											
Feb.			23.0	21.2	20.7		20.5			20.2	21.1
Mar.				22.0	22.0		21.5		20.7		21.6
Apr.			23.0	22.3	22.0						22.4
May			22.7	22.6	22.6		21.8		21.0		22.1
Jun.			23.3	22.7	23.0		21.8				22.7
Jul.			23.3	23.0	23.2	20.8	22.0		21.7		22.3
Aug.			23.5	22.7	23.5	20.8	21.7		21.8		22.3
Sept.			23.5	22.8	23.2		22.0		22.0		22.7
Oct.			23.8	22.7	23.0				22.0		22.9
Nov.				23.5		22.0			23.0		22.8
Dec.				23.7			21.8				22.8
5 years											
Feb.				25.2	23.7					22.4	23.8
Mar.				25.7	23.7					22.7	24.0
Apr.										22.7	22.7
May					24.5		24.0	23.7			24.1
Jun.			25.2		25.0	24.5	24.2	24.0		24.0	24.5
Jul.							24.8	24.3			24.6
Aug.				26.7			24.0	24.5	24.2	24.2	24.7
Sept.						24.5	24.2	24.0	24.0	24.0	24.1
Oct.				26.3			25.2	24.0			25.2
Nov.				27.0			25.5	25.0			25.8
Dec.				26.8	24.7			24.5			25.3
6 years											
Feb.					26.2						26.2
Mar.											
Apr.											
May											
Jun.											
Jul.								26.0			26.0
Aug.									26.2		26.2
Sept.											
Oct.								25.8			25.8
Nov.									26.3		26.3
Dec.											
7 years											
Feb.										27.2	27.2
Jul.										27.5	27.5




Figure 1. Shrimp in Div. 3 M: Towing positions in the Icelandic fleet on Flemish Cap in years 1993-1996.




Figure 2. Shrimp in Div. 3 M: Towing positions in the Icelandic fleet on Flemish Cap in years 1997-2000.




Figure 3. Shrimp in Div. 3 M: Towing positions in the Icelandic fleet on Flemish Cap in years 2001-2004.




Figure 4. The length frequency distribution of northern shrimp at Flemish Cap by months in 2003.




Figure 5. The length frequency distribution of northern shrimp at Flemish Cap by months in 2004.

February




Fig 6. The deviations of length frequencies of northern shrimp by years in February on the Flemish Cap from the mean length frequency of the years 1997-2004 in the same month.

March




Fig. 7. The deviations of length frequencies of northern shrimp by years in March on Flemish Cap from the mean length frequency distribution of the years 1994-2004 in the same month. 1994 and 1995 are data of Canada and other countries. Since 1996 data are solely from Iceland.

April





Fig 8. The deviations of length frequencies of northern shrimp by years in April on the Flemish Cap from the mean length frequency of the years 1993-2004 in the same month. 1993 through 1995 are data of Canada and other countries. Since 1996, data are solely from Iceland.

May



Carapace length mm

Fig 9. The deviations of length frequencies of northern shrimp by years in May on the Flemish Cap from the mean length frequency of the years 1993-2003 in the same month. 1993 through 1995 are data of Canada and other countries. Since 1996, data are solely from Iceland.

June





Fig 10. The deviations of length frequencies of northern shrimp by years in June on the Flemish Cap from the mean length frequency of the years 1993-2004 in the same month. 1993 through 1995 are data of Canada and other countries. Since 1996, data are solely from Iceland.

July



Carapace length mm

Fig 11. The deviations of length frequencies of northern shrimp by years in July on the Flemish Cap from the mean length frequency of the years 1993-2004 in the same month. 1993 through 1995 are data of Canada and other countries. Since 1996, data are solely from Iceland.

August




Fig 12. The deviations of length frequencies of northern shrimp by years in August on the Flemish Cap from the mean length frequency of the years 1996-2004 in the same month.




Fig 13. The deviations of length frequencies of northern shrimp by years in September on the Flemish Cap from the mean length frequency of the years 1993-2004 in the same month. 1993 through 1995 are data of Canada and other countries. Since 1996, data are solely from Iceland.




Fig 14. The deviations of length frequencies of northern shrimp by years in Oktober on the Flemish Cap from the mean length frequency of the years 1993-2003 in the same month.

November




Fig 15. The deviations of length frequencies of northern shrimp by years in November on the Flemish Ca from the mean length frequency of the years 1995-2003 in the same month.

December




Fig 16. The deviations of length frequencies of northern shrimp by years in December on the Flemish Cap from the mean length frequency of the years 1995-2003 in the same month.