

NOT TO BE CITED WITHOUT PRIOR
REFERENCE TO THE AUTHOR(S)

Northwest Atlantic



Fisheries Organization

Serial No. N5499

NAFO SCR Doc. 08/8

SCIENTIFIC COUNCIL MEETING – JUNE 2008

Atlantic cod and Yellowtail flounder indices from the Spanish Survey conducted in Divisions 3NO of the NAFO Regulatory Area

by

D. González-Troncoso and X. Paz

Instituto Español de Oceanografía,
P.O. Box 1552. Vigo, Spain.
e-mail: diana.gonzalez@vi.ieo.es

Abstract

Since 1995, Spain carries out a stratified random spring bottom trawl survey in Div. 3NO of the NAFO Regulatory Area. In 2001, the trawl vessel and gear were replaced; so, the time series indices were transformed. The transformed entire series of mean catches, biomass and length distribution for Atlantic cod (*Gadus morhua*) are presented for the period 1997-2000 and for Yellowtail flounder (*Limanda ferruginea*) for the period 1995-2000, and the both species no-transformed data for the years 2002-2007. For 2001, there are data from the two vessels. For Atlantic cod we can see a general decreasing in the biomass since the year 2002, although in 2006 the biomass is much higher because above all the catch of a simple town of almost 2 tons. In 2007 the biomass decreased, but the level is over the value in the period 2002-2005. For this species, an increase in the recruitment can be seen in 2004 and 2005, and in year 2007 the youngest length classes are much over the rest of the length classes. For Yellowtail flounder, there is no a clear trend since 1998; its indices are almost constant along this period.

Material and methods

The survey in Div. 3NO of NAFO Regulatory Area was initiated by Spain in 1995. Until 2001, the survey was carried out in Spring (May), on board the Spanish vessel C/V *Playa de Menduiña* (338 GT and 800 HP) using bottom trawl net type *Pedreira*. Since 2001, the R/V *Vizconde de Eza* replaced the C/V *Playa de Menduiña* as the research vessel for the survey, and *Campelen* net replaced *Pedreira* net as survey gear. The main specifications and geometry of these gears, as the rigging profile and the net plan, and a sheet with the resume of the main technical data of the survey are described in a previous paper (Walsh *et. al.*, 2001). Table 1 presents the number of valid tows, the depth strata covered and the dates of the survey series. The survey area was stratified following the standard stratification schemes (Bishop, 1994). Set number was allocated to strata proportionally to their size, with a minimum of two planned hauls per stratum and the trawl positions were chosen at random (Doubleday, 1981). Biomass indices were calculated by the swept area method (Cochran, 1997), assuming catchability factor of 1.

For Atlantic cod, the series are presented since 1997 because in years 1995 and 1996 the surveyed depth strata were only until 1000 meters, so they are not representative. As the strata where the Yellowtail flounder is presented were well surveyed, the series for this species are presented since 1995.

The catch from each haul was sorted by species and weighted. Random samples of each species catches were measured to total length to the nearest lower cm. Length distribution scaled from catches was estimated for the period 1997-2007 (Atlantic cod) and 1995-2007 (Yellowtail flounder) in two cm range. Data were grouping beginning with the pair number.

For each species, the haul mean catch, with its variance, and the stratified mean catches by stratum and year, with the annual variance, are presented, transformed until 2000 and no-transformed in the period 2002-2007. In the year 2001, there are data transformed from the former vessel with original data from the new vessel. Besides this, the biomass per stratum and year, with the annual variance, are presented, as the stratified mean catches per haul length distribution. To more information about the calculation of these indices, see González Troncoso *et al.*, 2004 and Paz *et al.*, 2004.

Results

Atlantic cod

Atlantic cod in Divisions 3NO has been under moratorium to directed fishing since 1994. According to the NAFO Scientific Council, the stock of Atlantic cod in Divisions 3NO declined dramatically during the mid-1980s, and the total biomass and the spawning biomass are currently at an extremely low level. Moreover, all recent year-classes have been weak (NAFO, 2007).

Mean Catches and Biomass

The Atlantic cod haul mean catches by stratum are presented in Table 2, included swept area, number of hauls and SD. Atlantic cod stratified mean catches per tow by stratum and year and their SD are presented in Table 3.

The entire time series (1997-2007) of biomass and their SD estimates for Atlantic cod are presented in Table 4. Estimated parameters a and b values of length-weight relationship are presented in Table 5.

We can see a great variation in the cod indices since 1997, but this is due to a few hauls in which the presence of cod was very high. For example, in 1998 and 2001, the C/V *Playa de Mendumía* made a more than seven tons cod catch in a single haul. Besides this, in 2001, the R/V *Vizconde de Eza* made two hauls with more than a ton of cod catches. But before year 2006, and apart from those hauls, the catches of cod were very poor. Since 2002, a decreasing in the biomass can be seen. In 2006, we can see a great increase in the biomass of this species. Although this increase is above all for a single catch of almost 2 tons, in general the catches of Atlantic cod in the survey of 2006 were over the mean. In 2007 we can see a decrease in the biomass over the 2006 biomass, but still remains greater than in the 2002-2005 period. The great value of the variance in some years is due to the tows with a large catch (Fig. 1 and 2).

Length Distribution

Table 6 and Figures 3 and 4 show the stratified mean catches per haul length distribution by year, besides the sampled size and its catch, for the period 1997-2007. The data have been grouped two by two, so we present the data every two cm. Except in 2001, 2006 and 2007, the modal values are very low. All lengths presence is very low, even it is very difficult to follow the modal values. In 2001 we have a good presence of individuals between 36 and 58 cm, probably due to the three hauls with great catches of this year, and in 2006 there is two modes in the length distribution, one around 30 cm and another one around 40 cm. There is no good recruitment until 2004, in which the individuals between 12 and 16 cm correspond to the greatest presence in the series, and in 2005 between 24 and 32, with a new mode between 12 and 16 cm, as in last year. In 2007 the youngest lengths dominate the length range with values of the lengths 12-16 that are between 2 and 4 times the abundance of the 48 cm length class, the following mode.

Yellowtail flounder

After a moratorium between 1994 and 1997, the Yellowtail flounder fishery is under TAC. According to the Report of NAFO Scientific Council Meeting, the stock size had a minimum in the mid 1990's, but since 1994 has steadily increased and now it is estimated to be at a level well above that of the mid-1980s (NAFO, 2006).

Mean Catches and Biomass

In Table 7 we present the haul mean catches by stratum for Yellowtail flounder, included swept area, number of hauls and SD. The stratified mean catches per tow by stratum and year and their SD are presented in Table 8 for this species.

The entire time series (1995-2007) of biomass by the swept area method and their SD estimates of Yellowtail flounder are presented in Table 9. The parameters a and b for the calculation of the length-weight relationship are presented in Table 10.

The Yellowtail flounder indices show no clear trend along the time (in the entire series). There was an increasing between 1995 and 1999 and since 2001 the indices are stabilised at a high level (Figures 5 and 6).

Length Distribution

The stratified mean catches per haul length distribution by sex and year, besides the sampled size and its catch, are presented in Table 11 and Figure 7 the period 1995-2007. The data have been grouped two by two, so we present the data every two cm. There is no presence of good recruitment last years. In Figure 8, we can see the evolution of a modal value since the beginning of the series, but, although there is a presence of juveniles in the lengths, this presence is very low.

References

- Bishop, C A.. 1994. Revisions and additions to stratification schemes used during research vessel surveys in NAFO subareas 2 and 3. NAFO SCR Doc. 94/43, Serial n° N2413, 23 pp.
- Cochran, W. G.. 1997. Sampling techniques. J. Wiley and Sons, N.Y., 428 pp.
- Doubleday, W. G.. 1981. Manual on groundfish surveys in the Northwest Atlantic. NAFO Sci. Coun. Studies, 2, 55.
- González Troncoso, D., X. Paz and C. González. 2004. Atlantic cod population indices obtained from the Spring surveys conducted by Spain in the NAFO Regulatory Area of Divisions 3NO, 1995-2003. NAFO SCR Doc. 04/12, Serial Number N4957, 21 pp.
- NAFO, 2006. Report of Scientific Council Meeting, 1-15 June 2006
- NAFO, 2007. Report of Scientific Council Meeting, 7-21 June 2007
- Paz, X., D. González Troncoso and E. Román. 2004. New time series for Yellowtail flounder from the comparative experience between the C/V *Playa de Mendumía* and the R/V *Vizconde de Eza* in the NAFO Regulatory Area of Divisions 3NO, 1995-2003. NAFO SCR Doc. 04/10, Serial Number N4955, 19 pp.
- Walsh, J.S., X. Paz and P. Durán. 2001. A preliminary investigation of the efficiency of Canadian and Spanish Survey bottom trawls on the Southern Bank. NAFO SCR Doc., 01/74, Serial n° N4453, 18 pp.

TABLE 1.- Spanish spring bottom trawl surveys on NAFO Div. 3NO: 1995-2007

Year	Vessel	Valid tows	Depth strata covered (m)	Dates
1995	C/V <i>Playa de Menduíña</i>	77	42-684	May 18-May 29
1996	C/V <i>Playa de Menduíña</i>	112	41-1135	May 07-May 24
1997	C/V <i>Playa de Menduíña</i>	128	42-1263	April 26-May 18
1998	C/V <i>Playa de Menduíña</i>	124	42-1390	May 06-May 26
1999	C/V <i>Playa de Menduíña</i>	114	41-1381	May 07-May 26
2000	C/V <i>Playa de Menduíña</i>	118	42-1401	May 07-May 28
2001 ^(*)	R/V <i>Vizconde de Eza</i>	83	36-1156	May 03-May 24
	C/V <i>Playa de Menduíña</i>	121	40-1500	May 05-May 23
2002	R/V <i>Vizconde de Eza</i>	125	38-1540	April 29-May 19
2003	R/V <i>Vizconde de Eza</i>	118	38-1666	May 11-June 02
2004	R/V <i>Vizconde de Eza</i>	120	43-1539	June 06-June 24
2005	R/V <i>Vizconde de Eza</i>	119	47-1485	June 10-June 29
2006	R/V <i>Vizconde de Eza</i>	120	45-1480	June 7-June 27
2007	R/V <i>Vizconde de Eza</i>	110	45-1374	May 29-June 19

(*) We took, for the calculation of the series, 83 hauls from the R/V *Vizconde de Eza* and 40 hauls from the C/V *Playa de Menduíña* (123 hauls in total)

TABLE 2.- Swept area, number of hauls and Atlantic cod mean catch (kg) and SD by stratum. Spanish Spring Surveys on NAFO Div. 3NO: 1997-2007. Swept area in square miles. n.s. means stratum not surveyed. 1997-2000 data are transformed C/V *Playa de Mendoña* data, and 2002-2007 data are original from R/V *Vizconde de Eza*. In 2001, there are data from the two vessels.

Stratum	1997				1998				1999				2000			
	Swept area	Tow number	A. cod Mean catch	A. cod SD	Swept area	Tow number	A. cod Mean catch	A. cod SD	Swept area	Tow number	A. cod Mean catch	A. cod SD	Swept area	Tow number	A. cod Mean catch	A. cod SD
353	0.0480	4	0.00	0.000	0.0465	4	0.00	0.004	0.0360	3	6.26	8.593	0.0356	3	8.59	9.984
354	0.0233	2	0.00	0.000	0.0356	3	17.67	29.046	0.0218	2	4.92	3.192	0.0356	3	18.44	27.099
355	0.0233	2	4.29	5.711	0.0221	2	27.05	3.662	0.0229	2	6.39	2.549	0.0233	2	94.83	76.209
356	0.0225	2	7.80	0.495	0.0221	2	6.23	0.247	0.0229	2	41.19	0.346	0.0225	2	16.34	17.172
357	0.0443	4	91.55	174.202	0.0240	2	7.45	0.742	0.0236	2	10.12	11.461	0.0124	1	9.15	-
358	0.0563	5	1.77	1.655	0.0236	3	4.46	4.030	0.0349	3	9.98	4.006	0.0341	3	184.88	194.829
359	0.0690	6	1.13	2.385	0.0698	6	0.39	0.858	0.0364	3	7.25	11.394	0.0469	4	18.26	17.367
360	0.3754	32	0.11	0.226	0.2561	25	0.22	0.700	0.2325	19	2.33	3.801	0.2396	20	2.16	3.561
374	0.0353	3	0.06	0.099	0.0353	3	0.00	0.000	0.0244	2	0.58	0.594	0.0240	2	0.00	0.000
375	0.0116	1	0.00	-	0.0345	3	0.78	0.403	0.0236	2	0.97	0.579	0.0244	2	0.00	0.000
376	0.1583	14	0.00	0.000	0.0930	10	0.20	0.187	0.1219	10	0.62	0.545	0.1200	10	0.90	1.852
377	0.0116	1	0.27	-	0.0229	2	1.89	2.375	0.0240	2	0.21	0.302	0.0229	2	0.02	0.027
378	0.0210	2	2.34	3.316	0.0120	2	3.46	0.940	0.0229	2	7.76	5.951	0.0233	2	10.65	11.169
379	0.0206	2	3.68	0.307	0.0356	3	8.30	5.847	0.0236	2	5.22	4.147	0.0225	2	41.12	54.683
380	0.0210	2	0.36	0.515	0.0113	2	2.33	1.361	0.0236	2	38.58	48.720	0.0236	2	8.21	3.236
381	0.0221	2	0.07	0.099	0.0229	2	0.21	0.187	0.0229	2	0.87	0.388	0.0236	2	1.74	0.730
382	0.0461	4	0.00	0.000	0.0229	3	0.32	0.336	0.0484	4	0.05	0.036	0.0499	4	0.71	0.561
721	0.0221	2	20.98	7.052	0.0203	2	0.61	0.866	0.0244	2	88.29	106.743	0.0236	2	28.34	17.122
722	0.0214	2	0.31	0.139	0.0101	2	0.00	0.000	0.0229	2	0.00	0.000	0.0218	2	0.90	1.277
723	0.0210	2	9.90	2.425	0.0233	2	4.39	3.736	0.0229	2	16.87	20.735	0.0248	2	22.02	12.010
724	0.0225	2	1.30	1.269	0.0206	2	1488.84	2101.820	0.0225	2	0.02	0.032	0.0233	2	0.70	0.341
725	0.0206	2	23.50	17.734	0.0086	1	30.86	-	0.0229	2	13.65	19.102	0.0210	2	4.34	3.857
726	n.s.	n.s.	n.s.	n.s.	0.0094	2	4.74	5.617	0.0225	2	0.81	0.492	0.0221	2	8.85	12.221
727	0.0094	1	0.12	-	0.0233	2	2.66	2.821	0.0236	2	9.20	4.701	0.0210	2	9.16	10.803
728	0.0214	2	1.17	0.569	0.0206	2	1.54	2.177	0.0233	2	0.00	0.000	0.0210	2	0.90	1.267
752	0.0218	2	0.00	0.000	0.0229	2	0.00	0.000	0.0233	2	0.00	0.000	0.0206	2	0.00	0.000
753	0.0214	2	0.00	0.000	0.0218	2	0.00	0.000	0.0229	2	0.00	0.000	0.0218	2	0.00	0.000
754	0.0330	3	0.00	0.000	0.0210	2	0.00	0.000	0.0206	2	0.00	0.000	0.0195	2	0.00	0.000
755	n.s.	n.s.	n.s.	n.s.	0.0206	2	0.00	0.000	0.0311	3	0.00	0.000	0.0431	4	0.00	0.000
756	0.0109	1	0.00	-	0.0225	2	0.32	0.449	0.0225	2	0.24	0.334	0.0203	2	0.36	0.257
757	0.0304	3	0.00	0.000	0.0206	2	0.00	0.000	0.0233	2	0.00	0.000	0.0214	2	0.00	0.000
758	0.0214	2	0.00	0.000	0.0105	2	0.00	0.000	0.0214	2	0.00	0.000	0.0210	2	0.00	0.000
759	n.s.	n.s.	n.s.	n.s.	0.0214	2	0.00	0.000	0.0218	2	0.00	0.000	0.0210	2	0.00	0.000
760	0.0105	1	0.00	-	0.0214	2	0.00	0.000	0.0225	2	0.00	0.000	0.0210	2	0.00	0.000
761	0.0315	3	0.00	0.000	0.0206	2	0.00	0.000	0.0210	2	0.00	0.000	0.0221	2	0.00	0.000
762	0.0308	3	0.00	0.000	0.0094	2	0.00	0.000	0.0210	2	0.00	0.000	0.0203	2	0.00	0.000
763	n.s.	n.s.	n.s.	n.s.	0.0218	2	0.00	0.000	0.0311	3	0.00	0.000	0.0416	4	1.08	2.170
764	0.0206	2	0.00	0.000	0.0218	2	0.00	0.000	0.0225	2	0.00	0.000	0.0218	2	0.00	0.000
765	0.0206	2	0.00	0.000	0.0098	2	0.00	0.000	0.0221	2	0.00	0.000	0.0203	2	0.00	0.000
766	0.0308	3	0.00	0.000	0.0191	2	0.00	0.000	0.0218	2	0.00	0.000	0.0214	2	0.00	0.000
767	n.s.	n.s.	n.s.	n.s.	0.0109	2	0.00	0.000	0.0214	2	0.00	0.000	0.0210	2	0.00	0.000

$$SD = \sqrt{\frac{\sum (x_i - \bar{x})^2}{n-1}}$$

TABLE 2 (cont.).- Swept area, number of hauls and Atlantic cod mean catch (kg) and SD by stratum. Spanish Spring Surveys on NAFO Div. 3NO: 1997-2007. Swept area in square miles. n.s. means stratum not surveyed. 1997-2000 data are transformed C/V *Playa de Mendoña* data, and 2002-2007 data are original from R/V *Vizconde de Eza*. In 2001, there are data from the two vessels.

Stratum	2001				2002				2003				2004			
	Swept area	Tow number	A. cod Mean catch	A. cod SD	Swept area	Tow number	A. cod Mean catch	A. cod SD	Swept area	Tow number	A. cod Mean catch	A. cod SD	Swept area	Tow number	A. cod Mean catch	A. cod SD
353	0.0341	3	7.33	8.145	0.0476	4	0.00	0.003	0.0334	3	0.00	0.000	0.0338	3	10.21	8.691
354	0.0338	3	16.07	3.315	0.0356	3	0.01	0.012	0.0338	3	7.63	13.221	0.0345	3	4.76	3.335
355	0.0240	2	56.11	64.898	0.0236	2	0.96	0.370	0.0229	2	3.02	2.390	0.0229	2	5.09	3.267
356	0.0240	2	149.60	76.650	0.0233	2	15.20	10.889	0.0225	2	15.61	1.605	0.0221	2	2.97	0.714
357	0.0244	2	27.20	36.062	0.0240	2	6.65	1.909	0.0229	2	5.28	7.460	0.0229	2	13.30	17.727
358	0.0345	3	3.42	2.592	0.0345	3	2.63	1.429	0.0338	3	207.22	260.186	0.0330	3	14.41	12.455
359	0.0803	7	176.35	433.935	0.0686	6	2.72	3.436	0.0791	7	1.03	1.522	0.0791	7	29.83	54.712
360	0.2423	20	11.36	27.470	0.2865	25	0.82	2.887	0.2254	20	1.14	2.952	0.2310	20	3.55	4.484
374	0.0240	2	0.00	0.000	0.0345	3	0.00	0.000	0.0225	2	0.00	0.000	0.0232	2	0.00	0.000
375	0.0338	3	0.00	0.000	0.0353	3	0.47	0.503	0.0330	3	0.48	0.826	0.0338	3	0.05	0.081
376	0.1155	10	0.04	0.119	0.1140	10	0.00	0.000	0.1125	10	0.65	1.987	0.1166	10	0.60	0.733
377	0.0229	2	0.00	0.000	0.0229	2	0.00	0.000	0.0225	2	1.25	1.768	0.0218	2	19.60	24.020
378	0.0236	2	11.98	15.726	0.0233	2	1.45	2.051	0.0225	2	19.18	19.141	0.0225	2	17.75	3.989
379	0.0229	2	9.54	9.001	0.0229	2	24.83	32.492	0.0229	2	4.35	0.481	0.0124	1	23.95	-
380	0.0206	2	6.00	2.895	0.0225	2	0.31	0.035	0.0229	2	1.09	0.976	0.0221	2	7.77	2.305
381	0.0236	2	0.66	0.891	0.0229	2	0.04	0.057	0.0229	2	0.00	0.000	0.0225	2	5.47	4.150
382	0.0469	4	0.12	0.145	0.0341	3	0.04	0.076	0.0454	4	0.00	0.000	0.0461	4	0.47	0.888
721	0.0248	2	4.85	6.859	0.0233	2	1.01	1.430	0.0225	2	9.40	13.287	0.0221	2	2.20	3.111
722	0.0233	2	0.00	0.000	0.0236	2	0.00	0.000	0.0221	2	1.73	2.447	0.0218	2	0.00	0.000
723	0.0240	2	676.15	932.179	0.0233	2	55.60	69.155	0.0229	2	0.65	0.919	0.0229	2	1.94	2.744
724	0.0353	3	6.16	10.254	0.0225	2	49.80	70.428	0.0225	2	10.46	14.786	0.0214	2	0.00	0.000
725	0.0116	2	1367.61	1856.733	0.0225	2	9.25	7.849	0.0229	2	2.17	3.062	0.0225	2	0.29	0.403
726	0.0116	2	1.83	2.593	0.0214	2	1122.95	1569.289	0.0225	2	0.00	0.000	0.0225	2	0.00	0.000
727	0.0225	2	10.40	4.810	0.0233	2	2.80	3.960	0.0218	2	7.45	9.405	0.0232	2	0.00	0.000
728	0.0229	2	0.00	0.000	0.0229	2	21.40	30.264	0.0225	2	0.00	0.000	0.0180	2	0.00	0.000
752	0.0210	2	0.00	0.000	0.0116	1	0.00	0.000	0.0229	2	0.00	0.000	0.0214	2	0.00	0.000
753	0.0214	2	0.00	0.000	0.0229	2	0.00	0.000	0.0229	2	0.00	0.000	0.0218	2	0.00	0.000
754	0.0195	2	0.00	0.000	0.0341	3	0.00	0.000	0.0218	2	0.00	0.000	0.0214	2	0.00	0.000
755	0.0416	4	0.00	0.000	0.0338	3	0.00	0.000	0.0221	2	0.00	0.000	0.0319	3	0.00	0.000
756	0.0113	2	0.04	0.057	0.0229	2	0.00	0.000	0.0221	2	0.00	0.000	0.0218	2	0.00	0.000
757	0.0233	2	0.00	0.000	0.0225	2	64.40	91.075	0.0221	2	0.00	0.000	0.0218	2	0.00	0.000
758	0.0218	2	0.00	0.000	0.0225	2	2.80	3.960	0.0221	2	0.00	0.000	0.0214	2	0.00	0.000
759	0.0221	2	0.00	0.000	0.0225	2	0.00	0.000	0.0113	1	0.00	-	0.0214	2	0.00	0.000
760	0.0229	2	0.00	0.000	0.0229	2	0.00	0.000	0.0218	2	0.00	0.000	0.0221	2	0.00	0.000
761	0.0225	2	0.00	0.000	0.0225	2	0.17	0.236	0.0225	2	0.00	0.000	0.0221	2	0.00	0.000
762	0.0116	2	0.00	0.000	0.0225	2	0.15	0.212	0.0225	2	0.00	0.000	0.0233	2	0.00	0.000
763	0.0330	3	0.00	0.000	0.0225	2	0.00	0.000	0.0311	3	0.00	0.000	0.0326	3	0.00	0.000
764	0.0240	2	0.00	0.000	0.0236	2	0.00	0.000	0.0221	2	0.00	0.000	0.0229	2	0.00	0.000
765	0.0113	2	0.00	0.000	0.0236	2	0.00	0.000	0.0113	1	0.00	-	0.0225	2	0.00	0.000
766	0.0203	2	0.00	0.000	0.0233	2	0.00	0.000	0.0225	2	0.00	0.000	0.0225	2	0.00	0.000
767	0.0225	2	0.00	0.000	0.0229	2	0.00	0.000	0.0229	2	0.00	0.000	0.0218	2	0.00	0.000

TABLE 2 (cont.).- Swept area, number of hauls and Atlantic cod mean catch (kg) and SD by stratum. Spanish Spring Surveys on NAFO Div. 3NO: 1997-2007. Swept area in square miles. n.s. means stratum not surveyed. 1997-2000 data are transformed C/V *Playa de Mendoña* data, and 2002-2007 data are original from R/V *Vizconde de Eza*. In 2001, there are data from the two vessels.

Stratum	2005					2006					2007				
	Swept area	Tow number	A. cod	Mean catch	SD	Swept area	Tow number	A. cod	Mean catch	SD	Swept area	Tow number	A. cod	Mean catch	SD
353	0.0353	3	4.20	3.962		0.0371	3	11.53	7.341		0.0364	3	0.14	0.138	
354	0.0353	3	6.76	8.311		0.0364	3	10.98	14.032		0.0364	3	16.81	14.624	
355	0.0225	2	1.97	0.255		0.0248	2	3.04	0.078		0.0240	2	41.34	12.820	
356	0.0233	2	1.43	1.478		0.0240	2	3.88	3.247		0.0240	2	0.96	1.351	
357	0.0233	2	3.98	4.603		0.0244	2	12.75	8.400		0.0360	3	1.42	1.323	
358	0.0349	3	22.75	17.967		0.0349	3	82.54	80.442		0.0368	3	113.84	43.776	
359	0.0814	7	57.31	134.609		0.0975	8	372.36	643.214		0.0855	7	3.17	4.658	
360	0.2325	20	2.47	4.698		0.2340	19	7.35	8.119		0.2378	20	2.42	4.606	
374	0.0229	2	0.11	0.148		0.0236	2	0.00	0.000		0.0240	2	0.00	0.000	
375	0.0349	3	0.00	0.000		0.0364	3	13.53	15.862		0.0364	3	1.71	1.646	
376	0.1174	10	0.76	0.963		0.1219	10	6.84	11.380		0.1185	10	0.68	1.167	
377	0.0233	2	61.19	64.955		0.0236	2	90.62	69.919		0.0240	2	698.56	987.885	
378	0.0225	2	8.59	10.087		0.0240	2	90.32	85.680		0.0233	2	85.98	23.723	
379	0.0236	2	5.70	7.078		0.0236	2	6.30	8.627		0.0240	2	3.13	0.394	
380	0.0229	2	27.53	24.784		0.0229	2	8.70	1.697		0.0240	2	4.20	5.945	
381	0.0233	2	3.63	3.765		0.0229	2	8.43	1.167		0.0240	2	2.19	0.354	
382	0.0458	4	0.97	0.639		0.0469	4	0.75	1.033		0.0484	4	0.00	0.000	
721	0.0229	2	0.00	0.000		0.0236	2	0.00	0.000		0.0116	1	0.00	-	
722	0.0233	2	0.00	0.000		0.0240	2	0.00	0.000		0.0225	2	0.00	0.000	
723	0.0233	2	0.00	0.000		0.0236	2	0.00	0.000		0.0240	2	3.15	4.455	
724	0.0225	2	0.00	0.000		0.0233	2	0.00	0.000		0.0233	2	0.00	0.000	
725	0.0236	2	1.47	2.073		0.0233	2	0.00	0.000		0.0225	2	11.89	11.823	
726	0.0113	1	0.00	-		0.0225	2	0.00	0.000		0.0229	2	0.00	0.000	
727	0.0229	2	0.00	0.000		0.0225	2	0.00	0.000		0.0240	2	0.00	0.000	
728	0.0109	1	0.00	-		0.0225	2	0.00	0.000		0.0225	2	0.00	0.000	
752	0.0236	2	0.00	0.000		0.0225	2	0.00	0.000		0.0225	2	0.00	0.000	
753	0.0225	2	0.00	0.000		0.0225	2	0.00	0.000		0.0225	2	0.00	0.000	
754	0.0225	2	0.00	0.000		0.0225	2	0.00	0.000		0.0225	2	0.00	0.000	
755	0.0450	4	0.00	0.000		0.0338	3	0.00	0.000		0.0338	3	0.00	0.000	
756	0.0233	2	0.00	0.000		0.0229	2	0.00	0.000		0.0225	2	0.00	0.000	
757	0.0225	2	0.00	0.000		0.0225	2	0.00	0.000		0.0229	2	0.00	0.000	
758	0.0225	2	0.00	0.000		0.0225	2	0.00	0.000		0.0225	2	0.00	0.000	
759	0.0229	2	0.00	0.000		0.0225	2	0.00	0.000	n.s.	n.s.	n.s.	n.s.	n.s.	
760	0.0229	2	0.00	0.000		0.0225	2	0.00	0.000	0.0223	2	0.00	0.000		
761	0.0221	2	0.00	0.000		0.0233	2	0.00	0.000	0.0225	2	0.00	0.000		
762	0.0225	2	0.00	0.000		0.0233	2	0.00	0.000	n.s.	n.s.	n.s.	n.s.	n.s.	
763	0.0334	3	0.00	0.000		0.0225	2	0.00	0.000	n.s.	n.s.	n.s.	n.s.	n.s.	
764	0.0233	2	0.00	0.000		0.0233	2	0.00	0.000	0.0225	2	0.00	0.000		
765	0.0229	2	0.00	0.000		0.0236	2	0.00	0.000	0.0225	2	0.00	0.000		
766	0.0229	2	0.00	0.000		0.0229	2	0.00	0.000	n.s.	n.s.	n.s.	n.s.	n.s.	
767	0.0113	1	0.00	-		0.0233	2	0.00	0.000	n.s.	n.s.	n.s.	n.s.	n.s.	

TABLE 3.- Stratified mean catches (Kg) by stratum and year and SD by year of Atlantic cod (1997-2007). n.s. means stratum not surveyed. 1997-2000 data are transformed C/V Playa de Menduiña data. 2002-2007 data are original from R/V *Vizconde de Eza*. In 2001, there are data from the two vessels.

Stratum	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007
353	0.00	0.59	1684.29	2310.56	1972.67	0.40	0.00	2746.49	1129.80	3100.67	36.76
354	0.00	4347.10	1209.44	4536.47	3954.04	1.64	1877.80	1172.11	1662.39	2700.42	4134.28
355	317.46	2001.63	472.57	7017.36	4152.14	71.15	223.48	376.66	145.78	224.59	3058.79
356	366.75	292.75	1935.74	768.05	7031.20	714.40	733.44	139.36	66.98	182.17	44.89
357	15014.55	1222.35	1659.07	1500.68	4460.80	1090.60	865.10	2180.38	651.90	2091.00	232.33
358	397.76	1002.53	2246.51	41597.12	768.75	592.50	46625.25	3241.50	5119.50	18570.75	25614.00
359	473.87	164.50	3052.91	7687.04	74245.15	1146.52	435.31	12557.95	24128.71	156764.14	1334.99
360	301.58	616.24	6478.57	6017.33	31605.14	2283.17	3169.28	9886.61	6869.14	20449.63	6724.01
374	12.23	0.00	124.31	0.00	0.00	0.00	0.00	0.00	22.47	0.00	0.00
375	0.00	211.79	261.73	0.00	0.00	126.47	129.18	12.65	0.00	3665.73	464.22
376	0.00	263.27	822.50	1202.94	50.03	0.00	864.70	801.87	1010.91	9129.90	911.39
377	26.59	188.96	21.35	1.92	0.00	0.00	125.00	1959.50	6119.00	9062.00	69855.95
378	325.88	481.53	1078.58	1480.09	1665.22	201.55	2665.33	2466.56	1194.36	12553.79	11950.53
379	390.21	880.31	553.41	4358.29	1010.71	2631.45	461.10	2538.70	603.67	667.80	331.94
380	34.94	223.39	3703.59	788.08	576.11	30.19	104.64	745.92	2642.40	835.20	403.58
381	10.08	30.36	125.22	250.68	95.74	5.76	0.00	787.90	523.08	1213.20	315.36
382	0.00	108.42	18.00	243.65	41.41	14.98	0.00	160.78	332.28	255.54	0.00
721	1363.56	39.80	5738.57	1842.35	315.25	65.75	610.68	143.00	0.00	0.00	0.00
722	26.16	0.00	0.00	75.84	0.00	0.00	145.32	0.00	0.00	0.00	0.00
723	1534.94	680.69	2614.28	3413.20	104803.25	8618.00	100.75	300.70	0.00	0.00	488.25
724	161.20	184615.64	2.82	87.21	764.25	6175.20	1296.42	0.00	0.00	0.00	0.00
725	2467.77	3240.64	1432.94	455.78	143598.88	971.25	227.33	29.93	153.93	0.00	1248.45
726	n.s	341.39	58.07	637.55	132.02	80852.04	0.00	0.00	0.00	0.00	0.00
727	11.42	255.30	883.49	879.12	998.37	268.80	715.20	0.00	0.00	0.00	0.00
728	91.43	120.09	0.00	69.87	0.00	1669.20	0.00	0.00	0.00	0.00	0.00
752	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
753	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
754	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
755	n.s	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
756	0.00	32.07	23.86	36.40	4.04	0.00	0.00	0.00	0.00	0.00	0.00
757	0.00	0.00	0.00	0.00	0.00	6568.80	0.00	0.00	0.00	0.00	0.00
758	0.00	0.00	0.00	0.00	0.00	277.20	0.00	0.00	0.00	0.00	0.00
759	n.s	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	n.s
760	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
761	0.00	0.00	0.00	0.00	0.00	28.56	0.00	0.00	0.00	0.00	0.00
762	0.00	0.00	0.00	0.00	0.00	31.80	0.00	0.00	0.00	0.00	n.s
763	n.s	0.00	0.00	283.12	0.00	0.00	0.00	0.00	0.00	0.00	n.s
764	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
765	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
766	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	n.s
767	n.s	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	n.s
TOTAL (\bar{Y})	23328.40	201361.35	36201.79	87540.73	382245.17	114437.38	61375.29	42248.54	52376.29	241466.52	127149.71
S.D.	2.50	19.47	3.50	8.46	36.96	11.07	5.93	4.09	5.06	23.35	13.47

TABLE 4.- Survey estimates (by the swept area method) of Atlantic cod biomass (t) and SD by stratum and year on NAFO Div. 3NO. n.s. means stratum not surveyed. 1997-2000 data are transformed C/V *Playa de Menduíña* data. 2002-2007 data are original from R/V *Vizconde de Eza*. In 2001, there are data from the two vessels.

Stratum	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007
353	0	0	140	195	173	0	0	244	96	251	3
354	0	366	111	382	351	0	167	102	141	223	341
355	27	181	41	604	346	6	20	33	13	18	255
356	33	26	169	68	586	61	65	13	6	15	4
357	1357	102	140	121	366	91	76	191	56	172	19
358	35	86	194	3657	67	52	4144	295	440	1597	2091
359	41	14	252	656	6476	100	39	1111	2076	12863	109
360	26	53	529	502	2609	199	281	856	591	1660	566
374	1	0	10	0	0	0	0	0	2	0	0
375	0	18	22	0	0	11	12	1	0	302	38
376	0	23	67	100	4	0	77	69	86	749	77
377	2	17	2	0	0	0	11	180	526	767	5821
378	31	41	95	127	141	17	237	219	106	1046	1028
379	38	74	47	387	88	230	40	205	51	57	28
380	3	20	314	67	56	3	9	67	231	73	34
381	1	3	11	21	8	1	0	70	45	106	26
382	0	10	1	20	4	1	0	14	29	22	0
721	123	4	471	156	25	6	54	13	0	0	0
722	2	0	0	7	0	0	13	0	0	0	0
723	146	59	229	276	8734	741	9	26	0	0	41
724	14	17902	0	8	65	549	115	0	0	0	0
725	239	376	125	43	12347	86	20	3	13	0	111
726	n.s.	33	5	58	11	7565	0	0	0	0	0
727	1	22	75	84	89	23	66	0	0	0	0
728	9	12	0	7	0	146	0	0	0	0	0
752	0	0	0	0	0	0	0	0	0	0	0
753	0	0	0	0	0	0	0	0	0	0	0
754	0	0	0	0	0	0	0	0	0	0	0
755	n.s.	0	0	0	0	0	0	0	0	0	0
756	0	3	2	4	0	0	0	0	0	0	0
757	0	0	0	0	0	584	0	0	0	0	0
758	0	0	0	0	0	25	0	0	0	0	0
759	n.s.	0	0	0	0	0	0	0	0	0	n.s.
760	0	0	0	0	0	0	0	0	0	0	0
761	0	0	0	0	0	3	0	0	0	0	0
762	0	0	0	0	0	3	0	0	0	0	n.s.
763	n.s.	0	0	27	0	0	0	0	0	0	n.s.
764	0	0	0	0	0	0	0	0	0	0	0
765	0	0	0	0	0	0	0	0	0	0	0
766	0	0	0	0	0	0	0	0	0	0	n.s.
767	n.s.	0	0	0	0	0	0	0	0	0	n.s.
TOTAL	2131	19444	3054	7576	32548	10502	5455	3712	4509	19921	10592
S.D.	1322	18206	655	2566	15903	7971	3016	848	1984	8109	5853

TABLE 5.- Length weight relationships in the calculation of Atlantic cod biomass. The equation is $Weight = a(l + 0.5)^b$
Spanish Spring Surveys on NAFO Div. 3NO: 1997-2007.

	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007
a	0.0102 Error = 0.2480	0.0061 Error = 0.0748	0.0048 Error = 0.0788	0.0060 Error = 0.0706	0.0048 Error = 0.0893	0.0057 Error = 0.1025	0.0046 Error = 0.0581	0.0052 Error = 0.0698	0.0052 Error = 0.0715	0.0058 Error = 0.0678	0.0059 Error = 0.0570
b	2.9387 Error = 0.0629	3.0671 Error = 0.0197	3.1313 Error = 0.0203	3.0822 Error = 0.0179	3.1198 Error = 0.0228	3.0783 Error = 0.0274	3.1370 Error = 0.0153	3.1107 Error = 0.0185	3.1238 Error = 0.0189	3.0965 Error = 0.0174	3.0762 Error = 0.0153
	R ² = 0.975 N = 431	R2 = 0.997 N = 687	R ² = 0.997 N = 430	R ² = 0.997 N = 877	R ² = 0.996 N = 488	R ² = 0.995 N = 678	R ² = 0.998 N = 516	R ² = 0.997 N = 656	R ² = 0.997 N = 612	R ² = 0.999 N = 1129	R2 = 0.998 N= 1011

TABLE 6.- Atlantic cod length distribution per haul mean catches by sex and year. Number per stratified mean catches. Spanish Spring Survey on NAFO 3NO: 1997-2007. Indet. means indeterminate. 1997-2000 data are transformed C/V *Playa de Mendoña* data. 2002-2007 data are original R/V *Vizconde de Eza* data. (*) indicates untransformed data.

Length (cm.)	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007
8	0.000	0.000	0.013	0.000	0.009	0.011	0.000	0.000	0.000	0.014	0.000
10	0.000	0.000	0.011	0.000	0.000	0.039	0.000	0.408	0.088	0.000	0.015
12	0.000	0.010	0.010	0.006	0.000	0.030	0.054	2.141	0.722	0.014	0.416
14	0.000	0.025	0.205	0.020	0.000	0.000	0.065	1.542	1.129	0.008	4.160
16	0.000	0.011	0.741	0.124	0.021	0.040	0.028	0.828	0.644	0.014	8.177
18	0.000	0.007	0.774	0.115	0.019	0.007	0.037	0.186	0.245	0.013	4.890
20	0.000	0.004	0.287	0.211	0.103	0.000	0.061	0.063	0.036	0.007	1.020
22	0.000	0.002	0.478	0.285	0.184	0.083	0.062	0.076	0.094	0.041	0.070
24	0.002	0.005	1.173	0.241	0.175	0.096	0.075	0.119	0.224	0.367	0.024
26	0.002	0.009	1.740	0.319	0.194	0.139	0.150	0.219	0.515	0.948	0.008
28	0.013	0.028	1.251	0.385	0.216	0.153	0.189	0.141	1.044	3.595	0.000
30	0.013	0.037	0.525	0.417	0.132	0.284	0.243	0.162	1.151	5.313	0.041
32	0.051	0.028	0.172	0.327	0.162	0.312	0.087	0.149	0.551	3.854	0.086
34	0.087	0.030	0.096	0.438	0.380	0.524	0.067	0.330	0.393	1.710	0.317
36	0.142	0.022	0.070	0.559	0.923	0.596	0.071	0.421	0.189	1.102	0.442
38	0.184	0.026	0.090	1.038	1.787	0.572	0.121	0.420	0.129	2.258	0.729
40	0.108	0.105	0.086	1.030	3.363	0.689	0.081	0.217	0.135	5.496	0.925
42	0.066	0.075	0.031	0.897	3.463	1.005	0.078	0.248	0.113	5.305	0.881
44	0.106	0.365	0.047	0.473	4.234	1.141	0.117	0.101	0.097	4.004	0.885
46	0.073	0.603	0.025	0.307	5.028	1.483	0.111	0.110	0.136	2.317	0.788
48	0.091	0.931	0.045	0.183	5.686	1.090	0.175	0.077	0.173	1.054	1.632
50	0.043	0.963	0.044	0.137	4.959	1.058	0.225	0.060	0.101	0.487	2.035
52	0.074	0.924	0.063	0.099	4.098	1.111	0.298	0.088	0.128	0.279	1.748
54	0.087	1.499	0.106	0.109	3.195	0.895	0.390	0.072	0.026	0.276	1.412
56	0.142	1.537	0.081	0.069	1.224	0.691	0.428	0.065	0.028	0.227	0.651
58	0.124	1.764	0.113	0.136	0.693	0.223	0.322	0.110	0.012	0.231	0.401
60	0.195	1.026	0.130	0.101	0.532	0.370	0.306	0.074	0.055	0.256	0.262
62	0.114	0.540	0.098	0.065	0.181	0.126	0.183	0.093	0.078	0.229	0.094
64	0.088	0.505	0.072	0.152	0.032	0.005	0.227	0.104	0.092	0.204	0.054
66	0.111	0.163	0.049	0.134	0.047	0.057	0.098	0.063	0.089	0.114	0.079
68	0.014	0.271	0.067	0.101	0.014	0.000	0.093	0.071	0.077	0.098	0.056
70	0.029	0.157	0.019	0.137	0.015	0.061	0.085	0.042	0.093	0.092	0.096
72	0.004	0.193	0.013	0.104	0.028	0.007	0.027	0.031	0.083	0.074	0.037
74	0.013	0.136	0.018	0.142	0.012	0.000	0.011	0.033	0.078	0.096	0.029
76	0.002	0.086	0.011	0.066	0.017	0.002	0.015	0.030	0.079	0.071	0.012
78	0.003	0.080	0.008	0.034	0.022	0.000	0.010	0.017	0.056	0.121	0.042
80	0.006	0.079	0.015	0.073	0.039	0.000	0.027	0.036	0.047	0.051	0.029
82	0.001	0.038	0.005	0.032	0.013	0.000	0.000	0.009	0.018	0.103	0.008
84	0.003	0.000	0.004	0.044	0.000	0.011	0.025	0.003	0.006	0.057	0.036
86	0.001	0.048	0.012	0.026	0.021	0.000	0.008	0.000	0.022	0.041	0.000
88	0.000	0.042	0.010	0.021	0.003	0.007	0.002	0.022	0.014	0.041	0.000
90	0.001	0.000	0.000	0.016	0.011	0.000	0.000	0.008	0.014	0.013	0.000
92	0.000	0.003	0.019	0.020	0.000	0.000	0.000	0.009	0.000	0.039	0.015
94	0.000	0.000	0.000	0.005	0.003	0.000	0.000	0.013	0.000	0.005	0.000
96	0.000	0.000	0.005	0.003	0.012	0.000	0.008	0.000	0.000	0.000	0.000
98	0.000	0.000	0.005	0.003	0.008	0.000	0.000	0.000	0.000	0.026	0.000
100	0.000	0.000	0.005	0.000	0.000	0.000	0.000	0.000	0.000	0.025	0.000
102	0.000	0.000	0.000	0.010	0.000	0.000	0.008	0.000	0.000	0.000	0.000
104	0.000	0.001	0.000	0.000	0.000	0.011	0.000	0.027	0.000	0.014	0.000
106	0.000	0.000	0.000	0.005	0.000	0.000	0.014	0.000	0.000	0.014	0.000
108	0.000	0.000	0.000	0.000	0.011	0.000	0.000	0.000	0.000	0.000	0.000
110	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
112	0.000	0.000	0.000	0.000	0.009	0.000	0.000	0.000	0.000	0.000	0.000
114	0.001	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
116	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.004
118	0.002	0.000	0.000	0.003	0.000	0.000	0.000	0.000	0.000	0.000	0.000
120	0.000	0.000	0.006	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
122	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
124	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
126	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
128	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
130	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
132	0.000	0.000	0.000	0.000	0.009	0.000	0.000	0.000	0.000	0.000	0.000
Total	1.997	12.378	8.847	9.220	41.290	12.930	4.684	9.035	9.005	40.718	32.605
Nº samples (*):	40	55	72	70	32	41	42	58	59	64	58
Nº Ind. (*):	742	967	2770	2753	1591	1030	539	939	1126	2909	2301
Sampled catch:	248	410	527	752	1107	776	654	554	778	2026	1115
Range (*):	24-118	12-104	9-121	13-118	8-132	9-104	12-106	10-105	11-91	7-104	9-114
Total catch:	572	3873	613	1274	3487	2806	846	554	794	3994	2182
Total hauls (*):	128	124	114	118	123	125	118	120	119	120	110

TABLE 7.- Swept area, number of hauls and Yellowtail flounder mean catch (kg) and SD by stratum. Spanish Spring Surveys on NAFO Div. 3NO: 1995-2007. Swept area in square miles. n.s. means stratum not surveyed. 1995-2000 data are transformed C/V *Playa de Menduña* data, and 2002-2007 data are original from R/V *Vizconde de Eza*. In 2001, there are data from the two vessels.

1995				1996				1997				1998				
Stratum	Swept area	Tow number	Y. flounder Mean catch	Swept area	Tow number	Y. flounder Mean catch	Y. flounder SD	Swept area	Tow number	Y. flounder Mean catch	Y. flounder SD	Swept area	Tow number	Y. flounder Mean catch	Y. flounder SD	
353	0.0353	3	5.82	4.105	0.0371	3	74.88	94.62	0.0480	4	12.55	14.26	0.0465	4	12.22	20.16
354	0.0353	3	1.78	3.089	0.0319	3	1.11	0.84	0.0233	2	1.41	1.56	0.0356	3	1.22	0.24
355	n.s.	n.s.	n.s.	n.s.	0.0221	2	0.25	0.35	0.0233	2	2.20	0.31	0.0221	2	0.13	0.18
356	n.s.	n.s.	n.s.	n.s.	0.0203	2	0.00	0.00	0.0225	2	0.32	0.46	0.0221	2	0.00	0.00
357	0.0109	1	0.00	-	0.0218	2	0.00	0.00	0.0443	4	0.00	0.00	0.0240	2	0.00	0.00
358	0.0319	3	0.00	0.000	0.0319	3	0.13	0.23	0.0563	5	0.02	0.04	0.0236	3	0.00	0.00
359	0.0345	3	1.35	2.336	0.0548	5	0.92	0.83	0.0690	6	0.08	0.14	0.0698	6	0.17	0.22
360	0.3563	31	20.44	40.707	0.3761	31	142.09	128.86	0.3754	32	80.92	155.59	0.2561	25	373.90	629.84
374	0.0225	2	0.00	0.000	0.0233	2	0.00	0.00	0.0353	3	0.00	0.00	0.0353	3	0.04	0.02
375	0.0225	2	1.48	1.875	0.0229	2	41.40	58.54	0.0116	1	0.20	-	0.0345	3	12.37	21.37
376	0.1729	15	35.06	58.691	0.1650	14	71.40	86.94	0.1583	14	162.35	179.83	0.0930	10	279.27	181.29
377	0.0221	2	0.00	0.000	0.0229	2	0.00	0.00	0.0116	1	0.00	-	0.0229	2	0.00	0.00
378	0.0435	4	0.00	0.000	0.0330	3	0.06	0.10	0.0210	2	0.00	0.00	0.0120	2	0.00	0.00
379	0.0221	2	0.00	0.000	0.0113	1	0.00	-	0.0206	2	0.00	0.00	0.0356	3	0.00	0.00
380	n.s.	n.s.	n.s.	n.s.	0.0221	2	0.00	0.00	0.0210	2	0.00	0.00	0.0113	2	0.00	0.00
381	n.s.	n.s.	n.s.	n.s.	0.0229	2	0.00	0.00	0.0221	2	0.00	0.00	0.0229	2	0.00	0.00
382	n.s.	n.s.	n.s.	n.s.	0.0338	3	0.00	0.00	0.0461	4	0.00	0.00	0.0229	3	0.00	0.00
721	n.s.	n.s.	n.s.	n.s.	0.0214	2	0.03	0.05	0.0221	2	0.75	1.06	0.0203	2	0.00	0.00
722	n.s.	n.s.	n.s.	n.s.	0.0206	2	0.00	0.00	0.0214	2	0.00	0.00	0.0101	2	0.00	0.00
723	n.s.	n.s.	n.s.	n.s.	0.0109	1	0.00	-	0.0210	2	0.00	0.00	0.0233	2	0.00	0.00
724	0.0105	1	0.00	-	0.0203	2	0.00	0.00	0.0225	2	0.00	0.00	0.0206	2	0.00	0.00
725	0.0334	3	0.00	0.000	0.0225	2	0.00	0.00	0.0206	2	0.00	0.00	0.0086	1	0.00	-
726	0.0214	2	0.00	0.000	0.0218	2	0.00	0.00	n.s.	n.s.	n.s.	n.s.	0.0094	2	0.00	0.00
727	n.s.	n.s.	n.s.	n.s.	0.0210	2	0.00	0.00	0.0094	1	0.00	-	0.0233	2	0.00	0.00
728	n.s.	n.s.	n.s.	n.s.	0.0218	2	0.00	0.00	0.0214	2	0.00	0.00	0.0206	2	0.00	0.00
752	n.s.	n.s.	n.s.	n.s.	0.0109	1	0.00	-	0.0218	2	0.00	0.00	0.0229	2	0.00	0.00
753	n.s.	n.s.	n.s.	n.s.	0.0199	2	0.00	0.00	0.0214	2	0.00	0.00	0.0218	2	0.00	0.00
754	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	0.0330	3	0.00	0.00	0.0210	2	0.00	0.00
755	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	0.0206	2	0.00	0.00
756	n.s.	n.s.	n.s.	n.s.	0.0210	2	0.00	0.00	0.0109	1	0.00	-	0.0225	2	0.00	0.00
757	n.s.	n.s.	n.s.	n.s.	0.0188	2	0.00	0.00	0.0304	3	0.00	0.00	0.0206	2	0.00	0.00
758	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	0.0214	2	0.00	0.00	0.0105	2	0.00	0.00
759	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	0.0214	2	0.00	0.00
760	n.s.	n.s.	n.s.	n.s.	0.0210	2	0.00	0.00	0.0105	1	0.00	-	0.0214	2	0.00	0.00
761	n.s.	n.s.	n.s.	n.s.	0.0199	2	0.00	0.00	0.0315	3	0.00	0.00	0.0206	2	0.00	0.00
762	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	0.0308	3	0.00	0.00	0.0094	2	0.00	0.00
763	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	0.0218	2	0.00	0.00
764	n.s.	n.s.	n.s.	n.s.	0.0210	2	0.00	0.00	0.0206	2	0.00	0.00	0.0218	2	0.00	0.00
765	n.s.	n.s.	n.s.	n.s.	0.0199	2	0.00	0.00	0.0206	2	0.00	0.00	0.0098	2	0.00	0.00
766	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	0.0308	3	0.00	0.00	0.0191	2	0.00	0.00
767	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	0.0109	2	0.00	0.00

$$SD = \sqrt{\frac{\sum (x_i - \bar{x})^2}{n-1}}$$

TABLE 7 (cont.).- Swept area, number of hauls and Yellowtail flounder mean catch (kg) and SD by stratum. Spanish Spring Surveys on NAFO Div. 3NO: 1995-2007. Swept area in square miles. n.s. means stratum not surveyed. 1995-2000 data are transformed C/V *Playa de Menduña* data, and 2002-2007 data are original from R/V *Vizconde de Eza*. In 2001, there are data from the two vessels.

Stratum	1999				2000				2001			
	Swept area	Tow number	Y. flounder Mean catch	Y. flounder SD	Swept area	Tow number	Y. flounder Mean catch	Y. flounder SD	Swept area	Tow number	Y. flounder Mean catch	Y. flounder SD
353	0.0360	3	150.18	182.44	0.0356	3	67.87	91.37	0.0341	3	61.42	102.797
354	0.0218	2	0.08	0.12	0.0356	3	1.79	1.93	0.0338	3	0.34	0.322
355	0.0229	2	0.00	0.00	0.0233	2	0.00	0.00	0.0240	2	0.00	0.000
356	0.0229	2	0.00	0.00	0.0225	2	0.00	0.00	0.0240	2	0.01	0.007
357	0.0236	2	0.00	0.00	0.0124	1	0.00	-	0.0244	2	0.00	0.000
358	0.0349	3	0.00	0.00	0.0341	3	0.00	0.00	0.0345	3	0.00	0.000
359	0.0364	3	0.34	0.47	0.0469	4	2.36	2.93	0.0803	7	1.42	2.836
360	0.2325	19	545.18	424.37	0.2396	20	391.18	331.64	0.2423	20	536.80	488.657
374	0.0244	2	74.16	103.18	0.0240	2	20.47	23.55	0.0240	2	238.75	111.369
375	0.0236	2	347.15	168.25	0.0244	2	153.36	2.06	0.0338	3	100.33	68.319
376	0.1219	10	551.60	165.61	0.1200	10	435.27	236.60	0.1155	10	443.12	196.619
377	0.0240	2	0.00	0.00	0.0229	2	0.05	0.06	0.0229	2	0.00	0.000
378	0.0229	2	0.00	0.00	0.0233	2	0.00	0.00	0.0236	2	0.00	0.000
379	0.0236	2	0.00	0.00	0.0225	2	0.00	0.00	0.0229	2	0.00	0.000
380	0.0236	2	0.00	0.00	0.0236	2	0.00	0.00	0.0206	2	0.00	0.000
381	0.0229	2	0.00	0.00	0.0236	2	0.00	0.00	0.0236	2	0.00	0.000
382	0.0484	4	0.00	0.00	0.0499	4	0.00	0.00	0.0469	4	0.02	0.030
721	0.0244	2	0.00	0.00	0.0236	2	0.00	0.00	0.0248	2	0.00	0.000
722	0.0229	2	0.00	0.00	0.0218	2	0.00	0.00	0.0233	2	0.00	0.000
723	0.0229	2	0.00	0.00	0.0248	2	0.00	0.00	0.0240	2	0.00	0.000
724	0.0225	2	0.00	0.00	0.0233	2	0.00	0.00	0.0353	3	0.00	0.000
725	0.0229	2	0.00	0.00	0.0210	2	0.00	0.00	0.0116	2	0.00	0.000
726	0.0225	2	0.00	0.00	0.0221	2	0.00	0.00	0.0116	2	0.00	0.000
727	0.0236	2	0.00	0.00	0.0210	2	0.00	0.00	0.0225	2	0.00	0.000
728	0.0233	2	0.00	0.00	0.0210	2	0.00	0.00	0.0229	2	0.00	0.000
752	0.0233	2	0.00	0.00	0.0206	2	0.00	0.00	0.0210	2	0.06	0.083
753	0.0229	2	0.00	0.00	0.0218	2	0.00	0.00	0.0214	2	0.00	0.000
754	0.0206	2	0.00	0.00	0.0195	2	0.00	0.00	0.0195	2	0.00	0.000
755	0.0311	3	0.00	0.00	0.0431	4	0.00	0.00	0.0416	4	0.00	0.000
756	0.0225	2	0.00	0.00	0.0203	2	0.00	0.00	0.0113	2	0.00	0.000
757	0.0233	2	0.00	0.00	0.0214	2	0.00	0.00	0.0233	2	0.00	0.000
758	0.0214	2	0.00	0.00	0.0210	2	0.00	0.00	0.0218	2	0.00	0.000
759	0.0218	2	0.00	0.00	0.0210	2	0.00	0.00	0.0221	2	0.00	0.000
760	0.0225	2	0.00	0.00	0.0210	2	0.00	0.00	0.0229	2	0.00	0.000
761	0.0210	2	0.00	0.00	0.0221	2	0.00	0.00	0.0225	2	0.00	0.000
762	0.0210	2	0.00	0.00	0.0203	2	0.00	0.00	0.0116	2	0.00	0.000
763	0.0311	3	0.00	0.00	0.0416	4	0.00	0.00	0.0330	3	0.00	0.000
764	0.0225	2	0.00	0.00	0.0218	2	0.00	0.00	0.0240	2	0.00	0.000
765	0.0221	2	0.00	0.00	0.0203	2	0.00	0.00	0.0113	2	0.00	0.000
766	0.0218	2	0.00	0.00	0.0214	2	0.00	0.00	0.0203	2	0.00	0.000
767	0.0214	2	0.00	0.00	0.0210	2	0.00	0.00	0.0218	2	0.00	0.000

$$SD = \sqrt{\frac{\sum (x_i - \bar{x})^2}{n-1}}$$

TABLE 7 (cont.).- Swept area, number of hauls and Yellowtail flounder mean catch (kg) and SD by stratum. Spanish Spring Surveys on NAFO Div. 3NO: 1995-2007. Swept area in square miles. n.s. means stratum not surveyed. 1995-2000 data are transformed C/V *Playa de Mendoña* data, and 2002-2007 data are original from R/V *Vizconde de Eza*. In 2001, there are data from the two vessels.

Stratum	2002				2003				2004			
	Swept area	Tow number	Y. flounder Mean catch	Y. flounder SD	Swept area	Tow number	Y. flounder Mean catch	Y. flounder SD	Swept area	Tow number	Y. flounder Mean catch	Y. flounder SD
353	0.0476	4	75.13	88.259	0.0334	3	11.15	19.307	0.0338	3	8.79	14.005
354	0.0356	3	0.17	0.289	0.0338	3	0.00	0.000	0.0345	3	0.62	1.065
355	0.0236	2	0.00	0.000	0.0229	2	0.00	0.000	0.0229	2	0.00	0.000
356	0.0233	2	0.00	0.000	0.0225	2	0.00	0.000	0.0221	2	0.00	0.000
357	0.0240	2	0.00	0.000	0.0229	2	0.00	0.000	0.0229	2	0.00	0.000
358	0.0345	3	0.00	0.000	0.0338	3	0.00	0.000	0.0330	3	0.26	0.442
359	0.0686	6	0.11	0.261	0.0791	7	0.00	0.000	0.0791	7	25.01	38.371
360	0.2865	25	340.23	356.687	0.2254	20	360.55	298.992	0.2310	20	403.19	333.463
374	0.0345	3	32.04	52.542	0.0225	2	16.13	8.238	0.0233	2	193.46	225.058
375	0.0353	3	48.61	68.927	0.0330	3	28.45	35.557	0.0338	3	543.04	155.015
376	0.1140	10	533.62	416.745	0.1125	10	391.60	257.289	0.1166	10	481.06	140.810
377	0.0229	2	0.00	0.000	0.0225	2	0.70	0.990	0.0218	2	0.00	0.000
378	0.0233	2	0.00	0.000	0.0225	2	0.00	0.000	0.0225	2	0.00	0.000
379	0.0229	2	0.00	0.000	0.0229	2	0.00	0.000	0.0124	1	0.00	-
380	0.0225	2	0.00	0.000	0.0229	2	0.00	0.000	0.0221	2	0.00	0.000
381	0.0229	2	0.00	0.000	0.0229	2	0.00	0.000	0.0225	2	0.00	0.000
382	0.0341	3	0.00	0.000	0.0454	4	0.00	0.000	0.0461	4	0.00	0.000
721	0.0233	2	0.00	0.000	0.0225	2	0.00	0.000	0.0221	2	0.00	0.000
722	0.0236	2	0.00	0.000	0.0221	2	0.00	0.000	0.0218	2	0.00	0.000
723	0.0233	2	0.00	0.000	0.0229	2	0.00	0.000	0.0229	2	0.00	0.000
724	0.0225	2	0.00	0.000	0.0225	2	0.52	0.735	0.0214	2	0.00	0.000
725	0.0225	2	0.00	0.000	0.0229	2	0.00	0.000	0.0225	2	0.00	0.000
726	0.0214	2	0.00	0.000	0.0225	2	0.00	0.000	0.0225	2	0.00	0.000
727	0.0233	2	0.00	0.000	0.0218	2	0.00	0.000	0.0233	2	0.00	0.000
728	0.0229	2	0.00	0.000	0.0225	2	0.00	0.000	0.0180	2	0.00	0.000
752	0.0116	1	0.00	-	0.0229	2	0.00	0.000	0.0214	2	0.00	0.000
753	0.0229	2	0.00	0.000	0.0229	2	0.00	0.000	0.0218	2	0.00	0.000
754	0.0341	3	0.00	0.000	0.0218	2	0.00	0.000	0.0214	2	0.00	0.000
755	0.0338	3	0.00	0.000	0.0221	2	0.00	0.000	0.0319	3	0.00	0.000
756	0.0229	2	0.00	0.000	0.0221	2	0.00	0.000	0.0218	2	0.00	0.000
757	0.0225	2	0.00	0.000	0.0221	2	0.00	0.000	0.0218	2	0.00	0.000
758	0.0225	2	0.00	0.000	0.0221	2	0.00	0.000	0.0214	2	0.00	0.000
759	0.0225	2	0.00	0.000	0.0113	1	0.00	-	0.0214	2	0.00	0.000
760	0.0229	2	0.00	0.000	0.0218	2	0.00	0.000	0.0221	2	0.00	0.000
761	0.0225	2	0.00	0.000	0.0225	2	0.00	0.000	0.0221	2	0.00	0.000
762	0.0225	2	0.00	0.000	0.0225	2	0.00	0.000	0.0233	2	0.00	0.000
763	0.0225	2	0.00	0.000	0.0311	3	0.00	0.000	0.0326	3	0.00	0.000
764	0.0236	2	0.00	0.000	0.0221	2	0.00	0.000	0.0229	2	0.00	0.000
765	0.0236	2	0.00	0.000	0.0113	1	0.00	-	0.0225	2	0.00	0.000
766	0.0233	2	0.00	0.000	0.0225	2	0.00	0.000	0.0225	2	0.00	0.000
767	0.0225	2	0.00	0.000	0.0229	2	0.00	0.000	0.0218	2	0.00	0.000

TABLE 7 (cont.).- Swept area, number of hauls and Yellowtail flounder mean catch (kg) and SD by stratum. Spanish Spring Surveys on NAFO Div. 3NO: 1995-2007. Swept area in square miles. n.s. means stratum not surveyed. 1995-2000 data are transformed C/V *Playa de Menduíña* data, and 2002-2007 data are original from R/V *Vizconde de Eza*. In 2001, there are data from the two vessels.

Stratum	2005				2006				2007			
	Swept area	Tow number	Y. flounder Mean catch	Y. flounder SD	Swept area	Tow number	Y. flounder Mean catch	Y. flounder SD	Swept area	Tow number	Y. flounder Mean catch	Y. flounder SD
353	0.0353	3	58.83	99.610	0.0371	3	71.98	122.954	0.0364	3	0.64	0.172
354	0.0353	3	0.21	0.188	0.0364	3	0.21	0.371	0.0364	3	0.16	0.283
355	0.0225	2	0.00	0.000	0.0248	2	0.00	0.000	0.0240	2	0.00	0.000
356	0.0233	2	0.00	0.000	0.0240	2	0.00	0.000	0.0240	2	0.00	0.000
357	0.0233	2	0.00	0.000	0.0244	2	0.00	0.000	0.0360	3	0.00	0.000
358	0.0349	3	0.00	0.000	0.0349	3	0.00	0.000	0.0368	3	0.00	0.000
359	0.0814	7	99.52	142.727	0.0975	8	169.33	359.779	0.0855	7	102.63	116.690
360	0.2325	20	342.14	223.566	0.2340	19	361.02	266.205	0.2378	20	349.70	307.902
374	0.0229	2	300.46	128.092	0.0236	2	610.03	73.518	0.0240	2	1057.60	455.094
375	0.0349	3	288.64	138.290	0.0364	3	287.65	109.715	0.0364	3	145.73	86.977
376	0.1174	10	500.53	238.908	0.1219	10	489.81	231.495	0.1185	10	460.24	203.990
377	0.0233	2	42.84	60.518	0.0236	2	6.09	8.605	0.0240	2	165.35	233.840
378	0.0225	2	0.00	0.000	0.0240	2	0.00	0.000	0.0233	2	0.00	0.000
379	0.0236	2	0.00	0.000	0.0236	2	0.00	0.000	0.0240	2	0.00	0.000
380	0.0229	2	0.00	0.000	0.0229	2	0.00	0.000	0.0240	2	0.00	0.000
381	0.0233	2	0.00	0.000	0.0229	2	0.00	0.000	0.0240	2	0.00	0.000
382	0.0458	4	0.00	0.000	0.0469	4	0.00	0.000	0.0484	4	0.00	0.000
721	0.0229	2	0.00	0.000	0.0236	2	0.00	0.000	0.0116	1	0.00	-
722	0.0233	2	0.00	0.000	0.0240	2	0.00	0.000	0.0225	2	0.00	0.000
723	0.0233	2	0.00	0.000	0.0236	2	0.18	0.247	0.0240	2	0.00	0.000
724	0.0225	2	0.00	0.000	0.0233	2	0.00	0.000	0.0233	2	0.00	0.000
725	0.0236	2	0.00	0.000	0.0233	2	0.00	0.000	0.0225	2	0.00	0.000
726	0.0113	1	0.00	-	0.0225	2	0.00	0.000	0.0229	2	0.00	0.000
727	0.0229	2	0.00	0.000	0.0225	2	0.00	0.000	0.0240	2	0.00	0.000
728	0.0109	1	0.00	-	0.0225	2	0.00	0.000	0.0225	2	0.00	0.000
752	0.0236	2	0.00	0.000	0.0225	2	0.00	0.000	0.0225	2	0.00	0.000
753	0.0225	2	0.00	0.000	0.0225	2	0.00	0.000	0.0225	2	0.00	0.000
754	0.0225	2	0.00	0.000	0.0225	2	0.00	0.000	0.0225	2	0.00	0.000
755	0.0450	4	0.00	0.000	0.0338	3	0.00	0.000	0.0338	3	0.00	0.000
756	0.0233	2	0.00	0.000	0.0229	2	0.00	0.000	0.0225	2	0.00	0.000
757	0.0225	2	0.00	0.000	0.0225	2	0.00	0.000	0.0229	2	0.00	0.000
758	0.0225	2	0.00	0.000	0.0225	2	0.00	0.000	0.0225	2	0.00	0.000
759	0.0229	2	0.00	0.000	0.0225	2	0.00	0.000	n.s.	n.s.	n.s.	n.s.
760	0.0229	2	0.35	0.488	0.0225	2	0.00	0.000	0.0233	2	0.00	0.000
761	0.0221	2	0.00	0.000	0.0233	2	0.00	0.000	0.0225	2	0.00	0.000
762	0.0225	2	0.00	0.000	0.0233	2	0.00	0.000	n.s.	n.s.	n.s.	n.s.
763	0.0334	3	0.00	0.000	0.0225	2	0.00	0.000	n.s.	n.s.	n.s.	n.s.
764	0.0233	2	0.00	0.000	0.0233	2	0.00	0.000	0.0225	2	0.00	0.000
765	0.0229	2	0.00	0.000	0.0236	2	0.00	0.000	0.0225	2	0.00	0.000
766	0.0229	2	0.00	0.000	0.0229	2	0.00	0.000	n.s.	n.s.	n.s.	n.s.
767	0.0113	1	0.00	-	0.0233	2	0.00	0.000	n.s.	n.s.	n.s.	n.s.

TABLE 8.- Stratified mean catches (Kg) by stratum and year and SD by year of Yellowtail flounder (1995-2007). n.s. means stratum not surveyed. 1995-2000 data are transformed C/V *Playa de Menduíña* data. 2002-2007 data are original from R/V *Vizconde de Eza*. In 2001, there are data from the two vessels.

Stratum	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007
353	1565.07	20142.03	3376.59	3288.11	40399.20	18255.85	16521.08	20208.63	2998.45	2364.96	15825.27	19363.88	173.06
354	438.70	0.00	346.30	299.00	20.56	439.52	83.64	41.00	0.00	151.29	52.07	52.64	40.18
355	n.s.	0.00	163.06	9.34	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
356	n.s.	0.00	15.24	0.00	0.00	0.00	0.24	0.00	0.00	0.00	0.00	0.00	0.00
357	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
358	0.00	30.09	4.21	0.00	0.00	0.00	0.00	0.00	0.00	57.45	0.00	0.00	0.00
359	567.78	385.71	34.00	72.73	143.12	994.90	597.82	44.91	0.00	10527.59	41896.12	71289.88	43209.03
360	56884.98	395448.50	225203.35	1040562.34	1517232.56	1088647.76	1493908.83	946847.84	1003413.43	1122077.77	952164.35	1004707.97	973222.06
374	0.00	0.00	0.00	9.54	15871.12	4379.59	51092.50	6856.85	3450.75	41400.44	64297.37	130545.35	226326.40
375	401.88	11218.18	54.37	3352.77	94076.82	41560.71	27190.33	13173.31	7709.95	147164.74	78220.54	77952.61	39493.73
376	46774.78	95247.02	216576.13	372549.36	735836.39	580653.95	591126.08	711849.08	522389.06	641736.71	667712.36	653412.94	613960.16
377	0.00	0.00	0.00	0.00	0.00	4.51	0.00	0.00	70.00	0.00	4283.75	608.50	16535.00
378	0.00	7.75	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
379	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
380	n.s.	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
381	n.s.	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
382	n.s.	0.00	0.00	0.00	0.00	0.00	5.16	0.00	0.00	0.00	0.00	0.00	0.00
721	n.s.	2.17	48.90	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
722	n.s.	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
723	n.s.	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	27.13	0.00
724	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	64.48	0.00	0.00	0.00	0.00
725	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
726	0.00	0.00	n.s.	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
727	n.s.	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
728	n.s.	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
752	n.s.	0.00	0.00	0.00	0.00	0.00	7.67	0.00	0.00	0.00	0.00	0.00	0.00
753	n.s.	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
754	n.s.	n.s.	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
755	n.s.	n.s.	n.s.	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
756	n.s.	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
757	n.s.	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
758	n.s.	n.s.	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
759	n.s.	n.s.	n.s.	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	n.s.
760	n.s.	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	53.13	0.00	0.00
761	n.s.	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
762	n.s.	n.s.	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	n.s.
763	n.s.	n.s.	n.s.	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	n.s.
764	n.s.	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
765	n.s.	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
766	n.s.	n.s.	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	n.s.
767	n.s.	n.s.	n.s.	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	n.s.
TOTAL (\bar{Y})	106633.19	522481.47	445822.16	1420143.19	2403579.77	1734936.80	2180533.35	1699021.61	1540096.13	1965480.95	1824504.95	1957960.89	1912959.62
S.D.	4.37	8.41	10.69	34.70	27.41	22.21	30.58	24.92	20.84	21.27	17.06	19.83	23.61

TABLE 9.- Survey estimates (by the swept area method) of Yellowtail flounder biomass (t) and SD by stratum and year on NAFO Div. 3NO. n.s. means stratum not surveyed. 1995-2000 data are transformed C/V *Playa de Menduña* data. 2002-2007 data are original from R/V *Vizconde de Eza*. In 2001, there are data from the two vessels.

Stratum	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007
353	133	1628	281	282	3367	1537	1452	1697	270	210	1347	1565	14
354	37	26	30	25	2	37	7	3	0	13	4	4	3
355	n.s.	2	14	0	0	0	0	0	0	0	0	0	0
356	n.s.	0	1	0	0	0	0	0	0	0	0	0	0
357	0	0	0	0	0	0	0	0	0	0	0	0	0
358	0	3	0	0	0	0	0	0	0	5	0	0	0
359	49	35	3	6	12	85	52	4	0	931	3604	5849	3538
360	4950	32593	19198	89742	123989	90863	123341	82622	89057	97150	81907	81579	81869
374	0	0	0	0	1302	365	4258	596	307	3561	5622	11051	18861
375	36	981	5	291	7964	3410	2417	1121	701	13081	6729	6429	3257
376	4059	8082	19160	32255	60376	48388	51175	62443	46435	55026	56887	53613	51811
377	0	0	0	0	0	0	0	0	6	0	368	52	1378
378	0	1	0	0	0	0	0	0	0	0	0	0	0
379	0	0	0	0	0	0	0	0	0	0	0	0	0
380	n.s.	0	0	0	0	0	0	0	0	0	0	0	0
381	n.s.	0	0	0	0	0	0	0	0	0	0	0	0
382	n.s.	0	0	0	0	0	0	0	0	0	0	0	0
721	n.s.	0	4	0	0	0	0	0	0	0	0	0	0
722	n.s.	0	0	0	0	0	0	0	0	0	0	0	0
723	n.s.	0	0	0	0	0	0	0	0	0	0	0	2
724	0	0	0	0	0	0	0	0	0	0	0	0	0
725	0	0	0	0	0	0	0	0	0	0	0	0	0
726	0	0	n.s.	0	0	0	0	0	0	0	0	0	0
727	n.s.	0	0	0	0	0	0	0	0	0	0	0	0
728	n.s.	0	0	0	0	0	0	0	0	0	0	0	0
752	n.s.	0	0	0	0	0	1	0	0	0	0	0	0
753	n.s.	0	0	0	0	0	0	0	0	0	0	0	0
754	n.s.	n.s.	0	0	0	0	0	0	0	0	0	0	0
755	n.s.	n.s.	n.s.	0	0	0	0	0	0	0	0	0	0
756	n.s.	0	0	0	0	0	0	0	0	0	0	0	0
757	n.s.	0	0	0	0	0	0	0	0	0	0	0	0
758	n.s.	n.s.	0	0	0	0	0	0	0	0	0	0	0
759	n.s.	n.s.	n.s.	0	0	0	0	0	0	0	0	0	n.s.
760	n.s.	0	0	0	0	0	0	0	0	0	5	0	0
761	n.s.	0	0	0	0	0	0	0	0	0	0	0	0
762	n.s.	n.s.	0	0	0	0	0	0	0	0	0	0	n.s.
763	n.s.	n.s.	n.s.	0	0	0	0	0	0	0	0	0	n.s.
764	n.s.	0	0	0	0	0	0	0	0	0	0	0	0
765	n.s.	0	0	0	0	0	0	0	0	0	0	0	0
766	n.s.	n.s.	0	0	0	0	0	0	0	0	0	0	n.s.
767	n.s.	n.s.	n.s.	0	0	0	0	0	0	0	0	0	n.s.
TOTAL	9264	43349	38697	122601	197012	144685	182704	148487	136775	169978	156472	160145	160731
S.D.	2484	6032	8527	31359	22938	19097	25847	23368	19287	18869	15271	16458	18852

TABLE 10.- Length weight relationships in the calculation of Yellowtail flounder biomass. The equation is $Weight = a(l + 0.5)^b$ ^b Spanish Spring Surveys on NAFO Div. 3NO: 1995-2007. To calculate the parameters for the indeterminate individuals, we used the total data (males + females + indeterminate individuals). *E* means Error.

		1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007
Males	a	0.0079 E = 0.2653	0.008 E = 0.0907	0.0081 E = 0.0936	0.0075 E = 0.1034	0.0084 E = 0.2119	0.0036 E = 0.0994	0.0081 E = 0.1248	0.0075 E = 0.0729	0.0121 E = 0.1109	0.0053 E = 0.1352	0.0027 E = 0.0882	0.0096 E = 0.0825	0.0074 E = 0.0655
	b	3.0416 E = 0.0799	3.0342 E = 0.0269	3.0197 E = 0.0281	3.0376 E = 0.0313	3.0098 E = 0.0610	3.2403 E = 0.0300	3.0176 E = 0.0374	3.0271 E = 0.0226	2.8978 E = 0.0348	3.1236 E = 0.0419	3.3274 E = 0.0274	2.9463 E = 0.0263	3.0190 E = 0.0201
		R2 = 0.984 N=137	R2 = 0.998 N=430	R2 = 0.997 N=556	R2 = 0.997 N=523	R2 = 0.994 N=56	R2 = 0.997 N=270	R2 = 0.995 N=271	R2 = 0.998 N=274	R2 = 0.995 N=316	R2 = 0.995 N=411	R2 = 0.997 N=311	R2 = 0.999 N=371	R2 = 0.999 N= 578
Females	a	0.0063 E = 0.1251	0.0056 E = 0.0632	0.0056 E = 0.0517	0.0067 E = 0.1290	0.0073 E = 0.2607	0.0026 E = 0.0914	0.006 E = 0.0841	0.0051 E = 0.0901	0.0061 E = 0.0995	0.0047 E = 0.0630	0.0027 E = 0.0634	0.0069 E = 0.1137	0.0043 E = 0.1973
	b	3.1083 E = 0.0367	3.1496 E = 0.0179	3.1382 E = 0.0152	3.0788 E = 0.0384	3.0577 E = 0.0739	3.3504 E = 0.0267	3.1122 E = 0.0249	3.1448 E = 0.0274	3.1079 E = 0.0307	3.1768 E = 0.0191	3.329 E = 0.0177	3.0584 E = 0.0347	3.1915 E = 0.0582
		R2 = 0.995 N=246	R2 = 0.999 N=735	R2 = 0.999 N=910	R2 = 0.994 N=682	R2 = 0.989 N=62	R2 = 0.998 N=344	R2 = 0.997 N=378	R2 = 0.997 N=343	R2 = 0.996 N=513	R2 = 0.999 N=547	R2 = 0.998 N=569	R2 = 0.997 N=507	R2 = 0.987 N= 731
Indet.	a	0.0088 E = 0.1109	0.006 E = 0.0656	0.006 E = 0.0580	0.0071 E = 0.0652	0.0078 E = 0.1656	0.0026 E = 0.0835	0.0092 E = 0.1075	0.006 E = 0.0402	0.0069 E = 0.1095	0.004 E = 0.0608	0.0025 E = 0.0523	0.0102 E = 0.1453	0.0068 E = 0.1078
	b	3.0144 E = 0.0330	3.1285 E = 0.0188	3.1166 E = 0.0171	3.0614 E = 0.0195	3.0406 E = 0.0477	3.3423 E = 0.0245	2.9883 E = 0.0329	3.0977 E = 0.0123	3.0737 E = 0.0337	3.2137 E = 0.0186	3.3552 E = 0.0148	2.9471 E = 0.0448	3.0606 E = 0.0327
		R2 = 0.996 N=391	R2 = 0.999 N=1181	R2 = 0.999 N=1466	R2 = 0.994 N=1211	R2 = 0.995 N=118	R2 = 0.999 N=614	R2 = 0.994 N=703	R2 = 0.999 N=620	R2 = 0.995 N=833	R2 = 0.999 N=969	R2 = 0.995 N=884	R2 = 0.995 N=887	R2 = 0.995 N= 1312

TABLE 11.- Yellowtail flounder length distribution. Estimated numbers per haul mean catches. Spanish Spring Survey on NAFO 3NO: 1995-2007. Indet. means indeterminate. 1995-2000 data are transformed C/V *Playa de Mendoña* data. 2002-2007 data are original R/V *Vizconde de Eza* data. In 2001, there are data from the two vessels. (*) indicates untransformed data.

Length (cm.)	1995				1996				1997				1998				
	Males	Females	Indet.	Total	Males	Females	Indet.	Total	Males	Females	Indet.	Total	Males	Females	Indet.	Total	
4	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	
6	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	
8	0.000	0.000	0.185	0.185	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	
10	0.000	0.000	0.456	0.456	0.000	0.000	0.498	0.498	0.000	0.000	0.000	0.000	0.000	0.000	0.071	0.071	
12	0.103	0.870	2.350	3.323	0.000	0.000	0.877	0.877	1.356	0.560	0.000	1.916	0.000	0.000	1.538	1.538	
14	1.557	1.441	2.842	5.840	0.000	0.048	2.711	2.759	0.155	0.819	0.000	0.974	0.121	0.157	0.000	0.278	
16	2.045	3.581	0.277	5.903	0.288	3.152	5.167	8.607	2.947	1.811	0.000	4.758	1.500	1.535	0.000	3.034	
18	2.649	3.358	0.031	6.038	2.334	15.279	3.167	20.780	5.076	4.415	0.000	9.491	8.365	5.129	0.000	13.495	
20	2.984	3.212	0.000	6.196	5.319	26.981	0.750	33.050	13.857	15.055	0.000	28.912	8.974	10.166	0.000	19.140	
22	4.807	6.015	0.000	10.823	8.522	32.231	0.065	40.818	28.296	23.048	0.000	51.345	25.957	20.452	0.000	46.409	
24	4.810	6.082	0.000	10.892	10.962	32.203	0.000	43.165	31.348	27.786	0.000	59.134	44.950	37.421	0.000	82.371	
26	2.340	2.446	0.000	4.786	9.552	16.875	0.000	26.427	24.015	26.970	0.000	50.985	72.376	60.520	0.000	132.896	
28	2.704	2.544	0.000	5.248	9.151	11.591	0.000	20.742	13.921	21.248	0.000	35.169	57.459	62.401	0.000	119.861	
30	2.588	4.738	0.000	7.325	7.206	9.915	0.000	17.122	6.159	10.349	0.000	16.508	32.472	56.275	0.000	88.747	
32	1.664	4.451	0.000	6.115	6.379	6.166	0.000	12.545	3.761	5.090	0.000	8.851	15.566	32.294	0.000	47.859	
34	1.290	3.070	0.000	4.361	5.565	6.928	0.000	12.493	1.894	2.803	0.000	4.698	5.840	22.613	0.000	28.453	
36	0.661	1.797	0.000	2.459	4.143	9.508	0.000	13.651	1.195	2.683	0.000	3.878	2.638	12.385	0.000	15.023	
38	0.475	1.395	0.000	1.870	2.083	6.687	0.000	8.771	0.485	2.407	0.000	2.892	2.475	8.439	0.000	10.914	
40	0.373	0.937	0.000	1.310	0.724	5.018	0.000	5.742	0.245	1.723	0.000	1.968	1.060	7.705	0.000	8.765	
42	0.059	0.588	0.000	0.647	0.694	3.305	0.000	4.000	0.099	0.801	0.000	0.899	0.065	3.260	0.000	3.324	
44	0.004	0.471	0.000	0.475	0.087	1.550	0.000	1.637	0.031	0.281	0.000	0.311	0.008	1.729	0.000	1.737	
46	0.004	0.081	0.000	0.085	0.081	0.969	0.000	1.050	0.006	0.044	0.000	0.049	0.000	0.600	0.000	0.600	
48	0.000	0.191	0.000	0.191	0.018	0.286	0.000	0.304	0.000	0.052	0.000	0.052	0.004	0.273	0.000	0.277	
50	0.000	0.027	0.000	0.027	0.000	0.045	0.000	0.045	0.000	0.018	0.000	0.018	0.000	0.000	0.000	0.000	
52	0.000	0.052	0.000	0.052	0.000	0.053	0.000	0.053	0.000	0.018	0.000	0.018	0.000	0.000	0.000	0.000	
54	0.000	0.005	0.000	0.005	0.000	0.039	0.000	0.039	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	
56	0.000	0.005	0.000	0.005	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	
Total	31.117	47.358	6.141	84.616	73.109	188.829	13.235	275.173	134.845	147.982	0.000	282.827	279.828	343.354	1.609	624.791	
Nº samples (*):					43				33				54				48
Nº Ind. (*):	1876	3003	81	4960	1837	4584	249	6670	3635	4469	0	8104	2848	3693	3	6544	
Sampled catch:					375				532				585				536
Range (*):					9-56				10-55				12-53				11-49
Total catch:					2731				5721				4956				12231
Total hauls (*):					77				112				128				124

TABLE 11 (cont.).- Yellowtail flounder length distribution. Estimated numbers per haul mean catches. Spanish Spring Survey on NAFO 3NO: 1995-2007. Indet. means indeterminate. 1995-2000 data are transformed C/V *Playa de Menduíña* data. 2002-2007 data are original R/V *Vizconde de Eza* data. In 2001, there are data from the two vessels. (*) indicates untransformed data.

Length (cm.)	1999				2000				2001			
	Males	Females	Indet.	Total	Males	Females	Indet.	Total	Males	Females	Indet.	Total
4	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
6	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.325	0.325
8	0.000	0.000	1.516	1.516	0.000	0.000	0.000	0.000	0.000	0.000	1.937	1.937
10	5.154	3.352	2.960	11.465	0.000	0.793	0.000	0.793	0.104	0.356	1.850	2.310
12	12.807	8.911	0.000	21.718	3.716	1.266	0.000	4.982	0.320	1.239	1.187	2.746
14	19.227	16.710	0.000	35.938	7.773	11.915	0.000	19.687	0.952	1.477	1.114	3.543
16	13.999	15.356	0.000	29.355	10.311	10.506	0.000	20.817	3.575	4.509	0.412	8.497
18	8.893	10.757	0.000	19.650	14.266	16.475	0.000	30.741	10.107	10.530	0.149	20.786
20	14.809	10.199	0.000	25.008	16.177	19.576	0.000	35.753	17.815	24.898	0.000	42.713
22	33.285	22.789	0.000	56.073	17.231	18.660	0.000	35.891	21.299	29.178	0.000	50.477
24	61.756	39.009	0.000	100.765	21.395	20.983	0.000	42.378	24.254	23.840	0.000	48.094
26	98.561	59.521	0.000	158.083	48.000	33.100	0.000	81.100	28.911	24.809	0.000	53.720
28	107.816	84.193	0.000	192.009	67.229	39.182	0.000	106.412	58.237	33.305	0.000	91.542
30	72.947	92.236	0.000	165.183	64.336	44.684	0.000	109.020	72.412	45.107	0.000	117.519
32	28.850	75.169	0.000	104.018	36.450	53.416	0.000	89.865	49.179	59.052	0.000	108.232
34	15.810	43.595	0.000	59.405	12.695	39.970	0.000	52.665	22.267	64.772	0.000	87.039
36	9.185	24.775	0.000	33.960	6.653	25.712	0.000	32.365	8.702	46.598	0.000	55.300
38	3.658	14.964	0.000	18.623	3.526	15.747	0.000	19.274	6.293	30.315	0.000	36.608
40	1.466	8.582	0.000	10.049	1.996	10.642	0.000	12.638	2.145	12.925	0.000	15.070
42	0.262	5.318	0.000	5.580	0.286	6.803	0.000	7.089	0.857	7.788	0.000	8.645
44	0.111	2.620	0.000	2.731	0.013	4.005	0.000	4.018	0.614	4.596	0.000	5.210
46	0.028	0.988	0.000	1.016	0.000	1.806	0.000	1.806	0.221	1.968	0.000	2.190
48	0.096	0.486	0.000	0.582	0.003	0.845	0.000	0.848	0.000	0.775	0.000	0.775
50	0.000	0.140	0.000	0.140	0.000	0.246	0.000	0.246	0.000	0.242	0.000	0.242
52	0.000	0.032	0.000	0.032	0.000	0.000	0.000	0.000	0.000	0.051	0.000	0.051
54	0.000	0.000	0.000	0.000	0.000	0.033	0.000	0.033	0.000	0.000	0.000	0.000
56	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Total	508.721	539.702	4.475	1052.898	332.057	376.364	0.000	708.421	328.265	428.326	6.975	763.567
Nº samples (*):				39				42				43
Nº Ind. (*):	4616	5076	6	9698	3323	4100	0	7423	3358	4684	80	8122
Sampled catch:				796				717				2298
Range (*):				8-52				11-54				6-53
Total catch:				17169				12742				16141
Total hauls (*):				114				118				123

TABLE 11 (cont.).- Yellowtail flounder length distribution. Estimated numbers per haul mean catches. Spanish Spring Survey on NAFO 3NO: 1995-2007. Indet. means indeterminate. 1995-2000 data are transformed C/V *Playa de Menduíña* data. 2002-2007 data are original R/V *Vizconde de Eza* data. In 2001, there are data from the two vessels. (*) indicates untransformed data.

Length (cm.)	2002				2003				2004			
	Males	Females	Indet.	Total	Males	Females	Indet.	Total	Males	Females	Indet.	Total
4	0.000	0.000	0.000	0.000	0.000	0.000	0.009	0.009	0.000	0.000	0.116	0.116
6	0.000	0.141	0.475	0.616	0.000	0.107	0.297	0.404	0.000	0.000	0.337	0.337
8	0.349	0.639	0.332	1.321	0.036	0.121	0.274	0.431	0.109	0.049	0.741	0.899
10	1.315	0.712	0.000	2.027	0.847	0.572	0.140	1.559	0.528	0.637	0.000	1.165
12	0.620	0.675	0.000	1.295	0.969	1.205	0.000	2.174	2.005	1.577	0.000	3.582
14	1.544	1.064	0.000	2.608	0.977	0.869	0.000	1.846	3.503	2.632	0.000	6.135
16	1.889	2.134	0.000	4.023	0.946	0.289	0.000	1.234	4.580	3.608	0.000	8.188
18	3.180	2.479	0.000	5.660	1.665	1.689	0.000	3.355	4.649	3.543	0.000	8.192
20	7.908	6.122	0.000	14.030	1.695	2.233	0.000	3.928	5.414	6.205	0.000	11.619
22	16.552	12.664	0.000	29.217	4.214	4.602	0.000	8.817	5.563	5.757	0.000	11.321
24	21.724	22.245	0.000	43.968	11.364	8.741	0.000	20.105	8.232	7.732	0.000	15.964
26	27.246	24.307	0.000	51.553	27.765	19.581	0.000	47.347	25.572	16.572	0.000	42.145
28	40.151	22.443	0.000	62.594	37.413	29.153	0.000	66.566	57.974	27.637	0.000	85.611
30	57.549	34.445	0.000	91.994	52.296	29.328	0.000	81.624	87.376	52.285	0.000	139.661
32	46.938	50.680	0.000	97.618	45.761	40.076	0.000	85.836	74.712	58.683	0.000	133.396
34	18.047	57.599	0.000	75.646	19.769	52.100	0.000	71.869	30.847	58.596	0.000	89.443
36	7.014	45.699	0.000	52.713	6.757	39.555	0.000	46.312	7.531	46.290	0.000	53.820
38	2.651	25.514	0.000	28.165	2.130	23.649	0.000	25.779	2.056	26.594	0.000	28.650
40	1.183	12.427	0.000	13.610	0.832	9.444	0.000	10.276	1.716	10.932	0.000	12.648
42	0.616	6.257	0.000	6.873	0.256	3.895	0.000	4.151	0.514	3.725	0.000	4.240
44	0.042	2.690	0.000	2.732	0.268	2.432	0.000	2.700	0.028	2.033	0.000	2.061
46	0.024	1.150	0.000	1.174	0.000	1.113	0.000	1.113	0.000	0.575	0.000	0.575
48	0.000	0.818	0.000	0.818	0.000	0.525	0.000	0.525	0.000	0.303	0.000	0.303
50	0.020	0.149	0.000	0.169	0.000	0.202	0.000	0.202	0.000	0.009	0.000	0.009
52	0.000	0.038	0.000	0.038	0.000	0.009	0.000	0.009	0.000	0.055	0.000	0.055
54	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
56	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Total	256.565	333.090	0.807	590.462	215.959	271.489	0.721	488.169	322.910	336.032	1.193	660.136
Nº samples (*):					43				37			45
Nº Ind. (*):	3419	4576	7	8002	2424	3254	12	5690	3703	4234	16	7953
Sampled catch:					2269				1864			2587
Range (*):					6-52				5-52			5-53
Total catch:					14385				11280			15117
Total hauls (*):					125				118			120

TABLE 11 (cont.).- Yellowtail flounder length distribution. Estimated numbers per haul mean catches. Spanish Spring Survey on NAFO 3NO: 1995-2007. Indet. means indeterminate. 1995-2000 data are transformed C/V *Playa de Menduíña* data. 2002-2007 data are original R/V *Vizconde de Eza* data. In 2001, there are data from the two vessels. (*) indicates untransformed data.

Length (cm.)	2005				2006				2007			
	Males	Females	Indet.	Total	Males	Females	Indet.	Total	Males	Females	Indet.	Total
4	0.000	0.000	0.000	0.000	0.060	0.000	0.000	0.060	0.000	0.000	0.000	0.000
6	0.000	0.013	0.192	0.205	0.000	0.000	0.079	0.079	0.000	0.000	0.103	0.103
8	0.269	0.018	0.054	0.341	0.187	0.162	0.245	0.594	0.000	0.000	0.000	0.000
10	1.725	0.467	0.051	2.243	0.686	0.384	0.276	1.346	0.041	0.059	0.000	0.101
12	2.353	1.877	0.000	4.229	2.026	1.734	0.000	3.760	0.536	0.449	0.000	0.985
14	4.728	3.053	0.000	7.780	3.645	3.862	0.000	7.507	1.148	0.578	0.000	1.725
16	4.674	3.630	0.000	8.304	5.776	6.009	0.000	11.785	2.222	2.551	0.000	4.773
18	3.334	3.348	0.000	6.682	5.989	5.547	0.000	11.536	5.728	4.614	0.000	10.342
20	4.905	4.847	0.000	9.752	9.721	8.196	0.000	17.917	9.024	7.293	0.000	16.317
22	8.934	6.836	0.000	15.770	10.735	10.545	0.000	21.280	13.286	14.190	0.000	27.476
24	8.930	7.162	0.000	16.092	11.073	12.977	0.000	24.050	17.380	19.046	0.000	36.426
26	15.997	8.451	0.000	24.447	13.117	13.439	0.000	26.556	20.689	18.113	0.000	38.802
28	34.840	17.504	0.000	52.344	26.251	15.412	0.000	41.663	35.157	19.170	0.000	54.327
30	75.001	34.103	0.000	109.105	64.180	25.059	0.000	89.238	75.144	25.235	0.000	100.379
32	70.556	58.866	0.000	129.423	74.126	52.415	0.000	126.541	76.329	50.253	0.000	126.582
34	28.072	62.961	0.000	91.032	38.379	67.737	0.000	106.116	42.232	68.548	0.000	110.780
36	8.105	48.672	0.000	56.777	11.021	63.706	0.000	74.727	12.733	61.691	0.000	74.424
38	1.965	26.547	0.000	28.512	3.046	39.877	0.000	42.923	3.973	41.839	0.000	45.812
40	0.908	11.697	0.000	12.606	0.981	17.493	0.000	18.474	1.430	20.920	0.000	22.350
42	0.172	4.746	0.000	4.918	0.081	5.709	0.000	5.789	0.213	6.891	0.000	7.104
44	0.050	2.020	0.000	2.070	0.072	2.190	0.000	2.262	0.000	2.454	0.000	2.454
46	0.000	1.128	0.000	1.128	0.000	1.341	0.000	1.341	0.071	1.043	0.000	1.114
48	0.000	0.200	0.000	0.200	0.000	0.560	0.000	0.560	0.000	0.367	0.000	0.367
50	0.000	0.030	0.000	0.030	0.000	0.231	0.000	0.231	0.000	0.107	0.000	0.107
52	0.000	0.000	0.000	0.000	0.000	0.012	0.000	0.012	0.000	0.120	0.000	0.120
54	0.000	0.079	0.000	0.079	0.000	0.091	0.000	0.091	0.000	0.000	0.000	0.000
56	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Total	275.518	308.254	0.297	584.069	281.150	354.688	0.601	636.440	317.336	365.532	0.103	682.971
Nº samples (*):				48				45				47
Nº Ind. (*):	4790	6556	6	11352	4404	6012	10	10426	5083	5533	1	10617
Sampled catch:				3784				3407				2761
Range (*):				6-55				5-54				7-52
Total catch:				14275				15424				15200
Total hauls (*):				119				120				110




FIGURE 1.- Atlantic cod stratified mean catches in Kg and \pm SD by year. Spanish Spring surveys on NAFO Div. 3NO: 1997-2007 (1997-2000 transformed data from C/V *Playa de Mendoña*; 2002-2007 original data from R/V *Vizconde de Eza*. In 2001, there are data from the two vessels).




FIGURE 2.- Atlantic cod biomass calculated by the swept area method in tons and \pm SD by year. Spanish Spring surveys on NAFO Div. 3NO: 1997-2007 (1997-2000 transformed data from C/V *Playa de Mendoña*; 2002-2007 original data from R/V *Vizconde de Eza*. In 2001, there are data from the two vessels).




FIGURE 3.- Atlantic cod length distribution (cm) on NAFO 3NO: 1997-2007 . Mean catches per tow numbers. 1997-2000 data are transformed data from C/V *Playa de Menduña*, and 2002-2007 data are original from R/V *Vizconde de Eza*. In 2001, there are data from the two vessels.




FIGURE 4.- Atlantic cod mean catches per tow length distribution (cm) on NAFO 3NO: 1997-2007.




FIGURE 5.- Yellowtail flounder stratified mean catches in Kg and \pm SD by year. Spanish Spring surveys on NAFO Div. 3NO: 1995-2007 (1995-2000 transformed data from C/V *Playa de Mendoña*; 2002-2007 original data from R/V *Vizconde de Eza*. In 2001, there are data from the two vessels).




FIGURE 6.- Yellowtail flounder biomass calculated by the swept area method in tons and \pm SD by year. Spanish Spring surveys on NAFO Div. 3NO: 1995-2007 (1995-2000 transformed data from C/V *Playa de Mendoña*; 2002-2007 original data from R/V *Vizconde de Eza*. In 2001, there are data from the two vessels).




FIGURE 7.- Yellowtail flounder length distribution (cm) on NAFO 3NO: 1995-2007. Mean catches per tow numbers. 1995-2000 data are transformed data from C/V *Playa de Menduña*, and 2002-2007 data are original from R/V *Vizconde de Eza*. In 2001, there are data from the two vessels




FIGURE 7 (Cont.).- Yellowtail flounder length distribution (cm) on NAFO 3NO: 1995-2007. Mean catches per tow numbers. 1995-2000 data are transformed data from C/V *Playa de Menduña*, and 2002-2007 data are original from R/V *Vizconde de Eza*. In 2001, there are data from the two vessels




FIGURE 8.- Yellowtail flounder mean catches per tow length distribution (cm) on NAFO 3NO: 1995-2007.