

NOT TO BE CITED WITHOUT PRIOR
REFERENCE TO THE AUTHOR(S)

Northwest Atlantic



Fisheries Organization

Serial No. N5546

NAFO SCR Doc. 08/44

SCIENTIFIC COUNCIL MEETING – JUNE 2008

Distribution and Abundance of Yellowtail Flounder (*Limanda ferruginea*) on the Grand Bank, NAFO Divisions 3LNO, from Canadian Bottom Trawl Survey Estimates from 1984-2007

by

D. Maddock Parsons and Bill Brodie

Northwest Atlantic Fisheries Centre, Science Branch
Department of Fisheries and Oceans, 80 East White Hills Road
St. John's, Newfoundland A1C 5X1

Abstract

Abundance and biomass indices of Grand Bank yellowtail flounder in NAFO Divisions 3LNO were derived from annual multi-species, random-stratified bottom trawl surveys conducted by Canada during the spring of 1984-2007 and during the fall from 1990 to 2007. The majority of the stock is found in depths less than 93 m and in Div. 3NO. Stock size and geographical range of yellowtail flounder declined from the mid-1980s to the mid-1990s, but recent surveys show the stock size has increased dramatically and has again expanded its northward range to re-occupy habitats on the northern Grand Bank. The 2006 spring estimate of biomass is the highest in the time series (504 kt) and the 2007 spring estimate (443 kt), is the second highest in the series. Biomass in the fall of 2006 is estimated to be 306 kt, a decrease from the 2005 value, but the 2007 estimate is the highest in the series at 482 kt.

Introduction

Annual multi-species, stratified-random bottom trawl surveys have been conducted by the Newfoundland region of the Canadian Dept of Fisheries and Oceans on the Grand Bank, in Div. 3LNO, during the spring (April-June) of each year since 1971. Since 1990, a second series of surveys has been carried out on the Grand Bank during the fall period, from October to December. However, since 1971 there have been two changes in survey gears and only one set of conversion factors has been developed for the 1984-1995 time series. Consequently only data from 1984 to 2007 will be presented here.

From both the spring and fall surveys, swept area abundance and biomass estimates are derived for yellowtail flounder (*Limanda ferruginea*) and serve as fishery-independent indices of stock size. Because catchability of the standard survey trawl is unknown and assumed to be ≤ 1.0 (Walsh, 1996), the indices are considered to be relative estimates of stock size.

The purpose of this paper is to update the results of these annual surveys. The last review of the survey results in any detail took place in 2006 (Walsh *et al.*, 2006) and because this stock is on a two year cycle for full assessment, attention will focused on monitoring annual changes in stock size and recent changes in stock size and temporal and spatial patterns of distribution since 2006 in relation to historical patterns.

Materials and Methods

Survey design

The stratification scheme is based on depth and shown in Fig. 1 (see Doubleday, 1981, for a review of procedures). The timing of the spring surveys, the frequency of fishing sets in the inshore strata (beginning in 1997) and the range of depths surveyed are shown in Table 1.

The 1984-2007 spring and the 1990-1994 fall surveys both covered depths from 45 to 731 m. Beginning in the fall of 1995 with the use of the new Campelen survey trawl, the coverage of the fall surveys extended to 1 500m. Due to mechanical problems with the CCG *Teleost* survey vessel, only sets in the deepwater strata of Div. 3L were fished in 1995 (Table 2). Surveys in 1999, and 2004-2007 had poor coverage in deep water strata (Table 2) (see Brodie and Stansbury, 2007) however, this is thought to have negligible effect on estimation of the relative abundance and biomass of yellowtail flounder, in most years, because the stock is found almost exclusively in depths less than 93 m. Nevertheless, the inclusion of these deepwater sets does lower the overall mean catch per tow by Division in those years. In the 2006 spring survey, there were also fewer sets in some strata in 3N, and stratum 373 was not surveyed. In recent years, this stratum contributed significant biomass to the total index and missing this stratum, and coupled with reduced coverage in other important strata, the 2006 estimates of abundance and biomass may not be comparable to estimates in other years. In addition, in years, 1995, 2002-2005, some northern portions of the surveys have overlapped into January of the following calendar year due to mechanical problems with the survey vessels. However, these delays are not expected to affect yellowtail flounder estimates because of its shallow water distribution in the southern section of the survey area.

Survey gears and vessels

From 1971 to 1982 the surveys of the Grand Bank were conducted by the 54 m side trawler, the FRV A. T. Cameron (ATC) using a two bridle Yankee 41.5 otter trawl rigged with rubber disk footgear. In 1983, this trawl was replaced by the three bridle Engel 145 Hi-Lift otter trawl rigged with large steel bobbin footgear and, at the same time, the A.T. Cameron was replaced by the 50 m stern trawler, the CCGS Wilfred Templeman (WT). Occasionally the W. Templeman's sister ship, the CCGS Alfred Needler (AN) took part in the surveys. In 1995, the old standard Engel trawl was replaced by a three bridle Campelen 1800 shrimp trawl rigged with 35 cm diameter rockhopper footgear. The Yankee and the Engel trawls were both towed at 3.5 kts, while the Campelen is towed at 3.0 kts (McCallum and Walsh, 1996). The Campelen trawl surveys of the Grand Bank began in the fall of 1995 aboard the CCGS Wilfred Templeman. The Campelen trawl also replaced the Yankee 41 shrimp trawl used in the annual fall juvenile groundfish surveys from 1985-94 (McCallum and Walsh, 1996). Beginning in the fall of 1996, the 63 m stern trawler, CCGS Teleost, began fishing mostly the deepwater survey sets of the annual fall surveys beyond 731 m in Div. 3LNO; however, shallower sets have also been fished when necessary (Table 2). In addition, the CCGS Alfred Needler has taken part in the fall surveys in 1996 and 2001. The Campelen trawl onboard the 2 other survey vessels is identical in construction and rigging as the one on the Wilfred Templeman. Since 1993, the geometry and performance of all bottom trawl surveys have been monitored by Scanmar trawl mounted acoustic instrumentation (Walsh and McCallum, 1995; McCallum and Walsh, 2001).

Time series

Conversion factors have been derived from comparative fishing trials to convert the 1984-95 spring and 1990-94 fall Engel trawl survey data into Campelen trawl units and were presented in Walsh *et al.* (1998a, 1998b). Survey data from 1971-82 time period has not been converted to Campelen trawl units and the unconverted time series can be found in the 1997 assessment paper (see Walsh *et al.*, 1997). Conversion factors into Campelen trawl units for yellowtail flounder have also been derived for the 1985-94 late summer-early fall juvenile groundfish series and the abundance and biomass data are found in a 2005 NAFO SCR paper (see Walsh, 2005). However, additional conversions of the database will be needed and consequently only annual spring and fall survey data from 1984 onward will be reported here.

Fishing and catch protocols

The Campelen carries out 15 minute tows using a towing speed of 3.0 kts and covers an average tow distance of 0.75 nautical miles. The catches are standardized to distance towed. The average wingspread used in estimating

swept area abundance indices is 16.84 m and the average swept area is estimated to be 24 950 m². After each set, all species in the catch are separated, counted and weighed. From each haul, the total catch or a sub-sample is taken to collect biological data on size, age, maturity and feeding for all commercial species.

Results

Spring Groundfish Surveys 1984-2007

Abundance and biomass trends:

Tables 4 to 9 give the survey catch rates by division in the form of stratified mean number and weight-per-tow by stratum. Tables 10-15 show abundance and biomass per stratum, along with confidence limits, for stock size in Div. 3L, 3N, and 3O, respectively, and table 28 gives combined estimates for Div. 3LNO from 1984-2007. Figure 2 show plots of the abundance and biomass estimates, as well as mean number and weight per tow, of surveys from 1984-2007. The high 1999 survey estimates point to a 'year effect' (Walsh *et al.*, 2000; STACFIS, 2000). Table 3 identifies large fishing sets that may contribute to variation seen around some of the estimates of stock size in a given year. Problems with the survey vessel in spring of 2006 resulted in reduced coverage, particularly in deep water of 3N and 3O. Only a few strata that were missed have had significant yellowtail catch in the past (notably strata 373) and it is felt that coverage of the area occupied by yellowtail flounder was sufficient to include estimates from this survey in the series.

In Div. 3L, there was a continuous decline in abundance and biomass from 1985 to "0.0 t" in 1995 (Tables 10, 13 and 28; Fig. 2). From 1996 to 1998, the stock showed a marginal increase to stabilize at an average biomass level of 500 t and then increased (by 5550%) to a level of 28 000 t in 1999 (Table 13; Fig. 2). From 2000-2002 the abundance and biomass declined dramatically and by 2002 the biomass was 600 t (1.6 million fish). From 2002-2006 the abundance and biomass have been variable but showed an overall increase to the highest estimates in the time series in 2006 at 251.5 million fish and 85.7 t biomass. In 2007, biomass and abundance estimates declined marginally, but remain second highest in the series. When the estimates are high most of the yellowtail flounder are generally found in stratum 363 and stratum 372.

In Div. 3N, in general, the majority of the stock was distributed in and around the Southeast Shoal area (strata 375, 376, 360 and 361 in Fig. 1), although in recent surveys, the abundance and biomass has been increasing in strata to the north of the Shoal, in particular strata 362 and 373 (Tables 11 and 14). The biomass index declined gradually from 168 000 t (435 million fish) in 1984 to 46 000 t (135 million fish) by 1994, a decline of 73% (Fig. 2). For the same period, the high abundance estimate of 478 million fish in 1989, was mainly due to the strong 1985 and 1986 year-classes which was not reflected in the biomass estimate for that survey. After a slight increase from 1994 to 1995, the survey biomass in 1996 jumped by 80% to 104 000 t (475 million fish) followed by a continued increase to a high of 238 000 t (965 million fish) in 1999 (Fig. 2). Since 1999, the survey abundance and biomass estimates have been variable, but increasing. The 2006 spring abundance and biomass estimates were the second highest and highest in the series, respectively (319 000 t; 1 billion fish), although stratum 373 was not surveyed in 2006, and fewer sets were conducted in several other strata, making it difficult to compare the index in 2006 to other years. In 2007 estimates remained high at 293 000 t and 954 million fish.

Variability around the 2001 estimate of abundance and biomass, both the highest estimates in their respective time series, was quite high (Fig. 2). Some recent estimates also have relatively wide confident limits. Some of this variation may be due to the several high catches. Over ½ of the estimate comes from sets in strata 360, 376, and 373. Similar to the 1999 survey where large catches were prevalent in strata 360, 361 362, 375 and 376, ten sets in 2001 had catches greater than 1 000 fish (5 in excess of 300 kg), in which one set caught 2400 (476 kg) in stratum 376 and another in stratum 373 caught 4800 (1.6 t) yellowtail flounder (Table 3). The biomass in stratum 373 contribute 79 000 t to the overall estimate of 298 000 t (Table 14). In 2003, there was also a few large catches (6> 900 fish; >200 kg) in strata 360, 373 and 376. In the 2000 and 2002 when the biomass was lower than in 1999, 2001 and 2003, there were a smaller number of incidences of large catches (2 and 4 >900 fish [>200 kg], respectively). In 2004 there were a few high catches in strata 360, 361 and 376 (4 >200 kg and >900 fish) and in 2005 strata 360, 361, 362, and 376 there were 4 sets greater than 200 kg. In 2006, 40% of the biomass estimate of 319 000 t is due to several sets in strata 360 that caught an average of 400kg and over 1300 fish per set. There is no one strata that dominates the estimate in 2007, however, and high catches are observed in several strata (360, 362, 373, 376; Table 3)

In Div. 3O, the abundance and biomass have shown a somewhat stable but slightly declining trend from 1984 to 1992 with an upward jump in 1993 before again declining (Tables 12 and 15; Fig. 2). The biomass index showed moderate fluctuations around an average value of 27 000 t (675 million fish) for the period 1984-92, increasing to 42 000 t (101 million fish) in 1993 and then declining to an average of 11 000 t during the 1994-95 period. The anomalous high estimate in 1993 may have been produced by the high catch rates in stratum 352 and is reflected in the high variability around the estimate. In 1996, the survey biomass dramatically increased by 492 % from 12 000 t (29 million fish) in 1995 to 71 000 t (162 million fish). Since 1996, estimates of biomass and abundance have been variable, but have shown an increasing trend and reach the series highs in 2006 (99 000 t) and 2007 (310 million fish) respectively. The discrepancy in trends between the two series is due to a shift in length frequency toward smaller fish in 2007 (Fig. 4). In this Division, most of the biomass is generally found in the two strata, 351 and 352 (see Fig. 1 for location) which borders Div. 3N. In 2005, for example, 83% of the biomass estimate is due to catch in strata 351 and 352. Whether some of the annual fluctuations are related to movement between Div. 3N and 3O is unknown.

In the spring surveys of Div. 3LNO because the majority of the survey abundance and biomass was found in Div. 3N and then total stock trends mimic that of Div. 3N. From 1989-1998 there have been negligible amounts in Div. 3L until the 1999 survey (Tables 10, 13 and 28; Fig. 2) and in the most recent 2 years, there has been an increase of 340% in the biomass estimate compared to the first two years of the survey.

Biomass in Div. 3LNO increased rapidly in the late 1990s from the lowest levels in the mid-1990s (Table 28). Between 1999 and 2007, abundance and biomass estimates have been variable but have continued an upward trend (Fig. 2). The 2001 survey estimate of abundance and biomass was the most variable, and like the 1999 estimate it had many sets with large catches, including one with 1.6 t (4 824 fish) in Div. 3N. These large catches probably contributed to the high variability around the estimate. Thus it is unlikely that this is a year effect as was seen in 1999 since the 2001 biomass was very low in Div. 3L and even showed a small decline in Div. 3O.

In 2006, the biomass reached its highest estimate in the time series at 504 000 t (1.6 billion fish) (Table 28). The upward trend was seen in all three divisions, in particular Div. 3L where the highest estimate in the time series was seen. Similar to 1999 and 2005 surveys, more yellowtail were again caught in the northern area of Div. 3N and in the southern area of Div. 3L in 2005 and 2006 when compared to 2002 (see Brodie *et. al.*, 2003). The majority of the biomass was located in and around strata 351 and 352 of Div. 3O and in strata 360, 361, 375 and 376, the Southeast Shoal area in Div. 3N and large catches (>400 kg) were taken in strata 360, 362, 373 and 376 (Table 3). It is more probable to say that 2002 was a negative anomaly as reflected in the lack of fish in the northern areas.

Fall Groundfish Surveys, 1990-2007

Abundance and biomass trends:

Tables 16-20 show the survey catch rates in the form of stratified mean number and weight-per -tow by stratum and division for the fall surveys, 1990-2007. Tables 22-27 show abundance and biomass per stratum, along with confidence intervals for all divisions. Figure 3 shows plots of the abundance and biomass estimates, mean numbers and weights per tow by division from 1990-2007. Overall estimates by division and for 3LNO combined are given in Table 29.

In Div. 3L, abundance and biomass were very low and variable without trend from 1990-1995, reaching an estimate close to zero in 1994 (Fig. 3). Noteworthy is that a "0.0" t biomass was also estimated for the 1995 spring series. From 1990 to 95 the abundance varied around an average level of 2 million fish and then tripled to 6 million fish in 1995 and 1996. The biomass varied around an average level of 1 000 t from 1990-1997 before increasing to about 26 000 t in 2001 (Table 28). Similarly the abundance rose from 6 million fish in 1996 to 75 million fish in 2001. A drop in both the abundance and biomass indices (of 56% and 46% respectively) in 2002 was followed by a general increase to the series high in 2007 (28 000 t; 91 million fish). Estimates have wide confidence limits since 2001, and may be due in part to most of the biomass occurring in 2 strata (363 and 372, which border 3N, account for upwards of 90% of biomass estimates in some years). These increases in biomass in Div. 3L are thought to be the result of an extension of the range of yellowtail flounder with increasing stock size. There are obvious within year differences in the amount of yellowtail flounder caught in this Division and this is reflected in the high variability around the estimates for 1999-2001 and 2003-2007.

In Div. 3N, the stock size, from 1990-92, fluctuated around an average value of 47 000 t before doubling in size in 1993 to 94 000 t (Table 28). Since then the stock increased to 369 000 t in 2001 followed by a decrease to 252 000 t in 2003 (Table 28; Fig. 3). Values have fluctuated around 250 000 t since then, and in 2007 increased to the series high at 378 000 t. Similarly, the survey abundance from 1990-94 fluctuated around an average size of 222 million fish before showing a strong upward trend in 1995 to 509 million fish and reaching 1.3 billion fish in 2001, representing an overall increase of 160% (Table 28; Fig 3). From 2001 the abundance has decreased to a level of 900 million fish. The large jump in stock biomass seen in the 2003 spring survey was not evident in the fall survey (81% vs. 10%, respectively) because the fall 2002 survey did not show a decline as was seen in the spring 2002. From 2001-2006, both the abundance and biomass estimates have varied around a level of 1.0 billion fish and 270 000 t biomass, respectively (Fig. 3). In 2007, estimates increased once more to the highest in the series at 1.5 billion fish and 378 000 t.

Much of the large increases seen in the 2001 survey was attributed to large catches in stratum 376 ranging with 5 sets from 420 to 1150 kg (2 000 to 4 000 fish; 2 sets >800 kg) contributing 33%, to the divisional total estimate and to the large confidence interval around both estimates of abundance and biomass for 2001 (Table 3). In 2004 and 2005 several sets on the Southeast Shoal, (strata 375 and 376) were above 400 kg (1 515-4 473 fish). In 2006 large sets were taken in strata 360 and 376, and in 2007, 11 sets met the criteria for large sets (>900 fish or >400 kg). Similar to the spring surveys strata 360, 361, 362, 373 and 376 account for most of the biomass in this Division.

In Div. 3O, both the abundance and biomass index showed no obvious trend from 1990-96, with abundance fluctuating around an average value of 55 million fish and biomass fluctuating around an average level of 20 000 t (Tables 24 and 28; Fig. 7 and 8). Then in 1997, the biomass level jumped by 205% to 26 000 t (159 million fish). From 1997-99 the biomass showed a slight decreasing trend before turning upward to 98 000 t by 2003. Since 1997 the abundance has been somewhat variable but with an upward trend moving from 159 million fish in 1997 to 334 million fish in 2003. Both the 2001 and 2003 estimates of abundance and biomass were highly variable, but the 2003 estimates of biomass and abundance, which are the highest in the time series, are not as variable as 2001 (Tables 24 and 28). Although slightly lower than 2003, the abundance and biomass estimates in 2004 to 2007 has been stabilize but variable at an average level of 206 million and 64 000 t, respectively.

A large catch of 1200 fish (463 kg) in the western stratum, 338, may have contributed to the large variability around the 2001 fall estimate. Even though the estimate of stock size was low in 2002 there were 3 large catches taken on the Southeast shoal strata. In 2003-2007, there were a few large catches (>1 000 fish) (Table 3 above) taken in the surveys (weight range of 200-400 kg). Similar to the spring surveys, most of the biomass in this division was found in strata 351 and 352 which borders Div. 3N.

In the fall survey of Div. 3LNO, similar to the spring surveys, the majority of the stock was found in Div. 3N. The abundance and biomass in this division has shown a general upward trend since the start of the surveys (Table 29 and Fig. 3). Since 1993, when the survey biomass was estimated to be 113 000 t (372 million fish), there has been an increasing upward trend to a high of 476 000 t (1.2 billion fish) in 2001, representing a 321% increase in stock biomass. The 2001 survey biomass estimate of 476 000 t showed a 42% increase in size over the 2000 estimate. In recent years the biomass in the Southeast Shoal's strata, 375 and 376, contribute significantly to the overall biomass: 34 % in 2000 and 55% in 2001 and the large catches in these strata contribute to the high variability around these two survey estimates. Since 2001 the biomass has decreased to 368 000 t (1.3 billion fish) in 2003 putting the 2002 and 2003 more in line with the 2000 estimates. The annual up and down pattern in the biomass and abundance from 1998 to 2003 evident in the spring surveys is not apparent in the fall surveys (Tables 15 and 27). From 2002 to 2006, the abundance and biomass were stable around an average level of 1.3 billion fish and 369 000 t, respectively. In 2007, estimates were the highest in the series for 3LNO combined (482 000 t; 1.8 million fish).

Distribution

Yellowtail flounder is concentrated mainly in Div. 3N and the bordering areas of Div. 3O and to a lesser extent the border of Div. 3LN, similar to most years in the time series. Figures 5 to 8 show the standard number and weight from the catches of individual fishing sets from the spring and fall Campelen surveys for 2004-2007.

In the 2004-2007 surveys, yellowtail flounder were most abundant on the Southeast Shoal and the strata immediate to the west in Div. 3N (Fig. 5 to 8) most of which straddle the Canadian 200 mile (360 km) limit and extend into the

Regulatory Area. This confirms earlier descriptions of distribution (Walsh, 1992; Brodie *et al.*, 1998; Walsh *et al.*, 1999; 2000; 2001a, c; Simpson and Walsh, 2003; Walsh *et al.*, 2004). Yellowtail flounder also appear to be more abundant in the Regulatory Area of Division 3N in the 1999-2005 surveys than in previous years and the northward distribution of the stock has extended to Div. 3L, similar to mid-1980s when the stock size was high (Simpson and Walsh 2003). Brodie *et al.* (1998) noted that the northward range extension of yellowtail flounder on the Grand Bank contracted with decreasing stock size during the mid to late 1980s and early 1990s so that the bulk of the stock was south of 45°N. Simpson and Walsh (2003) have shown that the observed range contraction of yellowtail flounder at low population levels represents selection for preferred habitats in the southern area of the Bank where depth and temperature are important covariates affecting the spatial pattern. Whereas during periods of large increases in stock size, the range of yellowtail flounder expands into less favourable habitats to north, depth but not temperature plays an influential role, and to a lesser extent westward in support of MacCall's Basin hypothesis. Recent tag returns from the 1998-2000 fishery have also confirmed the northward extension of the stock in recent years (Walsh *et al.*, 2001b; Walsh *et al.*, 2006).

Figure 9 shows a plot of the proportion of biomass north of 450 N from 1973 to 2007 and it is obvious that the range of the stock has extended northward since 1995. From 1996-2001, in 2003, and 2005-07, the proportion of biomass north of 450 N is higher in the spring than in the fall. The one obvious exception is the spring of 2002, when the proportion of biomass is much lower than in fall 2002, and is close to the low values in the early 1990s. The surrounding data suggest that the 2002 spring point is anomalous. The last 3 spring values are over 40%, and the last 3 fall values around 30%.

Colbourne and Walsh (2006) noted that in 1990-2005 surveys the centroid of the biomass of yellowtail flounder located within Div. 3NO was found over the Southeast Shoal of the Grand Bank. This area corresponds to some of the warmest bottom temperatures found anywhere on the Grand Banks. The authors reported that spring bottom temperatures in this region range from a minimum of 1-2°C during cold years (1990) to 3-4°C during warm years (1998 and 1999). Fall bottom temperatures are in general warmer than spring values ranging from 2-3°C in most years to maximum values of between 7-8°C during extreme years (1999). Since 1999, with the exception of 2002, survey catch rates of yellowtail have remained significantly higher than those before 1995. With the exception of 2003 spring bottom temperatures have also been higher than they were in the early 1990s. The cold temperature values observed in the spring of 2003 were anomalous and lasted from April to June and were above average during the remainder of the year (Colbourne *et al.*, 2004). However there is no indication that the temperature had a limiting factor in the northward distribution of yellowtail flounder in the 2002 spring survey (See Fig. 12); and 2004 fall survey (Fig. 12 and 13; Colbourne and Walsh, 2006). Temperature correlations have not been examined for the 2006 and 2007 survey data.

In summary, there was a steady increase in the abundance of yellowtail flounder coinciding with a northward expansion of the stock from 1995 up to 2007 that also coincided with an increasing trend in bottom temperatures (until 2006, at least). Colbourne and Walsh (2006) noted that these results indicate a temperature preference for yellowtail towards the warm water habitat of the Grand Banks. The 2004-2005 surveys showed increasing frequency of catches in the northern areas, especially Div. 3L in the spring (which continued in 2006 and 2007) with some catches being taken in less than 0°C in the fall of 2004. Temperature data has not been examined for the 2006 and 2007 Survey data in relation to yellowtail distribution. Vessel problems resulted in reduced coverage in the 2006 spring survey and estimates from that survey may not be comparable to those in other years.

References

- BRODIE, W. B., S. J. WALSH, and D. B. ATKINSON. 1998. The effect of stock abundance on range contraction of yellowtail flounder (*Pleuronectes ferruginea*) on the Grand Bank of Newfoundland in the Northwest Atlantic from 1975 to 1995. *J. Sea Res.*, **39**: 139-152
- BRODIE, W. B., S. J. WALSH, D. POWER, K. S. DWYER and M. F. VEITCH. 2003. Divisions 3LNO yellowtail flounder – interim monitoring update. NAFO SCR Doc. 03/52, Ser. No. N4870, 18p.
- BRODIE, W.B., AND D.E. STANSBURY. 2007. A brief description of the fall multispecies surveys in SA2 + Divisions 3KLMNO from 1995-2006. NAFO SCR Doc. 07/18, Ser. No. N5366, 24p.
- DOUBLEDAY, W. B. 1981 Manual on groundfish surveys in the Northwest Atlantic. *NAFO Sci. Coun. Studies*, **2**: 55 p.
- COLBOURNE, E. B., and S. J. WALSH. 2006. The distribution and abundance of yellowtail flounder (*Limanda ferruginea*) in relation to bottom temperatures in NAFO Divisions 3LNO based on multi-species surveys from 1990-2005. *NAFO SCR Doc.*, No. 23, Serial No. N5241, 16 p.
- COLBOURNE, E. B., C. FITZPATRICK, D. SENCIALL, P. STEAD, W. BAILEY, J. CRAIG, and C. BROMLEY. 2004. An assessment of physical oceanographic conditions in NAFO Sub-areas 2 and 3 for 2003. *NAFO SCR Doc.*, No. 15, Serial No. 4962, 26 p.
- KULKA, D. W., L. HENDRICKSON, N. BEZ, R. SCHLITZER, G. A. P. BLACK, and M. R. SIMPSON. 2005. Workshop on mapping and geostatistical methods for fisheries stock assessment. *NAFO Sci. Coun. Studies*, **39**: 43 p.
- MCCALLUM, B. R., and S. J. WALSH. 1996. Groundfish survey trawls used at the Northwest Atlantic Fisheries Centre, 1971-present. *NAFO SCR Doc.*, No. 50, Serial No. N27256, 18 p.
- MCCALLUM, B. R., and S. J. WALSH. 1999. Analysis of the performance of the Campelen 1800 shrimp trawl during annual Canadian bottom trawl surveys of Subarea 2J + Divisions 3KLMNO, and 3PS from 1995-1998. *NAFO SCR Doc.*, No. 4. Serial No. N4047, 13 p.
- MCCALLUM, B. R., and S. J. WALSH. 2001. Evaluating the success of the survey trawl standardization program at the Northwest Atlantic Fisheries Centre. *NAFO SCR Doc.*, No. 26, Serial No. N4399, 18 p.
- SIMPSON, M., and S. J. WALSH. 2003. Changes in the Spatial Structure of Grand Bank Yellowtail Flounder: Testing MacCall,s Basin Hypothesis. *J. Sea Res.*, **52**: xx to xx
- STACFIS. 2000. Appendix IV, Report of Standing Committee on Fisheries Science (STACFIS). Pp 122-134 In: Northwest Atlantic Fisheries Organization Scientific Council Reports 2000 ISSN-0250-6416, 303 p.
- WALSH, S. J. 1992. Factors influencing distribution of juvenile yellowtail flounder (*Limanda ferruginea*) on the Grand Bank of Newfoundland. *Neth. J. Sea Res.*, **29**: 193-203.
- WALSH, S. J. 1996 Efficiency of bottom sampling trawls in deriving survey abundance indices. In: Assessment of groundfish stocks based on bottom trawl survey results. *NAFO Sci. Coun. Studies*, **28**: 9-24.
- WALSH, S. J., and B. R. MCCALLUM. 1995. Survey trawl mensuration using acoustic trawl instrumentation. *ICES C.M. Doc.*, No. 1995/B:26, 20 p.
- WALSH, S. J., W. B. BRODIE, M. J. MORGAN, W. R. BOWERING, D. ORR, and M. VEITCH. 1997. An Assessment of the Grand Bank Yellowtail Flounder Stock in NAFO Divisions 3LNO. *NAFO SCR Doc.*, No. 72, Serial No. N2906, 54 p.

- WALSH, S. J., D. ORR, and W. B. BRODIE. 1998a. Conversion factors for yellowtail flounder survey indices derived from comparative fishing trials between the Engel 145 otter trawl and the Campelen 1800 shrimp trawl. *NAFO SCR Doc.*, No. 60, Serial No. N3052, 10 p.
- WALSH, S. J., W. B. BRODIE, M. VEITCH, D. ORR, C. MCFADDEN, and D. MADDOCK PARSONS. 1998b. An assessment of the Grand Bank yellowtail flounder stock in NAFO Divisions 3LNO. *NAFO SCR Doc.*, No. 72, Serial No. N3064, 78 p.
- WALSH, S. J., K. S. WHALEN and M. SIMPSON. 1999. Preliminary analysis of spatial and temporal variation in the distribution of juvenile yellowtail flounder on the Grand Bank: Investigating the methodology. *NAFO SCR Doc.*, No. 59, Serial No. N4118, 57 p.
- WALSH, S. J., M. F. VEITCH, M. J. MORGAN, W. R. BOWERING, and B. BRODIE. 2000. Distribution and abundance of yellowtail flounder (*Limanda ferruginea*) on the Grand Bank, NAFO Divisions 3LNO, as derived from annual Canadian bottom trawl surveys. *NAFO SCR Doc.*, No. 35, Serial No. N4264, 54 p.
- WALSH, S. J., M. J. MORGAN, M. J. MORGAN, and D. STANSBURY. 2001a. Distribution of juvenile yellowtail flounder, American plaice and Atlantic cod on the southern Grand Bank, NAFO Div. 3NO. *NAFO SCR Doc.*, No. 78, Serial No. N4457, 49 p.
- WALSH, S. J., M. J. MORGAN, W. B. BRODIE, K. S. DWYER, and L. MANSFIELD. 2001b A new tagging program for yellowtail flounder on the Grand Bank, NAFO Divs. 3LNO. *NAFO SCR Doc.*, No. 53, Serial No. N4431, 14 p.
- WALSH, S. J., M. F. VEITCH, and W. B. BRODIE. 2001c. Yellowtail flounder (*Limanda ferruginea*) distribution and abundance on the Grand Bank, NAFO Divisions 3LNO, 1984-2000. *NAFO SCR Doc.*, No. 50(revised), Serial No. N4428, 51 p.
- WALSH, S. J., M. F. VEITCH, W. B. BRODIE, and K. S. DWYER. 2004. The distribution and abundance of yellowtail flounder (*Limanda ferruginea*) on the Grand Bank, in NAFO Divisions 3LNO, from the Canadian annual bottom trawl surveys, from 1984-2003. *NAFO SCR Doc.*, No. 36, Serial No. N4986, 50 p.
- WALSH, S. J., and M. J. MORGAN. 2004. Observations of natural behaviour of yellowtail flounder derived from data storage tags. *ICES J. Mar. Sci.*, **61**: 1151-1156.
- WALSH, S. J. 2005. Conversion of the Canadian juvenile (Yankee trawl) groundfish survey time series for yellowtail flounder on the Grand Bank, NAFO Div. 3LNO, into Campelen trawl units. *NAFO SCR Doc.*, No. 48, Serial No. N5134, 11 p.
- WALSH, S. J., M. J. MORGAN, G. HAN, and J. CRAIG. 2006. Progress toward modeling tagging data to investigate spatial and temporal changes in habitat utilization of yellowtail flounder on the Grand Bank. *NAFO SCR Doc.*, No. 29, Serial No. N5248, 30 p.
- WALSH, S. J., W. B. BRODIE, and M. J. MORGAN. 2006. The 2006 Assessment of the Grand Bank Yellowtail Flounder Stock, NAFO Divisions 3LNO. *NAFO SCR Doc.*, No. 48, Serial No. N5274, 31p.

Table 1. Summary of Canadian RV Spring (Campelen time series) surveys from 1996-2007.

Year	Vessel	3L			3N		3O		3LNO	3LNO Survey timing	
		# sets	# inshore sets	Depth range (m)	# sets	Depth range (m)	# sets	Depth range (m)		Earliest set	Latest set
1996	W. Templeman	188		66-664	82	42-665	86	65-685	356	7-May	27-Jun
1997	W. Templeman	158		60-681	71	35-689	81	62-669	310	30-Apr	26-Jun
1998	W. Templeman	163		53-721	88	38-682	93	64-657	344	12-May	30-Jun
1999	W. Templeman	177		41-692	82	40-659	86	62-679	345	11-May	29-Jun
2000	W. Templeman	134		61-681	81	45-664	83	61-694	298	11-May	29-Jun
2001	W. Templeman	154		34-695	79	40-650	79	64-699	312	29-Apr	24-Jun
2002	W. Templeman	146		42-710	79	40-641	79	63-628	304	27-Apr	22-Jun
2003	W. Templeman	155		62-698	79	39-681	79	63-726	313	8-May	26-Jun
2004	W. Templeman	151		47-710	79	44-675	79	61-636	309	12-May	26-Jun
2005	W. Templeman	133		64-672	78	45-691	79	66-719	290	9-May	29-Jun
2006	W. Templeman A. Needler	141		60-701	4	68-77	3	75-84	148	10-Jun	30-Jun
					18	46-68	29	64-103	47	25-Jun	29-Jun
2007	Teleost W. Templeman	40		66-171	79	44-636	79	64-719	40	5-Jun	9-Jun
		97		61-702					255	3-May	12-Jul

Table 2. Summary of Canadian RV Fall (Campelen time series) surveys from 1995-2007.

Year	Vessel	3L			3N		3O		3LNO	3LNO Survey timing	
		# sets	# inshore sets	Depth range (m)	# sets	Depth range (m)	# sets	Depth range (m)		Earliest set	Latest set
1995	Teleost W. Templeman	5 165		733-1210 63-640	90	40-650	81	63-730	341	25-Jan 26-Sep	25-Jan 9-Dec
1996	Teleost W. Templeman A. Needler	31 180		805-1433 51-671	12 54	390-1147 37-309	24 19 15	68-690 65-139 63-304	335	26-Nov 9-Oct 24-Nov	17-Dec 12-Dec 5-Dec
1997	Teleost W. Templeman	71 134		161-1436 35-714	74	41-769	73	64-611	352	29-Nov 26-Sep	20-Dec 15-Dec
1998	Teleost W. Templeman	32 172		691-1437 34-675	78	42-1079	87	61-1076	381	28-Nov 10-Oct	16-Dec 15-Dec
1999	W. Templeman	169		63-1407	68	39-664	75	58-692	312	13-Oct	12-Dec
2000	Teleost W. Templeman	74 102		152-1430 42-447	23 70	747-1419 46-642	24 75	752-1424 62-654	368	11-Oct 11-Oct	18-Dec 14-Dec
2001	Teleost W. Templeman A. Needler	34 169 2		146-1457 38-702 187-203	24 70	739-1410 45-660	22 75	803-1391 67-703	396	22-Sep 6-Oct 24-Nov	20-Nov 6-Dec 24-Nov
2002	Teleost W. Templeman	30 176		763-1431 35-670	24 68	811-1429 44-675	24 75	775-1504 65-696	397	11-Oct 5-Oct	2-Dec 1-Dec
2003	Teleost W. Templeman	30 175		753-1446 32-702	70	43-727	8 75	761-1382 63-650	358	23-Sep 9-Oct	20-Jan 17-Dec
2004	Teleost W. Templeman	4 143		151-522 44-653	69	40-659	76	63-634	292	7-Dec 31-Oct	7-Dec 19-Dec
2005	Teleost W. Templeman A. Needler	7 120 57		803-1351 50-706 121-667	17 69	776-1445 42-633	24 75	754-1410 60-649	369	7-Oct 4-Oct 29-Oct	29-Jan 8-Dec 16-Nov
2006	Teleost W. Templeman	34 151		111-1401 61-641	69	46-650	74	63-674	328	19-Nov 30-Sep	18-Dec 19-Nov
2007	Teleost W. Templeman	48 120		81-1424 61-694	25 69	775-1419 48-652	24 75	753-1410 64-632	361	6-Oct 18-Oct	20-Dec 20-Dec

Table 3. Yellowtail flounder catches >900 fish or >400kg per survey set for Spring and Fall surveys from 1999-2007.

Year	Strata	Spring		Fall			
		3N		3N		3O	
		Number	Weight	Number	Weight	Number	Weight
1999	360	2131	666.5	972	252.7		
		1621	452.7				
	361	1140	145.8				
	362	1169	217.2	938	197.7		
		929	509.5	1772	240.5		
	375	1029	130.4				
	376	1540	267.0	1753	359.8		
2000	352				924	235.2	
	360			1060	238.0		
	362	1274	400.0				
	373	1147	450.6				
				1020	232.5		
	376			1392	682.4		
				2193	419.6		
2001	338				1192	463.4	
	360	1404	303.5				
		1043	225.8				
		1008	280.4				
	361	998	243.5	1275	376.0		
	362	1657	345.6	1433	424.2		
		1388	242.4				
2002	373	4824	1653.8				
	376	1080	305.9	4243	889.1		
		1370	293.2	2081	516.6		
		2384	493.5	3178	831.1		
	360	979	233.7	1642	742.0		
		1474	432.5	1190	315.2		
		1096	268.8				
2003		1200	376.4				
		1118	332.4				
	361			990	284.5		
				987	249.0		
	362			1289	295.7		
	375			1103	295.5		
	376			1365	297.7		
				1367	258.6		
	351				1345	255	
	353				993	340.31	
	360	922	288.5	1360	492.5		
		1105	290.7				
	361			1355	338.5		
				972	240.1		
	373	906	319.1				
	376	1976	393.6	1627	475.0		
		1004	305.1				
		1677	377.3				
	352						
	360						
	361						
	362						
	373						
	375						
	376						
	377						
	378						
	379						
	380						
	381						
	382						
	383						
	384						
	385						
	386						
	387						
	388						
	389						
	390						
	391						
	392						
	393						
	394						
	395						
	396						
	397						
	398						
	399						
	400						
	401						
	402						
	403						
	404						
	405						
	406						
	407						
	408						
	409						
	410						
	411						
	412						
	413						
	414						
	415						
	416						
	417						
	418						
	419						
	420						
	421						
	422						
	423						
	424						
	425						
	426						
	427						
	428						
	429						
	430						
	431						
	432						
	433						
	434						
	435						
	436						
	437						
	438						
	439						
	440						
	441						
	442						
	443						
	444						
	445						
	446						
	447						
	448						
	449						
	450						
	451						
	452						
	453						
	454						
	455						
	456						
	457						
	458						
	459						
	460						
	461						
	462						
	463						
	464						
	465						
	466						
	467						
	468						
	469						
	470						
	471						
	472						
	473						
	474						
	475						
	476						
	477						
	478						
	479						
	480						
	481						
	482						
	483						
	484						
	485						
	486						
	487						
	488						
	489						
	490						
	491						
	492						
	493						
	494						
	495						
	496						
	497						
	498						
	499						
	500						
	501						
	502						
	503						
	504						
	505						
	506						
	507						
	508						
	509						
	510						
	511						
	512						
	513						
	514						
	515						
	516						
	517						
	518						
	519						
	520						
	521						
	522						
	523						
	524						
	525						
	526						
	527						
	528						
	529						
	530						
	531						
	532						
	533						
	534						
	535						
	536						
	537						
	538						
	539						
	540						
	541						
	542						
	543						
	544						
	545						
	546						
	547						
	548						
	549						
	550						
	551						
	552						
	553						
	554						
	555						
	556						
	557						
	558						
	559						
	560						
	561						
	562						
	563						
	564						
	565						
	566						
	567						
	568						
	569						
	570						
	571						
	572						
	573					</td	

Table 4. Mean number per tow of Yellowtail flounder by stratum for Division 3L from Spring surveys 1984-2007.

Range	Str	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07			
30 - 56	784																	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0			
57 - 92	350	3.2	7.4	4.4	1.3	2.8	1.4	0.3	1.5	0.1	0.0	0.1	0.0	1.6	0.0	0.0	33.2	21.5	4.5	0.3	8.4	11.9	22.2	9.7	10.5			
	363	45.6	27.6	14.5	13.1	9.9	3.4	7.6	1.3	0.2	0.0	0.0	0.0	4.4	1.0	0.0	94.8	97.9	13.7	0.7	207.7	55.7	209.8	390.7	386.3			
	371	0.7	0.7	0.0	0.8	0.2	0.0	0.4	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	2.5	0.0	0.0	0.0	0.0	0.8	56.0	9.8	3.5			
	372	96.6	117.1	62.0	24.4	13.9	19.5	8.0	4.0	0.6	0.7	0.1	0.0	2.5	2.4	4.5	47.3	28.2	19.1	3.8	113.8	63.1	142.5	394.3	162.6			
	384	7.7	2.5	1.9	0.4	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.0	0.0	0.5	0.8	0.3	0.3	0.3	0.0	23.3	3.3				
	785																0.5	0.0	0.0	1.0			1.5					
93 - 183	328	0.0	0.0	0.0	0.1	0.0	0.0	0.1	0.2	0.0	0.0	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.0			
	341	0.0	0.2	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.2	0.0	0.2	0.2	1.3	3.8	3.6			
	342	0.0	0.0	0.0	0.0	0.0	0.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0			
	343	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0			
	348	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2			
	349	0.2	0.1	2.3	0.2	0.1	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	18.0	2.6	0.0	0.0	0.4	0.0	0.3	16.9	78.3					
	364	1.6	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2.9	0.5	0.0	0.0	0.0	0.0	1.9	22.3	0.3			
	365	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0			
	370	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0			
	385	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.1				
	390	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5			
	786														0.5	0.0	0.0											
	787														0.0	0.0	0.0											
	788														0.0													
	790														0.0													
	793														0.0													
	794														0.0			0.0										
	797														0.0			0.0										
	799																		0.0									
184 - 274	344	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
	347	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
	366	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
	369	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
	386	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
	389	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
	391	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
	795														0.0			0.6										
184 - 366	789														0.0			0.0										
	791														0.0			0.0										
	798														0.0			0.0										
275 - 366	345	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
	346	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
	368	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
	387	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
	388	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
	392	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
	796														0.0			0.0										
	800														0.0			0.0										
367 - 549	729	0.0													0.0			0.0										
	731	0.0													0.0			0.0										
	733	0.0													0.0			0.0										
	735	0.0													0.0			0.0										
	792														0.0			0.0										
550 - 731	730	0.0													0.0			0.0										
	732	0.0													0.0			0.0										
	734	0.0													0.0			0.0										
	736	0.0													0.0			0.0										
732 - 914	737														0.0													
	741														0.0													
	745														0.0													
	748														0.0													
3L mean # per tow		22.1	9.4	5.3	2.4	1.6	1.6	0.9	0.4	0.1	0.0	0.0	0.0	0.5	0.2	0.3	9.6	7.6	2.1	0.3	16.9	7.0	21.7	47.1	33.3			
Upper CI		39.3	14.6	7.8	3.6	2.4	2.6	1.6	0.7	0.1	0.1	0.0	0.0	0.7	0.4	0.8	15.6	11.3	4.2	0.6	30.2	13.0	37.5	64.2	48.1			
Lower CI		5.0	4.2	2.8	1.2	0.9	0.5	0.2	0.2	0.0	0.0	0.0	0.0	0.2	0.1	-0.2	3.6	3.9	0.0	0.0	3.6	1.0	5.8	30.1	18.4			

Table 5. Mean number per tow of Yellowtail flounder by stratum for Division 3N from Spring surveys 1984-2007.

Range	Str	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07
<=56	375	373.6	1656.6	4096.6	208.3	1185.7	82.3	259.5	21.5	340.3	135.7	29.0	139.7	603.3	487.2	411.6	476.4	301.6	213.4	395.0	286.2	240.0	302.3	463.9	
	376	91.5	220.3	162.3	719.6	125.7	521.3	764.1	183.7	35.0	2.3	10.8	67.7	1029.8	524.8	911.0	349.5	1145.8	243.8	730.8	830.0	768.7	851.7	522.9	
57 - 92	360	289.7	155.3	32.3	33.0	70	480.3	91.7	50.1	140.2	41.9	6.8	133.2	364.7	126.2	374.4	680.3	215.7	549.4	730.8	600.1	470.3	465.8	975.2	522.9
	361	338.6	171.0	101.4	130.1	166.6	142.3	293.3	242.9	63.6	237.9	45.1	276.7	453.6	427.2	455.7	586.7	54.0	639.2	375.3	526.2	472.4	415.1	443.5	386.0
93 - 183	362	227.1	74.4	159.9	103.3	73.3	50.9	79.4	53.7	7.5	86.8	2.3	6.6	169.3	210.5	300.0	307.9	519.1	522.6	55.6	263.2	307.9	456.0	773.8	406.8
	373	122.0	58.1	28.2	38.7	34.6	20.8	2.5	13.4	0.1	0.1	3.0	0.0	3.3	17.8	11.1	103.1	311.8	680.9	32.9	273.6	55.4	315.6	365.9	365.9
	374	59.7	38.5	14.8	7.6	4.2	1.8	0.4	1.0	0.0	0.0	0.0	0.0	0.0	10.7	5.8	248.7	225.5	88.3	31.3	77.9	225.3	254.0	240.0	679.7
	383	3.7	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	359	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	377	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	382	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
184 - 274	358	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	378	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	381	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	275 - 366	357	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	379	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	380	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
367 - 549	723	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	725	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	727	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
550 - 731	724	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	726	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	728	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
732 - 914	752	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	760	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
3N mean # per tow	189.7	104.6	100.0	128.1	58.9	208.4	133.1	111.7	79.3	60.4	51.5	51.5	198.0	233.2	240.4	402.1	289.6	496.4	220.0	381.0	287.4	342.4	660.7	397.1	
Upper Cl	251.2	135.1	141.7	206.4	86.3	335.7	103.6	165.4	127.0	103.6	89.1	89.1	349.9	324.1	499.8	356.3	660.0	287.5	467.6	357.7	421.1	872.1	490.9		
Lower Cl	128.2	74.1	54.3	116.5	59.0	57.9	31.6	17.1	13.8	14.1	11.6	11.6	156.7	304.4	222.8	272.8	182.6	294.5	217.2	262.9	449.3	212.1	490.9		

Table 6. Mean number per tow of Yellowtail flounder by stratum for Division 3O from Spring surveys 1984-2007.

Range	Str	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07	
57 - 92	330	1.0	14.8	5.0	1.5	1.1	2.0	1.2	9.2	0.0	0.1	0.0	0.0	1.8	0.6	0.5	0.6	47.2	1.6	6.0	10.3	8.9	8.4	14.7	36.4	
	331	50.0	62.3	5.3	26.5	9.0	25.0	0.0	2.0	5.5	0.5	1.5	5.3	1.0	69.8	43.5	30.0	17.0	86.5	45.5	36.5	45.5	25.5	25.5	144.8	
	338	30.0	22.2	10.6	4.1	48.9	13.2	11.3	17.1	18.0	13.0	5.0	10.0	66.0	68.1	54.3	63.7	43.2	148.8	28.7	30.2	15.6	21.0	53.6	53.6	144.8
	340	6.0	13.6	16.3	40.8	10.0	6.4	17.7	5.4	3.2	2.8	0.0	0.2	0.0	9.0	1.6	8.8	44.0	11.4	13.6	82.2	7.8	39.4	131.5	97.0	
	351	80.0	85.6	80.7	39.5	75.2	43.5	52.4	24.5	7.2	5.8	0.3	0.8	28.5	65.3	50.7	324.2	105.3	147.5	70.8	105.9	199.7	297.9	241.0	230.4	
	352	63.7	55.6	73.0	103.4	47.2	50.7	77.9	78.4	50.8	226.1	55.6	36.0	12.6	177.4	246.3	279.7	268.4	217.9	458.8	331.0	274.6	379.2	388.9	388.9	
	353	2.0	98.5	32.1	148.5	3.0	9.6	20.7	26.7	10.0	66.5	1.8	70.2	122.2	175.0	190.6	188.2	124.9	80.6	36.0	228.7	82.8	147.1	104.3	104.3	
93 - 183	329	0.0	0.0	0.0	0.1	0.0	0.0	0.4	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.3	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2	
	332	0.0	0.6	14.2	9.2	0.3	30.4	1.8	1.3	1.0	13.3	0.3	1.5	6.5	1.3	7.5	4.8	0.0	4.3	22.0	5.7	0.0	1.7	1.0	1.0	
	337	0.0	0.0	1.0	1.2	2.3	2.8	0.0	0.0	1.0	7.0	0.3	0.5	3.0	15.9	0.5	2.0	0.0	8.7	0.0	0.0	4.4	0.0	0.0	0.0	
	339	1.0	0.3	0.3	0.0	0.0	0.0	1.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.2	27.0	1.0	11.0	0.0	6.5	32.0	2.9	2.9	
	354	0.0	1.0	0.0	0.0	0.0	0.0	0.5	0.0	0.0	3.0	0.0	0.0	0.0	0.7	2.0	0.5	0.0	0.4	1.0	0.0	0.0	1.0	0.0	4.0	
	333	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	336	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
184 - 274	355	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	332	0.0	0.6	14.2	9.2	0.3	30.4	1.8	1.3	1.0	13.3	0.3	1.5	6.5	1.3	7.5	4.8	0.0	4.3	22.0	5.7	0.0	1.7	1.0	1.0	
	337	0.0	0.0	1.0	1.2	2.3	2.8	0.0	0.0	1.0	7.0	0.3	0.5	3.0	15.9	0.5	2.0	0.0	8.7	0.0	0.0	4.4	0.0	0.0	0.0	
	339	1.0	0.3	0.3	0.0	0.0	0.0	1.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.2	27.0	1.0	11.0	0.0	6.5	32.0	2.9	2.9	
	354	0.0	1.0	0.0	0.0	0.0	0.0	0.5	0.0	0.0	3.0	0.0	0.0	0.0	0.7	2.0	0.5	0.0	0.4	1.0	0.0	0.0	1.0	0.0	4.0	
	333	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	336	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
275 - 366	355	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	334	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	335	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	356	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	367 - 549	717	719	721	722	722	722	722	722	722	722	722	722	722	722	722	722	722	722	722	722	722	722	722	722	722
550 - 731	718	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	720	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
732 - 914	764	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	772	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
30 mean # per tow	25.8	34.2	28.5	36.9	24.2	18.9	23.9	19.7	11.0	39.8	8.5	8.5	63.3	54.6	60.5	105.4	73.1	77.3	63.1	95.3	93.2	89.0	169.8	121.4		
Upper CI	42.0	46.2	39.5	52.6	34.6	26.1	36.5	30.0	15.9	80.5	17.3	87.3	89.1	83.0	141.2	96.8	113.3	102.9	122.6	130.1	124.3	238.6	161.3			
Lower CI	9.6	22.1	17.5	21.2	13.9	11.8	11.2	9.3	6.1	-0.9	-17.3	-0.4	-39.4	69.7	49.4	41.2	23.3	68.0	56.3	53.7	100.8	81.4	81.4			

Table 7. Mean weight (kg) per tow of Yellowtail flounder by stratum for Division 3L from Spring surveys 1984-2007.

Range	Str	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07		
30 - 56	784																0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
57 - 92	350	1.4	3.5	2.0	0.6	1.4	0.6	0.2	0.7	0.1	0.0	0.1	0.0	0.7	0.0	0.0	16.3	8.4	2.1	0.0	2.6	4.6	7.9	3.7	3.9		
	363	22.2	12.6	6.9	6.3	4.5	1.6	3.4	0.6	0.1	0.0	0.0	0.0	2.2	0.5	0.0	51.6	43.6	5.3	0.1	77.6	23.7	84.0	144.1	135.2		
	371	0.4	0.3	0.0	0.4	0.1	0.0	0.1	0.0	0.0	0.0	0.0	0.1	0.0	0.0	1.4	0.0	0.0	0.0	0.0	0.0	0.3	20.9	4.4	1.2		
	372	46.5	48.2	28.7	11.2	6.2	9.9	4.0	2.0	0.3	0.3	0.1	0.0	1.1	0.7	1.4	24.2	12.0	7.0	1.5	43.0	23.8	50.9	121.9	55.8		
	384	3.7	1.5	1.2	0.2	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.0	0.2	0.3	0.3	0.0	0.2	0.0	11.8	1.8			
	785																0.0	0.0	0.2	0.0	0.8						
93 - 183	328	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0		
	341	0.0	0.1	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.0	0.1	0.3	1.2	1.1			
	342	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
	343	0.0	0.0	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
	348	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
	349	0.1	0.0	1.0	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	7.9	1.0	0.0	0.0	0.1	0.0	0.1	5.3	24.0			
	364	0.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.4	0.2	0.0	0.0	0.0	0.0	0.0	0.7	9.5	0.1		
	365	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
	370	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
	385	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.0			
	390	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2		
	786															0.2	0.0	0.0	0.0	0.0							
	787															0.0	0.0	0.0	0.0	0.0							
	788																0.0										
	790																	0.0									
	793																		0.0								
	794																		0.0								
	797																		0.0								
	799																			0.0							
184 - 274	344	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
	347	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
	366	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
	369	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
	386	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
	389	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
	391	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
184 - 366	795																			0.0	0.3						
	789																0.0				0.0						
	791																0.0										
	798																0.0				0.0						
275 - 366	345	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
	346	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
	368	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
	387	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
	388	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
	392	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
	796																0.0				0.0						
	800																0.0										
367 - 549	729	0.0															0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	731	0.0															0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	733	0.0															0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	735	0.0															0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	792																0.0	0.0	0.0	0.0							
550 - 731	730	0.0															0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	732	0.0															0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	734	0.0															0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	736	0.0															0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
732 - 914	737																0.0										
	741																0.0										
	745																0.0										
	748																0.0										
30 mean wt (kg) per tow		10.7	4.0	2.5	1.1	0.7	0.8	0.4	0.2	0.0	0.0	0.0	0.0	0.2	0.1	0.1	5.0	3.3	0.8	0.1	6.3	2.8	8.2	16.0	11.4		
Upper CI		19.0	6.0	3.6	1.7	1.1	1.3	0.8	0.4	0.1	0.1	0.0	0.0	0.3	0.1	0.2	8.0										

Table 8. Mean weight (kg) per tow of Yellowtail flounder by stratum for Division 3N from Spring surveys 1984-2007.

Table 9. Mean weight (kg) per tow of Yellowtail flounder by stratum for Division 3O from Spring surveys 1984-2007.

Table 10. Abundance (millions) of Yellowtail flounder by stratum for Division 3L from Spring surveys 1984-2007.

Range	Str	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07		
30 - 56	784																										
57 - 92	350	0.90	2.11	1.24	0.36	0.78	0.39	0.08	0.43	0.03	0.00	0.04	0.00	0.44	0.00	0.00	9.45	6.13	1.28	0.09	2.40	3.38	6.32	2.77	2.98		
	363	11.17	6.76	3.55	3.21	2.41	0.84	1.85	0.31	0.05	0.00	0.00	0.00	1.08	0.24	0.00	23.20	23.96	3.35	0.16	50.85	13.63	51.37	95.66	94.60		
	371	0.11	0.10	0.00	0.12	0.03	0.00	0.06	0.00	0.00	0.00	0.00	0.00	0.06	0.00	0.00	0.39	0.00	0.00	0.00	0.00	0.12	8.64	1.50	0.54		
	372	32.69	39.62	20.98	8.25	4.71	6.61	2.71	1.35	0.20	0.25	0.04	0.00	0.84	0.81	1.52	16.00	9.55	6.47	1.27	38.49	21.36	48.22	133.43	55.03		
	384	1.18	0.39	0.29	0.06	0.03	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.08	0.12	0.04	0.04	0.04	0.00	3.58	0.51			
	785																0.03	0.00									
93 - 183	328	0.00	0.00	0.00	0.03	0.00	0.00	0.03	0.00	0.00	0.00	0.00	0.00	0.00	0.03	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.04	0.00		
	341	0.00	0.05	0.00	0.04	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.26	0.00	0.04	0.04	0.27	0.82	0.78			
	342	0.00	0.00	0.00	0.00	0.00	0.05	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.04	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00			
	343	0.00	0.00	0.00	0.00	0.00	0.02	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00			
	348	0.00	0.00	0.04	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.07	0.05			
	349	0.05	0.02	0.66	0.05	0.04	0.00	0.00	0.00	0.00	0.00	0.04	0.00	0.03	0.00	0.00	5.23	0.76	0.00	0.12	0.00	0.10	4.93	22.77			
	364	0.62	0.00	0.00	0.03	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	1.11	0.19	0.00	0.00	0.00	0.00	8.65	0.13			
	365	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00			
	370	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00			
	385	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.03	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.08	0.04		
	390	0.05	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.11		
	786															0.01	0.00										
	787															0.00	0.00										
	788															0.00											
	790															0.00											
	793															0.00											
	794															0.00											
	797															0.00											
	799																0.00										
184 - 274	344	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
	347	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
	366	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
	369	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00			
	386	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00			
	389	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00			
	391	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00			
	795															0.00											
184 - 366	789															0.00											
	791															0.00											
	798															0.00											
275 - 366	345	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
	346	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
	368	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
	387	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
	388	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
	392	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
	796															0.00											
	800															0.00											
367 - 549	729	0.00														0.00											
	731	0.00														0.00											
	733	0.00														0.00											
	735	0.00														0.00											
	792															0.00											
550 - 731	730	0.00														0.00											
	732	0.00														0.00											
	734	0.00														0.00											
	736	0.00														0.00											
732 - 914	737															0.00											
	741															0.00											
	745															0.00											
	748															0.00											
3L Total		45.43	49.90	26.93	12.29	8.13	7.95	4.70	2.19	0.28	0.25	0.12	0.00	2.49	1.21	1.60	55.37	40.67	11.48	1.63	91.96	38.66	115.64	251.54	177.54		
Upper Cl		80.68	77.51	39.																							

Table 11. Abundance (millions) of Yellowtail flounder by stratum for Division 3N from Spring surveys 1984-2007.

Range	Str	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07		
<=56	375	81.87	36.29	89.76	45.63	25.97	18.02	56.87	4.71	74.58	29.73	6.36	30.61	132.21	201.74	104.40	78.67	66.09	46.76	86.56	62.72	52.59	66.25	101.61			
	376	18.87	45.42	33.47	48.39	25.91	201.46	107.49	157.57	37.88	7.22	0.46	2.23	13.98	51.94	150.10	54.81	154.08	279.99	88.80	226.12	246.99	133.58	191.71	151.41	175.63	
57 - 92	360	119.24	63.90	13.30	13.58	2.88	197.67	37.73	20.61	57.71	17.25	2.78	54.81	150.10	60.63	114.96	70.53	115.62	108.89	116.16	149.56	138.67	162.92	134.12	120.41	150.82	113.05
	361	86.31	43.59	25.85	33.17	42.46	36.27	74.77	61.91	16.22	2.60	30.08	0.81	0.21	58.70	72.98	103.99	176.03	179.95	181.15	19.26	91.25	106.74	158.07	268.22	144.17	98.39
93 - 143	362	78.74	25.78	55.44	35.81	25.41	17.65	27.52	18.24	2.60	30.08	0.81	0.21	58.70	72.98	103.99	176.03	179.95	181.15	19.26	91.25	106.74	158.07	268.22	144.17	98.39	
	373	42.29	20.14	9.78	13.41	11.99	7.23	0.87	4.63	0.03	0.04	1.04	0.00	0.21	58.70	72.98	103.99	176.03	179.95	181.15	19.26	91.25	106.74	158.07	268.22	144.17	98.39
	374	7.64	4.93	1.90	0.97	0.54	0.03	0.23	0.05	0.13	0.00	0.42	1.96	1.38	31.86	28.88	11.31	0.41	35.82	28.86	32.53	30.74	86.97	87.97	126.83	126.83	
	383	0.00	0.00	0.00	0.03	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.21	0.93	0.00	0.00		
	387	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
	382	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
184 - 274	358	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
	378	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
275 - 366	381	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
	357	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
	379	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
	380	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
367 - 549	723	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
	725	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
550 - 731	727	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
	726	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
	728	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
732 - 914	752	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
	756	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
3N Total	760	435.30	240.06	229.50	291.00	135.25	478.33	305.48	268.10	189.15	144.98	126.41	158.81	475.29	554.92	577.23	965.42	695.29	1119.91	528.33	914.86	690.11	822.00	1035.03	953.50		
Upper CI		576.47	310.03	459.56	197.30	473.01	72.55	186.44	373.39	139.00	70.30	218.84	218.84	832.53	611.79	835.57	895.46	690.37	1122.63	888.62	1012.86	1178.14	1178.14	1178.14	1178.14	1178.14	
Lower CI		294.14	170.08	133.85	122.45	170.08	133.85	122.45	170.08	133.85	122.45	170.08	133.85	277.31	736.35	730.89	535.00	655.00	366.25	521.40	631.15	703.92	728.44	728.44	728.44	728.44	

Table 12. Abundance (millions) of Yellowtail flounder by stratum for Division 3O from Spring surveys 1984-2007.

Table 13. Biomass (000 tons) of Yellowtail flounder by stratum for Division 3L from Spring surveys 1984-2007.

Range	Str	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07	
30 - 56	784																									
57 - 92	350	0.41	1.00	0.58	0.18	0.41	0.18	0.05	0.20	0.02	0.00	0.02	0.00	0.19	0.00	0.00	4.65	2.39	0.59	0.01	0.74	1.31	2.26	1.06	1.10	
	363	5.43	3.09	1.70	1.55	1.11	0.39	0.82	0.14	0.03	0.00	0.00	0.00	0.53	0.13	0.00	12.62	10.68	1.31	0.03	19.00	5.81	20.56	35.29	33.10	
	371		0.06	0.05	0.00	0.06	0.01	0.00	0.02	0.00	0.00	0.00	0.00	0.02	0.00	0.00	0.21	0.00	0.00	0.00	0.00	0.05	3.22	0.67	0.19	
	372	15.72	16.30	9.73	3.78	2.11	3.36	1.34	0.67	0.11	0.12	0.02	0.00	0.36	0.24	0.46	8.20	4.06	2.38	0.49	14.54	8.04	17.23	41.24	18.89	
	384		0.57	0.23	0.18	0.03	0.02	0.00	0.00	0.00	0.00	0.00	0.00	0.07	0.00	0.00	0.03	0.04	0.04	0.00	0.04	0.00	1.82	0.28		
	785																0.00	0.00	0.00	0.01						
93 - 183	328	0.00	0.00	0.00	0.01	0.00	0.00	0.02	0.03	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.02	0.00	
	341	0.00	0.02	0.00	0.02	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.10	0.00	0.01	0.01	0.07	0.26	0.25		
	342	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
	343	0.00	0.00	0.00	0.00	0.00	0.00	0.02	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
	348	0.00	0.00	0.03	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.01	
	349	0.02	0.01	0.29	0.03	0.03	0.00	0.00	0.00	0.00	0.01	0.00	0.02	0.00	0.00	0.00	2.30	0.28	0.00	0.03	0.00	0.03	1.54	6.98		
	364	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.54	0.08	0.00	0.00	0.00	0.00	0.26	3.68	0.05	
	365	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
	370	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
	385	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.06	0.01		
	390	0.03	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.05	
	786																									
	787																									
	788																									
	790																									
	793																									
	794																									
	797																									
	799																									
184 - 274	344	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
	347	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
	366	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
	369	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
	386	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
	389	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
	391	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
	795																									
184 - 366	789																0.00									
	791																0.00									
	798																0.00									
275 - 366	345	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
	346	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
	368	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
	387	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
	388	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
	392	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
	796																0.00									
	800																0.00									
367 - 549	729	0.00															0.00									
	731	0.00															0.00									
	733	0.00															0.00									
	735	0.00															0.00									
	792																0.00									
550 - 731	730	0.00															0.00									
	732	0.00															0.00									
	734	0.00															0.00									
	736	0.00															0.00									
732 - 914	737																0.00									
	741																0.00									
	745																0.00									
	748																0.00									
3L Total		21.86	21.08	12.58	5.79	3.75	3.97	2.25	1.06	0.15	0.12	0.05	0.00	1.13	0.46	0.47	28.52	17.53	4.42	0.59	34.33	15.31	43.62	85.66	60.91	
Upper Cl		38.90	32.00	18.31	8.50	5.40	6.75	4.03	1.76	0.36	0.32	0.11	0.11	1.74	0.79	1.28	46.20	25.76	8.87	1.24	59.19</td					

Table 14. Biomass (000 tons) of Yellowtail flounder by stratum for Division 3N from Spring surveys 1984-2007

Range	Str	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07
<=56	375	32.87	17.13	39.79	22.75	11.09	4.65	18.46	2.56	25.93	10.84	2.66	13.09	17.26	19.18	19.89	21.95	15.35	18.54	20.64	21.84	21.63	21.87	21.32	
57 - 92	360	6.19	13.77	16.22	2.61	25.09	14.63	2.98	4.62	1.05	0.12	0.57	1.12	25.50	20.54	30.98	15.02	52.28	10.39	54.84	43.15	40.47	44.17	43.32	
361	32.30	15.27	9.76	14.81	17.89	11.09	26.77	7.55	21.03	41.77	27.66	28.02	16.11	32.03	76.54	26.15	60.22	87.86	78.71	73.42	62.65	125.04	52.18		
362	30.10	11.14	21.23	13.98	12.16	8.54	10.50	8.45	1.00	14.16	0.50	0.10	28.93	33.67	31.20	32.94	41.93	26.21	41.71	33.63	26.15	31.45	30.30		
373	18.34	9.14	4.82	6.16	6.32	3.83	0.31	2.46	0.01	0.01	0.31	0.00	0.65	0.34	1.10	11.07	38.75	57.64	53.60	42.86	4.32	29.22	45.01		
374	3.85	2.70	1.14	0.55	0.29	0.01	0.07	0.02	0.08	0.00	0.00	0.14	0.91	0.38	0.15	8.84	9.52	3.06	33.12	43.40	14.74	49.40	37.67		
383	0.18	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.38		
359	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.37		
377	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	2.28		
382	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
184 - 274	358	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
378	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
381	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01		
275 - 366	357	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
379	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
380	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
367 - 549	723	725	727	724	726	728	729	730	731	732	733	734	735	736	737	738	739	740	741	742	743	744	745	746	
3N Total	167.68	88.20	95.13	77.54	51.39	78.33	70.73	69.07	49.62	50.75	46.34	57.88	103.94	121.30	143.67	238.45	197.33	287.92	147.32	280.20	216.75	263.66	319.06	292.78	
Upper CI	223.25	111.87	135.18	95.88	44.41	115.20	71.40	86.54	81.49	81.49	79.15	12.87	171.71	192.56	305.83	428.44	447.57	232.53	332.29	276.29	190.80	223.05	353.83	231.75	
Lower CI	112.12	55.88	61.58	31.38	10.35	115.24	12.88	11.96	11.96	11.96	11.96	11.96	78.16	70.89	91.79	174.23	116.23	188.26	228.16	157.20	199.80	223.05	231.75		

Table 15. Biomass (000 tons) of Yellowtail flounder by stratum for Division 3O from SORINA surveys 1984-2007.

Range	Str	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07	
57 - 92	330	0.18	1.91	0.74	0.21	0.16	0.32	0.19	1.14	0.00	0.03	0.00	0.00	0.26	0.06	0.04	0.09	0.19	0.53	0.57	0.85	0.86	1.93	3.42		
	331	1.36	1.85	0.18	0.29	0.93	0.04	0.09	0.18	0.01	0.03	0.00	0.00	0.10	0.00	0.00	0.17	0.63	0.41	2.43	0.77	0.77	0.69	0.69		
	338	3.32	2.76	1.42	0.50	5.10	1.67	1.45	1.33	2.12	1.38	0.71	1.28	0.03	0.03	0.00	0.00	5.54	7.20	4.73	14.07	3.29	3.00	2.38	4.97	
	340	0.68	1.55	1.77	4.31	1.10	0.76	2.01	0.63	0.37	0.35	0.00	0.01	0.00	0.00	0.00	0.19	0.98	4.21	1.03	0.74	4.78	0.54	3.86	10.20	6.90
	351	4.42	12.99	11.71	5.98	11.22	6.94	8.40	4.03	1.10	0.82	0.04	0.11	4.70	6.24	31.08	12.09	15.37	4.75	9.73	21.07	27.69	20.47	38.02		
	352	9.98	8.71	10.65	15.24	7.54	8.04	11.18	13.59	7.06	33.00	8.05	5.48	46.04	25.55	29.65	35.11	26.72	33.22	48.43	34.60	31.63	41.98	38.02		
	353	0.19	7.63	2.81	13.36	0.28	0.87	1.75	2.29	0.82	5.26	0.20	5.61	10.67	9.93	16.00	18.19	7.37	4.39	6.64	2.46	16.35	5.98	11.50	6.49	
93 - 183	329	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.05	0.00	0.02	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.02		
	332	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
	337	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
	339	0.05	0.02	0.01	0.00	0.00	0.00	0.00	0.03	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
	354	0.00	0.04	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
	336	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
	355	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
	334	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
	356	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
	367 - 549	717	719	721	720	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
	550 - 731	718	722	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
	732 - 914	764	772	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		
30 Total		28.18	37.50	30.47	41.24	25.85	21.47	25.13	23.30	11.60	42.40	9.24	12.74	70.56	53.19	58.04	98.72	72.15	63.65	51.60	71.98	75.79	81.48	99.11	89.31	
	Upper Cl	45.56	50.68	41.44	58.96	36.22	28.53	37.83	36.31	17.01	84.50	17.98	17.98	96.26	82.92	80.17	130.82	93.91	91.90	83.67	90.75	100.77	114.44	141.35	117.25	

Table 16. Mean number per tow of Yellowtail flounder by stratum for Division 3L from Fall Surveys 1990-2007.

Range	Str	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07
30 - 56	784							0.5	0.0	0.0	41.5	1.0	0.0	0.5	0.0	1.5			
57 - 92	350	5.9	0.7	0.5	0.0	0.1	0.4	0.3	0.0	0.4	1.3	3.1	12.4	18.4	29.3	17.3	2.9	3.3	0.5
	363	5.5	1.1	2.0	0.0	0.3	5.2	3.5	1.2	38.4	73.8	119.5	114.2	34.7	95.5	101.7	99.6	30.2	111.5
	371	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	1.3	0.3	0.3	0.3	0.8	0.0	
	372	3.9	4.8	3.8	7.7	0.0	6.4	16.9	17.2	10.2	6.5	18.0	125.8	55.8	79.9	98.4	29.0	136.2	187.0
	384	0.0	0.2	0.0	0.1	0.0	0.0	0.0	0.0	0.3	0.0	0.3	0.3	0.0	0.0	22.0	10.5	0.0	
	785							0.0	0.0	0.0	1.5	3.5	1.5	1.0	0.0	0.0			
93 - 183	328	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	341	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.0	0.0
	342	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	343	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	348	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3
	349	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.7	0.0	0.0	0.0	19.7	0.1
	364	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.5	0.6	0.0	0.2	0.2
	365	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	370	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	385	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	390	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	786							0.5	0.0	0.5	5.5	0.0	0.0	0.0	0.0	0.5			
	787							0.0	0.0	1.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	788							0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	790							0.0	0.0	0.0	0.0	0.0	0.0	1.0	0.0	0.0			
	793							0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.1	0.0			
	794							0.0	0.0	0.0		0.0	0.0	0.0	0.0	0.0	0.0		
	797							0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
	799							0.0	0.0	0.0	0.0	0.0	0.0	1.5	0.0	0.0			
184 - 274	344	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	347	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	366	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	369	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	386	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	389	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	391	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
184 - 366	789							0.0	0.0	0.0	0.0	0.5	0.4	0.0	0.5	0.0	0.0		
	791							0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
	798							0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
275 - 366	345	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	346	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	368	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	387	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	388	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	392	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	796							0.0	0.0	0.0	0.4	0.0	0.0	0.0	0.0	0.0	0.0	0.5	
367 - 549	800							0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
	729	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	731	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	733	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	735	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
550 - 731	730	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	732	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	734	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	736	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
732 - 914	737							0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	741							0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	745							0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	748							0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
915 - 1097	738							0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	742							0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	746							0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	749							0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
1098 - 1280	739							0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	743							0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	747							0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	750							0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
1281 - 1463	740							0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	744							0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	751							0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
3L mean # per tow		0.8	0.4	0.4	0.5	0.0	0.7	1.1	1.0	2.1	3.5	6.1	11.7	5.2	9.2	13.4	6.6	10.2	15.3
Upper CI		1.6	0.6	0.6	1.3	0.1	1.3	2.2	2.7	5.0	8.6	11.2	18.4	8.0	16.2	21.3	12.4	16.0	26.0
Lower CI		0.0	0.2	0.2	-0.3	0.0	0.1	-0.1	-0.7	-0.8	-1.6	1.0	5.0	2.4	2.3	5.4	0.8	4	

Table 17. Mean number per tow of Yellowtail flounder by stratum for Division 3N from Fall Surveys 1990-2007.

Range	Str	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07
<=56	375	40.7	58.0		76.5	329.8	398.5	216.7	212.6	310.9	372.8	460.5	643.3	545.5	488.5	1552.0	847.8	350.3	668.0
	376	323.3	342.8	323.0	674.8	206.3	711.6	831.3	873.3	782.2	722.5	2047.0	2539.0	1001.9	993.9	1099.3	3188.8	1443.8	1490.0
57 - 92	360	83.3	92.8	49.5	219.7	100.9	171.3	392.1	406.2	498.8	490.6	458.3	319.4	578.3	546.3	513.3	253.3	457.0	1112.3
	361	85.4	269.5	269.8	316.6	385.2	450.0	415.8	397.3	528.5	262.0	146.8	737.6	692.0	617.3	359.2	124.3	135.7	555.8
	362	47.6	60.7	6.7	1.9	6.8	245.0	75.6	307.3	139.4	572.0	202.7	571.4	434.7	339.1	536.3	250.3	279.0	231.1
	373	1.2	2.5	0.0	0.0	7.1	13.8	0.0	35.3	35.4	63.5	69.9	307.9	189.0	142.9	221.7	156.2	195.6	526.8
	374	0.0	1.0	0.0	0.0	0.0	30.0	18.0	15.7	182.3	130.3	202.3	108.3	64.7	192.3	155.0	362.3	493.8	
	383	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	3.5	0.0	1.5	178.2	
93 - 183	359	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	11.5	0.5	10.5	1.0	17.5	22.7	
	377	0.0	0.0	0.0	0.0	0.0	0.0	3.0	2.0	3.5	4.5	0.0	0.0	38.0	467.7	355.0	660.0		
	382	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	88.9	
184 - 274	358	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.0	0.0	0.0	
	378	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	6.5	0.0	0.0	18.7	
	381	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
275 - 366	357	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	10.0	0.0	0.0	
	379	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	380	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
367 - 549	723	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	725	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	727	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
550 - 731	724	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	726	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	728	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
732 - 914	752							0.0										0.0	
	756							0.0										0.0	
	760							0.0										0.0	
915 - 1097	753							0.0										0.0	
	757							0.0										0.0	
	761							0.0										0.0	
1098 - 1280	754							0.0										0.0	
	758							0.0										0.0	
	762							0.0										0.0	
1281 - 1463	755							0.0										0.0	
	759							0.0										0.0	
	763							0.0										0.0	
3N mean # per tow		65.9	92.1	86.4	137.7	108.0	212.0	215.0	256.7	241.2	312.4	320.3	489.5	361.7	364.8	485.5	446.1	339.1	526.6
Upper CI		108.2	151.7	198.7	227.0	179.3	294.2	302.9	321.4	313.7	423.2	494.3	673.2	462.3	464.7	611.3	575.2	463.4	634.8
Lower CI		23.6	32.5	-25.8	48.4	36.7	129.8	127.2	191.9	168.7	201.7	146.4	305.9	261.1	264.9	359.6	317.0	214.7	418.4

Table 18. Mean number per tow of Yellowtail flounder by stratum for Division 3O from Fall Surveys 1990-2007.

Range	Str	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07
57 - 92	330	1.3	0.1	1.3	3.3	0.1	8.2	0.2	7.3	1.7	23.8	3.3	20.0	8.3	22.3	18.0	15.2	10.2	55.3
	331	6.7	29.0	8.0	16.0	0.0	2.0	0.0	1.0	3.5	14.0	3.5	29.1	41.0	3.0	50.5	41.0	28.0	48.5
	338	8.5	20.0	2.0	8.8	0.3	97.0	0.5	38.2	31.2	35.8	78.0	260.0	6.4	72.2	2.7	10.1	76.5	8.6
	340	5.6	36.0	0.3	5.0	1.6	4.8	0.0	28.2	23.2	37.3	4.8	47.6	94.6	31.0	74.1	107.2	36.6	176.3
	351	36.9	15.9	1.8	35.3	7.0	15.8	11.6	107.3	207.4	135.3	272.6	171.1	170.8	446.1	247.7	114.9	191.9	286.9
	352	47.9	172.4	150.5	56.7	69.7	121.9	134.3	249.0	269.9	255.0	369.7	288.0	192.3	283.3	255.5	296.9	177.0	232.0
	353	28.0	0.0	0.0	8.7	0.0	8.7	7.0	82.8	0.5	73.5	30.0	70.0	53.0	253.0	23.8	46.8	42.9	38.3
93 - 183	329	1.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.0	0.2	0.2	0.0	0.2	0.0	0.0
	332	0.8	0.3	2.3	15.7	5.0	3.3	3.0	0.0	0.3	1.7	1.0	10.0	19.0	0.0	9.0	1.0	3.0	0.7
	337	0.0	1.0	0.0	0.0	0.0	0.0	19.0	1.3	5.3	0.3	0.9	0.0	0.0	0.0	3.4	16.0	7.3	0.0
	339	1.0	2.5	0.0	0.0	1.0	0.0	0.3	0.5	0.0	1.5	9.0	23.0	18.5	1.1	3.0	12.0	4.0	
	354	1.0	0.0	0.0	0.0	0.0	0.0	1.8	0.0	0.0	0.0	0.5	0.0	0.0	0.0	0.0	0.0	2.9	
184 - 274	333	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	336	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	355	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
275 - 366	334	0.0	0.0	0.0	0.0	0.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	335	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	356	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
367 - 549	717	0.0	0.0	3.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	719	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	721	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
550 - 731	718	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	720	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	722	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
732 - 914	764							0.0								0.0		0.0	
	768							0.0								0.0		0.0	
	772							0.0								0.0		0.0	
915 - 1097	765							0.0								0.0		0.0	
	769							0.0								0.0		0.0	
	773							0.0								0.0		0.0	
1098 - 1280	766							0.0								0.0		0.0	
	770							0.0											

Table 19. Mean weight (kg) per tow of Yellowtail flounder by stratum for Division 3L from Fall Surveys
1990-2007.

Range	Str	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07
30 - 56	784							0.0	0.0	0.0		4.0	0.1	0.0	0.1	0.0	0.1		
57 - 92	350	2.6	0.3	0.3	0.0	0.0	0.2	0.2	0.0	0.3	0.7	1.1	4.1	5.8	9.1	5.8	1.1	1.2	0.2
	363	2.7	0.5	0.9	0.0	0.1	2.3	1.5	0.6	15.9	36.3	41.8	41.1	12.7	35.1	35.9	32.3	10.8	35.0
	371	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.3	0.1	0.1	0.1	0.4	0.0
	372	1.9	2.3	1.7	3.3	0.0	1.9	5.4	3.3	3.6	1.4	5.3	41.9	25.5	21.8	34.6	12.0	46.3	57.1
	384	0.0	0.1	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.0	0.0	10.8	4.3	0.0
	785							0.0	0.0	0.0		0.4	0.2	0.2	0.4	0.0	0.0		
93 - 183	328	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	341	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0
	342	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	343	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	348	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1
	349	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.3	0.0	0.0	0.0	6.4	0.0
	364	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.3	0.0	0.1
	365	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	370	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	385	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	390	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	786							0.0	0.0	0.2		0.3	0.0	0.0	0.0	0.0	0.0	0.1	
	787							0.0	0.0	0.0		0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	788							0.0	0.0	0.0		0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	790							0.0	0.0	0.0		0.0	0.0	0.1	0.0	0.0			
	793							0.0	0.0	0.0		0.0	0.0	0.0	0.9	0.0			
	794							0.0	0.0	0.0		0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	797							0.0	0.0	0.0		0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	799							0.0	0.0	0.0		0.0	0.0	0.4	0.0	0.0			
184 - 274	344	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	347	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	366	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	369	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	386	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	389	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	391	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	795							0.0	0.0	0.0		0.0	0.0	0.0	0.0	0.0	0.0	0.0	
184 - 366	789							0.0	0.0	0.0		0.0	0.1	0.1	0.0	0.2	0.0	0.0	
	791							0.0	0.0	0.0		0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	798							0.0	0.0	0.0		0.0	0.0	0.0	0.0	0.0	0.0	0.0	
275 - 366	345	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	346	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	368	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	387	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	388	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	392	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	796							0.0	0.0	0.0		0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2
	800							0.0	0.0	0.0		0.0	0.0	0.0	0.0	0.0	0.0	0.0	
367 - 549	729	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	731	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	733	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	735	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	792							0.0	0.0	0.0		0.0	0.0	0.0	0.0	0.0	0.0	0.0	
550 - 731	730	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	732	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	734	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	736	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
732 - 914	737							0.0	0.0	0.0		0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	741							0.0	0.0	0.0		0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	745							0.0	0.0	0.0		0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	748							0.0	0.0	0.0		0.0	0.0	0.0	0.0	0.0	0.0	0.0	
915 - 1097	738							0.0	0.0	0.0		0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	742							0.0	0.0	0.0		0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	746							0.0	0.0	0.0		0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	749							0.0	0.0	0.0		0.0	0.0	0.0	0.0	0.0	0.0	0.0	
1098 - 1280	739							0.0	0.0	0.0		0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	743							0.0	0.0	0.0		0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	747							0.0	0.0	0.0		0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	750							0.0	0.0	0.0		0.0	0.0	0.0	0.0	0.0	0.0	0.0	
1281 - 1463	740							0.0	0.0	0.0		0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	744							0.0	0.0	0.0		0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	751							0.0	0.0	0.0		0.0	0.0	0.0	0.0	0.0	0.0	0.0	
3L mean wt (kg) per to		0.4	0.2	0.2	0.2	0.0	0.2	0.4	0.2	0.8	1.6	2.0	4.0	2.1	2.9	4.7	2.4	3.5	4.7
Upper CI		0.8	0.3	0.3	0.5	0.0	0.4	0.8	0.5	2.0	4.0	3.8	6.2	3.4	5.4	7.5	4.2	5.5	7.8
Lower CI		0.0	0.1	0.1	-0.1	0.0	0.0	-0.1	-0.4	-0.8	0.3	1.8	0.8	0.5	1.9	0.6	1.5	1.6	

Table 20. Mean weight (kg) per tow of Yellowtail flounder by stratum for Division 3N from Fall Surveys 1990-2007.

Range	Str	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07
<=56	375	14.6	23.1		36.4	142.0	67.7	54.8	70.1	87.1	112.2	115.8	177.8	149.1	142.0	347.5	195.8	95.5	190.5
	376	97.2	53.0	52.3	151.7	49.4	118.6	117.2	157.4	174.3	182.9	607.1	597.5	229.7	278.9	242.0	593.7	325.1	339.2
57 - 92	360	16.4	20.1	19.5	60.3	27.3	39.6	89.4	114.5	136.4	147.5	148.2	102.6	203.7	174.0	168.3	91.8	171.9	298.3
	361	37.3	77.0	95.3	116.9	161.0	133.7	122.5	142.9	146.3	69.6	40.7	234.5	185.3	153.4	83.3	30.7	33.3	135.0
	362	19.5	18.6	3.0	1.0	3.0	35.0	23.0	79.7	54.1	101.3	50.6	157.9	97.8	91.5	104.4	63.7	68.2	54.7
	373	0.6	1.4	0.0	0.0	2.5	2.8	0.0	12.2	15.6	20.5	23.4	119.2	66.4	51.2	79.2	42.8	66.3	151.8
	374	0.0	0.9	0.0	0.0	0.0	0.0	8.2	6.2	7.9	78.1	40.6	67.4	34.1	24.6	84.2	84.4	121.1	186.0
	383	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.0	0.7	82.0
93 - 183	359	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	5.4	0.2	4.5	0.3	7.9	6.9	
	377	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.4	0.4	1.0	1.1	0.0	0.0	14.7	196.8	176.1	228.3	
	382	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	25.6
184 - 274	358	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.0	0.0	0.0
	378	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2.7	0.0	0.0	6.6
	381	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
275 - 366	357	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	3.5	0.0	0.0	0.0
	379	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	380	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
367 - 549	723	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	725	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	727	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
550 - 731	724	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	726	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	728	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
732 - 914	752							0.0										0.0	
	756							0.0									0.0		0.0
	760							0.0									0.0		0.0
915 - 1097	753							0.0									0.0		0.0
	757							0.0									0.0		0.0
	761							0.0									0.0		0.0
1098 - 1280	754							0.0									0.0		0.0
	758							0.0									0.0		0.0
	762							0.0									0.0		0.0
1281 - 1463	755							0.0									0.0		0.0
	759							0.0									0.0		0.0
	763							0.0									0.0		0.0
3L mean wt (kg) per to		20.6	22.1	24.1	39.6	39.8	42.8	47.1	68.4	66.3	81.1	94.1	137.3	101.5	105.7	122.2	101.7	96.7	140.7
Upper CI		35.6	36.6	43.7	62.6	66.4	56.5	65.0	87.1	85.0	101.8	143.9	177.0	136.0	135.7	152.3	125.5	136.3	169.8
Lower CI		5.6	7.6	4.6	16.6	13.1	29.1	29.3	49.6	47.5	60.5	44.4	97.7	67.1	75.7	92.1	77.9	57.2	111.6

Table 21. Mean weight (kg) per tow of Yellowtail flounder by stratum for Division 3O from Fall Surveys 1990-2007.

Range	Str	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07
57 - 92	330	0.7	0.0	0.7	1.6	0.1	3.7	0.0	2.6	0.6	12.5	1.1	9.7	3.4	7.8	6.5	8.1	4.0	18.2
	331	3.8	14.9	4.6	8.8	0.0	0.6	0.0	0.3	1.2	1.9	1.1	6.9	12.7	1.5	20.0	14.9	12.3	17.3
	338	3.7	7.8	0.9	4.3	0.2	27.7	0.2	21.7	10.9	10.8	24.7	99.0	2.3	24.9	1.0	3.5	38.0	2.7
	340	2.7	16.8	0.2	1.3	0.8	2.0	0.0	10.9	9.2	11.0	2.1	13.8	38.8	9.0	28.5	33.7	11.0	68.6
	351	16.0	6.6	0.8	14.4	2.8	6.4	3.7	42.0	54.2	34.2	69.2	50.4	44.0	93.6	64.2	39.5	54.5	75.9
	352	19.6	59.2	51.3	23.5	26.1	38.6	42.8	74.6	80.2	66.1	102.8	76.4	62.6	108.8	75.4	107.3	43.4	69.3
	353	13.9	0.0	0.0	3.6	0.0	4.8	4.2	41.4	0.2	21.7	10.0	21.5	16.6	86.6	6.4	13.7	13.6	12.5
93 - 183	329	0.6	0.1	0.0	0.0	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.1	0.1	0.0	0.1	0.0	0.0
	332	0.4	0.2	1.0	7.3	2.6	0.9	1.7	0.0	0.0	0.5	0.4	3.4	6.2	0.0	2.9	0.4	1.1	0.4
	337	0.0	0.6	0.0	0.0	0.0	0.0	0.0	1.2	0.9	1.6	0.1	0.2	0.0	0.0	0.0	1.2	5.1	3.0
	339	0.5	1.1	0.0	0.0	0.5	0.0	0.1	0.3	0.0	0.6	2.6	8.2	4.9	0.4	0.7	4.0	1.1	0.0
	354	0.3	0.0	0.0	0.0	0.0	0.0	0.7	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0
184 - 274	333	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	336	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	355	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
275 - 366	334	0.0	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	335	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	356	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
367 - 549	717	0.0	0.0	1.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	719	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	721	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
550 - 731	718	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	720	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	722	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
732 - 914	764							0.0									0.0		0.0
	768							0.0									0.0		0.0
	772							0.0						0.0		0.0		0.0	0.0
915 - 1097	765							0.0									0.0		0.0
	769							0.0									0.0		0.0
	773																		

Table 22. Abundance (millions) of Yellowtail flounder by stratum for Division 3L from Fall surveys 1990-2007.

Range	Str	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07
30 - 56	784							0.02	0.00	0.00	1.53	0.04	0.00	0.02	0.00	0.06			
57 - 92	350	1.67	0.20	0.14	0.00	0.04	0.10	0.09	0.00	0.12	0.37	0.90	3.54	5.25	8.34	4.92	0.81	0.94	0.14
	363	1.35	0.27	0.48	0.00	0.06	1.27	0.86	0.29	9.41	18.08	29.26	27.95	8.49	23.38	24.91	24.40	7.40	27.31
	371	0.03	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.04	0.19	0.04	0.04	0.04	0.13	0.00
	372	1.32	1.63	1.30	2.61	0.00	2.18	5.72	5.83	3.45	2.20	6.09	42.57	18.87	27.03	33.29	9.81	46.08	63.29
	384	0.00	0.03	0.00	0.02	0.00	0.00	0.00	0.00	0.04	0.00	0.00	0.04	0.04	0.00	0.00	3.39	1.62	0.00
	785							0.00	0.00	0.00	0.10	0.22	0.10	0.06	0.00	0.00			
93 - 183	328	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
	341	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.05	0.00	0.00	
	342	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
	343	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
	348	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.08		
	349	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.04	0.21	0.00	0.00	0.00	5.73	0.04
	364	0.00	0.00	0.08	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.04	0.00	0.00	0.18	0.22	0.00	0.09
	365	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	370	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	385	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	390	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	786							0.01	0.00	0.01		0.06	0.00	0.00	0.00	0.00	0.01		
	787								0.00	0.00	0.08		0.00	0.00	0.00	0.00	0.00	0.00	0.00
	788								0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00
	790								0.00	0.00	0.00		0.00	0.00	0.01	0.00	0.00		
	793								0.00	0.00	0.00		0.00	0.00	0.00	0.01	0.00		
	794								0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00
	797								0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00
	799								0.00	0.00	0.00		0.00	0.00	0.01	0.00	0.00	0.00	0.00
184 - 274	344	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	347	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	366	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	369	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	386	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	389	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	391	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	795							0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
184 - 366	789							0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	791								0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00
	798								0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00
275 - 366	345	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	346	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	368	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	387	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	388	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	392	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01
	796								0.00	0.00	0.00		0.01	0.00	0.00	0.00	0.00	0.00	0.01
	800								0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00
367 - 549	729	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	731	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	733	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	735	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	792								0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00
550 - 731	730								0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00
	732	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	734	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	736	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
732 - 914	737								0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00
	741									0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00
	745									0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00
	748									0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00
915 - 1097	738								0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00
	742									0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00
	746									0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00
	749									0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00
1098 - 128	739									0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00
	743										0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00
	747										0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00
	750										0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00
1281 - 146	740										0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00
	744										0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00
	751										0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00
3L Total		4.37	2.12	2.00	2.63	0.10	3.55	6.68	6.12	13.11	20.64	37.95	74.50	33.14	58.89	63.37	38.78	61.90	90.96
Upper CI		8.68	3.29	3.13	6.62	0.29	6.77	14.10	16.91	31.62	50.49	69.41	117.20	51.25	103.00	101.17	72.83	96.85	154.47
Lower CI		0.07	0.96	0.87	-1.37	-0.09	0.33	-0.74	-4.67	-5.39	-9.20	6.49	31.79	15.04	14.78	25.57	4.73	26.95	27.44

Table 23. Abundance (millions) of Yellowtail flounder by stratum for Division 3N from Fall surveys 1990-2007.

Range	Str	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07
<=56	375	8.91	12.71	16.76	72.27	87.33	47.48	46.60	68.13	81.68	100.91	140.96	119.54	107.06	340.10	185.77	76.75	146.38	
	376	66.66	70.69	66.60	139.14	42.53	146.73	171.42	180.07	161.30	148.98	422.09	523.55	206.59	204.94	226.67	65.74	297.71	307.24
57 - 92	360	34.28	38.17	20.37	90.41	41.52	70.49	161.37	167.19	205.31	201.93	188.61	131.45	238.00	224.86	211.24	104.24	188.09	457.78
	361	21.76	68.70	68.76	80.70	98.18	114.71	105.99	101.27	134.72	66.78	37.42	188.01	176.39	157.36	91.56	31.70	34.59	141.67
	362	16.51	21.04	2.31	0.64	2.35	84.93	26.19	106.53	48.33	198.29	70.27	198.09	150.69	117.56	185.92	86.76	96.72	80.13
	373	0.41	0.87	0.00	0.00	2.48	4.78	0.00	12.23	12.28	22.01	24.22	106.72	65.52	49.55	76.84	54.15	67.80	182.60
	374	0.00	0.13	0.00	0.00	0.00	0.00	3.84	2.31	2.01	23.35	16.69	25.91	13.87	8.28	24.63	19.85	46.40	63.24
	383	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.32	0.00	0.14	16.52
93 - 183	359	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.67	0.03	0.61	0.06	1.01	1.32
	377	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.04	0.03	0.05	0.06	0.00	0.00	0.52	6.43	4.88	9.08	0.00
	382	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	7.91
184 - 274	358	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.02	0.00	0.00	0.00	0.00
	378	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.12	0.00	0.00	0.36	0.00
	381	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
275 - 366	357	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.23	0.00	0.00	0.00
	379	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	380	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
367 - 549	723	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	725	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	727	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
550 - 731	724	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	726	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	728	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
732 - 914	752																		
	756																		
	760																		
915 - 1097	753																		
	757																		
	761																		
1098 - 1280	754																		
	758																		
	762																		
1281 - 1463	755																		
	759																		
	763																		
3N Total		148.54	212.30	158.05	327.66	259.32	508.96	516.29	616.24	632.10	743.08	860.28	1314.70	971.27	869.63	1158.56	1146.73	814.09	1414.23
Upper CI		243.78	349.70	363.28	540.04	430.49	706.38	727.21	771.64	822.16	1006.50	1327.49	1807.98	1241.46	1107.71	1458.87	1478.60	1112.59	1074.89
Lower CI		53.30	74.91	-47.18	115.27	88.16	311.55	305.37	460.84	442.04	479.66	393.07	821.41	701.08	631.55	858.25	814.85	515.59	1153.29

Table 24. Abundance (millions) of Yellowtail flounder by stratum for Division 3O from Fall surveys 1990-2007.

Range	Str	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07
57 - 92	330	0.37	0.04	0.36	0.96	0.04	2.36	0.05	2.11	0.48	6.85	0.95	5.75	2.39	6.42	5.18	4.36	2.92	15.90
	331	0.42	1.82	0.50	1.00	0.00	0.13	0.00	0.06	0.22	0.88	0.22	1.83	2.57	0.19	3.17	2.57	1.76	3.04
	338	2.22	5.22	0.52	2.30	0.07	25.33	0.13	9.97	8.15	9.35	20.37	67.88	1.67	18.85	0.69	2.63	19.97	2.25
	340	1.31	8.50	0.06	1.18	0.38	1.12	0.00	6.66	5.48	8.80	1.13	11.24	22.33	7.32	17.49	25.31	8.46	41.62
	351	12.79	5.51	0.64	12.23	2.43	5.46	4.03	37.19	71.91	46.91	94.49	59.33	59.21	154.66	85.87	39.82	66.51	99.46
	352	17.00	61.19	53.41	20.13	24.73	43.25	47.66	88.37	95.80	90.50	131.19	102.20	68.24	100.53	90.67	105.36	62.82	82.34
	353	4.94	0.00	0.00	1.53	0.00	1.53	1.23	14.59	0.09	12.96	5.29	12.34	9.35	44.62	4.19	8.24	7.57	6.75
93 - 183	329	0.24	0.03	0.00	0.00	0.00	0.00	0.00	0.00	0.09	0.00	0.00	0.00	0.05	0.05	0.05	0.05	0.00	0.00
	332	0.12	0.04	0.34	2.26	0.72	0.48	0.43	0.00	0.05	0.24	0.14	1.44	2.74	0.00	1.30	0.14	0.43	0.10
	337	0.00	0.13	0.00	0.00	0.00	0.00	2.48	0.17	0.69	0.04	0.12	0.00	0.00	0.45	2.09	0.96	0.00	0.00
	339	0.08	0.20	0.00	0.00	0.08	0.00	0.03	0.04	0.00	0.12	0.72	1.85	1.49	0.09	0.24	0.97	0.32	0.00
	354	0.07	0.00	0.00	0.00	0.00	0.00	0.12	0.00	0.00	0.03	0.00	0.00	0.00	0.00	0.00	0.00	0.19	0.00
	333	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
184 - 274	336	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	355	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	334	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
275 - 366	335	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	356	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	334	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
367 - 549	717	0.00	0.00	0.04	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	719	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	721	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
550 - 731	718	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	720	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	722	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
732 - 914	764																		
	768																		
	772																		
915 - 1097	765																		
	769																		
	773																		
1098 - 1280	766																		
	770																		
	774																		
1281 - 1463	767																		
	771																		
	775																		
30 Total		39.55	82.67	55.83	41.58	28.48	79.65	56.16	159.17	182.95	176.54	254.06	262.73	170.41	334.12	209.10	190.80	172.55	26.96
Upper CI		59.03	130.44	126.94	69.29	54.54	128.90	93.49	214.23	261.96	240.30	364.29	430.49	240.59	504.40	328.59	304.72	253.17	364.81
Lower CI		20.07	34.91	-15.28	13.87	2.48	30.40	18.83	104.10	103.94	112.78	143.83	94.97	100.22	163.84	89.60	76.89	91.93	139.12

Table 25. Biomass (000 tons) of Yellowtail flounder by stratum for Division 3L from Fall surveys 1990-2007.

1000

Range	Str	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07
30 - 56	784						0.00	0.00	0.00		0.15	0.00	0.00	0.00	0.00	0.00	0.00		
57 - 92	350	0.75	0.09	0.07	0.00	0.01	0.04	0.05	0.00	0.09	0.20	0.30	1.18	1.66	2.59	1.65	0.33	0.35	0.05
	363	0.65	0.12	0.23	0.00	0.02	0.57	0.38	0.15	3.90	8.89	10.23	10.07	3.12	8.60	8.80	7.91	2.65	8.56
	371	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.02	0.05	0.01	0.01	0.02	0.06	0.00
	372	0.64	0.76	0.58	1.10	0.00	0.63	1.81	1.13	1.22	0.49	1.79	14.18	8.62	7.38	11.70	4.07	15.65	19.31
	384	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.01	0.01	0.00	0.00	1.66	0.66	0.00
	785						0.00	0.00	0.00		0.03	0.01	0.01	0.03	0.00	0.00	0.00		
93 - 183	328	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	341	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.02	0.00	0.00
	342	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	343	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	348	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.04	
	349	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.02	0.07	0.00	0.00	0.00	1.87	0.01
	364	0.00	0.00	0.06	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.02	0.00	0.00	0.06	0.11	0.00	0.03
	365	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	370	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	385	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	390	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	786						0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	787						0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	788						0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	790						0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	793						0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.01	0.00			
	794						0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	797						0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	799						0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
184 - 274	344	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	347	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	366	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	369	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	386	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	389	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	391	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	795						0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
184 - 366	789						0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	791						0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	798						0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
275 - 366	345	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	346	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	368	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	387	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	388	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	392	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	796						0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	800						0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
367 - 549	729	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	731	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	733	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	735	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	792						0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
550 - 731	730		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	732	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	734	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	736	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
732 - 914	737						0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	741						0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	745						0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	748						0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
915 - 1097	738						0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	742						0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	746						0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	749						0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1098 - 128	739						0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	743						0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	747						0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	750						0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1281 - 146	740						0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	744						0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	751						0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3L Total		2.06	0.99	0.94															

Table 26. Biomass (000 tons) of Yellowtail flounder by stratum for Division 3N from Fall surveys 1990-2007.

Range	Str	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07
<=56	375	3.20	5.05		7.98	31.11	14.83	12.01	15.36	19.08	24.58	25.38	38.96	32.66	31.12	76.15	42.90	20.94	41.74
	376	20.05	10.93	10.78	31.27	10.18	24.45	24.16	32.45	35.94	37.71	125.18	123.21	47.36	57.52	49.90	122.43	67.04	69.94
57 - 92	360	6.74	8.29	8.03	24.83	11.25	16.31	36.78	47.24	56.12	60.70	61.01	42.24	83.85	71.62	69.29	37.78	70.76	122.78
	361	9.51	19.62	24.30	29.81	41.03	34.09	31.23	36.44	37.28	17.74	10.38	59.76	47.22	39.10	21.23	7.81	8.48	34.41
	362	6.77	6.43	1.04	0.34	1.03	12.15	7.97	27.63	18.76	35.12	17.53	54.75	33.89	31.71	36.18	22.07	23.64	18.97
	373	0.21	0.49	0.00	0.00	0.87	0.98	0.00	4.22	5.41	7.09	8.11	41.31	23.00	17.76	27.44	14.84	22.98	52.64
	374	0.00	0.12	0.00	0.00	0.00	1.05	0.79	1.01	10.00	5.20	8.63	4.36	3.15	10.79	10.81	15.51	23.82	0.00
	383	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.08	0.00	0.06	7.61	0.00
93 - 183	359	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.31	0.01	0.26	0.02	0.45	0.00	0.00
	377	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.02	0.01	0.01	0.02	0.00	0.00	0.00	0.20	2.71	2.42	3.14
	382	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	2.27	0.00
184 - 274	358	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.00	0.00
	378	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.05	0.00	0.00	0.00	0.13
	381	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
275 - 366	357	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.08	0.00	0.00	0.00
	379	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	380	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
367 - 549	723	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	725	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	727	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
550 - 731	724	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	726	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	728	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
732 - 914	752																		
	756																		
	760																		
915 - 1097	753																		
	757																		
	761																		
1098 - 1254	754																		
	758																		
	762																		
1281 - 1464	755																		
	759																		
	763																		
3N Total		46.48	50.94	44.15	94.22	95.47	102.81	113.20	164.15	173.62	192.95	252.80	368.86	272.66	251.99	291.57	261.45	232.28	377.84
Upper CI		80.32	84.43	79.87	148.93	159.46	135.70	156.14	209.18	222.70	242.08	386.48	475.27	365.14	323.48	363.43	322.56	327.24	456.01
Lower CI		12.64	17.44	8.43	39.52	31.48	69.91	70.26	119.13	124.53	143.83	119.11	262.45	180.19	180.50	205.05	137.32	298.00	0.00

Table 27. Biomass (000 tons) of Yellowtail flounder by stratum for Division 3O from Fall surveys 1990-2007.

Range	Str	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07
57 - 92	330	0.20	0.01	0.19	0.47	0.03	1.06	0.00	0.76	0.17	3.59	0.32	2.78	0.97	2.23	1.87	2.32	1.16	5.23
	331	0.24	0.93	0.29	0.55	0.00	0.04	0.00	0.02	0.07	0.12	0.07	0.43	0.80	0.09	1.26	0.93	0.77	1.09
	338	0.97	2.04	0.24	1.11	0.06	7.24	0.05	5.67	2.85	2.81	6.45	25.86	0.59	6.51	0.26	0.91	9.93	0.71
	340	0.63	3.96	0.04	0.32	0.19	0.47	0.00	2.58	2.17	2.59	0.48	3.26	9.16	2.12	6.73	7.96	2.60	16.18
	351	5.55	2.30	0.29	4.99	0.97	2.23	1.27	14.55	18.79	11.87	23.99	17.49	15.25	32.46	22.27	13.69	18.88	26.30
	352	6.97	21.01	18.20	8.35	9.26	13.72	15.17	26.48	28.47	23.47	36.47	27.12	22.23	38.61	26.76	38.07	15.39	24.60
	353	2.45	0.00	0.00	0.63	0.00	0.84	0.74	7.30	0.03	3.83	1.76	3.80	2.92	15.27	1.13	2.42	2.40	2.20
93 - 183	329	0.14	0.02	0.00	0.00	0.00	0.00	0.00	0.05	0.00	0.00	0.00	0.00	0.03	0.03	0.00	0.01	0.00	0.00
	332	0.06	0.03	0.14	1.05	0.38	0.13	0.24	0.00	0.00	0.07	0.06	0.48	0.90	0.00	0.42	0.06	0.16	0.06
	337	0.00	0.08	0.00	0.00	0.00	0.00	1.32	0.11	0.21	0.02	0.03	0.00	0.00	0.00	0.16	0.67	0.40	0.00
	339	0.04	0.09	0.00	0.00	0.04	0.00	0.01	0.03	0.00	0.04	0.21	0.66	0.39	0.03	0.06	0.32	0.09	0.00
	354	0.02	0.00	0.00	0.00	0.00	0.00	0.05	0.00	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.07
184 - 274	333	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	336	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	355	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
275 - 366	334	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	335	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	356	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
367 - 549	717	0.00			0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	719	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	721	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
550 - 731	718		0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	720		0.00	0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	722	0.00		0.00	0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
732 - 914	764								0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	768								0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	772								0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
915 - 1092	765								0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	769								0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	773								0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1098 - 1268	766									0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	770									0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	774									0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1281 - 1462	767									0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	771									0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	775									0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3O Total		17.27	30.48	19.39	17.46	10.94	25.73	18.85	57.50	52.81	48.36	69.67	81.43	53.50	97.72	60.88	67.12	52.00	52.33
Upper CI		25.93	45.17	43.08	28.09	20.70	38.40	31.46	80.51	74.84	64.45	98.55	150.86	76.36	145.64	92.72	111.80	74.59	114.61
Lower CI		8.60	15.78	-4.29	6.84	1.19	13.07	6.24	34.48	30.79	32.28	40.79	12.00	30.63	49.80	29.05	22.43	29.42	38.44

Table 28. Estimates of abundance (millions), biomass ('000 tons), mean number and weight (kg) per tow for Spring surveys in NAFO Divisions 3LNO from 1984-2007.

	Spring Abundance (millions)				Spring Biomass ('000 tons)				Spring mean # per tow				Spring mean wt (kg) per tow			
	3L	3N	3O	3LNO	3L	3N	3O	3LNO	3L	3N	3O	3LNO	3L	3N	3O	3LNO
1984	45.4	435.3	63.5	544.2	21.9	167.7	28.2	217.7	22.1	189.7	25.8	79.9	10.7	73.1	11.4	32.0
1985	49.9	240.1	84.1	374.1	21.1	88.2	37.5	146.8	9.4	104.6	34.2	37.1	4.0	38.4	15.2	14.6
1986	26.9	229.5	70.1	326.5	12.6	95.1	30.5	138.2	5.3	100.0	28.5	33.3	2.5	41.5	12.4	14.1
1987	12.3	291.0	90.9	394.2	5.8	77.5	41.2	124.6	2.4	128.1	36.9	40.2	1.1	34.1	16.7	12.7
1988	8.1	135.3	59.7	203.1	3.7	51.4	25.8	81.0	1.6	58.9	24.2	20.7	0.7	22.4	10.5	8.2
1989	7.9	478.3	46.7	532.9	4.0	78.3	21.5	103.8	1.6	208.4	18.9	54.3	0.8	34.1	8.7	10.6
1990	4.7	305.5	57.3	367.4	2.2	75.7	25.1	103.1	0.9	133.1	23.9	37.7	0.4	33.0	10.5	10.6
1991	2.2	268.1	50.0	320.3	1.1	69.1	23.3	93.4	0.4	111.7	19.7	32.5	0.2	28.8	9.2	9.5
1992	0.3	189.2	28.0	217.4	0.2	49.6	11.6	61.4	0.1	79.3	11.0	21.2	0.0	20.8	4.6	6.0
1993	0.2	145.0	101.1	246.3	0.1	50.8	42.4	93.3	0.0	60.4	39.8	24.0	0.0	21.1	16.7	9.1
1994	0.1	126.4	21.9	148.4	0.0	46.3	9.2	55.6	0.0	51.5	8.5	14.1	0.0	18.9	3.6	5.3
1995	0.0	158.8	28.5	187.4	0.0	57.9	12.7	70.6	0.0	66.1	11.2	18.2	0.0	24.1	5.0	6.9
1996	2.5	475.3	161.7	639.4	1.1	103.9	70.6	175.6	0.5	198.0	63.3	62.1	0.2	43.3	27.6	17.1
1997	1.2	554.9	139.4	695.5	0.5	121.3	53.2	174.9	0.2	233.2	54.6	67.7	0.1	51.0	20.8	17.0
1998	1.6	577.2	154.5	733.3	0.5	143.7	58.0	202.2	0.3	240.4	60.5	69.9	0.1	59.8	22.7	19.3
1999	55.4	965.4	269.1	1289.9	28.5	238.5	98.7	365.7	9.6	402.1	105.4	120.4	5.0	99.3	38.7	34.1
2000	40.7	695.3	186.5	922.5	17.5	197.3	72.1	287.0	7.6	289.6	73.1	89.6	3.3	82.2	28.3	27.9
2001	11.5	1119.9	197.2	1328.5	4.4	297.9	63.6	366.0	2.1	466.4	77.3	126.6	0.8	124.1	24.9	34.9
2002	1.6	528.3	161.0	690.9	0.6	147.3	51.6	199.5	0.3	220.0	63.1	66.5	0.1	61.4	20.2	19.2
2003	92.0	914.9	243.2	1250.1	34.3	280.2	72.0	386.5	16.9	381.0	95.3	120.2	6.3	116.7	28.2	37.2
2004	38.7	690.1	237.9	966.7	15.3	216.7	75.8	307.9	7.0	287.4	93.2	92.0	2.8	90.3	29.7	29.3
2005	115.6	822.0	227.1	1164.8	43.6	263.7	81.5	388.8	21.7	342.4	89.0	113.2	8.2	109.8	31.9	37.8
2006	251.5	1035.0	295.9	1582.4	85.7	319.1	99.1	503.8	47.1	660.7	169.8	183.0	16.0	203.7	56.9	58.3
2007	177.5	953.5	309.7	1440.7	60.9	292.8	89.3	443.0	33.3	397.1	121.4	140.0	11.4	121.9	35.0	43.0

Table 29. Estimates of abundance (millions), biomass ('000 tons), mean number and weight (kg) per tow for Fall surveys in NAFO Divisions 3LNO from 1990-2007.

	Fall Abundance (millions)				Fall Biomass ('000 tons)				Fall mean # per tow				Fall mean wt (kg) per tow			
	3L	3N	3O	3LNO	3L	3N	3O	3LNO	3L	3N	3O	3LNO	3L	3N	3O	3LNO
1990	4.4	148.5	39.5	192.5	2.1	46.5	17.3	65.8	0.8	65.9	16.1	19.3	0.4	20.6	7.0	6.6
1991	2.1	212.3	82.7	297.1	1.0	50.9	30.5	82.4	0.4	92.1	33.1	29.3	0.2	22.1	12.2	8.1
1992	2.0	158.0	55.8	215.9	0.9	44.1	19.4	64.5	0.4	86.4	22.7	22.4	0.2	24.1	7.9	6.7
1993	2.6	327.7	41.6	371.9	1.1	94.2	17.5	112.8	0.5	137.7	16.4	37.4	0.2	39.6	6.9	11.3
1994	0.1	259.3	28.5	287.9	0.0	95.5	10.9	106.4	0.0	108.0	11.2	28.0	0.0	39.8	4.3	10.3
1995	3.6	509.0	79.6	592.2	1.2	102.8	25.7	129.8	0.7	212.0	31.2	57.2	0.2	42.8	10.1	12.5
1996	6.7	516.3	56.2	579.1	2.2	113.2	18.9	134.3	1.1	215.0	22.7	51.6	0.4	47.1	7.6	12.0
1997	6.1	616.2	159.2	781.5	1.3	164.2	57.5	222.9	1.0	256.7	62.7	69.1	0.2	68.4	22.7	19.7
1998	13.1	632.1	183.0	828.2	5.2	173.6	52.8	231.6	2.1	241.2	69.0	71.1	0.8	66.3	19.9	19.9
1999	20.6	743.1	176.5	940.3	9.6	193.0	48.4	250.9	3.5	312.4	71.4	87.8	1.6	81.1	19.6	23.4
2000	37.9	860.3	254.1	1152.3	12.5	252.8	69.7	335.0	6.1	320.3	91.5	98.8	2.0	94.1	25.1	28.7
2001	74.5	1314.7	262.7	1651.9	25.5	368.9	81.4	475.8	11.7	489.5	95.3	139.8	4.0	137.3	29.5	40.3
2002	33.1	971.3	170.4	1174.8	13.6	272.7	53.5	339.7	5.2	361.7	61.4	99.3	2.1	101.5	19.3	28.7
2003	58.9	869.6	334.1	1262.6	18.6	252.0	97.7	368.3	9.2	364.8	127.1	110.9	2.9	105.7	37.2	32.3
2004	63.4	1158.6	209.1	1431.0	22.2	291.6	60.9	374.7	13.4	485.5	81.9	147.8	4.7	122.2	23.9	38.7
2005	38.8	1146.7	190.8	1376.3	14.1	261.5	67.1	342.7	6.6	446.1	68.7	122.7	2.4	101.7	24.2	30.6
2006	61.9	814.1	172.5	1048.5	21.2	232.3	52.0	305.5	10.2	339.1	68.1	95.4	3.5	96.7	20.5	27.8
2007	91.0	1414.2	252.0	1757.2	28.0	377.8	76.5	482.4	15.3	526.6	90.8	154.0	4.7	140.7	27.6	42.3




Figure 1. Designation of strata in NAFO divisions 3LNO.




Figure 2. Abundance (millions), Biomass ('000 tons), Mean number and weight (kg) per tow for Yellowtail flounder in spring surveys by NAFO division and for 3LNQ combined from 1984-2007.




Figure 3. Abundance (millions), Biomass ('000 tons), Mean number and weight (kg) per tow for Yellowtail flounder in fall surveys in NAFO divisions 3LNO from 1990-2007.




Figure 4. Length distributions of Yellowtail flounder from the 2006 and 2007 Canadian Spring survey in NAFO division 3O.




Figure 5. Distribution of Yellowtail flounder (number per tow) in Canadian spring surveys of NAFO divisions 3LNO from 2004-2007.




Figure 6. Distribution of Yellowtail flounder (weight (kg) per tow) in Canadian spring surveys of NAFO divisions 3LNO from 2004-2007.




Figure 7. Distribution of Yellowtail flounder (number per tow) in Canadian fall surveys of NAFO divisions 3LNO from 2004-2007.




Figure 8. Distribution of Yellowtail flounder (weight (kg) per tow) in Canadian fall surveys of NAFO divisions 3LNO from 2004-2007.




Figure 9. Proportion of yellowtail flounder caught north of 45° N in Divs. 3LNO. All data up to 1995 are from spring surveys only.