

Northwest Atlantic



Fisheries Organization

Serial No. N5601

NAFO SCR Doc. 08/069

## NAFO/ICES WG PANDALUS ASSESSMENT GROUP – OCTOBER 2008

A preliminary estimate of Atlantic cod (*Gadus morhua*) biomass in West Greenland offshore waters (NAFO Subarea 1) for 2008 and recent changes in the spatial overlap with Northern shrimp (*Pandalus borealis*)

by

Kaj Sünksen

Greenland Institute of Natural Resources  
P.O. Box 570, 3900 Nuuk, Greenland

### Abstract

Catches of Atlantic cod in West Greenland waters were compared between the German ground fish survey and the Greenland survey for shrimp and fish. The analysis was restricted to years with sufficient area coverage and regions included in both surveys. The two survey's estimates of cod biomass were closely correlated ( $r^2 = 0.92$ ,  $P < 0.001$ ). Linear regression analysis using data from 16 years revealed that the biomass index of Atlantic cod biomass estimated from the Greenland survey for 2008 would correspond to 84 699 tons in the German survey. The biomass of Atlantic cod is still low compared with the 1980s, despite its moderate increase in the most recent years. The distribution is pronounced southern since 80 % of the biomass is found in NAFO Div. 1F. The impact of Atlantic cod on Northern shrimp through predation therefore appears currently to be small considering the low spatial overlap between the two species.

### Introduction

Hvingel (2002, 2003) has incorporated the effect of cod predation into a model for assessing the stock of Northern shrimp (*Pandalus borealis*) at West Greenland (Hvingel and Kingsley, 2002), and STACFIS and the Scientific Council of NAFO have adopted the results from this model since 2002 (Anon., 2004). No consistent time series of Atlantic cod biomass exist for the different stock components and therefore Hvingel and Kingsley (2002) constructed a cod biomass series for West Greenland based on VPA estimates of stock size for East and West Greenland combined for the period 1955 to 1991 and separate survey indices for East and West Greenland offshore waters available since 1982. From 1993 and forward the time series used in the model was derived from a German offshore cod survey in West Greenland waters alone. This survey is conducted each year in autumn (October-November) and its results are not available at the time the assessment for Northern shrimp is carried out. Instead, information on Atlantic cod abundance and biomass from a Greenland bottom trawl survey for shrimp and fish, which commenced in 1988 and which is carried out in summer (July-September), could be used to assess the actual impact of predation by Atlantic cod on the stock of Northern shrimp. The two surveys, however, do not survey exactly the same area and in some years, area coverage or data sampling has been insufficient in one or the other survey.

Wieland and Storr-Paulsen (2004) compared different time series of Atlantic cod biomass in West Greenland offshore waters and concluded that the approach used by Hvingel and Kingsley (2002) could lead to a substantial overestimation of the impact of predation by Atlantic cod on Northern shrimp. A new time series of Atlantic cod was constructed based on VPA estimates by Buch et al. (1994) and survey indices, which avoids some of the inconsistencies of the data series introduced by Hvingel & Kingsley (2002).

This paper presents an update of a comparison of Atlantic cod biomass estimates from the Greenland survey for shrimp and fish and the German ground fish survey from the last year (Sünksen, 2007). It provides a preliminary estimate of Atlantic cod biomass in West Greenland waters for 2008 that is comparable to the biomass indices from previous years for the area covered by the German survey. Furthermore, the paper examines changes in the spatial overlap between Atlantic cod and Northern shrimp and suggests an alternative time series of Atlantic cod biomass for consideration in the assessment of Northern shrimp.

### Materials and Methods

The Greenland bottom trawl survey for fish and shrimps covers the offshore areas at West Greenland and has been conducted from 1988-2008. The area has been expanded through time and since 1992 the survey has covered the area between 72°30'N and 59°15'N (NAFO Division 1A-1F) from the 3 nautical mile limit to the 600 m depth contour (Fig. 1) and the inshore area Disko Bay. The survey area is divided into NAFO Divisions, which are further subdivided into four depth strata ( $\leq 150$  m, 151-200 m, 201-400 m and 401-600 m). The survey was originally designed as a shrimp survey and sampling of fish data was not complete in the period 1988-1991. Since 1992 the sampling of fish has improved and it is now considered as a combined survey for shrimp and fish (Nygaard *et al.*, 2008). The survey is designed as a stratified random trawl survey with a minimum of two hauls per stratum. The sampling period is July to September and throughout the entire time period the survey has been conducted by the Greenland Institute of Natural Resources with the research trawler RV *Paamiut*. Shrimp trawls with a high (10-12 m) vertical opening have been used, which were equipped with a heavy bobbin chain ground gear until 2004 (Skjervøy trawl) or a rockhopper bobbin/rubber disc ground gear since 2005 (Cosmos 2600 trawl). Towing speed has always been about 2.5 knots (see Ziemer, 2008 for further details on the survey design).

The German survey is as a stratified random groundfish survey covering the shelf area outside the 3 nautical mile limit and the continental slope down to a depth of 400 m off East and West Greenland between 67°00'N and 59°00'N (ICES area 14 and NAFO Div. 1B-1F, Fig. 1). The Institute for Sea Fisheries, Germany, has conducted this survey annually since 1982. The primary target is cod, and the fishing gear used is a groundfish trawl rigged with a heavy ground gear. Towing speed is 4 knots. The survey provides swept area estimates of abundance (by age) and biomass (all ages pooled) for the East and the West Greenland offshore component, which form the primary basis for the evaluation of the status of the offshore cod stock (ICES, 2008).

Area coverage was incomplete in one or the other surveys the following years and areas:

- NAFO Div. 1A and 1BN have always been covered by the Greenland survey only,
- the southern NAFO Div. 1E and 1F were not included in the Greenland survey prior to 1992 and 1993, respectively,
- the northern areas, i.e. 1B and NAFO Div. 1C, were not covered in the German survey in 1995, in 2001 to 2003, and in 2005,
- the number of stations was insufficient in the German survey in NAFO Div. 1F in 1993, 1999 and 2002 and in NAFO Div. 1D in 1995 and 2002.

Consequently, new annual indices of Atlantic cod abundance and biomass were calculated for the Greenland survey for all years since 1992 excluding areas for which no corresponding data have been available from both surveys.

Final factors for converting cod catches from 2005 to 2008 made with the Cosmos trawl to Skjervøy catches from the period 1998-2004 are yet not available. Therefore the 2005 to 2008 estimates were converted preliminarily to the old trawl standard using a size-independent conversion of total catches and an adjustment for the smaller average swept area fished with the old trawl (Nygaard *et al.*, 2008):

$$\text{Cod}_{\text{Skjervøy}} = 1/1.78 * 1.15 * \text{Cod}_{\text{Cosmos}} = 0.65 * \text{Cod}_{\text{Cosmos}}$$

The major difference between the two trawls is the ground gear, and Lewy *et al.* (2004) reported a conversion factor of about 0.5 in such a case, which is close to the value used here.

The changes in the geographical distribution of Atlantic cod and Northern shrimp were examined using both data from the fishery and from the two surveys described above. Commercial catches reported by NAFO Division have

been available for Atlantic cod from Horsted (2000) for the period 1953-1991 and for Northern shrimp from Kingsley and Hvingel (2005) for the period 1975-2005. These values were used to calculate mean latitude of the catches based on the mid of the geographical boundaries of the NAFO statistical areas (Fig. 1) weighted by the aggregated catches. Here, the northern limit of NAFO Division 1A was set to 73°00' N and the southern limit of NAFO Division 1F was set to 59°30' N considering the main distributional range of the fishery for the two species.

Wieland et al. (2006) have recently applied a geostatistical tool, the global index of collocation (GIC) developed by Bez and Rivoirard (2000), to examine the spatial overlap of Atlantic cod and Northern shrimp based on geo-referenced survey data. These results were used to establish a regression between the GIC and the mean latitude of survey biomass of Atlantic cod. This relationship was used to estimate GIC values back in time, i.e. for the periods in which only commercial catch data by NAFO division and no survey data by set were available or the survey appeared to insufficient due to poor area coverage either in the German or the Greenland survey.

## Results and Discussion

Atlantic cod biomasses estimated from the Greenland survey were considerably below the estimates from the German survey in the overlapping years and area, and different trawl types and towing speeds might be the major causes for this. Despite pronounced differences between the two surveys at the upper levels of cod biomass, highly significant correlations ( $r^2 = 0.92$ ,  $P < 0.001$ ,  $n = 16$ ) between the two data series were found, and the equation obtained by linear regression on a log-log scale (Fig. 2) is:

$$\log(\text{Cod biomass}_{\text{German Survey}}) = 0.6792 + 0.9696 * \log(\text{Cod biomass}_{\text{Greenland Survey}})$$

For the most recent years, the survey estimates of stock biomass indicate a pronounced southern distribution of Atlantic cod. This is in 2008 confirmed by the Greenlandic survey where close to 80% of the biomass were found in NAFO Div. 1F which is south of 60°45' (Tab. 1).

Figure 3 shows the geographical distribution of Atlantic cod recorded in the Greenland survey for shrimp and fish in 2008. Apart from the dense aggregation found in 1F fairly high densities were also found in 1D and 1E. In addition, cod were found in many of the stations taken in NAFO Div. 1C and 1B. Overall estimates of cod biomass are 39 904 tons for the entire surveyed area and 37 240 tons for the reduced area that corresponds to the German survey. Taking into account the change of the trawl in the Greenland survey, the latter value converts to 24 093 tons.

The regression equation derived from the comparison of the German and the Greenland survey in the previous years gives an estimate of Atlantic cod biomass for 2008 of 84 699 tons in NAFO Div. 1B (south of 67°N) to 1F (Tab. 2). This value is however quite different from the German survey index in 2007 where a few large hauls caused a biomass somewhat higher than expected from the Greenland survey the same year. However these levels still indicate that the level of Atlantic cod biomass is still much below what has been observed in late 1980s (Fig. 4).

## Acknowledgements

Greenland Institute of Natural Resources would like to thank Dr. Manfred Stein, and Dr. Heino Fock involved in the German groundfish survey for their cooperation.

## References

- ANON. 2004. Scientific Council Meeting –October/November 2004. *NAFO SCS Doc.*, No. 20, Serial No. N5061.
- ZIEMER, N. AND H. SIEGSTAD, 2008. Results of the Greenland Bottom Trawl Survey for Northern shrimp (*Pandalus borealis*) off West Greenland (NAFO Subarea 1 and Division 0A), 1988-2008. *NAFO SCR Doc.*, No. 71 Serial No. N5603.
- BEZ, N., and J. RIVOIRARD. 2000. Indices of collocation between populations. In: Checkley, DM, Hunter, LR, Motos, L, van der Lingen, CD (eds.). Report of a workshop on the use of the Continuous Underway Fish Egg Sampler (CUFES) for mapping spawning habitats of pelagic fish. GLOBEC Report 14: 48-52.
- BUCH, E, S.A. HORSTED, and H. HOVGÅRD. 1994. Fluctuations in the occurrence of cod in Greenland waters and their possible causes. *ICES Marine Science Symposia* 198: 158-174.
- HORSTED, S.A. 2000. A review of the cod fisheries at Greenland, 1910-1995. *Journal of Northwest Atlantic Fishery Science* 28: 1-112.
- HVINGEL, C. 2002. Assessment, prediction and risk analysis of stock development: Shrimp off West Greenland, 2002. *NAFO SCR Doc.*, No. 157, Serial No. N4786.
- HVINGEL, C. 2003. Assessment, prediction and risk analysis: Stock development and production of northern shrimp off West Greenland. *NAFO SCR Doc.*, No. 73, Serial No. N4912.
- HVINGEL, C., and M. C. S. KINGSLEY. 2002. A framework for the development of management advice on a shrimp stock using a Bayesian approach. *NAFO SCR Doc.*, No. 158, Serial No. N4787.
- ICES. 2008. Report of the North Western Working Group. *ICES C.M. Doc.* 2008/ACOM:03, 617 pp.
- KINGSLEY, MCS. and C. HVINGEL. 2005. The fishery for Northern shrimp (*Pandalus borealis*) off West Greenland, 1970-2005. *NAFO SCR Doc.*, No. 83, Serial No. N5188.
- LEWY, P., J.R. NIELSEN and H. HOVGÅRD. 2004. Survey gear calibration independent of spatial fish distribution. *Can. J. Fish. Aquat. Sci.* 61:636-647
- NYGAARD, R., SÜNKSEN, K. and O. JØRGENSEN. 2008. Biomass and abundance of demersal fish stocks off West Greenland estimated from the Greenland shrimp survey, 1988-2007. *NAFO SCR Doc.*, No. 28, Serial No. N5528.
- WIELAND, K., and M. STORR-PAULSEN, 2004. A comparison of different time series of Atlantic cod (*Gadus morhua*) biomass at West Greenland and their potential use for the assessment of Northern Shrimp (*Pandalus borealis*) in NAFO Subarea 1. *NAFO SCR Doc.*, No. 71, Serial No. N5051.
- WIELAND, K., M. STORR-PAULSEN, and K. SÜNKSEN. 2007. Response in stock size and recruitment of northern shrimp (*Pandalus borealis*) to changes in predator biomass and distribution in West Greenland waters. *Journal of Northwest Atlantic Fishery Science* 39: 21-33.
- SÜNKSEN, K. 2007. A preliminary estimate of Atlantic Cod (*Gadus morhua*) biomass in West Greenland offshore waters (NAFO Subarea 1) for 2007 and recent changes in the spatial overlap with Northern shrimp (*Pandalus borealis*). *NAFO SCR Doc.*, No. 73, Serial No. N5459.

Table 1: Observed Cod biomass (tons) by survey and area. Greenland survey as observed, i.e. without conversion  
 -: not covered. +: covered but not to be included in the comparison.

Year	German survey					Greenlandic survey				
	Stratum					Stratum (without > 400 m depth)				
	1BS+1C	1D	1E	1F	Sum	1BS+1C	1D	1E	1F	Sum
1992	332	60	35	184	611	109	118	0	2	229
1993	224	83	55	-	362	30	39	124	+	193
1994	59	28	33	16	136	47	0	1	0	48
1995	-	-	38	17	55	+	+	3	4	7
1996	84	44	50	193	371	15	23	27	49	114
1997	40	24	111	110	285	55	0	0	0	55
1998	75	11	2	42	130	0	47	50	3	100
1999	51	13	129	-	193	6	17	1	+	24
2000	232	156	150	33	571	75	9	2	46	132
2001	-	327	2179	172	2678	+	178	98	100	376
2002	-	-	264	-	264	+	+	42	+	42
2003	-	248	1422	593	2263	+	453	118	46	555
2004	506	578	1329	3870	6283	183	680	685	305	1853
2005	-	1345	3617	20255	25217	+	1317	1047	38765	41129
2006	419	79099	4771	62207	146496	420	5180	545	19111	25256
2007	2098	3168	13621	140098	158985	223	582	471	21091	22367
2008						1051	2133	4844	29210	37240

Tab. 2: Time series of Atlantic cod biomass (tons) (\*: using mean VPA to survey ratio based on Wieland & Storr-Paulsen, (2004).

Year	Cod biomass			Spatial overlap GIC Cod - Shr observed	'estimated'	'effective' Cod biomass
	VPA (Buch et al. 1984)	Survey (ICES 2008)	VPA and survey combined *			
1950	4076.5		4076.5			
1951	3722.9		3722.9			
1952	3285.7		3285.7			
1953	2855.3		2855.3		0.692	1977.1
1954	2848.3		2848.3		0.714	2033.9
1955	2731.4		2731.4		0.703	1919.1
1956	2298.7		2298.7		0.693	1592.7
1957	2037.5		2037.5		0.684	1392.9
1958	1866.2		1866.2		0.674	1258.3
1959	1687.7		1687.7		0.719	1212.6
1960	1823.1		1823.1		0.706	1287.3
1961	1793.9		1793.9		0.704	1263.1
1962	1469.2		1469.2		0.716	1051.3
1963	1328.4		1328.4		0.686	911.2
1964	1327.6		1327.6		0.676	898.1
1965	1345.3		1345.3		0.706	950.2
1966	1254.9		1254.9		0.709	889.2
1967	1167.5		1167.5		0.683	797.4
1968	904.9		904.9		0.639	578.1
1969	637.4		637.4		0.611	389.7
1970	442.2		442.2		0.554	244.9
1971	372.8		372.8		0.587	218.7
1972	283.0		283.0		0.678	191.9
1973	179.5		179.5		0.643	115.4
1974	132.1		132.1		0.642	84.7
1975	108.5		108.5		0.629	68.2
1976	228.8		228.8		0.579	132.5
1977	251.7		251.7		0.574	144.5
1978	253.5		253.5		0.672	170.3
1979	217.3		217.3		0.670	145.6
1980	240.4		240.4		0.680	163.4
1981	178.3		178.3		0.619	110.4
1982	190.9	128.5	190.9		0.518	98.8
1983	134.0	82.4	134.0		0.461	61.7
1984	79.0	25.6	79.0		0.479	37.8
1985	51.7	35.7	51.7		0.482	25.0
1986	38.4	86.7	38.4		0.510	19.6
1987	466.7	638.6	466.7		0.604	282.1
1988	481.2	608.0	481.2		0.618	297.3
1989	403.2	333.9	403.2		0.370	149.1
1990		34.4	34.3		0.289	9.9
1991		5.1	5.1		0.313	1.6
1992		0.6	0.6		0.523	0.3
1993		0.4	0.4	0.633	0.658	0.2
1994		0.1	0.1	(0.356)	0.599	0.1
1995		0.1	0.1	(0.120)	0.483	0.0
1996		0.4	0.4	0.280		0.1
1997		0.3	0.3	0.490		0.1
1998		0.1	0.1	0.390		0.1
1999		0.2	0.2	0.496		0.1
2000		0.6	0.6	0.643		0.4
2001		2.7	2.7	0.462		1.2
2002		2.1	2.1	0.278		0.6
2003		2.3	2.3	0.398		0.9
2004		6.3	6.3	0.257		1.6
2005		25.2	25.2	0.074		1.9
2006		146.5	146.5	0.220		32.2
2007		159.0	159.0	0.139		22.1
2008			84.7	0.156		13.2




Fig. 1. Survey coverage for the Greenland fish and shrimp survey and the German ground fish survey in West Greenland offshore waters. (\*: NAFO divisions 1B and 1C were not covered by the German survey in 1995, in 2001 to 2003 and in 2005).




Fig. 2. Relationship between estimates of cod biomass in the overlapping area of the German groundfish survey and the Greenland Bottom Trawl Survey, 1992-2007. Dashed and dotted lines indicate limits of the 95% confidence and prediction intervals, respectively.






Fig. 3. Geographical distribution of cod density (in kg/km<sup>2</sup>) in the Greenland Bottom Trawl Survey in 2008. Numbers in parentheses denotes number of tows.




Fig. 4. Survey indices of Atlantic cod biomass for West Greenland offshore waters, 1982-2008 (1982-2007: original estimates from the German groundfish survey; 2008: estimate based on the Greenland Bottom Trawl Survey).