



SCIENTIFIC COUNCIL MEETING – JUNE 2009

Physical Oceanographic Conditions on the Scotian Shelf and in the eastern Gulf of Maine
(NAFO areas 4V,W,X) during 2008

B. Petrie, R. G. Pettipas, W. M. Petrie and V. V. Soukhovtsev

Department of Fisheries and Oceans, Maritimes Region
Ocean Sciences Division, Bedford Institute of Oceanography
P.O. Box 1006, Dartmouth, N.S. B2Y 4A2

Abstract

A review of the 2008 physical oceanographic conditions on the Scotian Shelf and in the eastern Gulf of Maine and adjacent offshore areas indicates that there was an overall balance between positive and negative anomalies. The climate index, a composite of 18 selected, normalized time series, had 12 variables within 0.5 standard deviations of their normal values and 3 each that were more than 0.5 SD above or below normal. The overall average of the 18 normalized anomalies was 0.06 ± 0.97 . However, there was systematic variability within the region: 7 of the 10 series from the eastern half (Halifax and eastward) were negative, with the 10 variables having an average value of -0.17 SD; whereas, 6 of the 8 from the western half were positive, with an average value of the 8 variables of +0.35 SD. Deep temperatures in Emerald and Georges Basins were 0.88 and 0.65 SD less than normal; Cabot Strait 200-300 m temperature was 0.45 SD below normal. This indicates colder than normal slope water conditions. These below normal temperatures were also reflected in the bottom temperatures in areas 4W and 4X which were 1.8 and 0.72 SD below normal.

Introduction

This document briefly describes air temperature, ice area and volume, and ocean temperature variability of Scotian Shelf and Gulf of Maine waters during 2008 (see Fig. 1 for the study area, Fig. 2 for area names of time series presented in document). The results are derived from data obtained at coastal and long-term monitoring stations, on annual ecosystem surveys and Atlantic Zone Monitoring Program (AZMP) missions, and from ships-of-opportunity and other research cruises. Most of the data are available in the BIO temperature and salinity (CLIMATE) database¹, which is updated several times per year from the national archive maintained by Integrated Science Data Management (ISDM), Department of Fisheries and Oceans (DFO) in Ottawa. Our analyses use data archived prior to 14 January 2009. Additional hydrographic data were obtained directly from DFO fisheries surveys.

Many of the products which have been presented previously in the overview are now available on the ISDM website devoted to the AZMP program². The products available include sections from the AZMP spring and fall surveys, time series of physical properties from fixed stations, climate indexes such as coastal temperature series, frontal positions, and bottom temperatures from ecosystem surveys. In addition, the availability of quality controlled data from sources such as CLIMATE allow individuals to develop the product that most suits their needs. For these reasons, we present an abbreviated overview to give a general idea of environmental variability in 2008.

¹http://www.mar.dfo-mpo.gc.ca/science/ocean/database/data_query.html

²<http://www.meds-sdmm.dfo-mpo.gc.ca/isdm-gdsi/azmp-pmza/index-eng.html>

In order to detect interannual variability and long-term trends of the time series presented, we have removed the potentially large seasonal cycle by determining the monthly differences, i.e. the anomalies, from the long-term means. In some cases, we present the standardized anomaly (anomaly divided by the standard deviation). When possible, long-term monthly and annual means, and standard deviations (SD) are based on 1971-2000 data.

Temperature and salinity conditions in the Scotian Shelf, Bay of Fundy and Gulf of Maine regions are determined by many processes: heat transfer between the ocean and atmosphere, inflow from the Gulf of St. Lawrence supplemented by flow from the Newfoundland Shelf, exchange with offshore slope waters, local mixing, freshwater runoff, direct precipitation and melting of sea-ice. The Nova Scotia Current is the dominant inflow, originating in the Gulf of St. Lawrence and entering the region through Cabot Strait (see inset, Fig. 1). The Current, whose path is strongly affected by topography, has a general southwestward drift over the Scotian Shelf and continues into the Gulf of Maine where it contributes to the counter-clockwise mean circulation. The properties of shelf waters are modified by mixing with offshore waters from the continental slope. These offshore waters are generally of two types, Warm Slope Water, with temperatures in the range of 8-13°C and salinities from 34.7-35.6, and Labrador Slope Water, with temperatures from 3.5°C to 8°C and salinities from 34.3 to 35. Shelf water properties have large seasonal cycles, east-west and inshore-offshore gradients, and vary with depth (Petrie et al. 1996).

Air Temperatures

Annual air temperature anomalies for six sites in the Scotian Shelf-Gulf of Maine region are shown in Fig. 3 and Table 1. In 2008, annual air temperature anomalies were positive at all sites and ranged from +0.2 to +0.8°C. The time series of annual anomalies indicates that most sites feature increasing temperatures over the long term with decadal scale variability superimposed. For shorter periods, this can lead to no trend or decreasing temperatures (Fig. 3). Linear trends from 1900 to present from Boston, Shearwater and Sydney correspond to changes of +1.7°C, +1.4°C and 1.0°C respectively; the trends from Yarmouth and Saint John are small, 0.5 and 0.2°C/century, corresponding to changes of 0.6 and 0.2°C for 1900-present; the corresponding change for the shorter Sable Island record (1915-present) is +1.1°C.

The anomalies for all 6 sites, summarized in Fig. 4(A), illustrate two points: for most years the anomalies have the same sign. i.e. the stacked bars and the scatter plot coincide. Since 1915 when all sites were operating, 72 of the 94 years had 4 or more stations with the annual anomalies having the same signs; for 49 years, all 6 stations had anomalies with the same sign. This indicates that the spatial scale of the air temperature patterns is greater than the largest spacing among sites. In fact, plotting the correlations between annual anomalies against station spacings yields an e-folding scale of 1800 km. Second, the time scale of the dominant variability has been changing from longer periods for 1915-1954 to shorter periods for 1955-2008 (Fig. 4C).

Table 1. Air temperature statistics 2008 for Scotian Shelf and Gulf of Maine (Standard Deviation SD)

Site	Annual Anomaly 2008 (°C)		Monthly SD 2008 (°C)	1971-2000 Annual	
	Observed	Normalized		Mean (°C)	SD (°C)
Sydney	+0.8	+1.1	1.3	5.53	0.76
Sable I.	+0.5	+0.7	1.1	7.61	0.68
Shearwater	+0.6	+0.8	1.0	6.67	0.67
Yarmouth	+0.7	+1.2	0.8	6.95	0.57
Saint John	+0.6	+0.9	0.8	5.02	0.67
Boston	+0.2	+0.3	0.9	10.88	0.57

Sea Ice

The greater part of sea ice on the Scotian Shelf originates in the Gulf of St. Lawrence and is transported through Cabot Strait pushed by northwesterly winds and ocean currents. Sydney Bight and the southern coast of Cape Breton are typically the only areas heavily affected by ice. In 2008, ice first appeared on the Scotian Shelf in late February, about 2 weeks later than normal (Fig. 5). Ice lasted until about mid-April, 2 weeks later than normal.

Over the winter, ice duration was near or slightly below normal. Note that in the duration maps, ice boundaries exceed those of first and last appearance. We consider an area that has not had ice in the current year but has had ice for 5 years of the 30 year climatological period to have a negative duration anomaly. Consequently, those areas are colour mapped as light red.

The ice areas and volumes are shown in Fig. 6 and compiled in Tables 2 and 3. The heaviest months, January to April, featured below normal coverage and volumes from January to March, and above normal values in April. Overall, the January to April coverage and volume were 0.84 (15th lowest in 40 year record) and 0.42 SD (19th lowest in 40 year record) below normal. Consider the amount of cooling that the melting of ice would cause on the Scotian Shelf. If $O(10 \text{ km}^3)$ of Gulf ice were melted (a rough estimate and probably a lower bound, see Table 3), then this would lower an 8 m thick layer of water on the open Shelf by about 1°C.

Table 2. Ice Area statistics, Scotian Shelf

Month	07-08 Ice Year (km ²)	1971-2000 Mean (km ²)	Anomaly (km ²)	Normalized
Jan	100	1600	-1500	-0.56
Feb	3100	14000	-10000	-1.04
Mar	7200	18000	-10000	-0.84
Apr	7300	6000	+1600	+0.31

Table 3. Ice Volume statistics, Scotian Shelf

Month	07-08 Ice Year (km ³)	1971-2000 Mean (km ³)	Anomaly (km ³)	Normalized
Jan	0.0	0.2	-0.2	-0.59
Feb	0.7	3.4	-2.7	-1.03
Mar	4.1	7.2	-3.1	-0.62
Apr	5.9	3.5	+2.3	+0.69

Remotely-Sensed Sea Surface Temperature

A 9 km resolution Pathfinder sea surface temperature database is maintained at BIO. In the following analysis, we substituted the 18 km resolution Multi-Channel Sea Surface Temperature (MCSST) data from 1999 for the Pathfinder observations from that year because there was serious degradation of the latter, particularly towards the end of the year. This deterioration of the Pathfinder data quality was not evident in other years nor was it found for the MCSST data. The Pathfinder dataset runs to June 2003 when this version of the data series was terminated. To provide data for June 2003 to present, we used the sea surface temperature data (1997-present) downloaded from the satellites by the remote sensing group in the Ocean Research & Monitoring Section (ORMS). Comparison of the Pathfinder and ORMS temperatures during the common time period led to a conversion equation $SST(\text{Pathfinder}) = 0.976 * SST(\text{ORMS}) + 0.46$ with an $r^2 = 0.98$. We adjusted the ORMS observations to bring them in line with the longer Pathfinder series. Anomalies were based on 1985-2008 averages.

Annual anomalies for 7 subareas in the Scotian Shelf-Gulf of Maine region were determined from the averages of monthly anomalies (Fig. 7, Table 4). In 2008, anomalies ranged from -0°C (-0.0 SD) over Georges Bank to +0.6°C (+1 SD) at Cabot Strait. Seven of the eight areas had positive anomalies, 6 were greater than or equal to +0.5 SD. The average anomaly was +0.6 SD over the region. All areas from Cabot Strait to the western Scotian Shelf and the Bay of Fundy show increasing temperature trends from 1985-2008 corresponding to temperature changes of 0.6°C (WSS) to 1.7°C (ESS); Lurcher Shoals and Georges Bank feature temperature decreases of 0.1°C

Table 4. SST anomalies, 2008 and long-term SST statistics.

Site	2008 SST Anomaly (°C)	2008 SST Anomaly Normalized	1985-2008 Mean Annual SST (°C)	1985-2008 SST Std. Dev. (°C)
Cabot Str.	+0.6	+1.0	6.11	0.65
ESS	+0.4	+0.6	7.32	0.71
CSS	+0.3	+0.5	8.82	0.66
Western Bank	+0.4	+0.6	9.26	0.70
WSS	+0.3	+0.6	8.57	0.56
Lurcher Shoal	+0.1	+0.1	7.77	0.57
Bay of Fundy	+0.4	+0.7	7.85	0.51
Georges Bank	-0.0	-0.0	10.59	0.56

Annual anomalies for 7 subareas in the Scotian Shelf-Gulf of Maine region were determined from the averages of monthly anomalies (Fig. 7, Table 4). In 2008, anomalies ranged from -0°C (-0.0 SD) over Georges Bank to +0.6°C (+1 SD) at Cabot Strait. Seven of the eight areas had positive anomalies, 6 were greater than or equal to +0.5 SD. The average anomaly was +0.6 SD over the region. All areas from Cabot Strait to the western Scotian Shelf and the Bay of Fundy show increasing temperature trends from 1985-2008 corresponding to temperature changes of 0.6°C (WSS) to 1.7°C (ESS); Lurcher Shoals and Georges Bank feature temperature decreases of 0.1°C

Coastal Temperatures and Salinities

Coastal sea surface temperatures have been collected at St. Andrews (New Brunswick) and Halifax (Nova Scotia) since the 1920s (Fig. 8). In 2008, the SST anomalies were +0.18°C (+0.37 SD) for St. Andrews and -0.30°C (-0.45 SD) for Halifax, increases of 0.22°C and 0.68°C from 2007 values.

Temperature and salinity measurements, for the most part sampled monthly, have been taken since 1924 at Prince 5, at the entrance to the Bay of Fundy (Fig. 1). It is the longest continuously operating hydrographic monitoring site in eastern Canada. Its waters are generally well-mixed from the surface to the bottom (90 m). The depth-averaged (0-90 m) temperature and salinity time series are shown in Fig. 8. In 2008, the annual temperature anomaly was +0.34°C (+0.76 SD) and the salinity anomaly was -0.16 (-0.65 SD). These represent changes of +0.26°C and -0.34 from the 2007 values.

Scotian Shelf and Gulf of Maine Temperatures from Long-term Stations

Drinkwater and Trites (1987) tabulated monthly mean temperatures and salinities from available bottle data for areas on the Scotian Shelf and in the eastern Gulf of Maine that generally correspond to topographic features such as banks and basins (Fig. 2). Petrie et al. (1996) updated their report using these same areas and all available hydrographic data. We present time series of annual mean and filtered (5 year running means) temperature anomalies at selected depths for five areas (Fig.9). The Cabot Strait temperatures represent the slope waters entering the Gulf of St. Lawrence along Laurentian Channel; the Misaine Bank series characterizes the near bottom temperatures on the eastern Scotian Shelf; the deep Emerald Basin anomalies represent the slope water intrusions onto the Shelf that are subsequently trapped in the deep inner basins; the Lurcher Shoals observations define the ocean climate in SW Nova Scotia and the shallow waters entering the Gulf of Maine via the Nova Scotia Current; finally, the Georges Basin series indicates the slope waters entering the Gulf of Maine through the Northeast Channel. Annual anomalies are based on the averages of monthly values; however, observations are not available for each month in each area. For Cabot Strait, Misaine Bank, Emerald Basin, Lurcher Shoals and Georges Basin, 2008 annual anomalies are based on observations from 5, 8, 5, 4 and 6 months.

In 2008, the annual anomalies were -0.18°C (-0.45 SD) for Cabot Strait 200-300 m, -0.20°C (-0.32 SD) for Misaine Bank 100 m, -0.88°C (-1.1 SD) for Emerald Basin 250 m, +0.09°C (+0.13 SD) for Lurcher Shoals 50 m, and -0.65°C (-1.2 SD) for Georges Basin 200 m. These values correspond to changes of -0.00°C, +0.20°C, +0.99°C, +1.20°C and +0.14°C, respectively over 2007 values.

Temperatures during the Summer Groundfish Surveys

The broadest spatial CTD coverage of the Scotian Shelf is obtained during the annual July DFO ecosystem survey which covers the Scotian Shelf from Cabot Strait to the Bay of Fundy. The deep water boundary of the survey is marked roughly by 200 m isobath along the shelf break at the Laurentian Channel, the outer Scotian Shelf, and the Northeast Channel into the Gulf of Maine towards the Bay of Fundy. A total of 167 CTD stations were taken during the 2008 survey and an additional 59 bottom temperature stations were obtained as part of the ITQ (Individual Transferable Quota) fleet survey. The groundfish survey takes 1 month to complete with the area west of Halifax sampled first and the area east of Halifax sampled second.

The temperatures from both surveys were combined and interpolated onto a 0.2° by 0.2° latitude-longitude grid using an objective analysis procedure known as optimal estimation. The interpolation method uses the 15 "nearest neighbours" with a horizontal length scale of 30 km and a vertical length scale of 15 m in the upper 30 m and 25 m at deeper depths. Data near the interpolation grid point are weighted proportionately more than those farther away. Temperatures were optimally estimated for 0, 50, 100 m and near bottom.

Bottom temperatures ranged from 3.3°C in area 4Vs to 6.9°C in 4X illustrating the substantial difference in the environmental conditions across the Shelf. The anomalies were least in the eastern areas 4Vn,s, greatest in 4W: $+0.0^\circ\text{C}$ ($+0.01$ SD) in 4Vn; -0.16°C (-0.22 SD) in 4Vs; -1.2°C (-1.8 SD) in 4W; and -0.49°C (-0.72 SD) in 4X (Fig. 10 A-D). Compared to 2007, bottom temperatures in areas 4Vn,s and 4X increased by 0.1, 0.3 and 0.6°C , whereas temperature decreased by 0.2°C in 4W.

The volume of the Cold Intermediate Layer (CIL), defined as waters with temperatures $<4^\circ\text{C}$, was estimated from the full depth CTD profiles for the region from Cabot Strait to Cape Sable (Fig. 10E). For the period 1970 to 1989, the number of CTD profiles per year was limited; therefore, 5 year blocks of data, e.g. 1970-1974, centre date 1972, were used as input for the procedure to map the irregularly spaced data onto a regular grid. The data were then incremented by 1 year and a new set of estimates made (i.e., 1970-74, 1971-75, ...). This procedure is similar to filtering the data for the 1970-89 period, effectively reducing the variance. Thus the long-term mean and particularly the SD (based on the 1972-2000 data in Fig. 10E) could be affected. We expect that the true SD is higher than the one derived here.

There is considerable variation in the volume of the CIL since 1998 (Fig. 10E). In 2008, the observed volume of 6600 km^3 was 1.6 SD greater than the long-term mean value of 5100 km^3 but down slightly from 2007.

Density Stratification

Stratification of the near surface layer influences physical and biological processes in the ocean such as vertical mixing, the ocean's response to wind forcing, the timing of the spring bloom, vertical nutrient fluxes and plankton speciation. Under increased stratification, there is a tendency for more primary production to be recycled within the upper mixed layer and hence less available for the deeper layers. We examined the variability in stratification by calculating the density (σ_t) difference between 0 and 50 m. The density differences were based on monthly mean density profiles calculated for each area in Fig. 2. The long-term monthly mean density gradients for 1971-2000 were estimated; these were subtracted from the individual monthly values to obtain monthly anomalies. Annual anomalies were estimated by averaging all available monthly anomalies within a calendar year. This could be misleading if, in a particular year, most data were collected in months when stratification was weak, while in another year, sampling occurred when stratification was strong. However, initial results, using normalized monthly anomalies, were qualitatively similar to the plots presented here. The annual anomalies and their 5-year running means were then calculated for an area-weighted combination of subareas 4-23 on the Scotian Shelf (Fig. 2, 11). A value of $0.01\text{ (kg m}^{-3}\text{)/m}$ represents a difference of 0.5 kg m^{-3} over 50 m.

The dominant feature is the long term trend of increasing stratification from the 1960s to present. Stratification on the Scotian Shelf in 2008 was the 4th strongest in 49 years and increased from its 2007 value. The average stratification parameter for areas 4-23 was above normal in 2008 by 1 SD.

Sea Level

Sea level is a primary variable in the Global Ocean Observing System. Relative sea level is measured with respect to a fixed reference point on land. Consequently, relative sea level consists of two major components: one due to true changes of sea level and a second caused by sinking or rising of the land. In Atlantic Canada, post-glacial rebound (PGR) is causing the area roughly south (north) of the north shore of the Gulf of St. Lawrence to sink (rise) in response to glacial retreat; this results in an apparent rise (fall) of sea level. Tushingham and Peltier (1991) estimate a PGR component of the sea level trend of 23 cm/century at Halifax.

Relative sea level at Halifax (1920-2008) is plotted as monthly means and as a filtered series using a 12-month running-mean filter (Fig. 12A). The linear trend of the monthly mean data has a positive slope of 31.9 (± 6.7) cm/century, lower than the value of 36.7 cm/century (1897-1980) given by Barnett (1984) but within the standard error (note Barnett does not give a standard error). In 2008, relative sea level at Halifax increased by 1.7 cm above the 2007 level. However, the interesting feature of the data is the long-term variation that has occurred since the 1920s. In Fig. 12(B) we show the differences of the annual sea level from the long-term trend. It is apparent that from the 1920s to the early 1970s, the trend was greater than the trend calculated using all of the data. In fact, the trend from 1920-72 was 40.4 (± 1.4) cm/century; for the period 1972-2008, the value was 20.3 (± 2.4) cm/century, i.e. close to the PGR value predicted by the Tushingham-Peltier model.

Summary

A graphical summary of many of the time series already shown indicates that the periods 1987-1993 and 2003-2004 were predominantly colder than normal and 1999-2000 was warmer than normal (Fig. 13). The period 1979-1986 also tends to be warmer than normal. In this figure, annual anomalies based on the 1971-2000 means have been normalized by dividing by the 1971-2000 standard deviations for each variable. The results are displayed as the number of standard deviations above (red) and below (blue) normal. In 2008, 14 of the 22 series shown were within 0.5 SD of their normal values. Of the 8 remaining series, 4 were more than 0.5 SD greater than normal and 4 were more than 0.5 SD below normal. Most of the series associated with the eastern and central Shelf had negative anomalies; most associated with the western Shelf were positive.

Eighteen selected variables of the mosaic plot are summarized as a combination bar and line-scatter plot in Fig. 14. This plot is an attempt to derive an overall climate index for the area. We have selected “profiles” for the eastern (Misaine), central (Emerald) and western (Lurcher) Scotian Shelf, the Bay of Fundy (Prince 5) and Georges Bank. In addition, we have included the spatially comprehensive but temporally limited July groundfish survey bottom temperatures (4Vn,s, 4W and 4X) and surface temperatures for Halifax and St. Andrews because of their long-term nature. The bar components are colour coded so that for any year the contribution of each variable can be determined and systematic spatial patterns seen. The height of each variable’s contribution to the bar depends on its magnitude. The positive components are stacked on the positive side, the negative components on the negative side. The sum of the normalized anomalies (difference between the positive and negative stacks) is shown as a black line connecting grey circles. This is a measure of whether the year tended to be colder or warmer than normal and can serve as an overall climate index.

The cold periods of 1987-1993 and 2003-2004 and the warm period of 1999-2000 are apparent. Systematic differences from the overall tendency within a given year are also evident. The overall index in 2008 averaged +0.1 (± 1) SD, i.e. essentially normal; 12 variables were within 0.5 standard deviations of their normal values and 3 each were more than 0.5 SD above or below normal. However, there was systematic variability within the region: 7 of the 10 series from the eastern half (Halifax and eastward) were negative, with the 10 variables having an average value of -0.17 SD; whereas, 6 of the 8 from the western half were positive, with an average value of the 8 variables of +0.35 SD. Deep temperatures in Emerald and Georges Basins were 0.88 and 0.65 SD less than normal; Cabot Strait 200-300 m temperature was 0.45 SD below normal. This indicates colder than normal slope water conditions. These below normal temperatures were also reflected in the bottom temperatures in areas 4W and 4X which were 1.8 and 0.72 SD below normal.

Acknowledgements

We wish to thank the many individuals who provided data or helped in the preparation of this paper, including: Mathieu Ouellet of the Marine Environmental Data Service in Ottawa; R. Losier of the Biological Station in St. Andrews, for providing St. Andrews and Prince 5 data. We also thank Eugene Colbourne, Manfred Stein and Peter Galbraith for their comments which improved the document.

References

- Barnett, T. 1984. The estimation of “global” sea level change: a problem of uniqueness. *J. Geophys. Res.* 89, 7980-7988.
- Drinkwater, K. F. and R. W. Trites 1987. Monthly means of temperature and salinity in the Scotian Shelf region. *Can. Tech. Rep. Fish. Aquat. Sci.* 1539: 101 p.
- Petrie, B., K. Drinkwater, D. Gregory, R. Pettipas, and A. Sandström 1996. Temperature and salinity atlas for the Scotian Shelf and the Gulf of Maine. *Can. Data Rep Hydrog. Ocean Sci.* 171: 398 p.
- Tushingham, A. and R. Peltier 1991. Ice 3-G: a new global model of late Pleistocene deglaciation based on geophysical predictions of post-glacial relative sea level change. *J. Geophys. Res.* 96, 4497-4523.




Fig. 1. The Scotian Shelf and the Gulf of Maine showing hydrographic stations, standard sections and topographic features. The dotted lines indicate the boundaries of the NAFO Subareas. Inset depicts major circulation features.




Fig. 2. Areas on the Scotian Shelf and eastern Gulf of Maine from Drinkwater and Trites (1987).




Fig. 3. Annual air temperature anomalies (dashed line) and 5-yr running means (solid line) at selected sites in Scotian Shelf-Gulf of Maine region.




Fig. 4. (A) The contributions of each of the annual temperature anomalies for 6 Scotian Shelf-Gulf of Maine sites are shown as a bar chart and their summation as a time series (grey circles, black line). (B) Variations of the correlations of the annual temperature anomalies with station separation. (C) The autocorrelation of the average annual temperature anomaly for the 6 sites.




Fig. 5. The time when ice first appeared during 2008 in days from the beginning of the year (left, top panel) and its anomaly from the 1971-2000 mean in days (left, bottom panel). Negative (positive) anomalies indicate earlier (later) than normal appearance. The time when ice was last seen in 2008 in days from the beginning of the year (centre, top panel) and its anomaly from the 1971-2000 mean in days (centre, bottom panel). Positive (negative) anomalies indicate later (earlier) than normal disappearance. The duration of ice in days (right, top panel) during 2008 and the anomalies from the 1971-2000 mean in days (right, bottom panel). Positive (negative) anomalies indicate durations longer (shorter) than the mean.




Fig. 6. Time series of the monthly mean ice area for the Scotian Shelf (top panel), the average ice area during the usual periods of advancement (January-March) and retreat (April-May) (middle panel) and the comparison of the monthly areas and volumes to the 1971-2000 means (bottom panel)




Fig. 7. Scotian Shelf-Gulf of Maine areas used for extraction of sea-surface temperature (upper panel). The annual sea surface temperature anomalies derived from satellite imagery compared to their long-term means (lower panel). Pathfinder estimates were used for September 1985-May 2003. Estimates for June 2003-December 2008 were from the remote sensing laboratory, Ocean Research and Monitoring Section of the Ecosystem Research Division at BIO. These values were adjusted by the regression $\text{Pathfinder} = 0.976 \cdot \text{ORMS} + 0.46$ based on a comparison between overlapping Pathfinder-ORMS data.




Fig. 8. The annual surface temperature anomalies (dashed line with circles) and their 5-year running means (heavy black line) for (A) Halifax Harbour and (B) St. Andrews; annual depth-averaged (0-90 m) (C) temperature and (D) salinity anomalies for the Prince 5 monitoring station at the mouth of the Bay of Fundy. Horizontal dashed lines mark ± 1 standard deviation.




Fig. 9. The annual mean temperature anomaly time series (dashed line with circles) and the 5 year running mean filtered anomalies (heavy solid line) on the Scotian Shelf and in the Gulf of Maine at (A) Cabot Strait 200-300 m; (B) Misaine Bank 100 m; (C) Emerald Basin 250 m; (D) Lurcher Shoals 50 m; and Georges Basin (200 m) (see Fig. 2).




Fig. 10. Time series of July bottom temperature anomalies (dashed lines with circles) and 5 year running mean filtered series (heavy line) for areas (A) 4Vn, (B) 4Vs, (C) 4W and (D) 4X. The horizontal dashed lines are ± 1 SD. (E) Time series of the Cold Intermediate Layer (CIL, defined as waters with $T < 4^{\circ}\text{C}$) volume on the Scotian Shelf based on the July ecosystem survey. The solid horizontal line is the long-term mean, the dashed horizontal lines are the mean ± 1 SD.




Fig. 11. The mean annual (dashed line with circles) and 5-yr running mean (heavy solid line) of the stratification index (0-50 m density gradient) averaged over the Scotian Shelf (areas 4-23 inclusive, see Fig. 2). The horizontal dashed lines are ± 1 SD. Standard error estimates for each annual value are also shown.




Fig. 12. (A) The time series of the monthly means and a 12 month running mean of the relative sea level elevations at Halifax, along with the linear trend (1920-2008). (B) Residual relative sea level (monthly observed values – linear trend, averaged to annual estimates).




Fig. 13. Normalized annual anomalies of temperatures at the bottom and discrete depths for the Scotian Shelf-Gulf of Maine region. These normalized, annual anomalies are based on the 1971-2000 means, divided by the standard deviation. The scale represents the number of standard deviations an anomaly is from normal; blue indicates below normal, red above normal.




Fig. 14. The contributions of each of the normalized anomalies are shown as a bar chart and their summation as a time series (grey circles, black line).