

Northwest Atlantic



Fisheries Organization

Serial No. N5723

NAFO SCR Doc. 09/062

NAFO/ICES WG PANDALUS ASSESSMENT GROUP – OCTOBER 2009

The Norwegian fishery for northern shrimp (*Pandalus borealis*)
in the Barents Sea and round Svalbard

by

Carsten Hvingel and Trond Thangstad

Institute of Marine Research
Box 6404, N-9294 Tromsø, Norway

Abstract

The resource of northern shrimp (*Pandalus borealis*) in the Barents Sea is assessed as one stock. The fishery is multinational. Catches have ranged between 26 and 128 ktons. Norway accounts for 70-92% of the landings. The fishery is managed by effort control. Discard of small shrimp and by-catch of other species is believed to be low.

Overall catches have declined from 83 ktons in 2000 to 26 ktons in 2008 due to reduced market prices for shrimp products, and a major restructuring of the fleet has taken place. Catches in 2009 is projected at 23 ktons. A standardised catch-per-unit-effort series indicate a slight decline in stock density from 2000 to 2004 and a large increase thereafter to a peak in 2006. The indices for 2007 to 2009 are down approx. 10% as compared to 2006. A standardised effort series indicate a declining trend in fishing mortality since 2000.

Introduction

The resource of northern shrimp (*Pandalus borealis*) in the Barents Sea (Fig. 1) within the Norwegian EEZ and in the Svalbard zone (ICES Div I and II) is for assessment purposes considered as one stock. Norwegian and Russian vessels exploit the stock in the entire area while vessels from other nations are restricted to the Svalbard fisheries zone.

The fishery was initiated in 1970 by Norwegian vessels. As the fishery developed, vessels from several nations joined and catches reached 128 ktons in 1984 (Fig. 3). During the recent 10-year period annual yields have varied between 23 and 83 ktons. Norwegian vessels accounted for around 70-92% of the total catches and vessels from Russia, Iceland, Greenland and the EU for the rest (Tab. 1).

The fishery is regulated by effort control: licences are required for the Russian and Norwegian vessels and the fleets operating in the Svalbard zone is regulated by number of effective fishing days and number of vessels by country. Minimum mesh size is 35mm. Other species and small shrimp are protected by mandatory sorting grids and by temporary closing of areas whith excessive by-catch of juvenile cod, haddock, Greenland halibut, redfish and shrimp<15mm carapace length (measured in catch samples taken by independent observers).

A major restructuring of the fleet towards fewer and larger vessels has taken place since the mid 1990s. The fleet is now largely composed of a group of large freeze or factory trawlers (>3000HP (HP=engine horsepower)) and a small group of <500HP vessels. Trawling is mainly performed using two or three trawls simultaneously.

The present paper updates available information derived from catch statistics, logbooks and catch sampling from the Norwegian trawl fishery for shrimp in the Barents Sea (ICES Div. I and II).

Materials and methods

Logbook data were analysed to show the spatial and temporal distribution of the fishery and fleet composition. Catch-per-unit-effort (CPUE) data from Norwegian vessels were used in multiplicative models to calculate standardised annual catch rate indices (Hvingel et al., 2000). A Standardised effort series was derived by dividing total catch by the standardised CPUE.

The CPUE indices included the following variables: (1) vessel fishing power grouped by engine size, (2) seasonal availability of shrimp, (3) spatial availability of shrimp, (4) gear type (single, double or triple trawl) and (5) annual mean CPUE. The calculations were done using the SAS statistical software (Anon., 1988). The area definition used is similar to the stratification used in the 1980-2004 survey (Hvingel, 2007). The multiplicative model was represented in logarithmic form as:

$$\ln(CPUE_{kjmhi}) = \ln(u) + \ln(V_k) + \ln(S_j) + \ln(A_m) + \ln(G_h) + \ln(Y_i) + e_{kjmhi}$$

Where $CPUE_{kjmhi}$ is the mean CPUE for vessel k , fishing in area m in month j during year i with gear type h ($k = 1, \dots, n$; $m = 1, \dots, a$; $j = 1, \dots, s$; $i = 1, \dots, y$; $h = 1, 2, 3$); $\ln(u)$ is overall mean $\ln(CPUE)$; V_k is the effect of the k^{th} vessel; S_j is the effect of the j^{th} month; A_m is effect of the m^{th} area; G_h is the effect of gear type h ; Y_i is the effect of the i^{th} year; e_{kjmhi} is the error term assumed to be normally distributed $N(0, \sigma^2/n)$ where n is the number of observations in the cell. The standardised CPUE indices are the antilog of the year coefficient.

The fishing powers, V , were estimated at individual vessel level. Previously (before 2008) vessels were grouped by engine power and each group was one level in the model. The change was made in order not to have the year effects (the biomass indices) confounded by the recent restructuring of the fleet. Supposedly the least effective vessels are the ones leaving the fishery, which again would make the fishing power of vessel groups increase. By using individual vessels as the unit of fishing power this concern is now accommodated.

Results

Spatial and seasonal distribution

The fishery is conducted mainly in the Hopen area (central Barents Sea) which, along with the Svalbard shelf, is considered the most important fishing ground (Fig. 1). The partial data for 2009 show less activity in the Hopen Deep while more effort is spent further east in "Smutthullet". Information from the industry point to high densities of shrimp in this area and area closing in the traditional Hopen Deep are due to juvenile redfish bycatch as the main reasons for the change in fishing pattern.

The fishery takes place in all months but may in certain years be restricted by ice conditions. The lowest intensity is generally seen in October through March, the highest in May to August (Fig. 2).

Landings

Since the early 1980s annual landings have varied in a cyclic manner with local minima and maxima separated by periods of 4-5 years (Fig. 3). Overall catches have ranged from 26 to 128 kt. The most recent peak was seen in 2000 at approximately 83 kt. Catches thereafter declined to 26 kt in 2008. Based on data until August (logbooks and information from the industry) the total catch of 2009 is estimated at 23 kt.

Discards and by-catch

Discard of shrimp is believed to be small as the fishery is not catch regulated. Small cod, haddock, Greenland halibut and redfish in the size range of 5-25 cm are caught as by-catch. The by-catch of small cod ranged between 2 and 67 million individuals/yr since 1997, while 1-9 million haddock/yr and 0.5 to 14 million Greenland halibut/yr was registered since 2000 (Table 3). Redfish bycatch have been low (<7 mill/yr) in recent years. Details on by-catch are reported to AFWG (ICES, 2008).

Fleet composition and gear

A major restructuring of the fleet towards fewer and larger vessels has taken place since the mid 1990s. An average vessel had at that time around 1000 HP. 15 year later this value had increased to more than 6000 HP (Fig. 4).

Until 1996 the fishery was conducted by using single trawls only. Double trawls were then introduced and in 2002 approximately 2/3 the total effort spent was by using two trawls simultaneously (Fig. 6). In 2000 a few vessels started to experiment with triple trawls: 50% of the effort in 2009 is accounted for by this fishing method.

Standardised CPUE

The fishery dependent index of stock biomass – the standardised CPUE – is indicative of shrimp greater than 16 mm cpl., i.e. of the older male and the female stock combined.

The standardised CPUE declined by 60% from a maximum in 1984 to the lowest value of the time series in 1987 (Fig. 7) (Tab. 2). Since then it has shown an overall increasing trend. A new peak was reached in 2006. The 2007 to 2009 mean values are all about 10% lower than the 2006-value, but is still above the average of the series.

New methods for the calculation of this index series was introduced in 2008 see Hvingel and Thangstad (2008).

Effort

Standardised effort has shown a declining trend since 2000 (Fig. 7).

References

- ANON. 1988. SAS/STAT User's Guide, Release 6.03 Edition. Cary, NC: SAS Institute Inc., 1988. 1028
- ICES 2006. Report of the Pandalus assessment working group 2005. ICES CM 2006/ACFM:10. ref G. 72 pp.
- HVINGEL, C. 2007. Research survey information regarding northern shrimp (*Pandalus borealis*) in the Barents Sea. NAFO SCR Doc. 07/75, Serial No. N5461.
- HVINGEL, C. AND THANGSTAD, T. 2008. The Norwegian fishery for northern shrimp (*Pandalus borealis*) in the Barents Sea. NAFO SCR Doc. 07/74, Serial No. N5460.
- HVINGEL, C., LASSEN, H. AND PARSONS, D. G. 2000. A biomass index for northern shrimp (*Pandalus borealis*) in Davis Strait based on multiplicative modelling of commercial catch-per-unit-effort data (1976 - 1997). J. Northw. Atl. Fish. Sci. 26: 25–31.
- ICES 2007. Report of the Arctic Fisheries Working Group 2007. ICES CM 2007/ACFM:25.

Table 1. Nominal landings ('000 tons) by nation (2009 catch is estimated based on data until August).

Year	Norway	Russia	Others	Total
1970	5.508	0	0	5.508
1971	5.116	0	0	5.116
1972	6.772	0	0	6.772
1973	6.921	0	0	6.921
1974	8.008	0.992	0	9.000
1975	8.197	0	0.002	8.199
1976	9.752	0.548	0	10.300
1977	6.78	12.774	4.854	24.408
1978	20.484	15.859	0	36.343
1979	25.435	10.864	0.39	36.689
1980	35.061	11.219	0	46.280
1981	32.713	10.897	1.011	44.621
1982	43.451	15.552	3.835	62.838
1983	70.798	29.105	4.903	104.806
1984	76.636	43.180	8.246	128.062
1985	82.123	32.104	10.262	124.489
1986	48.569	10.216	6.538	65.323
1987	31.353	6.690	5.324	43.367
1988	32.021	12.32	4.348	48.689
1989	47.064	12.252	3.432	62.748
1990	54.182	20.295	6.687	81.164
1991	39.272	29.434	6.156	74.862
1992	39.603	20.944	8.021	68.568
1993	33.109	22.397	0.806	56.312
1994	20.116	7.108	1.063	28.287
1995	19.337	3.564	2.319	25.220
1996	25.445	5.747	3.320	34.512
1997	29.079	1.493	5.164	35.736
1998	44.792	4.895	6.1031	55.790
1999	52.612	10.765	12.292	75.669
2000	55.333	19.596	8.2413	83.170
2001	43.031	5.846	8.659	57.536
2002	48.799	3.790	8.899	61.488
2003	34.172	2.186	1.599	37.957
2004	35.918	1.170	4.211	41.299
2005	36.966	0.933	3.519	41.418
2006	27.352	0.000	2.282	29.634
2007	25.403	0.009	3.765	29.177
2008	20.638	0.370	5.129	26.137
2009	19.000	0.000	4.000	23.000

Table 2. Realised catch-per-unit-effort (CPUE) and effort (hrs trawled), and standardised (se text) CPUE and effort as proxies for fishable biomass and fishing mortality respectively. Based on Norwegian logbook data. (2009 values are estimated based on data until August).

year	Absolute		Standardised	
	CPUE	Effort	CPUE	Effort
	kg/hr	'000 hrs	index	index
1980	186	189	1.00	1.00
1981	216	152	1.19	0.81
1982	198	219	1.15	1.18
1983	231	306	1.31	1.73
1984	250	306	1.38	2.00
1985	231	356	1.15	2.35
1986	154	315	0.68	2.08
1987	116	270	0.53	1.76
1988	113	282	0.57	1.84
1989	143	330	0.72	1.88
1990	150	361	0.74	2.38
1991	171	230	0.78	2.08
1992	211	188	0.90	1.64
1993	209	159	0.97	1.25
1994	173	116	0.80	0.76
1995	150	129	0.67	0.81
1996	191	133	0.84	0.89
1997	228	127	0.80	0.97
1998	294	153	0.97	1.24
1999	295	178	1.02	1.60
2000	283	195	0.90	1.99
2001	356	121	0.91	1.37
2002	412	119	0.90	1.48
2003	386	88	0.88	0.93
2004	402	89	0.75	1.18
2005	611	61	1.04	0.86
2006	754	36	1.14	0.56
2007	866	30	1.04	0.61
2008	806	26	1.04	0.54
2009	865	22	1.03	0.48

Table 3. Estimated bycatch of cod, haddock, Greenland halibut and redfish (million individuals)

Year	Cod	Redfish	Haddock	Gr. Halibut
1983	14.57	91		
1984	12.6	167		
1985	92.41	198		
1986	10.91	18		
1987	9.87	110		
1988	5.19	46		
1989	1.5	199		
1990	9.02	94		
1991	22.52	51		
1992	25.43	78		
1993	19.23	22		
1994	4.56	23		
1995	5.92	2		
1996	17.1	25		
1997	28.69	24		
1998	67.11	3		
1999	13.43	11		
2000	7.77	15	3.72	13.94
2001	12.87	14	1.75	7.57
2002	2.46	5	9.19	0.19
2003	15.03	0.61	5.52	0.59
2004	2.66	1.1	1.22	0.33
2005	6.46	2.01		
2006	4.9	5.09		
2007	2.52	6.49		
2008	2.24	5.78		
2009	1.99	5.11		




Fig. 1. Distribution of catches by Norwegian vessels 2001-2009 based on logbook information. (2009 only data until August)




Fig. 2. Shrimp in the Barents Sea: Seasonal distribution of fishing effort 2006-may 2009 and mean 1980-2009. Hours trawled in a month as a percentage of total effort of the year. Norwegian data.




Fig. 3. Shrimp in the Barents Sea: Total annual landings. The 2009 projected value is estimated based on data until August.




Fig. 4. Shrimp in the Barents Sea: Mean engine size (horse powers) behind an hour of trawled in the years 1980-2009.




Fig. 5. Shrimp in the Barents Sea: Percentage of total fishing effort spent by using single, double or triple trawls 2000-2009. Norwegian data.




Fig 6. Shrimp in the Barents Sea: Standardised CPUE, Norwegian data.




Fig 7. Shrimp in the Barents Sea: Standardised effort, Norwegian data.

Appendix 1. Output from GLM-run of the Barents Sea index. Gear 55=single trawl, gear 58=double trawl, gear 59= triple trawl. Strata definitions see Hvingel 2007. Vessels are individual vessel identification code.

Class Level Information												
Class	Levels	Values										
strata	8	A B C D E F G H										
gear	3	55 58 59										
vessel	426	Not listed										
year	30	1980 1981 1982 1983 1984 1985 1986 1987 1988 1989 1990 1991 1992 1993										
		1994 1995 1996 1997 1998 1999 2000 2001 2002 2003 2004 2005 2006 2007										
		2008 2009										
month	12	1 2 3 4 5 6 7 8 9 10 11 12										

Number of Observations Used 207627

The GLM Procedure

Dependent Variable: lncpue
Weight: effort

Source	DF	Sum of Squares	Mean Square	F Value	Pr > F
Model	474	970449.893	2047.363	537.40	<.0001
Error	207152	789204.034	3.810		
Corrected Total	207626	1759653.927			

R-Square	Coeff Var	Root MSE	lncpue Mean
0.551500	37.55549	1.951866	5.197286

Source	DF	Type III SS	Mean Square	F Value	Pr > F
strata	7	16724.7879	2389.2554	627.14	<.0001
year	29	145926.3504	5031.9431	1320.80	<.0001
gear	2	217.5188	108.7594	28.55	<.0001
vessel	425	219067.3880	515.4527	135.30	<.0001
month	11	57125.4226	5193.2202	1363.13	<.0001

Dependent Variable: lncpue

Parameter	Estimate	Standard Error	t Value	Pr > t
Intercept	4.027749332 B	0.34215141	11.77	<.0001
strata A	-0.088511638 B	0.00546846	-16.19	<.0001
strata B	0.065451972 B	0.00484634	13.51	<.0001
strata C	0.069427559 B	0.00452606	15.34	<.0001
strata D	0.022476822 B	0.00946694	2.37	0.0176
strata E	0.192760228 B	0.00379745	50.76	<.0001
strata F	0.035372629 B	0.00982695	3.60	0.0003
strata G	0.015430357 B	0.00558441	2.76	0.0057
strata H	0.000000000 B	.	.	.
year 1981	0.177648232 B	0.00976584	18.19	<.0001
year 1982	0.139329902 B	0.00855602	16.28	<.0001
year 1983	0.266437224 B	0.00828722	32.15	<.0001
year 1984	0.323727464 B	0.00850822	38.05	<.0001
year 1985	0.135445168 B	0.00862698	15.70	<.0001
year 1986	-0.389256770 B	0.00890524	-43.71	<.0001
year 1987	-0.629433387 B	0.00944777	-66.62	<.0001
year 1988	-0.556505609 B	0.00914027	-60.89	<.0001
year 1989	-0.326353005 B	0.00879713	-37.10	<.0001
year 1990	-0.307022359 B	0.00874859	-35.09	<.0001
year 1991	-0.251519488 B	0.00923786	-27.23	<.0001
year 1992	-0.101917933 B	0.00952711	-10.70	<.0001
year 1993	-0.026467826 B	0.00998416	-2.65	0.0080
year 1994	-0.221936390 B	0.01103743	-20.11	<.0001
year 1995	-0.400806923 B	0.01107060	-36.20	<.0001
year 1996	-0.175741863 B	0.01072983	-16.38	<.0001
year 1997	-0.223842007 B	0.01093584	-20.47	<.0001
year 1998	-0.031836516 B	0.01069412	-2.98	0.0029
year 1999	0.019467585 B	0.01052680	1.85	0.0644
year 2000	-0.102958364 B	0.01104378	-9.32	<.0001

year	2001	-0.094064542 B	0.01238420	-7.60	<.0001
year	2002	-0.107463211 B	0.01298159	-8.28	<.0001
year	2003	-0.126088160 B	0.01407456	-8.96	<.0001
year	2004	-0.283169391 B	0.01404215	-20.17	<.0001
year	2005	0.043653535 B	0.01586719	2.75	0.0059
year	2006	0.134584350 B	0.01841011	7.31	<.0001
year	2007	0.037386964 B	0.01989219	1.88	0.0602
year	2008	0.041861471 B	0.02750962	1.52	0.1281
year	2009	0.027905245 B	0.03597421	0.78	0.4379
year	1980	0.000000000 B	.	.	.
gear	55	-0.162751005 B	0.02730953	-5.96	<.0001
gear	58	-0.103841192 B	0.02584916	-4.02	<.0001
gear	59	0.000000000 B	.	.	.
vessel	Values not printed				
month	1	0.215322620 B	0.00919751	23.41	<.0001
month	2	0.165440565 B	0.00959051	17.25	<.0001
month	3	0.279146909 B	0.00897436	31.10	<.0001
month	4	0.205020681 B	0.00812038	25.25	<.0001
month	5	0.137988165 B	0.00771301	17.89	<.0001
month	6	0.131569407 B	0.00767751	17.14	<.0001
month	7	0.076352317 B	0.00772399	9.89	<.0001
month	8	0.015490721 B	0.00775569	2.00	0.0458
month	9	-0.157565724 B	0.00797300	-19.76	<.0001
month	10	-0.384902184 B	0.00864761	-44.51	<.0001
month	11	-0.184960315 B	0.00840760	-22.00	<.0001
month	12	0.000000000 B	.	.	.

Continues.....

Plot of STUDENT*ESTIMATE. Legend: A = 1 obs, B = 2 obs, etc.

