



NAFO/ICES PANDALUS ASSESSMENT GROUP—OCTOBER 2010

A preliminary estimate of Atlantic cod (*Gadus morhua*) biomass in West Greenland offshore waters (NAFO Subarea 1) for 2010 and recent changes in the spatial overlap with Northern shrimp (*Pandalus borealis*)

by

Kaj Sünksen, Anja Retzel and Nikoline Ziemer

Greenland Institute of Natural Resources
P.O. Box 570, 3900 Nuuk, Greenland

Abstract

Catches of Atlantic cod in West Greenland waters were compared between the German ground fish survey and the Greenland survey for shrimp and fish. The analysis was restricted to years with regions included in both surveys. The two survey's estimates of cod biomass were closely correlated ($r^2 = 0.86$, $P < 0.001$). Linear regression analysis using data from 18 years revealed that the biomass index of Atlantic cod biomass estimated from the Greenland survey for 2010 would correspond to 14 040 tons in the German survey. This 2010 offshore biomass of Atlantic cod is still very low compared to the biomass in 2006-2007 and the spatial overlap between Atlantic cod and Northern shrimp has decreased in 2010. Therefore the impact of cod on Northern shrimp through predation appears therefore currently to be small.

Introduction

Hvingel (2002, 2003) has incorporated the effect of cod predation into a model for assessing the stock of Northern shrimp (*Pandalus borealis*) at West Greenland (Hvingel and Kingsley, 2002), and STACFIS and the Scientific Council of NAFO have adopted the results from this model since 2002 (Anon., 2004). No consistent time series of Atlantic cod biomass exist for the different stock components and therefore Hvingel and Kingsley (2002) constructed a cod biomass series for West Greenland based on VPA estimates of stock size for East and West Greenland combined for the period 1955 to 1991 and separate survey indices for East and West Greenland offshore waters available since 1982. From 1993 and forward the time series used in the model was derived from a German offshore cod survey in West Greenland waters alone. This survey is conducted each year in autumn (October-November) and its results are not available at the time the assessment for Northern shrimp is carried out. Instead, information on Atlantic cod abundance and biomass from a Greenland bottom trawl survey for shrimp and fish, which commenced in 1988 and which is carried out in summer (July-September), could be used to assess the actual impact of predation by Atlantic cod on the stock of Northern shrimp. The two surveys, however, do not survey exactly the same area and in some years, area coverage or data sampling has been insufficient in one or the other survey.

Wieland and Storr-Paulsen (2004) compared different time series of Atlantic cod biomass in West Greenland offshore waters and concluded that the approach used by Hvingel and Kingsley (2002) could lead to a substantial overestimation of the impact of predation by Atlantic cod on Northern shrimp. A new time series of Atlantic cod was constructed based on VPA estimates by Buch et al. (1994) and survey indices, which avoids some of the inconsistencies of the data series introduced by Hvingel & Kingsley (2002).

This paper presents an update of a comparison of Atlantic cod biomass estimates from the Greenland survey for shrimp and fish and the German ground fish survey from the last year (Sünksen and Ziemer, 2009). It provides a

preliminary estimate of Atlantic cod biomass in West Greenland waters for 2010 that is comparable to the biomass indices from previous years for the area covered by the German survey. Furthermore, the paper examines changes in the spatial overlap between Atlantic cod and Northern shrimp and suggests an alternative time series of Atlantic cod biomass for consideration in the assessment of Northern shrimp.

Materials and Methods

The Greenland bottom trawl survey for fish and shrimps covers the offshore areas at West Greenland and has been conducted from 1988-2010. The area has been expanded through time and since 1992 the survey has covered the area between 72°30'N and 59°15'N (NAFO Division 1A-1F) from the 3 nautical mile limit to the 600 m depth contour (Fig. 1) and the inshore area Disko Bay. The survey area is divided into NAFO Divisions, which are further subdivided into four depth strata (≤ 150 m, 151-200 m, 201-400 m and 401-600 m). The survey was originally designed as a shrimp survey and sampling of fish data was not complete in the period 1988-1991. Since 1992 the sampling of fish has improved and it is now considered as a combined survey for shrimp and fish (Nygaard and Jørgensen, 2010). The survey is designed as a stratified random trawl survey with a minimum of two hauls per stratum. The sampling period is June to August and throughout the entire time period the survey has been conducted by the Greenland Institute of Natural Resources with the research trawler RV *Pâmiut*. Shrimp trawls with a high (10-12 m) vertical opening have been used, which were equipped with a heavy bobbin chain ground gear until 2004 (Skjervøy trawl) or a rockhopper bobbin/rubber disc ground gear since 2005 (Cosmos 2600 trawl). Towing speed has always been about 2.5 knots (see Ziemer, Kingsley & Siegstad 2010 for further details on the survey design).

The German survey is as a stratified random groundfish survey covering the shelf area outside the 3 nautical mile limit and the continental slope down to a depth of 400 m off East and West Greenland between 67°00'N and 59°00'N (ICES area 14 and NAFO Div. 1B-1F, Fig. 1). The Institute for Sea Fisheries, Germany, has conducted this survey annually since 1982. The primary target is cod, and the fishing gear used is a groundfish trawl rigged with a heavy ground gear. Towing speed is 4 knots. The survey provides swept area estimates of abundance (by age) and biomass (all ages pooled) for the East and the West Greenland offshore component, which form the primary basis for the evaluation of the status of the offshore cod stock (ICES, 2010).

Area coverage was incomplete in one or the other surveys the following years and areas:

- NAFO Div. 1A and 1BN have always been covered by the Greenland survey only,
- the northern areas, i.e. 1B and NAFO Div. 1C, were not covered in the German survey in 1995, in 2001 to 2003, and in 2005,
- In 1995 NAFO Div. 1D were not covered by the German survey.

Consequently, new annual indices of Atlantic cod abundance and biomass were calculated for the Greenland survey for all years since 1992 excluding areas for which no corresponding data have been available from both surveys.

Final factors for converting cod catches from 2005 to 2010 made with the Cosmos trawl to Skjervøy catches from the period 1998-2004 are yet not available. Therefore the 2005 to 2010 estimates were converted preliminarily to the old trawl standard using a size-independent conversion of total catches and an adjustment for the smaller average swept area fished with the old trawl (Nygaard and Jørgensen, 2010):

$$\text{Cod}_{\text{Skjervøy}} = 1/1.78 * 1.15 * \text{Cod}_{\text{Cosmos}} = 0.65 * \text{Cod}_{\text{Cosmos}}$$

The major difference between the two trawls is the ground gear, and Lewy *et al.* (2004) reported a conversion factor of about 0.5 in such a case, which is close to the value used here.

The changes in the geographical distribution of Atlantic cod and Northern shrimp were examined using both data from the fishery and from the two surveys described above. Commercial catches reported by NAFO Division have been available for Atlantic cod from Horsted (2000) for the period 1953-1991 and for Northern shrimp from Kingsley and Hvingel (2005) for the period 1975-2005. These values were used to calculate mean latitude of the catches based on the mid of the geographical boundaries of the NAFO statistical areas (Fig. 1) weighted by the aggregated catches. Here, the northern limit of NAFO Division 1A was set to 73°00' N and the southern limit of NAFO Division 1F was set to 59°30' N considering the main distributional range of the fishery for the two species.

Wieland et al. (2006) have applied a geostatistical tool, the global index of collocation (GIC) developed by Bez and Rivoirard (2000), to examine the spatial overlap of Atlantic cod and Northern shrimp based on geo-referenced survey data. These results were used to establish a regression between the GIC and the mean latitude of survey biomass of Atlantic cod. This relationship was used to estimate GIC values back in time, i.e. for the periods in which only commercial catch data by NAFO division and no survey data by set were available or the survey appeared to insufficient due to poor area coverage either in the German or the Greenland survey.

Results and Discussion

Atlantic cod biomasses estimated from the Greenland survey were considerably below the estimates from the German survey in the overlapping years and area, and different trawl types and towing speeds might be the major causes for this. Despite pronounced differences between the two surveys at the upper levels of cod biomass, highly significant correlations ($r^2 = 0.86$, $P < 0.001$, $n = 18$) between the two data series were found, and the equation obtained by linear regression on a log-log scale (Fig. 2) is:

$$\log(\text{Cod biomass}_{\text{German Survey}}) = 0.946 + 0.892 * \log(\text{Cod biomass}_{\text{Greenland Survey}})$$

For the most recent years, the survey estimates of stock biomass indicated a pronounced southern distribution of Atlantic cod. This is also confirmed by the 2010 results where the main concentrations were found in NAFO subarea 1D and 1F (Tab. 1 and Figure 3).

Overall estimates of cod biomass are 7600 tons for the entire surveyed area and 5991 tons for the reduced area that corresponds to the German survey. Taking into account the change of the trawl in the Greenland survey, the latter value converts to 3876 tons that is comparable with the old values obtained with the Skjervøy trawl.

The regression equation derived from the comparison of the German and the Greenland survey in the previous years gives an biomass estimate of 14 040 tons of Atlantic cod to be found by the German survey in fall 2010 in NAFO Div. 1B (south of 67°N) to 1F (Tab. 2 and Figure 4). Although the biomass has increased compared to the very low 2009 level the 2010 biomass must still be considered low compared to the 2006 and 2007 level where the 2003 year class were found offshore giving to rise to biomasses on 160 000 to 170 000 tons respectively.

Acknowledgements

Greenland Institute of Natural Resources would like to thank Dr. Heino Fock involved in the German groundfish survey for their cooperation.

References

- ANON. 2004. Scientific Council Meeting –October/November 2004. *NAFO SCS Doc.*, No. 20, Serial No. N5061.
- ZIEMER, N., KINGSLEY, M. and H. SIEGSTAD, 2010. Results of the Greenland Bottom Trawl Survey for Northern shrimp (*Pandalus borealis*) off West Greenland (NAFO Subarea 1 and Division 0A), 1988-2010. *NAFO SCR Doc.*, No. 57, Serial No. N5848.
- BEZ, N., and J. RIVOIRARD. 2000. Indices of collocation between populations. In: Checkley, DM, Hunter, LR, Motos, L, van der Lingen, CD (eds.). Report of a workshop on the use of the Continuous Underway Fish Egg Sampler (CUFES) for mapping spawning habitats of pelagic fish. GLOBEC Report 14: 48-52.
- BUCH, E, S.A. HORSTED, and H. HOVGÅRD. 1994. Fluctuations in the occurrence of cod in Greenland waters and their possible causes. *ICES Marine Science Symposia* 198: 158-174.
- HORSTED, S.A. 2000. A review of the cod fisheries at Greenland, 1910-1995. *Journal of Northwest Atlantic Fishery Science* 28: 1-112.
- HVINGEL, C. 2002. Assessment, prediction and risk analysis of stock development: Shrimp off West Greenland, 2002. *NAFO SCR Doc.*, No. 157, Serial No. N4786.
- HVINGEL, C. 2003. Assessment, prediction and risk analysis: Stock development and production of northern shrimp off West Greenland. *NAFO SCR Doc.*, No. 73, Serial No. N4912.
- HVINGEL, C., and M. C. S. KINGSLEY. 2002. A framework for the development of management advice on a shrimp stock using a Bayesian approach. *NAFO SCR Doc.*, No. 158, Serial No. N4787.

- ICES. 2010. Report of the North Western Working Group. *ICES C.M.* 2010/ACOM:07
- KINGSLEY, MCS. and C. HVIINGEL. 2005. The fishery for Northern shrimp (*Pandalus borealis*) off West Greenland, 1970-2005. *NAFO SCR Doc.*, No. 83, Serial No. N5188.
- LEWY, P., J.R. NIELSEN and H. HOVGÅRD. 2004. Survey gear calibration independent of spatial fish distribution. *Can. J. Fish. Aquat. Sci.* 61:636-647
- NYGAARD, R. and O. JØRGENSEN. 2010. Biomass and Abundance of Demersal Fish Stocks off West Greenland Estimated from the Greenland Shrimp Survey, 1988-2009. *NAFO SCR Doc.*, No. 30, Serial No. N5788.
- WIELAND, K. and M. STORR-PAULSEN, 2004. A comparison of different time series of Atlantic cod (*Gadus morhua*) biomass at West Greenland and their potential use for the assessment of Northern Shrimp (*Pandalus borealis*) in NAFO Subarea 1. *NAFO SCR Doc.*, No. 71, Serial No. N5051.
- WIELAND, K., M. STORR-PAULSEN, and K. SÜNKSEN. 2007. Response in stock size and recruitment of northern shrimp (*Pandalus borealis*) to changes in predator biomass and distribution in West Greenland waters. *Journal of Northwest Atlantic Fishery Science* 39: 21-33.
- SÜNKSEN, K. and N. Ziemer. 2009. A preliminary estimate of Atlantic Cod (*Gadus morhua*) biomass in West Greenland offshore waters (NAFO Subarea 1) for 2009 and recent changes in the spatial overlap with Northern shrimp (*Pandalus borealis*). *NAFO SCR Doc.*, No. 65, Serial No. N5726.

Tab. 1: Observed Cod biomass (tons) by survey and area. Greenland survey as observed, i.e. without conversion -: not covered. +: covered but not to be included in the comparison.

| Year | German survey Stratum | | | | | Greenlandic survey Stratum (without > 400 m depth) | | | | |
|------|--------------------------|-------|-------|--------|--------|---|------|------|-------|-------|
| | 1BS+1C | 1D | 1E | 1F | Sum | 1BS+1C | 1D | 1E | 1F | Sum |
| 1992 | 337 | 94 | 35 | 257 | 723 | 109 | 118 | 0 | 2 | 229 |
| 1993 | 260 | 108 | 72 | 0 | 440 | 30 | 39 | 124 | 5 | 193 |
| 1994 | 59 | 28 | 32 | 18 | 137 | 47 | 0 | 1 | 0 | 48 |
| 1995 | - | - | 39 | 19 | 58 | + | + | 3 | 4 | 7 |
| 1996 | 96 | 46 | 44 | 202 | 388 | 15 | 23 | 27 | 49 | 114 |
| 1997 | 42 | 22 | 105 | 106 | 275 | 55 | 0 | 0 | 0 | 55 |
| 1998 | 77 | 17 | 1 | 46 | 141 | 0 | 47 | 50 | 3 | 100 |
| 1999 | 53 | 17 | 143 | 77 | 290 | 6 | 17 | 1 | 0 | 24 |
| 2000 | 249 | 188 | 157 | 44 | 638 | 75 | 9 | 2 | 46 | 132 |
| 2001 | - | 361 | 2056 | 185 | 2602 | + | 178 | 98 | 100 | 376 |
| 2002 | - | 178 | 285 | 1952 | 2415 | + | 1489 | 42 | 150 | 42 |
| 2003 | - | 283 | 1535 | 758 | 2576 | + | 453 | 118 | 46 | 555 |
| 2004 | 508 | 639 | 1196 | 4245 | 6588 | 183 | 680 | 685 | 305 | 1853 |
| 2005 | - | 1457 | 3884 | 21850 | 27191 | + | 1317 | 1047 | 38765 | 41129 |
| 2006 | 479 | 91733 | 5320 | 64593 | 162125 | 420 | 5180 | 545 | 19111 | 25256 |
| 2007 | 2408 | 3644 | 15694 | 151432 | 173178 | 223 | 582 | 471 | 21091 | 22367 |
| 2008 | 1745 | 798 | 3035 | 35535 | 41113 | 1051 | 2133 | 4106 | 23719 | 31010 |
| 2009 | 37 | 1204 | 1049 | 1720 | 4010 | 458 | 314 | 361 | 308 | 1441 |
| 2010 | | | | | | 419 | 2058 | 787 | 2728 | 5991 |

Tab. 2: Time series of Atlantic cod biomass (tons) (*: using mean VPA to survey ratio based on Wieland & Storr-Paulsen, (2004).

| Year | Cod biomass | | | Spatial overlap GIC Cod - Shr observed | 'effective' Cod biomass |
|------|---------------------------|-----------------------|------------------------------|--|-------------------------------|
| | VPA (Buch et al. 1984) | Survey (ICES 2008) | VPA and survey combined * | | |
| 1950 | 4076.5 | | 4076.5 | | |
| 1951 | 3722.9 | | 3722.9 | | |
| 1952 | 3285.7 | | 3285.7 | | |
| 1953 | 2855.3 | | 2855.3 | 0.692 | 1977.1 |
| 1954 | 2848.3 | | 2848.3 | 0.714 | 2033.9 |
| 1955 | 2731.4 | | 2731.4 | 0.703 | 1919.1 |
| 1956 | 2298.7 | | 2298.7 | 0.693 | 1592.7 |
| 1957 | 2037.5 | | 2037.5 | 0.684 | 1392.9 |
| 1958 | 1866.2 | | 1866.2 | 0.674 | 1258.3 |
| 1959 | 1687.7 | | 1687.7 | 0.719 | 1212.6 |
| 1960 | 1823.1 | | 1823.1 | 0.706 | 1287.3 |
| 1961 | 1793.9 | | 1793.9 | 0.704 | 1263.1 |
| 1962 | 1469.2 | | 1469.2 | 0.716 | 1051.3 |
| 1963 | 1328.4 | | 1328.4 | 0.686 | 911.2 |
| 1964 | 1327.6 | | 1327.6 | 0.676 | 898.1 |
| 1965 | 1345.3 | | 1345.3 | 0.706 | 950.2 |
| 1966 | 1254.9 | | 1254.9 | 0.709 | 889.2 |
| 1967 | 1167.5 | | 1167.5 | 0.683 | 797.4 |
| 1968 | 904.9 | | 904.9 | 0.639 | 578.1 |
| 1969 | 637.4 | | 637.4 | 0.611 | 389.7 |
| 1970 | 442.2 | | 442.2 | 0.554 | 244.9 |
| 1971 | 372.8 | | 372.8 | 0.587 | 218.7 |
| 1972 | 283.0 | | 283.0 | 0.678 | 191.9 |
| 1973 | 179.5 | | 179.5 | 0.643 | 115.4 |
| 1974 | 132.1 | | 132.1 | 0.642 | 84.7 |
| 1975 | 108.5 | | 108.5 | 0.629 | 68.2 |
| 1976 | 228.8 | | 228.8 | 0.579 | 132.5 |
| 1977 | 251.7 | | 251.7 | 0.574 | 144.5 |
| 1978 | 253.5 | | 253.5 | 0.672 | 170.3 |
| 1979 | 217.3 | | 217.3 | 0.670 | 145.6 |
| 1980 | 240.4 | | 240.4 | 0.680 | 163.4 |
| 1981 | 178.3 | | 178.3 | 0.619 | 110.4 |
| 1982 | 190.9 | 138.0 | 190.9 | 0.518 | 98.8 |
| 1983 | 134.0 | 80.9 | 134.0 | 0.461 | 61.7 |
| 1984 | 79.0 | 28.3 | 79.0 | 0.479 | 37.8 |
| 1985 | 51.7 | 34.1 | 51.7 | 0.482 | 25.0 |
| 1986 | 38.4 | 75.9 | 38.4 | 0.510 | 19.6 |
| 1987 | 466.7 | 628.0 | 466.7 | 0.604 | 282.1 |
| 1988 | 481.2 | 646.7 | 481.2 | 0.618 | 297.3 |
| 1989 | 403.2 | 332.6 | 403.2 | 0.370 | 149.1 |
| 1990 | | 42.1 | 42.1 | 0.289 | 12.2 |
| 1991 | | 6.8 | 6.8 | 0.313 | 2.1 |
| 1992 | | 0.7 | 0.7 | 0.523 | 0.4 |
| 1993 | | 0.4 | 0.4 | 0.633 | 0.3 |
| 1994 | | 0.1 | 0.1 | (0.356) | 0.1 |
| 1995 | | 0.1 | 0.1 | (0.120) | 0.0 |
| 1996 | | 0.4 | 0.4 | 0.280 | 0.1 |
| 1997 | | 0.3 | 0.3 | 0.490 | 0.1 |
| 1998 | | 0.1 | 0.1 | 0.390 | 0.0 |
| 1999 | | 0.3 | 0.3 | 0.496 | 0.1 |
| 2000 | | 0.6 | 0.6 | 0.643 | 0.4 |
| 2001 | | 2.6 | 2.6 | 0.462 | 1.2 |
| 2002 | | 2.4 | 2.4 | 0.278 | 0.7 |
| 2003 | | 2.6 | 2.6 | 0.398 | 1.0 |
| 2004 | | 6.6 | 6.6 | 0.257 | 1.7 |
| 2005 | | 27.2 | 27.2 | 0.074 | 2.0 |
| 2006 | | 162.1 | 162.1 | 0.220 | 35.7 |
| 2007 | | 173.2 | 173.2 | 0.139 | 24.0 |
| 2008 | | 41.1 | 41.1 | 0.156 | 6.4 |
| 2009 | | 4.0 | 4.0 | 0.602 | 2.4 |
| 2010 | | | 14.0 | 0.315 | 4.4 |




Fig. 1. Survey coverage for the Greenland fish and shrimp survey and the German ground fish survey in West Greenland offshore waters. (*: NAFO divisions 1B and 1C were not covered by the German survey in 1995, in 2001 to 2003 and in 2005).




Fig. 2. Relationship between estimates of cod biomass in the overlapping area of the German groundfish survey and the Greenland Bottom Trawl Survey, 1992-2009. Dashed and dotted lines indicate limits of the 95% confidence and prediction intervals, respectively.




Fig. 3. Geographical distribution of cod density (in kg/km^2) in the Greenland Bottom Trawl Survey in 2010. Numbers in parentheses denotes number of tows.




Fig. 4. Survey indices of Atlantic cod biomass for West Greenland offshore waters, 1982-2010 (1982-2009: original estimates from the German groundfish survey; 2010: estimate based on the Greenland Bottom Trawl Survey).