Serial No. N5896



NOT TO BE CITED WITHOUT PRIOR REFERENCE TO THE SECRETARIAT

Fisheries Organization

NAFO SCR Doc. 11/014

SCIENTIFIC COUNCIL MEETING – JUNE 2011

Physical Oceanographic Conditions on the Scotian Shelf and in the eastern Gulf of Maine (NAFO areas 4V,W,X) during 2010

B. Petrie, R. G. Pettipas and D. Hebert

Department of Fisheries and Oceans, Maritimes Region Ocean Sciences Division, Bedford Institute of Oceanography P.O. Box 1006, Dartmouth, N.S. B2Y 4A2

Abstract

A review of the 2010 physical oceanographic conditions on the Scotian Shelf and in the eastern Gulf of Maine and adjacent offshore areas indicates that above normal conditions prevailed. The climate index, a composite of 18 selected, normalized time series, averaged +1.0 (\pm 0.8) SD with 13 of the 18 variables more than 0.5 SD above normal; compared to the other 41 years, 2010 ranks as the 4th warmest. The anomalies did not show a strong spatial variation. Bottom temperatures were above normal but not exceptionally so with anomalies for NAFO areas 4Vn, 4Vs, 4W, 4X and eastern Georges Bank of +0.2°C (0.5SD), +0.4°C (0.6SD), +0.6°C (0.8SD), +0.3°C (0.5SD), and +0.4°C (0.7SD) respectively.

Introduction

This document briefly describes air temperature, ice area and volume, and ocean temperature variability of Scotian Shelf and Gulf of Maine waters during 2010 (see Fig. 1 for the study area, Fig. 2 for area names of time series presented in document). The results are derived from data obtained at coastal and long-term monitoring stations, on annual ecosystem surveys and Atlantic Zone Monitoring Program (AZMP) missions, and from ships-of-opportunity and other research cruises. Most of the data are available in the BIO temperature and salinity (CLIMATE) database¹, which is updated several times per year from the national archive maintained by Integrated Science Data Management (ISDM), Department of Fisheries and Oceans (DFO) in Ottawa. Our analyses use data archived prior to 15 February 2010. Additional hydrographic data were obtained directly from DFO fisheries surveys.

Many of the products which have been presented previously in the overview are now available on the ISDM website devoted to the AZMP program². The products available include sections from the AZMP spring and fall surveys, time series of physical properties from fixed stations, climate indexes such as coastal temperature series, frontal positions, and bottom temperatures from ecosystem surveys. In addition, the availability of quality controlled data from sources such as CLIMATE allow individuals to develop the product that most suits their needs. For these reasons, we present an abbreviated overview to give a general idea of environmental variability in 2010.

In order to detect interannual variability and long-term trends of the time series presented, we have removed the potentially large seasonal cycle by determining the monthly differences, i.e. the anomalies, from the long-term means. In some cases, we present the standardized anomaly (anomaly divided by the standard deviation). When possible, long-term monthly and annual means, and standard deviations (SD) are based on 1981-2010 data.

¹http://www.mar.dfo-mpo.gc.ca/science/ocean/database/data_query.html

²http://www.meds-sdmm.dfo-mpo.gc.ca/isdm-gdsi/azmp-pmza/index-eng.html

Temperature and salinity conditions in the Scotian Shelf, Bay of Fundy and Gulf of Maine regions are determined by many processes: heat transfer between the ocean and atmosphere, inflow from the Gulf of St. Lawrence supplemented by flow from the Newfoundland Shelf, exchange with offshore slope waters, local mixing, freshwater runoff, direct precipitation and melting of sea-ice. The Nova Scotia Current is the dominant inflow, originating in the Gulf of St. Lawrence and entering the region through Cabot Strait (see inset, Fig. 1). The Current, whose path is strongly affected by topography, has a general southwestward drift over the Scotian Shelf and continues into the Gulf of Maine where it contributes to the counter-clockwise mean circulation. The properties of shelf waters are modified by mixing with offshore waters from the continental slope. These offshore waters are generally of two types, Warm Slope Water, with temperatures in the range of 8-13°C and salinities from 34.7-35.6, and Labrador Slope Water, with temperatures from 3.5°C to 8°C and salinities from 34.3 to 35. Shelf water properties have large seasonal cycles, east-west and inshore-offshore gradients, and vary with depth (Petrie et al. 1996).

Air Temperatures

Annual air temperature anomalies for six sites in the Scotian Shelf-Gulf of Maine region are shown in Fig. 3 and Table 1. In 2010, annual air temperature anomalies were positive at all sites and ranged from $\pm 1.1^{\circ}$ C (Sable Island) to $\pm 1.7^{\circ}$ C (Shearwater). These values represent an increase from 2009 observations and all are approximately 2 standard deviations above the long-term mean; furthermore, the time series of annual anomalies indicates that all sites feature increasing temperatures over the long term with decadal scale variability superimposed. For shorter periods, this can lead to no trend or decreasing temperatures (Fig. 3). Linear trends from 1900 to present from Boston, Shearwater, Sydney and Sable Island correspond to changes of $\pm 1.7^{\circ}$ C, $\pm 1.4^{\circ}$ C, $\pm 1.0^{\circ}$ C and 0.9° C/century respectively; the trends from Yarmouth and Saint John are smaller, 0.6 and 0.3° C/century.

The individual anomalies for all 6 sites and their sum, displayed in Fig. 4, illustrate three points. 1) In the 111 year time series shown, 2010 overall was the warmest year for the region as a whole. 2) For most years the anomalies have the same sign. i.e. the stacked bars and the scatter plot coincide. Since 1900, 88 of the 111 years had 4 or more stations with the annual anomalies having the same signs; for 56 years, all 6 stations had anomalies with the same sign. This indicates that the spatial scale of the air temperature patterns is greater than the largest spacing among sites. In fact, plotting the correlations between annual anomalies against station separations yields an e-folding scale 1800 km (Petrie and Pettipas., 2010). The e-folding scale (=1/k) is determined from a least squares fit of an exponential function 1*exp(-kx) to the correlations among the six time series; it represents the distance at which the correlation equals 1/e, where e is the base of natural logarithms. 3) The time scale of the dominant variability has been changing from longer periods for 1900-1954 to shorter periods for 1955-2010. For the earlier period, a lagged autocorrelation analysis gives an e-folding scale of 3 years; whereas, for the later period, the scale was 0.6 years, i.e. less than the sampling interval of the series. In this case, the e-folding scale is based on the autocorrelation of the average annual temperature anomaly for the 6 sites (Petrie and Pettipas, 2010).

Site	Annual Anomaly 2010 (°C)		Monthly SD	1981-2010 Annual	
			2010 (°C)		
	Observed	Normalized		Mean (°C)	SD (°C)
Sydney	+1.5	+1.9	1.4	5.87	0.81
Sable I.	+1.1	+1.7	0.7	7.88	0.68
Shearwater	+1.7	+2.3	0.9	6.99	0.74
Yarmouth	+1.3	+2.1	0.7	7.16	0.62
Saint John	+1.6	+2.2	1.3	5.19	0.74
Boston	+1.2	+2.0	1.3	10.91	0.60

 Table 1. Air temperature statistics 2010 for Scotian Shelf and Gulf of Maine

Sea Ice

The greater part of sea ice on the Scotian Shelf originates in the Gulf of St. Lawrence and is transported through Cabot Strait pushed by northwesterly winds and ocean currents. Sydney Bight and the southern coast of Cape Breton are typically the only areas heavily affected by ice. In 2010, there was nearly no ice on the Scotian

Shelf from December 2009 until the end of the season in May 2010 (Fig. 5). Over the winter, ice duration was below normal. Note that in the duration maps, ice boundaries exceed those of first and last appearance. We consider an area that has not had ice in the current year but has had ice for 5 years of the 30 year climatological period to have a negative duration anomaly. Consequently, those areas are colour mapped as light red.

The ice areas and volumes are shown in Fig. 6 and compiled in Tables 2 and 3. The heaviest months, January to April, featured below normal coverage and volumes. Overall, the January to April 2010 coverage and ice volume were the second lowest in the 42 year long record. Only 1969 had lesser coverage and volume; the difference between these 2 years is within the uncertainty of the observations. Note that the ice statistics are based on the period 1971-2000 because the update to the new base period of 1981-2010 is a lengthy, complex task; in the coming year the base period will be revised to 1981-2010.

Month	09-10 Ice Year	1971-2000 Mean	Anomaly (km ²)	Normalized
	(km^2)	(km^2)		
Jan	90	1700	-1600	-0.6
Feb	60	13900	-13800	-1.4
Mar	10	18000	-18000	-1.5
Apr	2	5800	-5800	-1.1

 Table 2. Ice area statistics, Scotian Shelf

Table 5. ree volume statistics, beotian blief					
Month	09-10 Ice Year	1971-2000 Mean	Anomaly (km ³)	Normalized	
	(km^3)	(km^3)			
Jan	< 0.01	0.25	-0.25	-0.1	
Feb	< 0.01	3.5	-3.4	-0.7	
Mar	< 0.01	7.4	-7.3	-2.1	
Apr	< 0.01	3.6	-3.6	-11	

Table 3. Ice volume statistics. Scotian Shelf

Remotely-Sensed Sea Surface Temperature

A 9 km resolution Pathfinder sea surface temperature database is maintained at BIO. The Pathfinder dataset runs from January 1985 to December 2009; to provide data for 2010, we used the sea surface temperature data downloaded from the satellites by the remote sensing group in the Ocean Research & Monitoring Section (ORMS). Comparison of the Pathfinder and ORMS temperatures during the common time period led to a conversion equation SST(Pathfinder) = 0.976*SST(ORMS)+0.46 with an $r^2=0.98$. We adjusted the ORMS observations to bring them in line with the longer Pathfinder series. Anomalies were based on 1985-2009 averages.

Annual anomalies for 7 subareas in the Scotian Shelf-Gulf of Maine region were determined from the averages of monthly anomalies (Fig. 7a, Table 4). Periods of 1 year and longer accounted on average for 38% of the overall variance determined from the monthly anomalies. The annual anomalies during 2010 ranged from -0.4°C (-0.7 SD) over Georges Bank to $+0.9^{\circ}$ C (+1.3 SD) on the eastern Scotian Shelf. Six of the eight areas had positive anomalies; all 6 were greater than or equal to +0.78 SD. Over the lengths of the records all areas show increasing temperature trends, based on a linear least squares fit, corresponding to temperature changes from a lowest value 0.3° C (Georges Bank) to a highest value of 1.4° C (ESS).

The overall coherent variability of the annual temperature anomalies in the seven regions suggested a principal component analysis might be revealing. The leading mode, PCA1, captured 71% of the variance and all loadings had similar amplitudes, meaning roughly equal contributions from each series; PCA2 accounted for an additional 19% of the variance with negative loadings in the eastern half of the region, changing to positive values roughly to the west of the central Scotian Shelf (Fig. 7b).

Site	2010 SST	2010 SST	1985-2010	1985-2010	ΔT (°C)
	Anomaly	Anomaly	Mean Annual	SST	1985-2010
	(°C)	Normalized	SST (°C)	Std. Dev.	
				(°C)	
Cabot Str.	+0.9	+1.5	6.16	0.59	1.2
ESS	+0.9	+1.3	7.37	0.70	1.4
CSS	+0.7	+1.1	8.86	0.63	1.3
Western Bank	+0.6	+0.6	9.30	0.65	1.1
WSS	+0.5	+0.8	8.60	0.59	0.9
Lurcher Shoal	-0.1	-0.2	7.77	0.59	0.4
Bay of Fundy	+0.4	+0.8	7.88	0.55	0.8
Georges Bank	-0.4	-0.7	10.56	0.54	0.3

Table 4. SST anomalies, 2010 and long-term SST statistics.

Coastal Temperatures and Salinities

Coastal sea surface temperatures have been collected at St. Andrews (New Brunswick) and Halifax (Nova Scotia) since the 1920s (Fig. 8). In 2010, the SST anomalies were $\pm 1.0^{\circ}$ C (± 1.8 SD) for St. Andrews and $\pm 0.4^{\circ}$ C (± 0.54 SD) for Halifax, increases of 0.9°C and 0.5°C from 2009 values.

Temperature and salinity measurements, for the most part sampled monthly, have been taken since 1924 at Prince 5, at the entrance to the Bay of Fundy (Fig. 1). It is the longest continuously operating hydrographic monitoring site in eastern Canada. Its waters are generally well-mixed from the surface to the bottom (90 m). The depth-averaged (0-90 m) temperature, salinity and density anomaly time series are shown in Fig. 8(C-E). In 2010, the annual temperature anomaly was $+0.97^{\circ}$ C (+1.8 SD) and the salinity anomaly was -0.07 (-0.31 SD). These represent changes of $+0.88^{\circ}$ C and -0.11 from the 2009 values. The density anomaly variability is largely accounted for by the salinity anomaly (73%) and secondarily by the temperature anomaly (27%).

Scotian Shelf and Gulf of Maine Temperatures from Long-term Stations

Drinkwater and Trites (1987) tabulated monthly mean temperatures and salinities from available bottle data for areas on the Scotian Shelf and in the eastern Gulf of Maine that generally correspond to topographic features such as banks and basins (Fig. 2). Petrie et al. (1996) updated their report using these same areas and all available hydrographic data. We present time series of annual mean and filtered (5 year running means) temperature anomalies at selected depths for five areas (Fig. 9). The Cabot Strait temperatures represent the slope waters entering the Gulf of St. Lawrence along Laurentian Channel; the Misaine Bank series characterizes the near bottom temperatures on the eastern Scotian Shelf; the deep Emerald Basin anomalies represent the slope water intrusions onto the Shelf that are subsequently trapped in the deep inner basins; the Lurcher Shoals observations define the ocean climate in SW Nova Scotia and the shallow waters entering the Gulf of Maine via the Nova Scotia Current; finally, the Georges Basin series indicates the slope waters entering the Gulf of Maine through the Northeast Channel. Annual anomalies are based on the averages of monthly values; however, observations are not available for each month in each area. For Cabot Strait, Misaine Bank, Emerald Basin, Lurcher Shoals and Georges Basin, 2010 annual anomalies are based on observations from 6, 7, 6, 3 and 6 months.

In 2010, the annual anomalies were $\pm 0.27^{\circ}$ C (± 0.8 SD) for Cabot Strait 200-300 m, $\pm 0.43^{\circ}$ C (± 0.7 SD) for Misaine Bank 100 m, $\pm 0.63^{\circ}$ C (± 0.8 SD) for Emerald Basin 250 m, $\pm 0.09^{\circ}$ C (± 0.1 SD) for Lurcher Shoals 50 m, and $\pm 0.95^{\circ}$ C (± 1.8 SD) for Georges Basin 200 m. These values correspond to changes of $\pm 0.8^{\circ}$ C, $\pm 0.0^{\circ}$ C, $\pm 1.9^{\circ}$ C, $\pm 1.1^{\circ}$ C and $\pm 1.8^{\circ}$ C, respectively over 2009 values.

Temperatures during the Summer Groundfish Surveys

The broadest spatial CTD coverage of the Scotian Shelf is obtained during the annual July DFO ecosystem survey which covers the Scotian Shelf from Cabot Strait to the Bay of Fundy. The deep water boundary of the survey is marked roughly by 200 m isobath along the shelf break at the Laurentian Channel, the outer Scotian Shelf, and the

Northeast Channel into the Gulf of Maine towards the Bay of Fundy. A total of 195 CTD stations were taken during the 2010 survey and an additional 171 bottom temperature stations were obtained as part of the ITQ (Individual Transferable Quota) fleet survey. The groundfish survey takes 1 month to complete with the area west of Halifax sampled first and the area east of Halifax sampled second.

The temperatures from both surveys were combined and interpolated onto a 0.2° by 0.2° latitude-longitude grid using an objective analysis procedure known as optimal estimation. The interpolation method uses the 15 "nearest neighbours" with a horizontal length scale of 30 km and a vertical length scale of 15 m in the upper 30 m and 25 m at deeper depths. Data near the interpolation grid point are weighted proportionately more than those farther away. Temperatures were optimally estimated for 0, 50, 100 m and near bottom.

Bottom temperatures ranged from 3.7° C in area 4Vs to 7.6° C in 4X illustrating the difference in the environmental conditions across the Shelf. The anomalies were positive for these NAFO areas in 2010: +0.21°C (+0.50 SD) in 4Vn; +0.43°C (+0.61 SD) in 4Vs; +0.55°C (+0.74 SD) in 4W; and +0.33°C (+0.46 SD) in 4X (Fig. 10 A-D). Compared to 2009, bottom temperatures increased in areas 4Vn, 4W and 4X by 0.32, 0.26 and 0.33°C; temperature remained the same in area 4Vs.

The volume of the Cold Intermediate Layer (CIL), defined as waters with temperatures $<4^{\circ}$ C, was estimated from the full depth CTD profiles for the region from Cabot Strait to Cape Sable (Fig. 10E). For the period 1970 to 1989, the number of CTD profiles per year was limited; therefore, 5 year blocks of data, e.g. 1970-1974, centre date 1972, were used as input for the procedure to map the irregularly spaced data onto a regular grid. The data were then incremented by 1 year and a new set of estimates made (i.e., 1970-74, 1971-75, ...). This procedure is similar to filtering the data for the 1970-89 period, effectively reducing the variance. Thus the long-term mean and particularly the SD (based on the 1972-2000 data in Fig. 10E) could be affected. We expect that the true SD is higher than the one derived here.

There is considerable variation in the volume of the CIL since 1998 (Fig. 10E). In 2010, the observed volume of 4840 km³ was 0.6 SD less than the long-term mean value of 5510 km³ and a slight, 100 km³ decrease from 2009.

Density Stratification

Stratification of the near surface layer influences physical and biological processes in the ocean such as vertical mixing, the ocean's response to wind forcing, the timing of the spring bloom, vertical nutrient fluxes and plankton speciation. Under increased stratification, there is a tendency for more primary production to be recycled within the upper mixed layer and hence less available for the deeper layers. We examined the variability in stratification by calculating the density (sigma-t) difference between 0 and 50 m. The density differences were based on monthly mean density profiles calculated for each area in Fig. 2. The long-term monthly mean density gradients for 1981-2010 were estimated; these were subtracted from the individual monthly values to obtain monthly anomalies. Annual anomalies were estimated by averaging all available monthly anomalies within a calendar year. This could be misleading if, in a particular year, most data were collected in months when stratification was weak, while in another year, sampling occurred when stratification was strong. However, initial results, using normalized monthly anomalies, were qualitatively similar to the plots presented here. The annual anomalies and their 5-year running means were then calculated for an area-weighted combination of subareas 4-23 on the Scotian Shelf (Fig. 2, 11). A value of 0.01 (kg m⁻³)/m represents a difference of 0.5 kg m⁻³ over 50 m.

The dominant feature is the period from about 1950 to 1990 which featured generally below average stratification in contrast to the past 20 years which is characterized by above normal values. Stratification on the Scotian Shelf in 2010 strengthened compared to 2009; overall 2010 ranked as the third most stratified in the 64 year record.

Sea Level

Sea level is a primary variable in the Global Ocean Observing System. Relative sea level is measured with respect to a fixed reference point on land. Consequently, relative sea level consists of two major components: one due to true changes of sea level and a second caused by sinking or rising of the land. In Atlantic Canada, post-glacial rebound (PGR) is causing the area roughly south (north) of the north shore of the Gulf of St. Lawrence to sink (rise) in response to glacial retreat; this results in an apparent rise (fall) of sea level. Tushingham and Peltier (1991) estimate a PGR component of the sea level trend of 23 cm/century at Halifax.

Relative sea level at Halifax (1920-2010) is plotted as monthly means and as a filtered series using a 5-year running-mean filter (Fig. 12A). The linear trend of the monthly mean data has a positive slope of 32.6 (\pm 0.7) cm/century, lower than the value of 36.7 cm/century (1897-1980) given by Barnett (1984). In 2010, relative sea level at Halifax increased substantially by 6.9 cm above the 2009 level. However, the interesting feature of the data is the long-term variation that has occurred since the 1920s. In Fig. 12(B) we show the differences of the annual sea level from the long-term trend. It is apparent that from the 1920s to the early 1970s, the trend was greater than the trend calculated using all of the data. In fact, the trend from 1920-72 was 40.4 (\pm 1.4) cm/century; for the period 1972-2010, the value was 26.0 (\pm 4.6) cm/century, i.e. close to the PGR value predicted by the Tushingham-Peltier model. Several potential causes of this decadal scale variability have been examined; we still do not understand the cause of these changes.

Summary

A graphical summary of many of the time series already shown indicates that the periods 1987-1993 and 2003-2004 were predominantly colder than normal and 1999-2000 was warmer than normal (Fig. 13). The period 1979-1986 also tends to be warmer than normal. In this figure, annual anomalies based on the 1981-2010 means have been normalized by dividing by the 1981-2010 standard deviations for each variable. The results are displayed as the number of standard deviations above (red) and below (blue) normal. In 2010, 20 of the 22 series shown had positive anomalies; 6 variables were within 0.5 SD of their normal values and were concentrated in the western end of the region. Of the 16 remaining series, all were more than 0.5 SD greater than normal. In 2010, the average (median) normalized anomaly was 1.0 (0.75), the fourth (eighth) highest in the 41 year series. The standard deviation of the normalized anomalies was 0.77, the 24th highest in the series. These statistics indicate that 2010 was an exceptional warm year with a fairly uniform distribution of anomalies throughout the region.

Eighteen selected variables of the mosaic plot are summarized as a combination bar and line-scatter plot in Fig. 14. This plot is an attempt to derive an overall climate index for the area. We have selected "profiles" for the eastern (Misaine), central (Emerald) and western (Lurcher) Scotian Shelf, the Bay of Fundy (Prince 5) and Georges Bank. In addition, we have included the spatially comprehensive but temporally limited July groundfish survey bottom temperatures (4Vn,s, 4W and 4X) and surface temperatures for Halifax and St. Andrews because of their long-term nature. The bar components are colour coded so that for any year the contribution of each variable can be determined and systematic spatial patterns seen. The height of each variable's contribution to the bar depends on its magnitude. The positive components are stacked on the positive side, the negative components on the negative side. The sum of the normalized anomalies (difference between the positive and negative stacks) is shown as a black line connecting grey circles. This is a measure of whether the year tended to be colder or warmer than normal and can serve as an overall climate index.

The cold periods of 1987-1993 and 2003-2004 and the warm period of 1999-2000 are apparent. Systematic differences from the overall tendency within a given year are also evident. The overall index in 2010 averaged +1.0 (± 0.8) SD with 13 of the 18 variables more than 0.5 SD above normal; compared to the other 41 years, 2010 ranks as the 4th warmest. The anomalies did not show a strong spatial distribution in 2010. The leading mode of a principal component analysis of the 18 series captured 40% of the variance with all loadings having the same sign. The loadings of 16 of the 18 variables were strong (0.18 to 0.30) with weak contributions from the Emerald Basin 250 m (0.02) and Misaine Bank 0 m (0.1) series. The temporal variability of the leading mode was essentially the same ($r^2 = 0.99$) as shown by the sum of the anomalies (Fig. 14).

Acknowledgements

We wish to thank the many individuals who provided data for this paper, including: Mathieu Ouellet of the Marine Environmental Data Service in Ottawa; Paul McCurdy of the Biological Station in St. Andrews, for providing St. Andrews and Prince 5 data. We also thank Eugene Colbourne and Gary Maillet for their comments which improved the document.

References

- Barnett, T. 1984. The estimation of "global" sea level change: a problem of uniqueness. J. Geophys. Res. 89, 7980-7988.
- Drinkwater, K. F. and R. W. Trites 1987. Monthly means of temperature and salinity in the Scotian Shelf region. Can. Tech. Rep. Fish. Aquat. Sci. 1539: 101 p.
- Petrie, B., K. Drinkwater, D. Gregory, R. Pettipas, and A. Sandström 1996. Temperature and salinity atlas for the Scotian Shelf and the Gulf of Maine. Can. Data Rep Hydrog. Ocean Sci. 171: 398 p.
- Petrie, B. and R. G. Pettipas 2010. Physical Oceanographic Conditions on the Scotian Shelf and in the eastern Gulf of Maine (NAFO areas 4V,W,X) during 2009. NAFO SCR Doc. 10/12, 22 p.
- Tushingham, A. and R. Peltier 1991. Ice 3-G: a new global model of late Pleistocene deglaciation based on geophysical predictions of post-glacial relative sea level change. J. Geophys. Res. 96, 4497-4523.



Fig. 1. The Scotian Shelf and the Gulf of Maine showing hydrographic stations, standard sections and topographic features. The dotted lines indicate the boundaries of the NAFO Subareas. Inset depicts major circulation features.



Fig. 2. Areas on the Scotian Shelf and eastern Gulf of Maine from Drinkwater and Trites (1987).



Fig. 3. Annual air temperature anomalies (dashed line) and 5-yr running means (solid line) at selected sites in Scotian Shelf-Gulf of Maine region.



Fig. 4. The contributions of each of the annual temperature anomalies for 6 Scotian Shelf-Gulf of Maine sites are shown as a bar chart and their summation as a time series (grey circles, black line).



Fig. 5. The time when ice first appeared during 2010 in days from the beginning of the year (left, top panel) and its anomaly from the 1971-2000 mean in days (left, bottom panel). Negative (positive) anomalies indicate earlier (later) than normal appearance. The time when ice was last seen in 2010 in days from the beginning of the year (centre, top panel) and its anomaly from the 1971-2000 mean in days (centre, bottom panel). Positive (negative) anomalies indicate later (earlier) than normal disappearance. The duration of ice in days (right, top panel) during 2010 and the anomalies from the 1971-2000 mean in days (right, bottom panel). Positive (negative) anomalies indicate durations longer (shorter) than the mean.



Fig. 6. Time series of the monthly mean ice area for the Scotian Shelf (top panel), the average ice area during the usual periods of advancement (January-March) and retreat (April-May) (middle panel) and the comparison of the monthly areas and volumes to the 1971-2000 means (bottom panel)



Fig. 7a. Scotian Shelf-Gulf of Maine areas used for extraction of sea-surface temperature (upper panel). The annual sea surface temperature anomalies derived from satellite imagery compared to their long-term means (lower panel). Pathfinder estimates were used for September 1985-December 2009. Estimates for 2010 were from the remote sensing laboratory, Ocean Research and Monitoring Section of the Ecosystem Research Division at BIO. These values were adjusted by the regression Pathfinder=0.976*ORMS+0.46 based on a comparison between overlapping Pathfinder-ORMS data.



Figure 7b. PCA 1 (71% of variance) and PCA2 (19%) loadings from a principal components analysis of the annual mean temperature anomalies (Fig. 7a, lower panel) for the seven Scotian Shelf and Gulf of Maine regions (Fig. 7a, upper panel).



Fig. 8. The annual surface temperature anomalies (dotted line with circles) and their 5-year running means (heavy black line) for (A) Halifax Harbour and (B) St. Andrews; annual depth-averaged (0-90 m) (C) temperature, (D) salinity and density anomalies for the Prince 5 monitoring station at the mouth of the Bay of Fundy.



Fig. 9. The annual mean temperature anomaly time series (dotted line with circles) and the 5 year running mean filtered anomalies (heavy solid line) on the Scotian Shelf and in the Gulf of Maine at (A) Cabot Strait 200-300 m; (B) Misaine Bank 100 m; (C) Emerald Basin 250 m; (D) Lurcher Shoals 50 m; and Georges Basin (200 m) (see Fig. 2).



Fig. 10. Time series of July bottom temperature anomalies (dashed lines with circles) and 5 year running mean filtered series (heavy line) for areas (A) 4Vn, (B) 4Vs, (C) 4W and (D) 4X. (E) Time series of the Cold Intermediate Layer (CIL, defined as waters with T<4°C) volume on the Scotian Shelf based on the July ecosystem survey. The dashed horizontal line is the long-term mean.</p>



Fig. 11. The mean annual anomaly (dashed line with circles) and 5-yr running mean (heavy solid line) of the stratification index (0-50 m density gradient) averaged over the Scotian Shelf (areas 4-23 inclusive, see Fig. 2). Standard error estimates for each annual anomaly value are also shown.



Fig. 12. (A) The time series of the monthly means (grey) and a 5 year running mean (black) of the relative sea level elevations at Halifax, along with the linear trend (1920-2010, red). (B) Residual relative sea level (monthly observed values – linear trend, averaged to annual estimates).



Fig. 13. Normalized annual anomalies of temperatures at the bottom and discrete depths for the Scotian Shelf-Gulf of Maine region. These normalized, annual anomalies are based on the 1981-2010 means, divided by the standard deviation. The scale represents the number of standard deviations an anomaly is from normal; blue indicates below normal, red above normal.



Fig. 14. The contributions of each of the normalized anomalies are shown as a bar chart and their summation as a time series (grey circles, black line).