

NOT TO BE CITED WITHOUT PRIOR  
REFERENCE TO THE AUTHOR(S)

Northwest Atlantic



Fisheries Organization

Serial No. N6168

NAFO SCR Doc. 13/017

## SCIENTIFIC COUNCIL MEETING – JUNE 2013

Results for the Atlantic cod, roughhead grenadier, redfish, thorny skate and black dogfish of the Spanish Survey in the NAFO Div. 3L for the period 2003-2012

by

Esther Román, Ángeles Armesto and Diana González-Troncoso

Instituto Español de Oceanografía  
P.O. Box 1552. Vigo, Spain  
e-mail: [esther.roman@vi.ieo.es](mailto:esther.roman@vi.ieo.es)

### Abstract

Since 2003, a stratified random spring bottom trawl survey was conducted by Spain in Division 3L of NAFO Regulatory Area (Flemish Pass). The surveys were carried out by the R/V "Vizconde de Eza" using bottom trawl net type *Campelen*. Entire series of mean catches, biomass and length distribution for Atlantic cod, roughhead grenadier, redfish, thorny skate and black dogfish are presented for the period 2003-2012.

KEYWORDS: Survey, Flemish Pass, Atlantic Cod, Roughhead grenadier, Redfish, Thorny skate, Black dogfish.

### Material and Methods

The Spanish surveys in Div. 3L of NAFO Regulatory Area (Flemish Pass) were initiated by Spain in 2003. The Research vessel "Vizconde de Eza" has carried out the entire surveys series following the same procedures and using the same bottom trawl gear *Campelen 1800*. In 2003, the survey was carried out in spring (June) and it did not cover all strata adequately (69% of the total area prospected in 2006-2012). In 2004, the survey was carried out in August, for a period of nine days, and it covered only the 96%. In 2005, it was not possible to perform the survey due to problems with the winch of the ship; and in 2006, for the first time, an adequate prospecting survey was conducted in Division 3L with over 100 valid hauls. Table 1 shows the number of valid tows, the depth and number of covered strata and the dates of the survey series. To know more details about the technical specifications of the surveys, see Román *et al.*, 2013.

The catch from each haul was sorted out and weighted by species and a randomly selected sample of each species was taken in order to measure it and obtain the length distribution. In 2003 and 2004 the Atlantic cod samples were not sorted out by sex. There are two species of redfish in Division 3L (*Sebastes mentella* and *S. fasciatus*); the external characteristics of both species are very similar, which makes it difficult to distinguish between them and, as a result, they are treated together.

For Atlantic cod, redfish, thorny skate and black dogfish each individual of the sample was measured to the total length to the nearest lower cm and data are given in 2 cm intervals. However, roughhead grenadier individuals were measured from tip of snout to base of first anal-fin ray to the lower ½ cm., in 0.5 cm intervals, as adopted by NAFO in June 1980 (Atkinson, 1991) as a standard measurement for roundnose and roughhead grenadiers; length is presented as pre-anal-fin length (AFL) and data are given in 1 cm intervals.

It is presented the mean catch per haul, the stratified mean catch per haul and the biomass with their variance per year in the period 2003-2012. Length distribution in number per haul stratified mean catches per length, sex and year for these species are presented too. The following formula was used to obtain the biomass from length distribution: Weight=a(Length+0.5)<sup>b</sup>.

## Results

### **Atlantic Cod (*Gadus morhua* Linnaeus, 1758)**

NAFO manages 3 cod stocks in 3L, 3M and 3NO. After a dramatic decline of cod during the eighties and nineties, fishing bans were imposed in the 1990s. In recent assessment all stocks remain at a very low level although spawning biomass has increased in recent years. In 2010, after a decade long moratorium, a cod fishery on the Flemish Cap (Div. 3M) was re-opened but the moratoria (no directed fishery) continues for Div. 3NO and Div. 3L. (NAFO, 2012).

#### Mean catches and biomass

Table 2 shows the swept area, the tow number, the mean catches and their variance per haul by stratum for Atlantic cod. Table 3 and Figure 1 present the stratified mean catches by stratum and year with their total variance. The entire time series (2003-2012) of biomass and their total variance for Atlantic cod are presented in Table 4 and Figure 2. Estimated parameters values of length-weight relationship are presented in Table 5.

Figure 3 shows a map with the distribution of Atlantic cod catches per haul in 2012 Spanish 3L survey.

Atlantic cod indices show a great variation, due to a few hauls in which the presence of cod was very high, however there is no clear trend along the whole period (2003-2012). Stratified mean catch and biomass decreased from 2003 to 2004; then, the values of these indices increased in 2006 and declined briefly again in 2007. A great increase is shown in 2008 but this was due to a single haul in which the presence of cod was very high (1298.5 kg). The great value of the variance in some years is produced by the tows with a large catch. In 2009 declined again and since then an increasing trend in the biomass can be seen. In 2011 the biomass reaches the highest value in the time series. The highest values in the estimated biomass have been observed in the shallow strata, in a range of depth from 93 to 274 meters. In 2012, the biomass decreases at the same level than in 2008.

#### Length distribution

Table 6 presents the length distribution of stratified mean catches per haul for this species, by sex and year, with the number of samples in which there were length measurements, the sampled catch, the total number of individuals measured in each sample and the range of lengths achieved, as well as the total catch of this species and the total hauls made in the survey. In Figures 4 and 5 the evolution throughout the period can be followed.

In this period, individuals between 12 and 25 cm can be seen although in 2004 there was no presence of individuals below 24 cm. In general all lengths presence is very low, even it is very difficult to follow the modal values. In 2008 we have a good presence of individuals between 26 and 33 cm, probably due to the haul with great catch of that year, 29 cm is the mode in the length distribution. In 2009 the dominant lengths were between 36 and 41 cm (mode = 37 cm). In 2010 the mode was 44 cm with the dominant length between 40 and 47 cm. In 2011 the mode observed was 51 cm and the dominant lengths were between 47 and 55 cm. and in 2012 the dominant lengths were between 34 and 56 cm (mode = 46 cm)

### **Roughhead grenadier (*Macrourus berglax* Lacépède, 1802)**

Roughhead grenadier is not a regulated species. There is no directed fishery for this species and most catches are taken as by-catch in Greenland halibut fishery in Subareas 2 and 3. Roughhead grenadier is taken mainly in Div. 3LMN of NAFO Regulatory Area. The highest level of observed catches was reached in 1998. The biomass of this species presents a decreasing trend in the last years (NAFO, 2012).

#### Mean catches and biomass

Roughhead grenadier haul mean catches by stratum are presented in Table 7; swept area, number of hauls and SD are also shown in this table. Stratified mean catches per tow by stratum and year and their variance are presented in Table 8. The entire time series (2003-2012) of biomass and their SD estimates of this species are shown in Table 9 and length-weight relationships are shown in Table 5.

The indices of roughhead grenadier show no clear trend throughout the whole period, with an increasing in 2004 compared to 2003 and then remains stable (2006-2007). In 2008 the biomass increased, reaching the highest value of the series, but declining afterwards in the period 2009-2012. Biomass reaches the lowest value in the time series in the 2012 (Fig. 6 and 7).

Figure 3 shows a map with the distribution of roughhead grenadier catches per haul in 2012 Spanish 3L survey.

#### Length distribution

Table 10 shows the stratified mean catches per haul length distribution, for roughhead grenadier, by sex and year, with the number of samples in which there was length measurements, the sampled catch, the total number of individuals measured in these samples and the range of lengths found. The total catch of this species and the total hauls made in the survey are shown too. In Figures 5 and 8 the evolution along the years can be followed. A slight recruitment can be seen in all period. Females attain larger lengths than males in all years.

#### **Redfish (*Sebastes spp.* Cuvier, 1829)**

There are two species of redfish that have been commercially fished in Div. 3LN, *Sebastes fasciatus* (Acadian redfish) and *S. mentella* (deepwater redfish). The external characteristics are very similar, making them difficult to distinguish, and as a consequence they are reported collectively as "redfish" in the commercial fishery. The redfish stocks in 3LN, 3M, 3O, as well as those in Subarea 2 and Div. 1F+3K are managed by NAFO. There was a moratorium on 3LN stocks (no directed fishery) from 1998-2009 and the fishery was reopened in 2010. The stock biomass, female spawning biomass and abundance is higher in 2006 than in the early 1990s (NAFO, 2012).

#### Mean catches and biomass

Table 11 shows the swept area, the tow number, the mean catches per haul and year and their variance for redfish. Table 12 and Figure 9 present the stratified mean catches per stratum with the total variance per year.

Figure 3 shows a map with the distribution of redfish catches per haul in 2012 Spanish 3L survey.

Table 13 and Figure 10 show the biomass estimate per swept area per stratum and their total variance by year and also the estimated abundance. Redfish shows a great annual variability probably due to its pelagic habitat. The redfish biomass indices decreased in 2004, 2007 and 2011 and increased in 2006, 2008 and 2009 with a sharp increase in 2010. In 2012, the redfish indices show the greater increasing reaching the highest value of the series (this was due to some hauls in which the presence of redfish was very high). The length-weight relationships are presented in Table 5.

#### Length distribution

Table 14 presents the length distribution of the stratified mean catches per haul for redfish, by sex and year, with the number of samples in which there was length measurements, the sampled catch, the total number of individuals measured in these samples and the range of lengths found. The total catch of this species and the total hauls made in the survey are also shown. In Figures 5 and 11 the evolution along the years can be followed. The highest proportions of small individuals in the catches (smaller than 20 cm) were found in 2007 and 2008.

#### **Thorny skate (*Amblyraja radiata* Donovan, 1808)**

Commercial catches of skates comprise a mix of skate species. However, thorny skate dominates, comprising about 95% of the skate species taken in the Canadian and EU-Spain catches. Thus, the skate fishery on the Grand Banks can be considered a fishery for thorny skate. In 2005, NAFO Fisheries Commission established a TAC of 13 500 t for thorny skate in Div. 3LN. In 2010 and 2011, the TAC for Div. 3LN has been reduced to 12 000 t.

Catches for NAFO Div. 3LN increased in the mid-1980s with the beginning of fishery targeting thorny skate. Catches in 2010 were the lowest recorded in Div. 3LNOPS since the starting of the fishery (NAFO, 2012).

#### Mean catches and biomass

Table 15 shows the swept area, the tow number, the mean catches per haul and year and their variance for thorny skate. Table 16 presents the length-weight relationships. Table 17 and Figure 12 present the stratified mean catches per stratum with the total variance per year. Table 18 and Figure 13 present the biomass per swept area by stratum and year, their total variance per year and the abundance index. The indices of the thorny skate decreased from 2003

to 2004, increased in 2006-2007 and decreased again in the period 2008-2011. In 2012 the indices of the thorny skate increased again. Figure 3 shows a map with the distribution of thorny skate catches per haul in 2012 Spanish 3L survey.

#### Length distribution

Table 19 presents the stratified mean catches per haul length distribution for this species, by sex and year, with the number of samples in which there was length measurements, the sampled catch, the total number of individuals measured in these samples and the range of lengths achieved, as well as the total catch of this species and the total hauls made in the survey. In Figures 14 and 15, the evolution along the years can be followed. The highest proportion of small thorny skate in the catches was in 2007.

#### **Black dogfish (*Centroscyllium fabricii* Reinhardt, 1825)**

Black dogfish is present in all Divisions, but is more abundant in Div. 3NO and in depths greater than 900 m. Black dogfish is not a regulated species and commercial catches of this species are mainly a by-catch of the Greenland halibut fishery in Div. 3LMNO (González-Costas *et al.*, 2006).

#### Mean catches and biomass

Black dogfish haul mean catches by stratum are presented in Table 20, including swept area, number of hauls and SD. Stratified mean catches per tow by stratum and year and their variance are presented in Table 21. The entire time series (2003-2012) of biomass and their SD estimates of black dogfish are shown in Table 22. Length-weight relationships are presented in Table 16.

The abundance and biomass present the same trend as mean catches. Biomass estimated from the 3L survey displays an increasing trend since 2004 until 2007 and decreased in 2008, 2009 and 2012, increasing again in 2010 and 2011 being the second and third values of the time series (Fig. 16 and 17). In 2003, the catches occurred only in two strata (745 and 749), in which the catches were much different, what explain why the variance in that year is so large.

Figure 3 shows a map with the distribution of black dogfish catches per haul in 2012 Spanish 3L survey.

#### Length distribution

Table 23 presents the length distribution of the stratified mean catches per haul for black dogfish, by sex and year, with the number of samples in which there was length measurements, the sampled catch, the total number of individuals measured in these samples and the range of lengths met. The total catch of this species and the total hauls made in the survey are shown too. In Figures 15 and 18 the evolution throughout the years can be followed.

There is no presence of small individual (smaller 38 cm). Size compositions are mainly between 50 and 80 cm of length. In 2012 the observed mode was 62 cm and the dominant lengths were between 56 and 67 cm.

#### **References**

- ATKINSON, D. B. 1991. Relationships Between Pre-anal Fin Length and Total Length of Roughhead Grenadier (*Macrourus berglax* Lacépède) in the Northwest Atlantic. *J. Northw. Atlan. Fish. Sci.*, **11**: 7-9
- GONZÁLEZ-COSTAS, F., D. GONZÁLEZ-TRONCOSO, M. CASAS, and G. RAMILLO. 2006. Spiny Dogfish (*Squalus acanthias*) and Black Dogfish (*Centroscyllium fabricii*) Spanish Data (Surveys and Fishery) in NAFO Divisions 3LMNO. *NAFO SCR Doc.*, No. 30, Serial No. N5250, 10 p.
- NAFO, 2012. Report of Scientific Council Meeting, 1-14 June 2012.
- ROMÁN, E., C. GONZÁLEZ-IGLESIAS and D. GONZÁLEZ-TRONCOSO. 2013. Results for the Spanish Survey in the NAFO Regulatory Area of Division 3L for the period 2003-2012. *NAFO SCR Doc.*, No. 16, Serial No N6167.

**TABLE 1.-** Spanish bottom trawl surveys in NAFO Division 3L for the period 2003-2012.

Year	Vessel	Valid tows	Depth strata covered (m)	Surveyed strata (no.)	Dates
2003	R/V " <i>Vizconde de Eza</i> "	39	118-1100	17	June 2 - June 6, June 29
2004	R/V " <i>Vizconde de Eza</i> "	50	141-1452	23	August 7 - August 15
2005	-	-	-	-	-
2006	R/V " <i>Vizconde de Eza</i> "	100	116-1449	24	July 31 - August 18
2007	R/V " <i>Vizconde de Eza</i> "	94	119-1449	24	July 23 - August 11
2008	R/V " <i>Vizconde de Eza</i> "	100	105-1455	24	July 24 - August 11
2009	R/V " <i>Vizconde de Eza</i> "	98	111-1458	24	July 25 - August 12
2010	R/V " <i>Vizconde de Eza</i> "	97	119-1462	24	July 25 - August 14
2011	R/V " <i>Vizconde de Eza</i> "	89	115-1419	24	August 10 - August 24
2012	R/V " <i>Vizconde de Eza</i> "	98	112-1478	24	July 30 - August 18

**TABLE 2.-** Swept area, number of hauls and **Atlantic cod** mean catch (Kg) and SD (\*\*) by stratum. Spanish Survey on NAFO Div. 3L in the period 2003-2012, on board R/V "Vizconde de Eza". (\*) In 2003, the data correspond to 69% of the total area prospected in 2006-2012.

Stratum	2003 (*)				2004				2006				2007				2008			
	Swept area	Tow No.	Mean catch	SD	Swept area	Tow No.	Mean catch	SD	Swept area	Tow No.	Mean catch	SD	Swept area	Tow No.	Mean catch	SD	Swept area	Tow No.	Mean catch	SD
385	0.0225	2	0.062	0.040	0.0229	2	0.450	0.636	0.0229	2	1.783	2.521	0.0225	2	0.835	1.181	0.0229	2	6.051	6.537
387	0.0229	2	4.390	1.004	0.0214	2	1.885	1.888	0.0225	2	0.395	0.559	0.0225	2	1.992	1.105	0.0435	4	5.386	5.633
388	0.0334	3	7.870	6.987	0.0105	1	1.313	-	0.0566	5	7.028	5.142	0.0563	5	7.434	7.400	0.0559	5	18.665	19.454
389	0.0454	4	0.844	1.573	0.0225	2	0.510	0.721	0.0795	7	10.582	14.986	0.0900	8	4.162	4.621	0.0780	7	30.523	18.566
390	0.0563	5	0.000	0.000	0.0345	3	0.000	0.000	0.1249	11	0.081	0.249	0.1350	12	1.369	1.251	0.1395	12	8.682	15.848
391	0.0338	3	0.167	0.289	0.0218	2	0.000	0.000	0.0450	4	14.338	13.278	0.0450	4	11.183	15.378	0.0454	4	342.268	637.574
392	0.0116	1	0.400	-	0.0214	2	13.219	17.991	0.0229	2	2.045	1.506	0.0225	2	13.985	7.779	0.0221	2	0.000	0.000
729	0.0210	2	1.260	1.782	0.0221	2	0.000	0.000	0.0338	3	0.000	0.000	0.0338	3	0.000	0.000	0.0338	3	0.000	0.000
730	0.0221	2	0.000	0.000	0.0221	2	0.000	0.000	0.0326	3	0.000	0.000	0.0225	2	0.000	0.000	0.0323	3	0.000	0.000
731	0.0229	2	22.405	13.329	0.0233	2	0.496	0.530	0.0341	3	0.000	0.000	0.0338	3	0.510	0.883	0.0330	3	0.130	0.225
732	0.0113	1	0.000	-	0.0210	2	0.000	0.000	0.0334	3	0.000	0.000	0.0338	3	0.000	0.000	0.0446	4	0.000	0.000
733	n.s. n.s.	n.s.	n.s.		0.0330	3	0.000	0.000	0.0454	4	0.000	0.000	0.0338	3	0.427	0.739	0.0431	4	0.000	0.000
734	n.s. n.s.	n.s.	n.s.		0.0304	3	0.000	0.000	0.0225	2	0.000	0.000	0.0225	2	0.000	0.000	0.0221	2	0.000	0.000
741	0.0113	1	0.000	-	0.0323	3	0.000	0.000	0.0218	2	0.000	0.000	0.0225	2	0.000	0.000	0.0210	2	0.000	0.000
742	0.0116	1	0.000	-	0.0120	1	0.000	-	0.0229	2	0.000	0.000	0.0225	2	0.000	0.000	0.0210	2	0.000	0.000
743	n.s. n.s.	n.s.	n.s.		0.0188	2	0.000	0.000	0.0225	2	0.000	0.000	0.0225	2	0.000	0.000	0.0203	2	0.000	0.000
744	n.s. n.s.	n.s.	n.s.		0.0101	1	0.000	-	0.0229	2	0.000	0.000	0.0218	2	0.000	0.000	0.0221	2	0.000	0.000
745	0.0341	3	0.000	0.000	0.0319	3	0.000	0.000	0.0686	6	0.000	0.000	0.0675	6	0.000	0.000	0.0555	5	0.000	0.000
746	0.0446	4	0.000	0.000	0.0338	3	0.000	0.000	0.0675	6	0.000	0.000	0.0664	6	0.000	0.000	0.0638	6	0.000	0.000
747	n.s. n.s.	n.s.	n.s.		0.0308	3	0.000	0.000	0.1230	11	0.000	0.000	0.1238	11	0.000	0.000	0.1069	10	0.000	0.000
748	0.0109	1	0.000	-	0.0199	2	0.000	0.000	0.0326	3	0.000	0.000	0.0338	3	0.000	0.000	0.0218	2	0.000	0.000
749	0.0221	2	0.000	0.000	0.0221	2	0.000	0.000	0.0229	2	0.000	0.000	0.0113	1	0.000	-	0.0214	2	0.000	0.000
750	n.s. n.s.	n.s.	n.s.		0.0180	2	0.000	0.000	0.1005	9	0.000	0.000	0.0679	6	0.000	0.000	0.0844	8	0.000	0.000
751	n.s. n.s.	n.s.	n.s.		n.s. n.s.	n.s.	n.s.	n.s.	0.0454	4	0.000	0.000	0.0225	2	0.000	0.000	0.0413	4	0.000	0.000

$$(**) SD = \frac{\sum (x_i - \bar{x})}{n-1}$$

**TABLE 2 (cont).-** Swept area, number of hauls and **Atlantic cod** mean catch (Kg) and SD (\*\*) by stratum. Spanish Survey on NAFO Div. 3L in the period 2003-2012, on board R/V "Vizconde de Eza". (\*) In 2003, the data correspond to 69% of the total area prospected in 2006-2012.

Stratum	2009				2010				2011				2012			
	Swept area	Tow number	Mean catch	SD	Swept area	Tow number	Mean catch	SD	Swept area	Tow number	Mean catch	SD	Swept area	Tow number	Mean catch	SD
385	0.0225	2	5.285	3.514	0.0225	2	0.775	1.096	0.0229	2	93.750	118.723	0.0225	2	4.820	2.871
387	0.0439	4	23.204	40.440	0.0458	4	3.433	2.594	0.0450	4	36.505	32.228	0.0450	4	6.760	4.899
388	0.0555	5	7.413	3.853	0.0570	5	61.988	121.458	0.0563	5	15.241	14.829	0.0570	5	162.020	264.788
389	0.0803	7	40.874	54.955	0.0795	7	150.908	266.990	0.0675	6	26.796	42.096	0.0799	7	34.169	26.422
390	0.1373	12	22.441	43.094	0.1249	11	37.143	51.671	0.1009	9	217.889	231.959	0.1354	12	43.245	27.872
391	0.0458	4	65.264	62.051	0.0454	4	144.075	119.143	0.0458	4	150.275	91.993	0.0458	4	44.280	47.163
392	0.0229	2	0.063	0.089	0.0225	2	70.680	89.265	0.0229	2	3.268	3.129	0.0225	2	13.470	4.992
729	0.0341	3	0.000	0.000	0.0338	3	0.000	0.000	0.0338	3	0.000	0.000	0.0338	3	0.000	0.000
730	0.0338	3	0.000	0.000	0.0334	3	0.000	0.000	0.0334	3	0.000	0.000	0.0338	3	0.000	0.000
731	0.0341	3	0.000	0.000	0.0338	3	0.247	0.225	0.0334	3	0.000	0.000	0.0341	3	0.000	0.000
732	0.0450	4	0.000	0.000	0.0450	4	0.000	0.000	0.0454	4	0.000	0.000	0.0454	4	0.000	0.000
733	0.0450	4	0.000	0.000	0.0450	4	0.000	0.000	0.0454	4	0.545	0.642	0.0454	4	0.000	0.000
734	0.0218	2	0.000	0.000	0.0225	2	0.000	0.000	0.0225	2	0.000	0.000	0.0233	2	0.000	0.000
741	0.0221	2	0.000	0.000	0.0225	2	0.000	0.000	0.0218	2	0.000	0.000	0.0218	2	0.000	0.000
742	0.0214	2	0.000	0.000	0.0225	2	0.000	0.000	0.0225	2	0.000	0.000	0.0206	2	0.000	0.000
743	0.0203	2	0.000	0.000	0.0225	2	0.000	0.000	0.0221	2	0.000	0.000	0.0206	2	0.000	0.000
744	0.0210	2	0.000	0.000	0.0229	2	0.000	0.000	0.0221	2	0.000	0.000	0.0221	2	0.000	0.000
745	0.0559	5	0.000	0.000	0.0563	5	0.000	0.000	0.0446	4	0.000	0.000	0.0570	5	0.000	0.000
746	0.0668	6	0.000	0.000	0.0679	6	0.000	0.000	0.0566	5	0.000	0.000	0.0675	6	0.000	0.000
747	0.1118	10	0.000	0.000	0.1125	10	0.000	0.000	0.0893	8	0.000	0.000	0.1121	10	0.000	0.000
748	0.0229	2	0.000	0.000	0.0225	2	0.000	0.000	0.0221	2	0.000	0.000	0.0225	2	0.000	0.000
749	0.0225	2	0.000	0.000	0.0229	2	0.000	0.000	0.0221	2	0.000	0.000	0.0221	2	0.000	0.000
750	0.0791	7	0.000	0.000	0.0900	8	0.000	0.000	0.0668	6	0.000	0.000	0.0885	8	0.000	0.000
751	0.0338	3	0.000	0.000	0.0225	2	0.000	0.000	0.0334	3	0.000	0.000	0.0218	2	0.000	0.000

$$(**) SD = \frac{\sum (x_i - \bar{x})}{n-1}$$

**TABLE 3.-** Stratified mean catches (Kg) of **Atlantic cod** by stratum and year (2003-2012) and SD. Research Vessel *Vizconde de Eza*. n.s. means stratum not surveyed. In 2003: the data correspond to 69% of the total area prospected in 2006-2012.

Stratum	Survey									
	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
385	7.26	53.10	-	210.34	98.53	713.96	623.63	91.45	11062.50	568.76
387	1123.84	482.56	-	101.12	509.82	1378.75	5940.16	878.72	9345.28	1730.43
388	2809.59	468.74	-	2509.00	2653.87	6663.55	2646.51	22129.72	5441.04	57841.14
389	429.34	259.59	-	5386.31	2118.59	15536.35	20804.94	76812.24	13639.08	17391.88
390	0.00	0.00	-	65.94	1115.80	7076.10	18289.28	30271.32	177579.44	35245.01
391	47.00	0.00	-	4043.18	3153.47	96519.44	18404.45	40629.15	42377.55	12486.96
392	58.00	1916.68	-	296.53	2027.75	0.00	9.14	10248.60	473.79	1953.15
729	234.36	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00
730	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00
731	4839.48	107.03	-	0.00	110.16	28.08	0.00	53.28	0.00	0.00
732	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00
733	n.s	0.00	-	0.00	99.84	0.00	0.00	0.00	127.59	0.00
734	n.s	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00
741	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00
742	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00
743	n.s	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00
744	n.s	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00
745	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00
746	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00
747	n.s	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00
748	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00
749	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00
750	n.s	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00
751	n.s	n.s	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00
TOTAL	9548.87	3287.70	-	12612.40	11887.83	127916.23	66718.10	181114.48	260046.27	127217.33
SD	2.13	0.53	-	1.94	1.83	19.72	10.28	27.92	40.09	19.61
	0.57	0.30	-	0.55	0.42	13.89	2.75	9.17	10.15	6.72

**TABLE 4.**- Survey estimates (by the swept area method) of **Atlantic cod** biomass (t.) by stratum and year and their SD on NAFO Div. 3L (R/V *Vizconde de Eza*). n.s. means stratum not surveyed. In 2003, the data correspond to 69% of the total area prospected in 2006-2012.

Stratum	Survey									
	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
385	1	5	-	18	9	62	55	8	967	51
387	98	45	-	9	45	127	542	77	831	154
388	253	45	-	222	236	596	238	1941	484	5074
389	38	23	-	474	188	1394	1815	6763	1212	1524
390	0	0	-	6	99	609	1599	2667	15844	3124
391	4	0	-	359	280	8509	1609	3582	3705	1092
392	5	179	-	26	180	0	1	911	41	174
729	22	0	-	0	0	0	0	0	0	0
730	0	0	-	0	0	0	0	0	0	0
731	423	9	-	0	10	3	0	5	0	0
732	0	0	-	0	0	0	0	0	0	0
733	n.s.	0	-	0	9	0	0	0	11	0
734	n.s.	0	-	0	0	0	0	0	0	0
741	0	0	-	0	0	0	0	0	0	0
742	0	0	-	0	0	0	0	0	0	0
743	n.s.	0	-	0	0	0	0	0	0	0
744	n.s.	0	-	0	0	0	0	0	0	0
745	0	0	-	0	0	0	0	0	0	0
746	0	0	-	0	0	0	0	0	0	0
747	n.s.	0	-	0	0	0	0	0	0	0
748	0	0	-	0	0	0	0	0	0	0
749	0	0	-	0	0	0	0	0	0	0
750	n.s.	0	-	0	0	0	0	0	0	0
751	n.s.	n.s.	-	0	0	0	0	0	0	0
TOTAL	844	306	-	1114	1057	11300	5859	15953	23095	11192
SD	222	180	-	315	245	7745	1556	5265	5833	3877

**Table 5.**- Length-weight relationships in the calculation of biomass, for Division 3L (out ZEE Canada), 2003-2012 for **Atlantic cod**, **roughhead grenadier** and **redfish**. The equation is Weight=a(Length+0.5)<sup>b</sup>. To calculate the parameters for the indeterminate individuals, we used the total data (males+females+indeterminate individuals).

Atlantic cod							Roughhead grenadier							Redfish						
Year	Sex	L-W Equations		N	r <sup>2</sup>	Sex	L-W Equations		N	r <sup>2</sup>	Sex	L-W Equations		N	r <sup>2</sup>					
2003	All	$W = 0.0059 L^{3.0965}$	161	0.9875		All	$W = 0.0766 L^{3.0029}$	478	0.9872		All	$W = 0.0037 L^{3.3842}$	238	0.9902						
	Males	- - -				Males	$W = 0.0482 L^{3.1908}$	172	0.9772		Males	$W = 0.0103 L^{3.0686}$	95	0.9787						
	Females	- - -				Females	$W = 0.0824 L^{2.9761}$	290	0.9913		Females	$W = 0.0060 L^{3.2380}$	90	0.993						
2004	All	$W = 0.0045 L^{3.2037}$	58	0.9805		All	$W = 0.0791 L^{3.0113}$	1066	0.9896		All	$W = 0.0083 L^{3.1377}$	237	0.9808						
	Males	- - -				Males	$W = 0.0085 L^{2.9868}$	458	0.9866		Males	$W = 0.0161 L^{2.9333}$	97	0.9877						
	Females	- - -				Females	$W = 0.0788 L^{3.0119}$	597	0.9906		Females	$W = 0.0190 L^{2.8927}$	117	0.9881						
2006	All	$W = 0.0057 L^{3.3142}$	308	0.9854		All	$W = 0.0773 L^{3.0264}$	1645	0.9817		All	$W = 0.0096 L^{3.1034}$	920	0.9835						
	Males	$W = 0.0043 L^{3.2188}$	142	0.9808		Males	$W = 0.0664 L^{3.0810}$	655	0.9748		Males	$W = 0.0100 L^{3.0871}$	444	0.9843						
	Females	$W = 0.0069 L^{3.0874}$	166	0.9896		Females	$W = 0.0893 L^{2.9794}$	975	0.986		Females	$W = 0.0091 L^{3.1221}$	471	0.9811						
2007	All	$W = 0.0055 L^{3.1370}$	225	0.983		All	$W = 0.0885 L^{2.9691}$	1950	0.9895		All	$W = 0.0080 L^{3.1588}$	881	0.9842						
	Males	$W = 0.0061 L^{3.1114}$	107	0.991		Males	$W = 0.0946 L^{2.9435}$	754	0.9859		Males	$W = 0.0140 L^{2.9836}$	432	0.9858						
	Females	$W = 0.0047 L^{3.1750}$	118	0.9735		Females	$W = 0.0877 L^{2.9727}$	1165	0.9897		Females	$W = 0.0133 L^{3.0115}$	392	0.9868						
2008	All	$W = 0.0083 L^{3.0479}$	819	0.9856		All	$W = 0.1237 L^{2.8681}$	1773	0.9871		All	$W = 0.0142 L^{2.9849}$	699	0.9701						
	Males	$W = 0.0083 L^{3.0493}$	403	0.9855		Males	$W = 0.1174 L^{2.8868}$	754	0.9832		Males	$W = 0.0337 L^{2.7219}$	338	0.9343						
	Females	$W = 0.0084 L^{3.0467}$	416	0.9856		Females	$W = 0.1144 L^{2.8938}$	1024	0.988		Females	$W = 0.0314 L^{2.7511}$	340	0.9412						
2009	All	$W = 0.0084 L^{3.0256}$	684	0.9824		All	$W = 0.0903 L^{2.9583}$	1457	0.9911		All	$W = 0.0083 L^{3.1392}$	818	0.9854						
	Males	$W = 0.0089 L^{3.0085}$	296	0.9824		Males	$W = 0.0847 L^{2.9803}$	540	0.9871		Males	$W = 0.0135 L^{2.9882}$	354	0.9738						
	Females	$W = 0.0083 L^{3.0299}$	388	0.9821		Females	$W = 0.0927 L^{2.9505}$	899	0.9904		Females	$W = 0.0174 L^{2.9204}$	389	0.9763						
2010	All	$W = 0.0086 L^{3.0302}$	756	0.980		All	$W = 0.1006 L^{2.9369}$	1539	0.991		All	$W = 0.0110 L^{3.0593}$	808	0.9859						
	Males	$W = 0.0076 L^{3.0636}$	364	0.980		Males	$W = 0.0909 L^{2.9770}$	547	0.984		Males	$W = 0.0153 L^{2.9565}$	372	0.9754						
	Females	$W = 0.0095 L^{3.0027}$	392	0.979		Females	$W = 0.1071 L^{2.9152}$	947	0.990		Females	$W = 0.0161 L^{2.9484}$	397	0.9706						
2011	All	$W = 0.0090 L^{3.0101}$	1421	0.9874		All	$W = 0.0962 L^{2.9550}$	1545	0.9899		All	$W = 0.0105 L^{3.0803}$	1218	0.9882						
	Males	$W = 0.0102 L^{2.9790}$	682	0.9852		Males	$W = 0.1018 L^{2.9403}$	543	0.9796		Males	$W = 0.0129 L^{3.0158}$	529	0.9836						
	Females	$W = 0.0082 L^{3.0334}$	739	0.9892		Females	$W = 0.1169 L^{2.8873}$	913	0.9884		Females	$W = 0.0109 L^{3.0768}$	559	0.9855						
2012	All	$W = 0.0106 L^{2.9627}$	878	0.982		All	$W = 0.1070 L^{2.9148}$	1607	0.988		All	$W = 0.0126 L^{3.0228}$	978	0.9847						
	Males	$W = 0.0109 L^{2.9573}$	403	0.982		Males	$W = 0.1008 L^{2.9374}$	609	0.980		Males	$W = 0.0135 L^{2.9979}$	476	0.9856						
	Females	$W = 0.0123 L^{2.9243}$	474	0.980		Females	$W = 0.1081 L^{2.9117}$	934	0.988		Females	$W = 0.0157 L^{2.9616}$	491	0.9806						

**TABLE 6.- Atlantic cod** length distribution per haul mean catches by sex and year. Number per stratified mean catches. Spanish Summer Survey on NAFO 3L: 2004-2012 (R/V *Vizconde de Eza*). Indet. means indeterminate..

Length (cm.)	2004				2006				2007				2008			
	M	F	I	T	M	F	I	T	M	F	I	T	M	F	I	T
12	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.01	0.00	0.01	0.00	0.01	0.01	0.02	0.00	0.04
14	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.01	0.01	0.00	0.00	0.01
16	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.02	0.00	0.00	0.02	0.08	0.03	0.00	0.11
18	0.00	0.00	0.00	0.00	0.00	0.03	0.00	0.03	0.04	0.03	0.00	0.07	0.19	0.15	0.00	0.34
20	0.00	0.00	0.00	0.00	0.03	0.00	0.00	0.03	0.04	0.00	0.00	0.04	0.14	0.13	0.00	0.27
22	0.00	0.00	0.00	0.00	0.05	0.02	0.00	0.08	0.02	0.01	0.00	0.03	0.12	0.19	0.00	0.31
24	0.00	0.00	0.02	0.02	0.08	0.10	0.00	0.18	0.01	0.01	0.00	0.02	1.21	1.36	0.00	2.56
26	0.00	0.00	0.03	0.03	0.09	0.16	0.00	0.25	0.01	0.00	0.00	0.01	5.14	6.23	0.00	11.37
28	0.00	0.00	0.06	0.06	0.09	0.19	0.00	0.27	0.02	0.04	0.00	0.06	8.51	10.05	0.00	18.56
30	0.00	0.00	0.15	0.15	0.13	0.19	0.00	0.32	0.05	0.02	0.00	0.07	6.60	7.42	0.00	14.02
32	0.00	0.00	0.03	0.03	0.20	0.11	0.00	0.30	0.05	0.06	0.00	0.12	2.99	3.61	0.00	6.60
34	0.00	0.00	0.05	0.05	0.15	0.10	0.00	0.25	0.07	0.06	0.00	0.14	1.94	0.81	0.00	2.74
36	0.00	0.00	0.09	0.09	0.12	0.11	0.00	0.23	0.07	0.13	0.00	0.21	0.83	0.78	0.00	1.61
38	0.00	0.00	0.15	0.15	0.11	0.12	0.00	0.23	0.14	0.17	0.00	0.31	0.32	0.35	0.00	0.67
40	0.00	0.00	0.10	0.10	0.05	0.12	0.00	0.17	0.11	0.14	0.00	0.25	0.14	0.29	0.00	0.43
42	0.00	0.00	0.10	0.10	0.12	0.07	0.00	0.18	0.10	0.14	0.00	0.24	0.06	0.37	0.00	0.43
44	0.00	0.00	0.06	0.06	0.13	0.10	0.00	0.23	0.11	0.07	0.00	0.18	0.13	0.05	0.00	0.19
46	0.00	0.00	0.03	0.03	0.11	0.13	0.00	0.24	0.02	0.13	0.00	0.15	0.09	0.29	0.00	0.37
48	0.00	0.00	0.01	0.01	0.03	0.09	0.00	0.12	0.07	0.04	0.00	0.12	0.07	0.24	0.00	0.31
50	0.00	0.00	0.03	0.03	0.03	0.05	0.00	0.08	0.02	0.03	0.00	0.05	0.06	0.09	0.00	0.16
52	0.00	0.00	0.00	0.00	0.02	0.05	0.00	0.08	0.02	0.05	0.00	0.07	0.22	0.07	0.00	0.29
54	0.00	0.00	0.01	0.01	0.00	0.04	0.00	0.04	0.05	0.02	0.00	0.07	0.04	0.06	0.00	0.10
56	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.02	0.04	0.00	0.06	0.04	0.02	0.00	0.06
58	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.01	0.03	0.03	0.00	0.06	0.19	0.03	0.00	0.22
60	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.01	0.00	0.02	0.02	0.02	0.00	0.04
62	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.01	0.00	0.02	0.05	0.03	0.00	0.09
64	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	0.01	0.05	0.00	0.06
66	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.04	0.00	0.00	0.04
68	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.01
70	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.01
72	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
74	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01
76	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00
78	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
80	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
82	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
84	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
86	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
88	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
90	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
92	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Total	0.00	0.00	0.92	0.92	1.57	1.82	0.00	3.38	1.13	1.30	0.00	2.43	29.27	32.78	0.00	62.05
Nº samples:					9			22				32				34
Nº Ind.:	-	-	55	55	143	167	0	310	107	119	0	226	739	827	0	1566
Sampled catch:					34			176				168				1814
Range:					24-55			13-79				12-76				12-74
Total catch:					34			176				168				1814
Total valid hauls:					50			100				94				100

**TABLE 6 (cont.).- Atlantic cod** length distribution per haul mean catches by sex and year. Number per stratified mean catches. Spanish Summer Survey on NAFO 3L: 2004-2012 (R/V *Vizconde de Eza*). Indet. means indeterminate.

Length (cm.)	2009				2010				2011				2012			
	M	F	I	T	M	F	I	T	M	F	I	T	M	F	I	T
12	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.01
14	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
16	0.01	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
18	0.00	0.00	0.00	0.00	0.03	0.00	0.00	0.03	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00
20	0.02	0.01	0.00	0.03	0.00	0.00	0.00	0.00	0.01	0.01	0.00	0.03	0.00	0.00	0.00	0.00
22	0.06	0.06	0.00	0.12	0.01	0.00	0.00	0.01	0.04	0.05	0.00	0.08	0.01	0.00	0.00	0.01
24	0.08	0.05	0.00	0.13	0.07	0.07	0.00	0.13	0.07	0.16	0.00	0.23	0.02	0.02	0.00	0.04
26	0.12	0.12	0.00	0.24	0.21	0.24	0.00	0.45	0.37	0.31	0.00	0.68	0.07	0.04	0.00	0.11
28	0.14	0.18	0.00	0.32	0.49	0.88	0.00	1.37	0.46	0.66	0.00	1.12	0.01	0.00	0.00	0.01
30	0.20	0.15	0.00	0.36	0.99	1.06	0.00	2.05	0.58	0.71	0.00	1.29	0.02	0.01	0.00	0.03
32	0.39	0.37	0.00	0.77	1.34	1.23	0.00	2.57	0.67	0.78	0.00	1.45	0.05	0.03	0.00	0.08
34	0.66	1.04	0.00	1.70	0.87	1.07	0.00	1.95	0.81	0.72	0.00	1.53	0.06	0.04	0.00	0.10
36	1.11	1.16	0.00	2.26	1.27	1.35	0.00	2.62	0.68	0.75	0.00	1.43	0.15	0.03	0.00	0.18
38	1.09	1.42	0.00	2.51	1.31	1.44	0.00	2.75	0.71	0.75	0.00	1.46	0.14	0.06	0.00	0.20
40	0.92	1.07	0.00	1.99	1.65	2.08	0.00	3.72	0.76	1.09	0.00	1.85	0.15	0.07	0.00	0.22
42	0.49	0.76	0.00	1.25	1.91	2.12	0.00	4.02	0.95	0.86	0.00	1.82	0.39	0.33	0.00	0.72
44	0.28	0.47	0.00	0.75	1.79	2.52	0.00	4.31	0.99	1.29	0.00	2.28	0.44	0.58	0.00	1.03
46	0.15	0.37	0.00	0.52	1.60	2.24	0.00	3.85	1.18	1.61	0.00	2.79	0.68	0.98	0.00	1.66
48	0.04	0.15	0.00	0.18	1.17	1.48	0.00	2.65	1.41	2.14	0.00	3.55	0.73	0.82	0.00	1.55
50	0.08	0.14	0.00	0.22	0.51	0.95	0.00	1.46	2.26	2.42	0.00	4.68	0.71	1.08	0.00	1.79
52	0.07	0.13	0.00	0.20	0.28	0.43	0.00	0.71	1.86	2.21	0.00	4.07	0.75	0.85	0.00	1.60
54	0.07	0.08	0.00	0.15	0.18	0.31	0.00	0.49	1.34	2.00	0.00	3.34	0.91	0.97	0.00	1.88
56	0.09	0.11	0.00	0.20	0.05	0.21	0.00	0.25	0.71	1.05	0.00	1.75	0.64	0.97	0.00	1.61
58	0.01	0.13	0.00	0.14	0.12	0.13	0.00	0.26	0.49	0.62	0.00	1.11	0.63	0.79	0.00	1.42
60	0.02	0.07	0.00	0.09	0.16	0.06	0.00	0.22	0.36	0.32	0.00	0.68	0.48	0.62	0.00	1.10
62	0.03	0.04	0.00	0.07	0.05	0.07	0.00	0.12	0.08	0.22	0.00	0.30	0.45	0.54	0.00	0.99
64	0.01	0.06	0.00	0.07	0.05	0.01	0.00	0.06	0.09	0.06	0.00	0.15	0.55	0.48	0.00	1.03
66	0.01	0.03	0.00	0.04	0.02	0.05	0.00	0.07	0.07	0.05	0.00	0.12	0.22	0.22	0.00	0.45
68	0.01	0.02	0.00	0.03	0.04	0.01	0.00	0.05	0.02	0.09	0.00	0.11	0.16	0.33	0.00	0.48
70	0.01	0.03	0.00	0.04	0.01	0.00	0.00	0.01	0.00	0.05	0.00	0.05	0.10	0.19	0.00	0.29
72	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	0.05	0.17	0.00	0.22
74	0.01	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.01	0.01	0.00	0.02	0.02	0.12	0.00	0.14
76	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.04	0.04	0.00	0.08
78	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.06	0.00	0.07
80	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
82	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
84	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00
86	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01
88	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
90	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.03	0.00	0.03
92	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Total	6.18	8.25	0.00	14.44	16.19	20.07	0.00	36.25	17.01	21.02	0.00	38.03	8.65	10.47	0.01	19.14
Nº samples:				32				36				34				35
Nº Ind.:	580	781	0	1361	1014	1265	0	2279	1147	1440	0	2587	603	693	1	1297
Sampled catch:				957				2509				3141				1809
Range:				13-77				12-93				19-85				5-82
Total catch:				957				2509				3141				1809
Total valid hauls:				98				97				89				98

**TABLE 7.-** Swept area, number of hauls and **roughhead grenadier** mean catch (Kg) and SD (\*\*) by stratum. Spanish Survey on NAFO Div. 3L in the period 2003-2012, on board R/V "Vizconde de Eza". (\*) In 2003, the data correspond to 69% of the total area prospected in 2006-2012.

Stratum	2003 (*)				2004				2006				2007				2008			
	Swept area	Tow No.	Mean catch	SD	Swept area	Tow No.	Mean catch	SD	Swept area	Tow No.	Mean catch	SD	Swept area	Tow No.	Mean catch	SD	Swept area	Tow No.	Mean catch	SD
385	0.0225	2	0.000	0.000	0.0229	2	0.000	0.000	0.0229	2	0.000	0.000	0.0225	2	0.000	0.000	0.0229	2	0.000	0.000
387	0.0229	2	0.000	0.000	0.0214	2	59.987	23.598	0.0225	2	34.790	20.520	0.0225	2	45.990	51.746	0.0435	4	20.320	11.817
388	0.0334	3	0.000	0.000	0.0105	1	43.300		0.0566	5	26.406	7.803	0.0563	5	37.663	22.136	0.0559	5	15.056	11.298
389	0.0454	4	0.000	0.000	0.0225	2	1.875	2.652	0.0795	7	1.426	2.642	0.0900	8	3.075	8.697	0.0780	7	19.007	23.458
390	0.0563	5	0.560	1.252	0.0345	3	0.007	0.012	0.1249	11	0.000	0.000	0.1350	12	0.000	0.000	0.1395	12	0.580	1.338
391	0.0338	3	0.017	0.029	0.0218	2	0.018	0.025	0.0450	4	178.123	304.579	0.0450	4	86.525	171.255	0.0454	4	248.947	142.328
392	0.0116	1	3.900	-	0.0214	2	200.650	255.195	0.0229	2	118.025	159.347	0.0225	2	129.950	138.805	0.0221	2	58.175	54.836
729	0.0210	2	37.750	4.596	0.0221	2	29.475	17.501	0.0338	3	25.164	23.944	0.0338	3	26.490	13.222	0.0338	3	19.943	6.923
730	0.0221	2	101.050	37.972	0.0221	2	33.715	0.544	0.0326	3	53.270	7.021	0.0225	2	81.378	33.061	0.0323	3	35.119	29.483
731	0.0229	2	3.510	1.824	0.0233	2	10.450	5.162	0.0341	3	10.512	3.252	0.0338	3	14.333	7.365	0.0330	3	14.333	10.000
732	0.0113	1	34.400	-	0.0210	2	39.490	7.594	0.0334	3	22.164	9.200	0.0338	3	11.151	3.253	0.0446	4	21.545	3.045
733	n.s. n.s.	n.s.	n.s.		0.0330	3	15.553	5.137	0.0454	4	23.450	16.806	0.0338	3	19.104	14.162	0.0431	4	23.939	36.979
734	n.s. n.s.	n.s.	n.s.		0.0304	3	65.850	106.305	0.0225	2	39.315	9.638	0.0225	2	23.400	8.202	0.0221	2	30.580	20.182
741	0.0113	1	8.700	-	0.0323	3	1.055	1.342	0.0218	2	17.557	23.112	0.0225	2	4.650	6.166	0.0210	2	10.359	10.390
742	0.0116	1	24.400	-	0.0120	1	4.700	-	0.0229	2	20.933	7.015	0.0225	2	14.493	2.011	0.0210	2	16.861	11.943
743	n.s. n.s.	n.s.	n.s.		0.0188	2	26.245	6.017	0.0225	2	10.574	6.353	0.0225	2	29.666	25.928	0.0203	2	25.509	13.847
744	n.s. n.s.	n.s.	n.s.		0.0101	1	2.550	-	0.0229	2	15.365	15.111	0.0218	2	33.965	0.375	0.0221	2	58.670	15.570
745	0.0341	3	17.547	10.764	0.0319	3	5.800	2.722	0.0686	6	8.238	5.438	0.0675	6	3.624	1.509	0.0555	5	14.284	7.402
746	0.0446	4	63.800	71.784	0.0338	3	26.205	21.151	0.0675	6	41.767	29.972	0.0664	6	34.607	22.333	0.0638	6	30.720	16.486
747	n.s. n.s.	n.s.	n.s.		0.0308	3	43.627	13.999	0.1230	11	42.307	40.112	0.1238	11	62.510	26.732	0.1069	10	28.717	25.198
748	0.0109	1	55.980	-	0.0199	2	22.515	18.547	0.0326	3	67.920	73.796	0.0338	3	33.533	16.455	0.0218	2	217.340	286.322
749	0.0221	2	145.200	23.193	0.0221	2	45.900	51.336	0.0229	2	25.930	31.919	0.0113	1	28.700	-	0.0214	2	47.452	11.670
750	n.s. n.s.	n.s.	n.s.		0.0180	2	56.750	36.416	0.1005	9	16.866	18.117	0.0679	6	19.516	24.114	0.0844	8	11.937	6.673
751	n.s. n.s.	n.s.	n.s.		n.s. n.s.	n.s.	n.s.	n.s.	0.0454	4	4.253	3.543	0.0225	2	24.445	7.983	0.0413	4	9.038	8.141

$$(**) SD = \sqrt{\frac{\sum (x_i - \bar{x})^2}{n-1}}$$

**TABLE 7 (cont).**- Swept area, number of hauls and **roughhead grenadier** mean catch (Kg) and SD (\*\*) by stratum. Spanish Survey on NAFO Div. 3L in the period 2003-2012, on board R/V "Vizconde de Eza". (\*) In 2003, the data correspond to 69% of the total area prospected in 2006-2012.

Stratum	2009				2010				2011				2012			
	Swept area	Tow number	Mean catch	SD	Swept area	Tow number	Mean catch	SD	Swept area	Tow number	Mean catch	SD	Swept area	Tow number	Mean catch	SD
385	0.0225	2	0.000	0.000	0.0225	2	0.000	0.000	0.0229	2	1.010	1.428	0.0225	2	0.000	0.000
387	0.0439	4	30.045	16.013	0.0458	4	14.399	12.704	0.0450	4	11.304	9.250	0.0450	4	16.012	11.119
388	0.0555	5	27.627	27.428	0.0570	5	17.174	8.563	0.0563	5	5.022	4.969	0.0570	5	14.019	22.081
389	0.0803	7	31.105	63.627	0.0795	7	8.231	10.443	0.0675	6	4.711	3.126	0.0799	7	11.893	9.022
390	0.1373	12	4.648	14.283	0.1249	11	1.071	3.295	0.1009	9	2.856	7.168	0.1354	12	0.000	0.000
391	0.0458	4	72.878	56.298	0.0454	4	169.525	25.560	0.0458	4	153.179	92.811	0.0458	4	21.670	8.743
392	0.0229	2	60.934	78.701	0.0225	2	35.050	15.203	0.0229	2	83.417	29.674	0.0225	2	73.339	76.293
729	0.0341	3	9.991	5.382	0.0338	3	10.817	4.348	0.0338	3	3.398	2.102	0.0338	3	23.722	12.954
730	0.0338	3	75.453	99.963	0.0334	3	26.400	4.084	0.0334	3	66.456	55.464	0.0338	3	27.264	5.665
731	0.0341	3	4.980	1.654	0.0338	3	10.508	7.656	0.0334	3	2.002	1.506	0.0341	3	5.244	2.400
732	0.0450	4	8.289	3.314	0.0450	4	16.060	6.489	0.0454	4	2.393	2.786	0.0454	4	3.022	2.324
733	0.0450	4	19.108	13.978	0.0450	4	8.785	9.702	0.0454	4	6.622	8.721	0.0454	4	9.322	10.885
734	0.0218	2	28.777	12.760	0.0225	2	65.625	48.826	0.0225	2	8.413	1.874	0.0233	2	20.968	0.803
741	0.0221	2	11.334	6.316	0.0225	2	14.350	3.606	0.0218	2	7.707	9.880	0.0218	2	5.764	2.452
742	0.0214	2	3.425	1.803	0.0225	2	3.870	1.987	0.0225	2	14.545	14.221	0.0206	2	6.851	3.796
743	0.0203	2	13.278	13.438	0.0225	2	30.937	37.283	0.0221	2	18.488	1.660	0.0206	2	5.421	7.609
744	0.0210	2	8.208	6.495	0.0229	2	13.319	1.031	0.0221	2	6.254	3.743	0.0221	2	8.725	9.086
745	0.0559	5	3.787	2.256	0.0563	5	7.959	3.864	0.0446	4	2.802	4.240	0.0570	5	1.932	1.671
746	0.0668	6	23.474	20.537	0.0679	6	13.030	7.624	0.0566	5	8.981	7.193	0.0675	6	14.447	14.048
747	0.1118	10	33.180	25.868	0.1125	10	36.785	18.008	0.0893	8	22.273	17.958	0.1121	10	19.457	7.563
748	0.0229	2	92.330	127.477	0.0225	2	50.350	51.548	0.0221	2	25.955	33.074	0.0225	2	106.350	134.562
749	0.0225	2	13.700	9.334	0.0229	2	20.482	26.189	0.0221	2	27.713	30.670	0.0221	2	9.800	8.061
750	0.0791	7	16.895	14.145	0.0900	8	12.763	11.150	0.0668	6	9.292	4.047	0.0885	8	18.823	14.451
751	0.0338	3	88.193	144.495	0.0225	2	22.150	8.980	0.0334	3	14.880	6.137	0.0218	2	34.850	33.022

$$(**) SD = \sqrt{\frac{\sum (x_i - \bar{x})^2}{n-1}}$$

**TABLE 8.-** Stratified mean catches (Kg) of **roughhead grenadier** by stratum and year (2003-2012) and SD. Research Vessel *Vizconde de Eza*. n.s. means stratum not surveyed. In 2003: the data correspond to 69% of the total area prospected in 2006-2012.

Stratum	Survey									
	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
385	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	119.18	0.00
387	0.00	15356.54	-	8906.24	11773.44	5201.92	7691.52	3686.21	2893.70	4099.14
388	0.00	15458.10	-	9426.94	13445.76	5374.85	9862.70	6131.05	1792.71	5004.78
389	0.00	954.38	-	725.69	1565.18	9674.64	15832.37	4189.80	2397.73	6053.54
390	456.40	5.43	-	0.00	0.00	472.70	3787.71	872.79	2327.28	0.00
391	4.70	4.94	-	50230.55	24400.05	70203.05	20551.46	47806.05	43196.41	6110.94
392	565.50	29094.25	-	17113.63	18842.75	8435.38	8835.43	5082.25	12095.47	10634.08
729	7021.50	5482.35	-	4680.44	4927.20	3709.46	1858.39	2011.90	632.09	4412.23
730	17178.50	5731.55	-	9055.90	13834.26	5970.29	12827.07	4488.00	11297.58	4634.82
731	758.16	2257.20	-	2270.52	3095.93	3095.93	1075.61	2269.73	432.36	1132.78
732	7946.40	9122.19	-	5119.88	2575.96	4976.90	1914.82	3709.74	552.67	698.08
733	n.s.	3639.48	-	5487.30	4470.26	5601.67	4471.16	2055.69	1549.49	2181.41
734	n.s.	10075.05	-	6015.20	3580.20	4678.66	4402.88	10040.63	1287.19	3208.03
741	870.00	105.53	-	1755.70	465.00	1035.90	1133.40	1435.00	770.65	576.35
742	1561.60	300.80	-	1339.68	927.55	1079.10	219.20	247.68	930.85	438.46
743	n.s.	1338.50	-	539.27	1512.97	1300.93	677.18	1577.79	942.89	276.45
744	n.s.	168.30	-	1014.09	2241.69	3872.22	541.70	879.05	412.73	575.85
745	6106.24	2018.40	-	2866.88	1261.09	4970.83	1317.95	2769.59	975.10	672.20
746	25009.60	10272.36	-	16372.53	13565.94	12042.24	9201.61	5107.56	3520.47	5663.35
747	n.s.	31585.71	-	30630.47	45257.17	20791.04	24022.61	26632.56	16125.29	14086.51
748	8900.82	3579.89	-	10799.28	5331.80	34557.06	14680.47	8005.65	4126.85	16909.65
749	18295.20	5783.40	-	3267.18	3616.20	5978.95	1726.20	2580.67	3491.84	1234.80
750	n.s.	31553.00	-	9377.25	10850.99	6636.90	9393.86	7096.23	5166.44	10465.52
751	n.s.	n.s.	-	973.82	5597.91	2069.59	20196.12	5072.35	3407.52	7980.65
TOTAL	94674.62	183887.34	-	197968.44	193139.30	221730.20	176221.39	153747.96	120444.46	107049.61
( $\bar{y}$ )	21.16	29.38	-	30.52	29.77	34.18	27.17	23.70	18.57	16.50
SD	3.38	5.27	-	7.41	4.86	6.12	4.97	1.71	2.51	2.92

**TABLE 9.-** Survey estimates (by the swept area method) of **roughhead grenadier** biomass (t.) by stratum and year and their SD on NAFO Div. 3L (R/V *Vizconde de Eza*).  
n.s. means stratum not surveyed. In 2003, the data correspond to 69% of the total area prospected in 2006-2012.

Stratum	Survey									
	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
385	0	0	-	0	0	0	0	0	10	0
387	0	1437	-	792	1047	478	701	322	257	364
388	0	1472	-	832	1195	481	889	538	159	439
389	0	85	-	64	139	868	1381	369	213	531
390	41	0	-	0	0	41	331	77	208	0
391	0	0	-	4465	2169	6189	1797	4214	3777	534
392	49	2722	-	1496	1675	763	772	452	1058	945
729	669	496	-	416	438	330	163	179	56	392
730	1553	518	-	833	1230	555	1140	403	1016	412
731	66	194	-	200	275	281	95	202	39	100
732	706	869	-	460	229	446	170	330	49	62
733	n.s.	331	-	484	397	520	397	183	137	192
734	n.s.	995	-	535	318	423	405	893	114	276
741	77	10	-	161	41	99	102	128	71	53
742	134	25	-	117	82	103	21	22	83	43
743	n.s.	143	-	48	134	128	67	140	85	27
744	n.s.	17	-	89	206	350	52	77	37	52
745	537	190	-	251	112	448	118	246	87	59
746	2242	913	-	1455	1226	1133	827	451	311	503
747	n.s.	3082	-	2739	4023	1945	2150	2367	1445	1256
748	818	360	-	993	474	3178	1284	712	373	1503
749	1654	523	-	286	321	559	153	226	316	112
750	n.s.	3506	-	840	959	629	831	631	464	946
751	n.s.	n.s.	-	86	498	201	1795	451	306	734
TOTAL	8546	17887	-	17641	17190	20148	15641	13612	10672	9535
SD	1340	3240	-	4271	2799	3534	2844	972	1466	1676

**TABLE 10.- Roughhead grenadier** length distribution per haul mean catches by sex and year. Number per stratified mean catches. Spanish Summer Survey on NAFO 3L: 2004-2012 (R/V *Vizconde de Eza*). Indet. means indeterminate.

Length (cm.)	2004				2006				2007				2008				
	M	F	I	T	M	F	I	T	M	F	I	T	M	F	I	T	
1.5	0.00	0.00	0.02	0.02	0.00	0.00	0.04	0.04	0.00	0.02	0.01	0.03	0.00	0.00	0.02	0.02	
2.5	0.02	0.02	0.06	0.10	0.07	0.04	0.02	0.13	0.00	0.04	0.15	0.19	0.00	0.03	0.09	0.13	
3.5	0.77	0.21	0.18	1.15	0.61	0.16	0.14	0.91	0.40	0.17	0.70	1.26	0.28	0.08	1.42	1.78	
4.5	0.09	0.07	0.00	0.16	0.14	0.00	0.00	0.14	0.08	0.06	0.02	0.16	0.11	0.01	0.03	0.15	
5.5	0.36	0.42	0.00	0.77	0.12	0.15	0.00	0.27	0.34	0.21	0.02	0.57	0.10	0.13	0.01	0.24	
6.5	0.72	1.08	0.00	1.80	0.91	0.71	0.00	1.63	0.94	0.75	0.00	1.69	0.69	0.64	0.03	1.36	
7.5	0.32	0.39	0.00	0.71	0.62	0.48	0.00	1.10	0.28	0.33	0.00	0.61	0.24	0.38	0.00	0.62	
8.5	0.80	0.92	0.00	1.72	0.46	0.50	0.00	0.97	0.54	0.68	0.01	1.23	0.39	0.46	0.00	0.85	
9.5	1.05	1.12	0.00	2.17	0.95	0.87	0.00	1.82	0.60	0.81	0.00	1.42	0.74	0.58	0.00	1.31	
10.5	0.87	0.55	0.00	1.42	0.87	0.98	0.00	1.84	0.84	0.55	0.00	1.39	0.87	0.77	0.00	1.63	
11.5	1.03	1.13	0.00	2.16	1.36	1.26	0.00	2.62	1.21	1.12	0.00	2.32	1.19	1.32	0.00	2.51	
12.5	1.47	1.27	0.00	2.74	1.83	1.78	0.01	3.61	1.13	1.22	0.00	2.35	1.07	1.20	0.00	2.26	
13.5	1.64	1.05	0.00	2.69	1.66	1.75	0.01	3.41	1.46	1.45	0.00	2.91	1.58	1.36	0.00	2.93	
14.5	2.35	1.70	0.00	4.05	1.91	1.77	0.00	3.67	1.89	1.71	0.00	3.60	2.16	1.77	0.00	3.94	
15.5	3.43	2.57	0.00	6.00	2.21	1.64	0.00	3.85	1.54	1.47	0.00	3.01	2.61	2.21	0.00	4.82	
16.5	4.60	3.44	0.00	8.03	2.19	1.86	0.00	4.04	1.74	1.56	0.00	3.29	2.60	2.67	0.00	5.26	
17.5	3.98	3.20	0.00	7.18	3.45	1.88	0.01	5.34	1.97	1.45	0.00	3.41	1.92	1.97	0.00	3.89	
18.5	3.59	2.73	0.00	6.32	2.99	2.03	0.00	5.02	1.85	1.38	0.00	3.23	1.60	1.74	0.00	3.34	
19.5	1.98	2.90	0.00	4.88	1.73	2.94	0.00	4.66	1.57	1.57	0.00	3.14	1.36	1.77	0.00	3.13	
20.5	1.09	1.73	0.00	2.82	0.91	2.50	0.00	3.41	0.98	1.70	0.00	2.67	0.82	1.89	0.00	2.71	
21.5	0.13	1.13	0.00	1.26	0.51	2.60	0.00	3.11	0.40	2.38	0.00	2.78	0.37	1.71	0.00	2.09	
22.5	0.22	0.75	0.00	0.97	0.10	1.73	0.00	1.83	0.15	2.18	0.00	2.32	0.10	1.82	0.00	1.91	
23.5	0.00	0.57	0.00	0.57	0.03	1.44	0.00	1.47	0.05	1.90	0.00	1.95	0.03	1.83	0.00	1.86	
24.5	0.00	0.69	0.00	0.69	0.01	0.94	0.00	0.95	0.00	1.49	0.00	1.49	0.00	2.28	0.00	2.29	
25.5	0.01	0.43	0.00	0.45	0.00	0.84	0.00	0.84	0.01	1.18	0.00	1.20	0.00	1.87	0.00	1.87	
26.5	0.00	0.60	0.00	0.60	0.00	0.63	0.00	0.63	0.00	1.05	0.00	1.05	0.00	1.53	0.00	1.53	
27.5	0.00	0.15	0.00	0.15	0.00	0.25	0.00	0.25	0.00	0.69	0.00	0.69	0.00	0.88	0.00	0.88	
28.5	0.00	0.30	0.00	0.30	0.00	0.31	0.00	0.31	0.01	0.37	0.00	0.38	0.00	0.62	0.00	0.62	
29.5	0.00	0.24	0.00	0.24	0.00	0.20	0.00	0.20	0.01	0.35	0.00	0.37	0.00	0.58	0.00	0.58	
30.5	0.00	0.21	0.00	0.21	0.00	0.10	0.00	0.10	0.00	0.28	0.00	0.28	0.00	0.15	0.00	0.15	
31.5	0.00	0.24	0.00	0.24	0.00	0.13	0.00	0.13	0.00	0.21	0.00	0.21	0.00	0.11	0.00	0.11	
32.5	0.00	0.04	0.00	0.04	0.00	0.09	0.00	0.09	0.00	0.07	0.00	0.07	0.00	0.07	0.00	0.07	
33.5	0.00	0.01	0.00	0.01	0.00	0.04	0.00	0.04	0.00	0.03	0.00	0.03	0.00	0.03	0.00	0.03	
34.5	0.00	0.09	0.00	0.09	0.00	0.03	0.00	0.03	0.00	0.08	0.00	0.08	0.00	0.00	0.00	0.00	
35.5	0.00	0.08	0.00	0.08	0.00	0.01	0.00	0.01	0.00	0.05	0.00	0.05	0.00	0.01	0.00	0.01	
36.5	0.00	0.01	0.00	0.01	0.00	0.05	0.00	0.05	0.00	0.04	0.00	0.04	0.00	0.02	0.00	0.02	
37.5	0.00	0.06	0.00	0.06	0.00	0.01	0.00	0.01	0.00	0.02	0.00	0.02	0.00	0.00	0.00	0.00	
38.5	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.02	0.00	0.02	
39.5	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
40.5	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.02	0.00	0.02	0.00	0.00	0.00	0.00	
41.5	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	
Total	30.53	32.08	0.26	62.86	25.64	32.68	0.23	58.54	19.99	30.65	0.90	51.54	20.84	34.48	1.59	56.91	
Nº samples:					43				83				71				87
Nº Ind.:	1188	1359	17	2564	2107	2423	25	4555	1589	2246	69	3904	2022	3019	176	5217	
Sampled catch:					1579				2985				2712				3287
Range:					2-37.5				1.5-39				2-41				1.5-42.5
Total catch:					1579				2985				2712				3287
Total valid hauls:					50				100				94				100

**TABLE 10 (cont).- Roughhead grenadier** length distribution per haul mean catches by sex and year. Number per stratified mean catches. Spanish Summer Survey on NAFO 3L: 2003-2012 (R/V *Vizconde de Eza*). Indet. means indeterminate.

Length (cm.)	2009				2010				2011				2012				
	M	F	I	T	M	F	I	T	M	F	I	T	M	F	I	T	
1.5	0.00	0.00	0.01	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
2.5	0.01	0.00	0.13	0.15	0.03	0.00	0.26	0.29	0.02	0.00	0.17	0.19	0.00	0.00	0.24	0.24	
3.5	0.27	0.16	1.01	1.44	0.07	0.05	0.33	0.46	0.00	0.01	1.41	1.42	0.00	0.02	1.01	1.04	
4.5	0.07	0.00	0.05	0.12	0.04	0.09	0.01	0.15	0.03	0.03	0.07	0.14	0.10	0.03	0.03	0.16	
5.5	0.12	0.13	0.00	0.25	0.29	0.20	0.00	0.48	0.06	0.16	0.01	0.24	0.22	0.23	0.00	0.44	
6.5	0.38	0.45	0.00	0.83	0.58	0.59	0.00	1.17	0.24	0.30	0.02	0.56	0.92	0.88	0.00	1.80	
7.5	0.11	0.23	0.00	0.35	0.26	0.22	0.00	0.47	0.22	0.18	0.00	0.40	0.38	0.35	0.00	0.74	
8.5	0.25	0.30	0.00	0.54	0.28	0.36	0.00	0.64	0.44	0.34	0.00	0.78	0.38	0.28	0.00	0.66	
9.5	0.38	0.51	0.00	0.89	0.54	0.43	0.00	0.97	0.29	0.46	0.00	0.75	0.44	0.53	0.00	0.98	
10.5	0.56	0.52	0.00	1.08	0.76	0.66	0.00	1.42	0.31	0.42	0.00	0.72	0.57	0.42	0.00	0.99	
11.5	0.56	0.99	0.00	1.55	0.95	0.89	0.00	1.83	0.50	0.29	0.00	0.79	0.68	0.60	0.00	1.28	
12.5	1.24	0.91	0.00	2.15	1.26	1.10	0.00	2.37	0.62	0.63	0.00	1.25	0.65	0.59	0.00	1.24	
13.5	1.33	1.44	0.00	2.77	1.84	1.74	0.00	3.59	0.81	0.79	0.00	1.61	0.79	0.74	0.00	1.53	
14.5	1.58	1.53	0.00	3.11	2.46	2.38	0.00	4.85	1.48	1.13	0.00	2.61	1.26	0.91	0.00	2.17	
15.5	1.92	1.90	0.00	3.81	2.29	2.10	0.00	4.40	2.22	1.37	0.00	3.59	1.52	1.13	0.00	2.65	
16.5	1.96	1.80	0.00	3.76	2.32	2.49	0.00	4.80	2.24	1.41	0.00	3.65	1.63	1.02	0.00	2.65	
17.5	1.71	1.96	0.00	3.67	1.89	2.35	0.00	4.24	1.35	1.79	0.00	3.14	1.54	1.46	0.00	2.99	
18.5	1.31	1.52	0.00	2.83	1.35	2.30	0.00	3.65	1.31	1.99	0.00	3.30	1.06	1.38	0.00	2.45	
19.5	0.97	1.24	0.00	2.22	0.75	1.78	0.00	2.52	0.58	1.78	0.00	2.36	0.64	1.19	0.00	1.83	
20.5	0.59	1.22	0.00	1.81	0.36	1.26	0.00	1.62	0.16	1.26	0.00	1.42	0.29	1.25	0.00	1.55	
21.5	0.30	1.23	0.00	1.53	0.16	1.20	0.00	1.36	0.06	0.85	0.00	0.91	0.09	0.96	0.00	1.05	
22.5	0.15	1.21	0.00	1.37	0.04	0.85	0.00	0.89	0.06	0.66	0.00	0.72	0.01	0.98	0.00	0.99	
23.5	0.01	1.33	0.00	1.35	0.04	0.93	0.00	0.96	0.00	0.58	0.00	0.58	0.01	0.61	0.00	0.63	
24.5	0.00	1.25	0.00	1.25	0.00	0.56	0.00	0.56	0.01	0.73	0.00	0.74	0.00	0.70	0.00	0.70	
25.5	0.01	1.18	0.00	1.19	0.00	0.80	0.00	0.80	0.00	0.58	0.00	0.58	0.00	0.49	0.00	0.49	
26.5	0.00	1.19	0.00	1.19	0.00	0.56	0.00	0.56	0.00	0.63	0.00	0.63	0.00	0.45	0.00	0.45	
27.5	0.00	0.82	0.00	0.82	0.00	0.44	0.00	0.44	0.00	0.50	0.00	0.50	0.00	0.44	0.00	0.44	
28.5	0.00	0.52	0.00	0.52	0.00	0.38	0.00	0.38	0.00	0.37	0.00	0.37	0.00	0.23	0.00	0.23	
29.5	0.00	0.46	0.00	0.46	0.00	0.23	0.00	0.23	0.00	0.17	0.00	0.17	0.00	0.10	0.00	0.10	
30.5	0.00	0.27	0.00	0.27	0.00	0.11	0.00	0.11	0.00	0.10	0.00	0.10	0.00	0.08	0.00	0.08	
31.5	0.00	0.23	0.00	0.23	0.00	0.09	0.00	0.09	0.00	0.03	0.00	0.03	0.00	0.16	0.00	0.16	
32.5	0.00	0.14	0.00	0.14	0.00	0.06	0.00	0.06	0.00	0.04	0.00	0.04	0.00	0.02	0.00	0.02	
33.5	0.00	0.09	0.00	0.09	0.00	0.06	0.00	0.06	0.00	0.03	0.00	0.03	0.00	0.02	0.00	0.02	
34.5	0.00	0.08	0.00	0.08	0.00	0.06	0.00	0.06	0.00	0.01	0.00	0.01	0.00	0.02	0.00	0.02	
35.5	0.00	0.03	0.00	0.03	0.00	0.02	0.00	0.02	0.00	0.05	0.00	0.05	0.00	0.01	0.00	0.01	
36.5	0.00	0.03	0.00	0.03	0.00	0.05	0.00	0.05	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	
37.5	0.00	0.01	0.00	0.01	0.00	0.04	0.00	0.04	0.00	0.02	0.00	0.02	0.00	0.02	0.00	0.02	
38.5	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.03	0.00	0.03	0.00	0.03	0.00	0.03	
39.5	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
40.5	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
41.5	0.00	0.01	0.00	0.01	0.00	0.02	0.00	0.02	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
Total	15.78	26.93	1.21	43.93	18.58	27.44	0.61	46.63	13.01	19.73	1.68	34.43	13.19	18.35	1.29	32.83	
Nº samples:					81				84				83				82
Nº Ind.:	1409	2319	105	3833	1486	1997	65	3548	1037	1506	140	2683	1077	1413	113	2603	
Sampled catch:					2541				2234				1710				1508
Range:					2.0-41.5				2.5-42				2.5-39				2.5-38.5
Total catch:					2543				2234				1710				1508
Total valid hauls:					98				97				89				98

**TABLE 11.-** Swept area, number of hauls and **redfish** mean catch (Kg) and SD (\*\*) by stratum. Spanish Survey on NAFO Div. 3L in the period 2003-2012, on board R/V "Vizconde de Eza". (\*) In 2003, the data correspond to 69% of the total area prospected in 2006-2012.

Stratum	2003 (*)				2004				2006				2007				2008			
	Swept Tow area No.	Mean catch	SD	Swept Tow area No.	Mean catch	SD	Swept Tow area No.	Mean catch	SD	Swept Tow area No.	Mean catch	SD	Swept Tow area No.	Mean catch	SD	Swept Tow area No.	Mean catch	SD		
385	0.0225	2	0.001	0.001	0.0229	2	0.005	0.007	0.0229	2	0.000	0.000	0.0225	2	0.041	0.027	0.0229	2	0.495	0.644
387	0.0229	2	1.715	1.110	0.0214	2	56.000	51.619	0.0225	2	113.685	116.171	0.0225	2	80.400	34.083	0.0435	4	185.125	58.384
388	0.0334	3	6.453	6.142	0.0105	1	11.800	-	0.0566	5	66.040	32.355	0.0563	5	162.078	100.787	0.0559	5	212.750	142.882
389	0.0454	4	0.801	0.912	0.0225	2	33.050	44.901	0.0795	7	46.008	84.876	0.0900	8	10.723	18.542	0.0780	7	385.331	509.833
390	0.0563	5	0.580	1.242	0.0345	3	0.000	0.000	0.1249	11	0.188	0.318	0.1350	12	0.173	0.473	0.1395	12	0.922	2.280
391	0.0338	3	0.087	0.085	0.0218	2	1.435	1.718	0.0450	4	7.135	5.793	0.0450	4	6.013	6.351	0.0454	4	1093.130	1444.102
392	0.0116	1	46.300	-	0.0214	2	1222.320	1712.075	0.0229	2	4367.190	5741.976	0.0225	2	959.650	350.230	0.0221	2	209.150	15.203
729	0.0210	2	88.800	73.963	0.0221	2	310.250	239.780	0.0338	3	202.167	262.943	0.0338	3	128.889	184.792	0.0338	3	618.467	508.067
730	0.0221	2	231.080	64.389	0.0221	2	55.550	72.761	0.0326	3	145.923	148.390	0.0225	2	367.737	518.964	0.0323	3	29.790	42.861
731	0.0229	2	39.365	8.252	0.0233	2	79.550	68.236	0.0341	3	19.053	7.921	0.0338	3	37.100	28.646	0.0330	3	132.967	154.885
732	0.0113	1	72.200	-	0.0210	2	42.025	55.119	0.0334	3	5.638	7.067	0.0338	3	12.115	13.539	0.0446	4	11.975	11.596
733	n.s.	n.s.	n.s.	n.s.	0.0330	3	111.667	109.389	0.0454	4	72.600	47.167	0.0338	3	115.667	70.383	0.0431	4	132.600	203.165
734	n.s.	n.s.	n.s.	n.s.	0.0304	3	5.383	7.029	0.0225	2	12.328	3.921	0.0225	2	24.728	28.585	0.0221	2	22.485	27.457
741	0.0113	1	2240.000	-	0.0323	3	0.255	0.255	0.0218	2	0.000	0.000	0.0225	2	0.000	0.000	0.0210	2	0.555	0.049
742	0.0116	1	0.000	-	0.0120	1	0.331	-	0.0229	2	0.000	0.000	0.0225	2	0.300	0.424	0.0210	2	0.000	0.000
743	n.s.	n.s.	n.s.	n.s.	0.0188	2	2.090	2.956	0.0225	2	0.000	0.000	0.0225	2	0.000	0.000	0.0203	2	0.000	0.000
744	n.s.	n.s.	n.s.	n.s.	0.0101	1	0.000	-	0.0229	2	0.000	0.000	0.0218	2	0.479	0.677	0.0221	2	0.000	0.000
745	0.0341	3	1753.100	3028.408	0.0319	3	0.000	0.000	0.0686	6	0.119	0.221	0.0675	6	0.380	0.450	0.0555	5	0.364	0.664
746	0.0446	4	0.000	0.000	0.0338	3	0.000	0.000	0.0675	6	0.118	0.185	0.0664	6	0.000	0.000	0.0638	6	0.000	0.000
747	n.s.	n.s.	n.s.	n.s.	0.0308	3	0.200	0.346	0.1230	11	0.000	0.000	0.1238	11	0.000	0.000	0.1069	10	0.012	0.039
748	0.0109	1	2.700	-	0.0199	2	0.440	0.622	0.0326	3	0.130	0.225	0.0338	3	0.830	1.050	0.0218	2	4.290	6.067
749	0.0221	2	0.000	0.000	0.0221	2	0.000	0.000	0.0229	2	0.000	0.000	0.0113	1	0.000	-	0.0214	2	0.000	0.000
750	n.s.	n.s.	n.s.	n.s.	0.0180	2	0.000	0.000	0.1005	9	0.000	0.000	0.0679	6	0.000	0.000	0.0844	8	0.000	0.000
751	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	0.0454	4	0.000	0.000	0.0225	2	0.000	0.000	0.0413	4	0.000	0.000

$$(**) SD = \frac{\sum (x_i - \bar{x})}{n-1}$$

**TABLE 11 (cont).**- Swept area, number of hauls and **redfish** mean catch (Kg) and SD (\*\*) by stratum. Spanish Survey on NAFO Div. 3L in the period 2003-2012, on board R/V "Vizconde de Eza". (\*) In 2003, the data correspond to 69% of the total area prospected in 2006-2012

Stratum	2009				2010				2011				2012			
	Swept area	Tow number	Mean catch	SD	Swept area	Tow number	Mean catch	SD	Swept area	Tow number	Mean catch	SD	Swept area	Tow number	Mean catch	SD
385	0.0225	2	0.275	0.389	0.0225	2	0.000	0.000	0.0229	2	0.205	0.290	0.0225	2	0.000	0.000
387	0.0439	4	568.427	761.003	0.0458	4	278.625	163.544	0.0450	4	471.900	592.192	0.0450	4	456.188	146.956
388	0.0555	5	1686.275	2522.618	0.0570	5	922.261	770.678	0.0563	5	400.680	561.867	0.0570	5	3649.824	2735.118
389	0.0803	7	321.423	836.313	0.0795	7	3449.476	9037.325	0.0675	6	314.072	337.845	0.0799	7	5366.450	13039.715
390	0.1373	12	0.086	0.182	0.1249	11	0.005	0.011	0.1009	9	0.298	0.893	0.1354	12	0.307	0.723
391	0.0458	4	243.571	371.869	0.0454	4	2337.331	4421.647	0.0458	4	270.078	524.098	0.0458	4	1317.264	848.814
392	0.0229	2	797.546	42.491	0.0225	2	480.100	211.425	0.0229	2	7489.781	7767.171	0.0225	2	4138.815	2411.128
729	0.0341	3	50.830	11.765	0.0338	3	284.767	335.507	0.0338	3	1405.563	2154.649	0.0338	3	1491.733	2440.054
730	0.0338	3	167.600	193.999	0.0334	3	147.447	167.733	0.0334	3	98.992	73.752	0.0338	3	214.100	203.592
731	0.0341	3	37.000	30.152	0.0338	3	89.033	43.263	0.0334	3	45.227	32.987	0.0341	3	37.000	4.590
732	0.0450	4	8.311	9.503	0.0450	4	16.665	14.441	0.0454	4	12.480	9.605	0.0454	4	7.236	4.921
733	0.0450	4	59.725	53.776	0.0450	4	174.368	45.484	0.0454	4	255.160	236.623	0.0454	4	129.800	140.677
734	0.0218	2	16.220	17.367	0.0225	2	5.945	3.868	0.0225	2	7.888	0.972	0.0233	2	9.015	1.393
741	0.0221	2	0.903	0.012	0.0225	2	0.000	0.000	0.0218	2	0.500	0.707	0.0218	2	0.700	0.990
742	0.0214	2	0.000	0.000	0.0225	2	0.000	0.000	0.0225	2	0.208	0.294	0.0206	2	0.000	0.000
743	0.0203	2	5.575	7.884	0.0225	2	0.000	0.000	0.0221	2	0.000	0.000	0.0206	2	0.000	0.000
744	0.0210	2	0.000	0.000	0.0229	2	0.133	0.188	0.0221	2	0.858	1.213	0.0221	2	0.000	0.000
745	0.0559	5	0.000	0.000	0.0563	5	0.436	0.632	0.0446	4	0.745	1.007	0.0570	5	0.348	0.506
746	0.0668	6	0.043	0.106	0.0679	6	0.053	0.131	0.0566	5	0.000	0.000	0.0675	6	0.000	0.000
747	0.1118	10	0.000	0.000	0.1125	10	0.000	0.000	0.0893	8	0.379	1.071	0.1121	10	0.000	0.000
748	0.0229	2	1.576	2.228	0.0225	2	0.000	0.000	0.0221	2	0.595	0.134	0.0225	2	0.000	0.000
749	0.0225	2	0.000	0.000	0.0229	2	0.000	0.000	0.0221	2	0.000	0.000	0.0221	2	0.000	0.000
750	0.0791	7	0.230	0.609	0.0900	8	0.184	0.520	0.0668	6	0.242	0.592	0.0885	8	0.039	0.110
751	0.0338	3	0.000	0.000	0.0225	2	0.000	0.000	0.0334	3	0.000	0.000	0.0218	2	0.000	0.000

$$(**) SD = \sqrt{\frac{\sum (x_i - \bar{x})^2}{n-1}}$$

**TABLE 12.-** Stratified mean catches (Kg) of **redfish** by stratum and year (2003-2012) and SD. Research Vessel *Vizconde de Eza*. n.s. means stratum not surveyed. In 2003: the data correspond to 69% of the total area prospected in 2006-2012.

Stratum	Survey									
	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
385	0.12	0.59	-	0.00	4.84	58.35	32.45	0.00	24.19	0.00
387	439.04	14336.00	-	29103.36	20582.40	47392.00	145517.18	71328.00	120806.40	116784.00
388	2303.84	4212.60	-	23576.28	57861.85	75951.75	602000.18	329247.18	143042.76	1302987.17
389	407.58	16822.45	-	23418.22	5458.01	196133.55	163604.53	1755783.21	159862.48	2731523.05
390	472.70	0.00	-	153.59	141.00	751.23	70.36	3.93	242.69	250.00
391	24.44	404.67	-	2012.07	1695.53	308262.66	68687.02	659127.27	76161.86	371468.38
392	6713.50	177236.40	-	633242.55	139149.25	30326.75	115644.17	69614.50	1086018.17	600128.18
729	16516.80	57706.50	-	37603.00	23973.29	115034.80	9454.32	52966.60	261434.78	277462.40
730	39283.60	9443.50	-	24806.97	62515.29	5064.30	28492.00	25065.93	16828.70	36397.00
731	8502.84	17182.80	-	4115.52	8013.60	28720.80	7992.00	19231.20	9768.96	7992.00
732	16678.20	9707.78	-	1302.46	2798.49	2766.23	1919.90	3849.62	2882.88	1671.40
733	n.s	26130.00	-	16988.40	27066.00	31028.40	13975.65	40802.00	59707.44	30373.20
734	n.s	823.65	-	1886.11	3783.31	3440.21	2481.66	909.59	1206.79	1379.30
741	224000.00	25.50	-	0.00	0.00	55.50	90.25	0.00	50.00	70.00
742	0.00	21.18	-	0.00	19.20	0.00	0.00	0.00	13.31	0.00
743	n.s	106.59	-	0.00	0.00	0.00	284.33	0.00	0.00	0.00
744	n.s	0.00	-	0.00	31.58	0.00	0.00	8.78	56.63	0.00
745	610078.80	0.00	-	41.47	132.24	126.74	0.00	151.73	259.26	121.10
746	0.00	0.00	-	46.39	0.00	0.00	16.99	20.91	0.00	0.00
747	n.s	144.80	-	0.00	0.00	8.98	0.00	0.00	274.22	0.00
748	429.30	69.96	-	20.67	131.97	682.11	250.50	0.00	94.61	0.00
749	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00
750	n.s	0.00	-	0.00	0.00	0.00	127.88	102.17	134.37	21.68
751	n.s	n.s	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00
TOTAL	925850.76	334374.97		798317.04	353357.83	845804.35	1160641.36	3028212.59	1938870.48	5478628.86
	206.94	53.43		123.06	54.47	130.38	178.92	466.81	298.89	844.56
SD	136.03	28.87		90.99	11.94	36.35	69.07	285.47	130.15	396.90

**TABLE 13.-** Survey estimates (by the swept area method) of **redfish** biomass (t.) by stratum and year and their SD on NAFO Div. 3L (R/V *Vizconde de Eza*). n.s. means stratum not surveyed. In 2003, the data correspond to 69% of the total area prospected in 2006-2012.

Stratum	Survey									
	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
385	0	0	-	0	0	5	3	0	2	0
387	38	1341	-	2587	1830	4358	13267	6236	10738	10381
388	207	401	-	2082	5143	6797	54234	28881	12715	114297
389	36	1495	-	2062	485	17602	14271	154597	14210	239382
390	42	0	-	14	13	65	6	0	22	22
391	2	37	-	179	151	27175	6005	58105	6659	32478
392	578	16584	-	55365	12369	2741	10111	6188	94952	53345
729	1573	5216	-	3342	2131	10225	831	4708	23239	24663
730	3551	854	-	2281	5557	471	2533	2253	1513	3235
731	743	1478	-	362	712	2611	703	1709	878	703
732	1483	925	-	117	249	248	171	342	254	147
733	n.s.	2375	-	1498	2406	2878	1242	3627	5263	2678
734	n.s.	81	-	168	336	311	228	81	107	119
741	19911	2	-	0	0	5	8	0	5	6
742	0	2	-	0	2	0	0	0	1	0
743	n.s.	11	-	0	0	0	28	0	0	0
744	n.s.	0	-	0	3	0	0	1	5	0
745	53633	0	-	4	12	11	0	13	23	11
746	0	0	-	4	0	0	2	2	0	0
747	n.s.	14	-	0	0	1	0	0	25	0
748	39	7	-	2	12	63	22	0	9	0
749	0	0	-	0	0	0	0	0	0	0
750	n.s.	0	-	0	0	0	11	9	12	2
751	n.s.	n.s.	-	0	0	0	0	0	0	0
TOTAL	81837	30825		70066	31410	75567	103675	266754	170632	481469
SD	50717	17163		50718	6885	20435	40871	164597	72507	229026

**TABLE 14.- Redfish** length distribution per haul mean catches by sex and year. Number per stratified mean catches. Spanish Summer Survey on NAFO 3L: 2004-2012 (R/V *Vizconde de Eza*). Indet. means indeterminate.

Length (cm.)	2004				2006				2007				2008			
	M	F	I	T	M	F	I	T	M	F	I	T	M	F	I	T
4	0.00	0.00	0.04	0.04	0.00	0.01	0.00	0.01	0.00	0.00	0.04	0.04	0.00	0.00	0.16	0.16
6	0.00	0.00	3.68	3.68	0.10	0.05	2.83	2.98	0.00	0.00	17.45	17.45	0.00	0.00	8.19	8.19
8	0.24	0.09	5.05	5.39	0.90	1.28	13.68	15.86	0.01	0.19	26.86	27.06	0.00	0.00	17.35	17.35
10	3.48	2.03	1.69	7.20	2.18	1.28	1.82	5.28	1.45	2.17	1.64	5.26	0.81	0.21	57.74	58.76
12	4.86	3.30	4.93	13.09	3.00	3.27	0.12	6.40	4.45	3.71	0.53	8.69	3.70	2.13	17.78	23.62
14	3.92	1.83	2.16	7.91	11.25	8.43	0.00	19.68	3.44	1.80	0.01	5.25	8.31	3.62	0.11	12.04
16	8.87	5.17	0.26	14.30	20.69	19.49	0.00	40.18	5.97	3.81	0.00	9.77	19.39	18.88	0.00	38.27
18	11.87	12.09	0.00	23.96	14.29	13.66	0.00	27.95	11.85	13.08	0.00	24.92	66.37	46.99	0.05	113.41
20	23.04	20.03	0.00	43.07	23.65	11.01	0.00	34.66	25.50	15.85	0.00	41.35	96.85	63.72	0.00	160.57
22	19.46	18.47	0.00	37.93	41.88	31.01	0.00	72.89	36.00	30.40	0.00	66.41	81.51	63.44	0.00	144.94
24	30.92	12.78	0.00	43.70	40.39	44.21	0.00	84.60	19.89	32.60	0.00	52.48	49.16	50.05	0.00	99.21
26	35.91	14.43	0.00	50.34	9.50	58.30	0.00	67.79	7.34	11.29	0.00	18.63	25.59	33.03	0.00	58.62
28	16.80	12.42	0.00	29.22	8.69	64.05	0.00	72.74	4.69	6.69	0.00	11.39	22.11	21.05	0.00	43.16
30	5.36	5.65	0.00	11.01	6.12	47.61	0.00	53.73	4.33	5.57	0.00	9.90	10.25	9.73	0.00	19.99
32	0.52	2.70	0.00	3.22	4.13	23.73	0.00	27.86	5.48	7.42	0.00	12.90	3.50	4.98	0.00	8.48
34	0.16	0.59	0.00	0.76	0.72	3.74	0.00	4.47	2.66	2.82	0.00	5.48	1.11	2.86	0.00	3.96
36	0.30	0.37	0.00	0.66	0.12	2.15	0.00	2.27	0.20	0.96	0.00	1.16	0.49	0.68	0.00	1.18
38	0.12	0.05	0.00	0.17	0.08	1.05	0.00	1.12	0.05	0.13	0.00	0.18	0.06	0.29	0.00	0.35
40	0.00	0.00	0.00	0.00	0.02	0.01	0.00	0.03	0.02	0.03	0.00	0.06	0.01	0.12	0.00	0.13
42	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.01	0.03	0.00	0.04	0.01	0.11	0.00	0.12
44	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.01	0.00	0.00	0.00	0.00
46	0.00	0.00	0.00	0.00	0.00	0.03	0.00	0.03	0.00	0.00	0.00	0.00	0.00	0.13	0.00	0.13
48	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
50	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.01	0.00	0.00	0.00	0.00
52	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00
54	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Total	165.84	111.99	17.82	295.65	187.71	334.38	18.45	540.54	133.36	138.57	46.53	318.46	389.23	322.03	101.39	812.65
Nº samples:					28			48				51				52
Nº Ind.:	1903	1662	409	3974	3205	3089	1205	7499	2669	2360	2016	7045	3957	3147	1372	8476
Sampled catch:					3970			11080				4675				12283
Range:					5-39			5-48				5-53				5-47
Total catch:					3970			11080				4675				12283
Total valid hauls:					50			100				94				100

**TABLE 14 (cont).- Redfish** length distribution per haul mean catches by sex and year. Number per stratified mean catches. Spanish Summer Survey on NAFO 3L: 2003-2012 (R/V *Vizconde de Eza*). Indet. means indeterminate.

Length (cm.)	2009				2010				2011				2012			
	M	F	I	T	M	F	I	T	M	F	I	T	M	F	I	T
4	0.00	0.00	0.01	0.01	0.00	0.00	0.05	0.05	0.00	0.00	0.20	0.20	0.00	0.00	0.02	0.02
6	0.00	0.00	1.44	1.44	0.00	0.00	3.06	3.06	0.00	0.00	5.36	5.36	0.00	0.00	11.79	11.79
8	0.00	0.00	7.73	7.73	0.00	0.00	5.23	5.23	0.00	0.00	6.74	6.74	0.06	0.82	16.26	17.13
10	0.12	0.14	6.53	6.79	0.20	0.00	4.23	4.43	0.14	0.08	5.23	5.45	3.18	4.43	12.65	20.26
12	0.78	0.36	8.74	9.87	0.21	0.12	3.91	4.24	1.55	1.30	2.51	5.36	9.66	4.04	4.21	17.91
14	3.23	2.04	5.53	10.80	2.31	8.76	2.81	13.87	2.58	2.02	1.26	5.86	5.06	2.67	0.94	8.68
16	46.42	22.66	0.79	69.87	52.93	20.23	0.33	73.50	13.55	6.58	0.00	20.13	12.20	5.77	0.00	17.97
18	133.26	137.85	0.00	271.11	362.56	228.57	0.00	591.13	54.39	33.52	0.00	87.90	134.16	83.98	0.00	218.14
20	115.15	92.22	0.08	207.45	557.56	698.41	0.00	1255.97	141.06	124.18	0.00	265.25	635.81	404.59	0.00	1040.41
22	117.95	120.09	0.00	238.03	260.01	387.04	0.00	647.05	115.55	123.27	0.00	238.82	783.26	916.84	0.00	1700.11
24	67.44	106.44	0.00	173.88	91.63	122.89	0.00	214.51	165.60	80.38	0.00	245.98	279.36	676.30	0.00	955.66
26	15.72	82.79	0.00	98.51	53.99	95.89	0.00	149.88	110.11	66.27	0.00	176.37	118.77	229.31	0.00	348.08
28	9.27	17.36	0.00	26.62	21.46	66.19	0.00	87.65	33.80	104.64	0.00	138.43	23.11	113.92	0.00	137.02
30	2.75	10.77	0.00	13.52	8.10	14.77	0.00	22.87	5.54	79.03	0.00	84.57	6.96	74.74	0.00	81.70
32	2.46	4.50	0.00	6.96	4.85	10.51	0.00	15.36	2.92	27.91	0.00	30.82	3.54	30.04	0.00	33.58
34	2.23	2.06	0.00	4.29	2.69	4.84	0.00	7.54	1.12	17.35	0.00	18.48	3.37	6.71	0.00	10.08
36	0.60	1.49	0.00	2.10	1.25	2.39	0.00	3.64	1.18	5.13	0.00	6.31	1.21	2.74	0.00	3.96
38	0.15	0.03	0.00	0.19	0.60	1.72	0.00	2.31	0.21	0.67	0.00	0.88	1.21	1.64	0.00	2.85
40	0.32	0.37	0.00	0.70	0.06	0.95	0.00	1.01	0.01	0.05	0.00	0.06	0.06	0.46	0.00	0.53
42	0.00	0.04	0.00	0.04	0.06	1.79	0.00	1.85	0.02	0.00	0.00	0.02	0.00	0.01	0.00	0.01
44	0.00	0.31	0.00	0.31	0.00	0.11	0.00	0.11	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01
46	0.00	0.00	0.00	0.00	0.00	0.05	0.00	0.05	0.00	0.05	0.00	0.05	0.09	0.06	0.00	0.15
48	0.00	0.30	0.00	0.30	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.11	0.00	0.11
50	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
52	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.04	0.00	0.04	0.03	0.00	0.00	0.03
54	0.00	0.00	0.00	0.00	0.00	0.03	0.00	0.03	0.00	0.00	0.00	0.00	0.05	0.03	0.00	0.08
Total	517.84	601.84	30.85	1150.53	1420.47	1665.26	19.63	3105.35	649.33	672.46	21.31	1343.10	2021.16	2559.23	45.88	4626.26
Nº samples:				51				48				51				49
Nº Ind.:	3016	2723	558	6297	3216	3082	1178	7476	3017	3572	443	7032	3715	3954	502	8171
Sampled catch:				16615				42525				27586				76987
Range:				5-49				5-55				5-52				5-61
Total catch:				16615				42526				27586				76988
Total valid hauls:				98				97				89				98

**TABLE 15.-** Swept area, number of hauls and **thorny skate** mean catch (Kg) and SD (\*\*) by stratum. Spanish Survey on NAFO Div. 3L in the period 2003-2012, on board R/V "Vizconde de Eza". (\*) In 2003, the data correspond to 69% of the total area prospected in 2006-2012.

Stratum	2003 (*)				2004				2006				2007				2008			
	Swept area	Tow No.	Mean catch	SD	Swept area	Tow No.	Mean catch	SD	Swept area	Tow No.	Mean catch	SD	Swept area	Tow No.	Mean catch	SD	Swept area	Tow No.	Mean catch	SD
385	0.0225	2	0.000	0.000	0.0229	2	7.050	9.970	0.0229	2	6.044	4.588	0.0225	2	30.260	11.653	0.0229	2	37.608	26.315
387	0.0229	2	5.295	4.957	0.0214	2	10.700	2.263	0.0225	2	16.438	16.599	0.0225	2	32.485	2.143	0.0435	4	26.276	17.380
388	0.0334	3	13.273	13.347	0.0105	1	16.700	-	0.0566	5	44.186	24.414	0.0563	5	31.096	13.246	0.0559	5	37.148	12.932
389	0.0454	4	5.984	5.117	0.0225	2	10.900	13.294	0.0795	7	32.979	14.712	0.0900	8	25.861	11.704	0.0780	7	33.065	8.029
390	0.0563	5	0.190	0.425	0.0345	3	1.997	1.730	0.1249	11	5.529	7.479	0.1350	12	7.366	7.441	0.1395	12	5.044	7.191
391	0.0338	3	1.723	1.509	0.0218	2	64.250	65.125	0.0450	4	151.088	51.460	0.0450	4	100.658	56.818	0.0454	4	190.795	35.749
392	0.0116	1	10.050	-	0.0214	2	62.300	0.141	0.0229	2	149.500	165.604	0.0225	2	330.100	170.554	0.0221	2	159.247	95.534
729	0.0210	2	54.955	31.176	0.0221	2	140.375	186.712	0.0338	3	49.261	27.663	0.0338	3	164.760	243.624	0.0338	3	34.265	25.540
730	0.0221	2	71.400	60.670	0.0221	2	0.000	0.000	0.0326	3	4.348	7.532	0.0225	2	0.000	0.000	0.0323	3	0.000	0.000
731	0.0229	2	38.705	25.873	0.0233	2	18.510	22.330	0.0341	3	46.757	62.791	0.0338	3	57.448	64.552	0.0330	3	9.140	13.870
732	0.0113	1	76.200	-	0.0210	2	0.000	0.000	0.0334	3	2.015	1.851	0.0338	3	0.000	0.000	0.0446	4	0.727	1.454
733	n.s.	n.s.	n.s.	n.s.	0.0330	3	9.363	5.299	0.0454	4	14.573	8.911	0.0338	3	6.427	8.497	0.0431	4	14.693	15.502
734	n.s.	n.s.	n.s.	n.s.	0.0304	3	0.000	0.000	0.0225	2	0.000	0.000	0.0225	2	0.000	0.000	0.0221	2	0.000	0.000
741	0.0113	1	0	-	0.0323	3	0.000	0.000	0.0218	2	0.000	0.000	0.0225	2	0.000	0.000	0.0210	2	0.000	0.000
742	0.0116	1	0	-	0.0120	1	0.000	-	0.0229	2	0.000	0.000	0.0225	2	0.000	0.000	0.0210	2	0.000	0.000
743	n.s.	n.s.	n.s.	n.s.	0.0188	2	0.000	0.000	0.0225	2	0.000	0.000	0.0225	2	0.000	0.000	0.0203	2	0.000	0.000
744	n.s.	n.s.	n.s.	n.s.	0.0101	1	0.000	-	0.0229	2	0.000	0.000	0.0218	2	0.000	0.000	0.0221	2	0.000	0.000
745	0.0341	3	22.077	21.917	0.0319	3	0.000	0.000	0.0686	6	0.000	0.000	0.0675	6	0.000	0.000	0.0555	5	0.000	0.000
746	0.0446	4	2.318	3.703	0.0338	3	0.000	0.000	0.0675	6	0.000	0.000	0.0664	6	0.000	0.000	0.0638	6	0.000	0.000
747	n.s.	n.s.	-	-	0.0308	3	0.000	0.000	0.1230	11	0.000	0.000	0.1238	11	0.000	0.000	0.1069	10	0.000	0.000
748	0.0109	1	65.220	-	0.0199	2	0.000	0.000	0.0326	3	0.837	1.449	0.0338	3	0.000	0.000	0.0218	2	0.000	0.000
749	0.0221	2	8.060	6.067	0.0221	2	0.000	0.000	0.0229	2	0.000	0.000	0.0113	1	0.000	-	0.0214	2	0.000	0.000
750	n.s.	n.s.	n.s.	n.s.	0.0180	2	1.375	1.945	0.1005	9	0.393	1.180	0.0679	6	0.000	0.000	0.0844	8	0.000	0.000
751	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	0.0454	4	0.000	0.000	0.0225	2	0.000	0.000	0.0413	4	0.000	0.000

$$(**) SD = \frac{\sum (x_i - \bar{x})}{n-1}$$

**TABLE 15 (cont).**- Swept area, number of hauls and **thorny skate** mean catch (Kg) and SD (\*\*) by stratum. Spanish Survey on NAFO Div. 3L in the period 2003-2012, on board R/V "Vizconde de Eza". (\*) In 2003, the data correspond to 69% of the total area prospected in 2006-2012.

Stratum	2009				2010				2011				2012			
	Swept area	Tow number	Mean catch	SD	Swept area	Tow number	Mean catch	SD	Swept area	Tow number	Mean catch	SD	Swept area	Tow number	Mean catch	SD
385	0.0225	2	22.855	12.155	0.0225	2	4.230	5.204	0.0229	2	40.870	7.722	0.0225	2	38.670	8.358
387	0.0439	4	20.590	15.584	0.0458	4	22.350	21.258	0.0450	4	5.241	5.174	0.0450	4	7.559	6.290
388	0.0555	5	33.480	11.888	0.0570	5	34.932	33.326	0.0563	5	9.356	7.705	0.0570	5	42.734	32.557
389	0.0803	7	12.954	7.076	0.0795	7	27.170	24.762	0.0675	6	11.893	10.892	0.0799	7	14.376	12.301
390	0.1373	12	14.043	24.187	0.1249	11	12.900	8.972	0.1009	9	20.264	12.350	0.1354	12	18.599	15.739
391	0.0458	4	31.899	30.002	0.0454	4	24.041	14.994	0.0458	4	32.718	28.277	0.0458	4	38.843	29.385
392	0.0229	2	41.322	31.215	0.0225	2	36.728	7.462	0.0229	2	40.537	19.861	0.0225	2	178.990	196.916
729	0.0341	3	38.090	23.526	0.0338	3	6.453	5.814	0.0338	3	4.906	5.481	0.0338	3	35.344	8.527
730	0.0338	3	0.000	0.000	0.0334	3	0.012	0.021	0.0334	3	1.467	2.540	0.0338	3	3.670	6.357
731	0.0341	3	22.847	22.201	0.0338	3	11.114	11.389	0.0334	3	4.470	5.812	0.0341	3	3.263	2.986
732	0.0450	4	7.100	11.428	0.0450	4	0.000	0.000	0.0454	4	0.000	0.000	0.0454	4	0.000	0.000
733	0.0450	4	4.315	6.530	0.0450	4	5.573	4.374	0.0454	4	2.899	3.869	0.0454	4	5.995	4.874
734	0.0218	2	0.000	0.000	0.0225	2	0.000	0.000	0.0225	2	0.000	0.000	0.0233	2	0.010	0.014
741	0.0221	2	0.000	0.000	0.0225	2	0.000	0.000	0.0218	2	0.000	0.000	0.0218	2	0.000	0.000
742	0.0214	2	0.000	0.000	0.0225	2	0.011	0.016	0.0225	2	0.000	0.000	0.0206	2	0.000	0.000
743	0.0203	2	1.395	1.973	0.0225	2	0.000	0.000	0.0221	2	0.000	0.000	0.0206	2	0.000	0.000
744	0.0210	2	0.000	0.000	0.0229	2	0.000	0.000	0.0221	2	0.000	0.000	0.0221	2	0.000	0.000
745	0.0559	5	0.000	0.000	0.0563	5	0.650	1.453	0.0446	4	0.000	0.000	0.0570	5	0.004	0.008
746	0.0668	6	0.000	0.000	0.0679	6	0.000	0.000	0.0566	5	0.000	0.000	0.0675	6	0.000	0.000
747	0.1118	10	0.000	0.000	0.1125	10	0.000	0.000	0.0893	8	0.424	1.199	0.1121	10	0.000	0.000
748	0.0229	2	0.000	0.000	0.0225	2	0.000	0.000	0.0221	2	0.000	0.000	0.0225	2	0.000	0.000
749	0.0225	2	0.000	0.000	0.0229	2	0.000	0.000	0.0221	2	0.000	0.000	0.0221	2	0.000	0.000
750	0.0791	7	0.000	0.000	0.0900	8	0.000	0.000	0.0668	6	0.000	0.000	0.0885	8	0.000	0.000
751	0.0338	3	0.000	0.000	0.0225	2	0.000	0.000	0.0334	3	0.000	0.000	0.0218	2	0.000	0.000

$$(**) SD = \sqrt{\frac{\sum (x_i - \bar{x})^2}{n-1}}$$

**Table 16.-** Length-weight relationships in the calculation of biomass, for Division 3L (out ZEE Canada), 2003-2012 for **thorny skate and black dogfish**. The equation is Weight=a(Length+0.5)<sup>b</sup>. To calculate the parameters for the indeterminate individuals, we used the total data (males+females+indeterminate individuals).

Thorny skate				Black dogfish								
Year	Sex	L-W Equations	N	r <sup>2</sup>	Sex	L-W Equations	N	r <sup>2</sup>	Sex	L-W Equations	N	r <sup>2</sup>
2003	All	$W = 0.0050 L^{3.1712}$	305	0.9896		$W = 0.0081 L^{2.8882}$	20	0.9637				
	Males	$W = 0.0051 L^{3.1619}$	141	0.9906		$W = 0.1143 L^{2.2194}$	5	0.9382				
	Females	$W = 0.0048 L^{3.1855}$	164	0.9888		$W = 0.0072 L^{2.9265}$	15	0.9782				
2004	All	$W = 0.0067 L^{3.1187}$	186	0.9661		$W = 0.0025 L^{3.1608}$	113	0.9592				
	Males	$W = 0.0054 L^{3.1684}$	94	0.97		$W = 0.0272 L^{2.5776}$	58	0.8969				
	Females	$W = 0.0086 L^{3.0629}$	92	0.9639		$W = 0.0013 L^{3.3314}$	55	0.9809				
2006	All	$W = 0.0084 L^{3.0587}$	491	0.983		$W = 0.0011 L^{3.3758}$	283	0.9216				
	Males	$W = 0.0103 L^{3.0011}$	210	0.9847		$W = 0.0071 L^{2.9000}$	99	0.9233				
	Females	$W = 0.0061 L^{3.1402}$	281	0.9814		$W = 0.0008 L^{3.4608}$	184	0.9363				
2007	All	$W = 0.0080 L^{3.0609}$	539	0.9848		$W = 0.0008 L^{3.4421}$	362	0.9155				
	Males	$W = 0.0091 L^{3.0242}$	255	0.9868		$W = 0.0099 L^{2.8281}$	147	0.9029				
	Females	$W = 0.0072 L^{3.0929}$	284	0.9839		$W = 0.0006 L^{3.5445}$	215	0.9373				
2008	All	$W = 0.0071 L^{3.0883}$	598	0.9884		$W = 0.0014 L^{3.3183}$	279	0.9006				
	Males	$W = 0.0077 L^{3.0618}$	282	0.9903		$W = 0.0087 L^{2.8575}$	160	0.8956				
	Females	$W = 0.0064 L^{3.1175}$	316	0.9867		$W = 0.0008 L^{3.4541}$	119	0.9283				
2009	All	$W = 0.0072 L^{3.0862}$	283	0.9864		$W = 0.0007 L^{3.4922}$	236	0.9246				
	Males	$W = 0.0093 L^{3.0231}$	171	0.9848		$W = 0.0132 L^{2.7605}$	75	0.8865				
	Females	$W = 0.0057 L^{3.1507}$	112	0.9881		$W = 0.0007 L^{3.5184}$	161	0.9465				
2010	All	$W = 0.0060 L^{3.1361}$	290	0.9906		$W = 0.0019 L^{3.2510}$	299	0.9506				
	Males	$W = 0.0060 L^{3.1285}$	149	0.9892		$W = 0.0137 L^{2.7559}$	130	0.9408				
	Females	$W = 0.0056 L^{3.1630}$	141	0.9927		$W = 0.0012 L^{3.3617}$	169	0.9637				
2011	All	$W = 0.0031 L^{3.2899}$	218	0.9937		$W = 0.0020 L^{3.2316}$	455	0.9518				
	Males	$W = 0.0036 L^{3.2468}$	136	0.9941		$W = 0.0059 L^{2.9580}$	171	0.9493				
	Females	$W = 0.0024 L^{3.3657}$	82	0.9941		$W = 0.0014 L^{3.3220}$	284	0.9568				
2012	All	$W = 0.0065 L^{3.1140}$	352	0.9918		$W = 0.0019 L^{3.2460}$	242	0.9531				
	Males	$W = 0.0085 L^{3.0429}$	219	0.9925		$W = 0.0107 L^{2.8100}$	116	0.9571				
	Females	$W = 0.0040 L^{3.2467}$	133	0.9933		$W = 0.0010 L^{3.4151}$	126	0.9718				

**TABLE 17.-** Stratified mean catches (Kg) of **thorny skate** by stratum and year (2003-2012) and SD. Research Vessel *Vizconde de Eza*. n.s. means stratum not surveyed. In 2003: the data correspond to 69% of the total area prospected in 2006-2012.

Stratum	Survey									
	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
385	0.00	831.90	-	713.19	3570.68	4437.69	2696.89	499.14	4822.66	4563.06
387	1355.52	2739.20	-	4208.00	8316.16	6726.59	5271.04	5721.60	1341.76	1935.04
388	4738.58	5961.90	-	15774.40	11101.27	13261.69	11952.50	12470.58	3340.16	15256.04
389	3045.60	5548.10	-	16786.09	13163.25	16830.16	6593.66	13829.31	6053.28	7317.60
390	154.85	1627.28	-	4506.21	6003.36	4110.66	11444.98	10513.50	16515.07	15158.46
391	485.98	18118.50	-	42606.68	28385.42	53804.19	8995.45	6779.63	9226.41	10953.66
392	1457.25	9033.50	-	21677.50	47864.50	23090.82	5991.69	5325.49	5877.79	25953.48
729	10221.63	26109.75	-	9162.48	30645.36	6373.35	7084.74	1200.20	912.52	6573.92
730	12138.00	0.00	-	739.22	0.00	0.00	0.00	2.04	249.33	623.90
731	8360.28	3998.16	-	10099.44	12408.84	1974.24	4934.88	2400.70	965.52	704.74
732	17602.20	0.00	-	465.47	0.00	167.94	1640.10	0.00	0.00	0.00
733	n.s.	2191.02	-	3410.14	1503.84	3438.05	1009.71	1304.02	678.31	1402.83
734	n.s.	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	1.53
741	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00
742	0.00	0.00	-	0.00	0.00	0.00	0.00	0.70	0.00	0.00
743	n.s.	0.00	-	0.00	0.00	0.00	71.15	0.00	0.00	0.00
744	n.s.	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00
745	7682.68	0.00	-	0.00	0.00	0.00	0.00	226.20	0.00	1.32
746	908.46	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00
747	n.s.	0.00	-	0.00	0.00	0.00	0.00	0.00	306.80	0.00
748	10369.98	0.00	-	133.03	0.00	0.00	0.00	0.00	0.00	0.00
749	1015.56	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00
750	n.s.	764.50	-	218.69	0.00	0.00	0.00	0.00	0.00	0.00
751	n.s.	n.s.	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00
TOTAL	79536.57	76923.81	-	130500.54	162962.67	134215.36	67686.78	60273.11	50289.61	90445.57
SD	17.78	12.29	-	20.12	25.12	20.69	10.43	9.29	7.75	13.94
	2.41	4.54	-	3.27	5.19	1.92	1.44	1.30	0.98	3.36

**TABLE 18.-** Survey estimates (by the swept area method) of **thorny skate** biomass (t.) by stratum and year and their SD on NAFO Div. 3L (R/V *Vizconde de Eza*). n.s. means stratum not surveyed. In 2003, the data correspond to 69% of the total area prospected in 2006-2012.

Stratum	Survey									
	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
385	0	73	-	62	317	388	240	44	422	406
387	119	256	-	374	739	619	481	500	119	172
388	426	568	-	1393	987	1187	1077	1094	297	1338
389	268	493	-	1478	1170	1510	575	1218	538	641
390	14	142	-	397	534	354	1001	926	1473	1344
391	43	1666	-	3787	2523	4743	786	598	807	958
392	125	845	-	1895	4255	2087	524	473	514	2307
729	973	2360	-	814	2724	567	623	107	81	584
730	1097	0	-	68	0	0	0	0	22	55
731	731	344	-	888	1103	179	434	213	87	62
732	1565	0	-	42	0	15	146	0	0	0
733	n.s.	199	-	301	134	319	90	116	60	124
734	n.s.	0	-	0	0	0	0	0	0	0
741	0	0	-	0	0	0	0	0	0	0
742	0	0	-	0	0	0	0	0	0	0
743	n.s.	0	-	0	0	0	7	0	0	0
744	n.s.	0	-	0	0	0	0	0	0	0
745	675	0	-	0	0	0	0	20	0	0
746	81	0	-	0	0	0	0	0	0	0
747	n.s.	0	-	0	0	0	0	0	28	0
748	954	0	-	12	0	0	0	0	0	0
749	92	0	-	0	0	0	0	0	0	0
750	n.s.	85	-	20	0	0	0	0	0	0
751	n.s.	n.s.	-	0	0	0	0	0	0	0
TOTAL	7164	7031	-	11531	14486	11968	5982	5310	4448	7991
SD	942	2642	-	1887	2993	1124	808	740	560	2008

**TABLE 19.-** Thorny skate length distribution per haul mean catches by sex and year. Number per stratified mean catches. Spanish Summer Survey on NAFO 3L: 2004-2012 (R/V *Vizconde de Eza*). Indet. means indeterminate.

Length (cm.)	2004				2006				2007				2008				
	M	F	I	T	M	F	I	T	M	F	I	T	M	F	I	T	
10	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
12	0.00	0.00	0.00	0.00	0.04	0.01	0.00	0.05	0.16	0.08	0.00	0.24	0.09	0.11	0.00	0.20	
14	0.04	0.00	0.00	0.04	0.09	0.00	0.00	0.09	0.08	0.12	0.00	0.21	0.12	0.09	0.00	0.20	
16	0.00	0.00	0.00	0.00	0.04	0.03	0.00	0.07	0.05	0.10	0.00	0.15	0.03	0.03	0.00	0.06	
18	0.02	0.00	0.00	0.02	0.03	0.02	0.00	0.06	0.13	0.12	0.00	0.25	0.04	0.01	0.00	0.05	
20	0.00	0.02	0.00	0.02	0.03	0.03	0.00	0.06	0.18	0.10	0.00	0.28	0.09	0.01	0.00	0.10	
22	0.00	0.00	0.00	0.00	0.00	0.03	0.00	0.03	0.15	0.19	0.00	0.34	0.02	0.01	0.00	0.03	
24	0.00	0.00	0.00	0.00	0.03	0.01	0.00	0.05	0.13	0.14	0.00	0.27	0.02	0.02	0.00	0.04	
26	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.13	0.16	0.00	0.30	0.08	0.07	0.00	0.14	
28	0.00	0.00	0.00	0.00	0.02	0.01	0.00	0.03	0.10	0.06	0.00	0.16	0.02	0.05	0.00	0.08	
30	0.00	0.00	0.00	0.00	0.00	0.03	0.00	0.03	0.08	0.05	0.00	0.12	0.04	0.05	0.00	0.10	
32	0.01	0.02	0.00	0.04	0.03	0.01	0.00	0.05	0.08	0.06	0.00	0.13	0.07	0.05	0.00	0.12	
34	0.01	0.00	0.00	0.01	0.01	0.03	0.00	0.05	0.09	0.03	0.00	0.12	0.05	0.04	0.00	0.10	
36	0.01	0.00	0.00	0.01	0.02	0.01	0.00	0.03	0.06	0.05	0.00	0.11	0.03	0.05	0.00	0.08	
38	0.00	0.00	0.00	0.00	0.00	0.04	0.00	0.04	0.05	0.06	0.00	0.11	0.01	0.03	0.00	0.04	
40	0.04	0.05	0.00	0.10	0.05	0.03	0.00	0.08	0.02	0.01	0.00	0.03	0.05	0.01	0.00	0.06	
42	0.07	0.07	0.00	0.13	0.00	0.03	0.00	0.03	0.03	0.06	0.00	0.09	0.02	0.05	0.00	0.07	
44	0.10	0.25	0.00	0.35	0.01	0.03	0.00	0.05	0.04	0.04	0.00	0.08	0.01	0.02	0.00	0.03	
46	0.14	0.14	0.00	0.28	0.09	0.08	0.00	0.17	0.05	0.09	0.00	0.14	0.03	0.06	0.00	0.09	
48	0.17	0.12	0.00	0.30	0.10	0.08	0.00	0.18	0.05	0.09	0.00	0.14	0.02	0.01	0.00	0.03	
50	0.26	0.24	0.00	0.50	0.13	0.17	0.00	0.30	0.12	0.13	0.00	0.25	0.06	0.03	0.00	0.09	
52	0.35	0.26	0.00	0.61	0.22	0.13	0.00	0.35	0.09	0.15	0.00	0.24	0.07	0.08	0.00	0.15	
54	0.33	0.14	0.00	0.47	0.27	0.37	0.00	0.64	0.21	0.24	0.00	0.44	0.08	0.09	0.00	0.17	
56	0.32	0.23	0.00	0.55	0.22	0.24	0.00	0.45	0.19	0.34	0.00	0.53	0.03	0.13	0.00	0.16	
58	0.32	0.24	0.00	0.56	0.22	0.46	0.00	0.67	0.30	0.27	0.00	0.57	0.12	0.22	0.00	0.34	
60	0.41	0.20	0.00	0.61	0.36	0.39	0.00	0.75	0.27	0.59	0.00	0.86	0.22	0.28	0.00	0.50	
62	0.11	0.15	0.00	0.26	0.22	0.53	0.00	0.76	0.46	0.76	0.00	1.22	0.29	0.35	0.00	0.65	
64	0.18	0.19	0.00	0.37	0.41	0.54	0.00	0.95	0.42	0.62	0.00	1.04	0.35	0.45	0.00	0.81	
66	0.22	0.14	0.00	0.37	0.34	0.39	0.00	0.72	0.34	0.54	0.00	0.88	0.39	0.45	0.00	0.84	
68	0.12	0.05	0.00	0.17	0.17	0.41	0.00	0.58	0.37	0.64	0.00	1.02	0.32	0.44	0.00	0.76	
70	0.05	0.02	0.00	0.08	0.19	0.22	0.00	0.41	0.25	0.38	0.00	0.62	0.25	0.37	0.00	0.62	
72	0.06	0.00	0.00	0.06	0.08	0.13	0.00	0.21	0.18	0.24	0.00	0.43	0.19	0.15	0.00	0.34	
74	0.00	0.00	0.00	0.00	0.09	0.07	0.00	0.16	0.12	0.13	0.00	0.25	0.26	0.16	0.00	0.42	
76	0.00	0.00	0.00	0.00	0.08	0.05	0.00	0.13	0.04	0.05	0.00	0.10	0.10	0.13	0.00	0.23	
78	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.03	0.03	0.00	0.06	0.09	0.03	0.00	0.12	
80	0.00	0.00	0.00	0.00	0.01	0.01	0.00	0.02	0.01	0.00	0.00	0.01	0.07	0.00	0.00	0.07	
82	0.08	0.00	0.00	0.08	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.01	0.05	0.02	0.00	0.07	
84	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.03	0.00	0.00	0.03	
86	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.02	0.00	0.00	0.02	
88	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.01	
Total	3.44	2.56	0.00	5.99	3.61	4.66	0.00	8.27	5.10	6.71	0.00	11.81	5.10	6.71	0.00	11.81	
Nº samples:					18				42				43				
Nº Ind.:	170	135	0	305	312	420	0	732	457	621	0	1078	457	621	0	1078	
Sampled catch:					617				1832				2325				2325
Range:					14-83				13-81				12-82				12-82
Total catch:					682				1832				2325				2325
Total valid hauls:					50				100				94				94

**TABLE 19.-** Thorny skate length distribution per haul mean catches by sex and year. Number per stratified mean catches. Spanish Summer Survey on NAFO 3L: 2003-2012 (R/V *Vizconde de Eza*). Indet. means indeterminate.

Length (cm.)	2009				2010				2011				2012			
	M	F	I	T	M	F	I	T	M	F	I	T	M	F	I	T
10	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00
12	0.11	0.05	0.00	0.16	0.05	0.05	0.00	0.10	0.09	0.03	0.00	0.11	0.09	0.05	0.00	0.14
14	0.06	0.07	0.00	0.13	0.08	0.07	0.00	0.15	0.06	0.08	0.00	0.14	0.07	0.05	0.00	0.11
16	0.02	0.02	0.00	0.04	0.00	0.03	0.00	0.03	0.01	0.00	0.00	0.01	0.01	0.00	0.00	0.01
18	0.00	0.05	0.00	0.05	0.01	0.02	0.00	0.03	0.00	0.01	0.00	0.01	0.03	0.00	0.00	0.03
20	0.01	0.03	0.00	0.04	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	0.04	0.01	0.00	0.05
22	0.00	0.00	0.00	0.00	0.00	0.02	0.00	0.02	0.00	0.00	0.00	0.00	0.03	0.03	0.00	0.06
24	0.02	0.02	0.00	0.05	0.01	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.05	0.06	0.00	0.11
26	0.01	0.00	0.00	0.01	0.01	0.02	0.00	0.03	0.00	0.00	0.00	0.00	0.04	0.03	0.00	0.07
28	0.02	0.01	0.00	0.03	0.00	0.01	0.00	0.01	0.00	0.03	0.00	0.03	0.03	0.11	0.00	0.14
30	0.02	0.03	0.00	0.06	0.04	0.02	0.00	0.07	0.02	0.01	0.00	0.03	0.08	0.01	0.00	0.09
32	0.00	0.01	0.00	0.01	0.00	0.03	0.00	0.03	0.01	0.00	0.00	0.01	0.04	0.04	0.00	0.08
34	0.01	0.01	0.00	0.02	0.01	0.04	0.00	0.05	0.00	0.01	0.00	0.01	0.04	0.04	0.00	0.08
36	0.00	0.00	0.00	0.00	0.04	0.02	0.00	0.07	0.02	0.01	0.00	0.03	0.06	0.06	0.00	0.12
38	0.02	0.01	0.00	0.03	0.02	0.01	0.00	0.03	0.00	0.02	0.00	0.02	0.06	0.04	0.00	0.10
40	0.02	0.00	0.00	0.02	0.00	0.01	0.00	0.01	0.01	0.01	0.00	0.03	0.07	0.05	0.00	0.12
42	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	0.06	0.01	0.00	0.07
44	0.01	0.04	0.00	0.05	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	0.02	0.02	0.00	0.05
46	0.00	0.01	0.00	0.01	0.01	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.02	0.03	0.00	0.05
48	0.01	0.02	0.00	0.03	0.01	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.01
50	0.05	0.01	0.00	0.06	0.00	0.01	0.00	0.01	0.01	0.00	0.00	0.01	0.02	0.05	0.00	0.07
52	0.02	0.02	0.00	0.04	0.01	0.02	0.00	0.03	0.00	0.00	0.00	0.00	0.05	0.03	0.00	0.08
54	0.05	0.05	0.00	0.09	0.00	0.02	0.00	0.02	0.00	0.01	0.00	0.01	0.02	0.01	0.00	0.03
56	0.02	0.15	0.00	0.17	0.02	0.04	0.00	0.07	0.00	0.01	0.00	0.01	0.01	0.00	0.00	0.01
58	0.13	0.09	0.00	0.22	0.08	0.09	0.00	0.17	0.02	0.05	0.00	0.07	0.03	0.04	0.00	0.07
60	0.16	0.08	0.00	0.24	0.03	0.12	0.00	0.16	0.06	0.09	0.00	0.15	0.03	0.05	0.00	0.08
62	0.23	0.24	0.00	0.47	0.08	0.10	0.00	0.18	0.03	0.08	0.00	0.10	0.07	0.10	0.00	0.18
64	0.23	0.14	0.00	0.36	0.12	0.16	0.00	0.28	0.15	0.06	0.00	0.20	0.10	0.11	0.00	0.21
66	0.25	0.18	0.00	0.43	0.21	0.18	0.00	0.38	0.13	0.09	0.00	0.23	0.10	0.14	0.00	0.24
68	0.28	0.18	0.00	0.47	0.19	0.23	0.00	0.42	0.19	0.12	0.00	0.31	0.28	0.22	0.00	0.50
70	0.19	0.07	0.00	0.26	0.21	0.07	0.00	0.28	0.17	0.15	0.00	0.32	0.30	0.09	0.00	0.38
72	0.17	0.09	0.00	0.25	0.13	0.08	0.00	0.21	0.18	0.03	0.00	0.22	0.37	0.12	0.00	0.49
74	0.19	0.01	0.00	0.20	0.11	0.05	0.00	0.16	0.16	0.05	0.00	0.21	0.23	0.03	0.00	0.26
76	0.02	0.03	0.00	0.06	0.09	0.03	0.00	0.12	0.11	0.00	0.00	0.11	0.19	0.07	0.00	0.26
78	0.04	0.03	0.00	0.07	0.09	0.01	0.00	0.10	0.05	0.00	0.00	0.05	0.21	0.03	0.00	0.24
80	0.01	0.00	0.00	0.01	0.03	0.00	0.00	0.03	0.04	0.00	0.00	0.04	0.14	0.01	0.00	0.15
82	0.01	0.00	0.00	0.01	0.02	0.00	0.00	0.02	0.01	0.00	0.00	0.01	0.09	0.01	0.00	0.10
84	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.01	0.07	0.00	0.00	0.07
86	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.02	0.00	0.00	0.02
88	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.01	0.04	0.00	0.00	0.04	0.01	0.00	0.00	0.01
Total	2.41	1.76	0.00	4.17	1.73	1.62	0.00	3.35	1.59	0.96	0.00	2.55	3.19	1.74	0.00	4.93
Nº samples:				44				46				39				44
Nº Ind.:	211	156	0	367	159	145	0	304	136	82	0	218	266	151	0	417
Sampled catch:				996.2				853				663				1309
Range:				12-82				12-88				11-88				12-88
Total catch:				996.2				853				663				1309
Total valid hauls:				98				97				89				98

**TABLE 20.-** Swept area, number of hauls and **black dogfish** mean catch (Kg) and SD (\*\*) by stratum. Spanish Survey on NAFO Div. 3L in the period 2003-2012, on board R/V "Vizconde de Eza". (\*) In 2003, the data correspond to 69% of the total area prospected in 2006-2012.

Stratum	2003 (*)				2004				2006				2007				2008			
	Swept area	Tow No.	Mean catch	SD	Swept area	Tow No.	Mean catch	SD	Swept area	Tow No.	Mean catch	SD	Swept area	Tow No.	Mean catch	SD	Swept area	Tow No.	Mean catch	SD
385	0.0225	2	0.000	0.000	0.0229	2	0.000	0.000	0.0229	2	0.000	0.000	0.0225	2	0.000	0.000	0.0229	2	0.000	0.000
387	0.0229	2	0.000	0.000	0.0214	2	0.000	0.000	0.0225	2	0.000	0.000	0.0225	2	0.000	0.000	0.0435	4	0.000	0.000
388	0.0334	3	0.000	0.000	0.0105	1	0.000	-	0.0566	5	0.000	0.000	0.0563	5	0.000	0.000	0.0559	5	0.000	0.000
389	0.0454	4	0.000	0.000	0.0225	2	0.000	0.000	0.0795	7	0.000	0.000	0.0900	8	0.000	0.000	0.0780	7	0.000	0.000
390	0.0563	5	0.000	0.000	0.0345	3	0.000	0.000	0.1249	11	0.000	0.000	0.1350	12	0.000	0.000	0.1395	12	0.000	0.000
391	0.0338	3	0.000	0.000	0.0218	2	0.000	0.000	0.0450	4	0.000	0.000	0.0450	4	0.000	0.000	0.0454	4	0.000	0.000
392	0.0116	1	0.000	-	0.0214	2	0.000	0.000	0.0229	2	0.000	0.000	0.0225	2	0.000	0.000	0.0221	2	0.000	0.000
729	0.0210	2	0.000	0.000	0.0221	2	0.000	0.000	0.0338	3	0.000	0.000	0.0338	3	0.000	0.000	0.0338	3	0.000	0.000
730	0.0221	2	0.000	0.000	0.0221	2	2.175	3.076	0.0326	3	3.690	6.391	0.0225	2	19.488	26.067	0.0323	3	27.367	47.400
731	0.0229	2	0.000	0.000	0.0233	2	0.000	0.000	0.0341	3	0.000	0.000	0.0338	3	0.000	0.000	0.0330	3	0.000	0.000
732	0.0113	1	0.000	-	0.0210	2	0.000	0.000	0.0334	3	0.000	0.000	0.0338	3	0.000	0.000	0.0446	4	0.000	0.000
733	n.s.	n.s.	n.s.	n.s.	0.0330	3	0.000	0.000	0.0454	4	0.000	0.000	0.0338	3	0.000	0.000	0.0431	4	0.000	0.000
734	n.s.	n.s.	n.s.	n.s.	0.0304	3	0.000	0.000	0.0225	2	0.000	0.000	0.0225	2	0.000	0.000	0.0221	2	0.000	0.000
741	0.0113	1	0.000	-	0.0323	3	0.000	0.000	0.0218	2	0.000	0.000	0.0225	2	0.000	0.000	0.0210	2	0.000	0.000
742	0.0116	1	0.000	-	0.0120	1	0.000	-	0.0229	2	0.000	0.000	0.0225	2	0.000	0.000	0.0210	2	0.000	0.000
743	n.s.	n.s.	n.s.	n.s.	0.0188	2	0.626	0.862	0.0225	2	0.000	0.000	0.0225	2	0.000	0.000	0.0203	2	0.000	0.000
744	n.s.	n.s.	n.s.	n.s.	0.0101	1	0.000	-	0.0229	2	0.725	1.025	0.0218	2	1.663	0.541	0.0221	2	0.880	0.198
745	0.0341	3	0.007	0.012	0.0319	3	0.000	0.000	0.0686	6	0.000	0.000	0.0675	6	0.000	0.000	0.0555	5	0.000	0.000
746	0.0446	4	0.000	0.000	0.0338	3	0.000	0.000	0.0675	6	9.033	10.572	0.0664	6	9.171	6.742	0.0638	6	6.142	1.917
747	n.s.	n.s.	n.s.	n.s.	0.0308	3	4.067	3.591	0.1230	11	3.656	2.707	0.1238	11	6.015	5.815	0.1069	10	5.894	5.184
748	0.0109	1	0.000	-	0.0199	2	36.980	52.298	0.0326	3	15.713	18.383	0.0338	3	35.817	40.266	0.0218	2	80.800	114.268
749	0.0221	2	219.750	310.773	0.0221	2	17.300	5.515	0.0229	2	91.125	124.599	0.0113	1	229.700	-	0.0214	2	35.410	19.827
750	n.s.	n.s.	n.s.	n.s.	0.0180	2	2.800	3.960	0.1005	9	6.213	9.605	0.0679	6	13.979	28.671	0.0844	8	12.366	21.347
751	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	0.0454	4	1.103	1.497	0.0225	2	4.405	0.191	0.0413	4	3.780	2.765

$$(**) SD = \frac{\sum (x_i - \bar{x})}{n-1}$$

**TABLE 20 (cont).**- Swept area, number of hauls and **black dogfish** mean catch (Kg) and SD (\*\*) by stratum. Spanish Survey on NAFO Div. 3L in the period 2003-2012, on board R/V "Vizconde de Eza". (\*) In 2003, the data correspond to 69% of the total area prospected in 2006-2012.

Stratum	2009				2010				2011				2012			
	Swept area	Tow number	Mean catch	SD	Swept area	Tow number	Mean catch	SD	Swept area	Tow number	Mean catch	SD	Swept area	Tow number	Mean catch	SD
385	0.0225	2	0.000	0.000	0.0225	2	0.000	0.000	0.0229	2	0.000	0.000	0.0225	2	0.000	0.000
387	0.0439	4	0.000	0.000	0.0458	4	0.000	0.000	0.0450	4	0.000	0.000	0.0450	4	0.000	0.000
388	0.0555	5	0.000	0.000	0.0570	5	0.000	0.000	0.0563	5	0.000	0.000	0.0570	5	0.000	0.000
389	0.0803	7	0.000	0.000	0.0795	7	0.000	0.000	0.0675	6	0.000	0.000	0.0799	7	0.000	0.000
390	0.1373	12	0.000	0.000	0.1249	11	0.000	0.000	0.1009	9	0.000	0.000	0.1354	12	0.000	0.000
391	0.0458	4	0.000	0.000	0.0454	4	0.000	0.000	0.0458	4	0.000	0.000	0.0458	4	0.000	0.000
392	0.0229	2	0.000	0.000	0.0225	2	0.000	0.000	0.0229	2	0.000	0.000	0.0225	2	0.000	0.000
729	0.0341	3	0.000	0.000	0.0338	3	0.000	0.000	0.0338	3	0.000	0.000	0.0338	3	0.000	0.000
730	0.0338	3	30.959	51.654	0.0334	3	19.640	25.019	0.0334	3	3.646	6.315	0.0338	3	10.040	17.053
731	0.0341	3	0.000	0.000	0.0338	3	0.000	0.000	0.0334	3	0.000	0.000	0.0341	3	0.000	0.000
732	0.0450	4	0.000	0.000	0.0450	4	0.300	0.600	0.0454	4	0.000	0.000	0.0454	4	0.000	0.000
733	0.0450	4	0.000	0.000	0.0450	4	0.000	0.000	0.0454	4	0.000	0.000	0.0454	4	0.000	0.000
734	0.0218	2	0.000	0.000	0.0225	2	0.000	0.000	0.0225	2	0.000	0.000	0.0233	2	0.000	0.000
741	0.0221	2	0.000	0.000	0.0225	2	0.000	0.000	0.0218	2	0.000	0.000	0.0218	2	0.000	0.000
742	0.0214	2	0.000	0.000	0.0225	2	0.000	0.000	0.0225	2	0.000	0.000	0.0206	2	0.000	0.000
743	0.0203	2	1.835	1.082	0.0225	2	0.000	0.000	0.0221	2	0.000	0.000	0.0206	2	0.000	0.000
744	0.0210	2	0.430	0.608	0.0229	2	0.000	0.000	0.0221	2	0.612	0.865	0.0221	2	0.000	0.000
745	0.0559	5	0.000	0.000	0.0563	5	0.000	0.000	0.0446	4	0.705	1.410	0.0570	5	0.000	0.000
746	0.0668	6	3.939	5.074	0.0679	6	4.817	2.936	0.0566	5	7.160	9.335	0.0675	6	6.004	4.804
747	0.1118	10	6.653	4.933	0.1125	10	5.965	5.925	0.0893	8	5.204	3.122	0.1121	10	4.889	4.861
748	0.0229	2	12.240	17.310	0.0225	2	83.545	40.807	0.0221	2	135.930	187.058	0.0225	2	25.190	35.624
749	0.0225	2	131.090	156.143	0.0229	2	148.715	196.837	0.0221	2	114.000	69.141	0.0221	2	70.633	84.905
750	0.0791	7	9.146	7.225	0.0900	8	0.848	1.376	0.0668	6	1.711	2.351	0.0885	8	4.283	6.729
751	0.0338	3	5.343	4.636	0.0225	2	1.870	1.414	0.0334	3	3.076	2.976	0.0218	2	9.550	5.388

$$(**) SD = \sqrt{\frac{\sum(x_i - \bar{x})^2}{n-1}}$$

**TABLE 21.-** Stratified mean catches (Kg) of **black dogfish** by stratum and year (2003-2012) and SD. Research Vessel *Vizconde de Eza*. n.s. means stratum not surveyed.  
In 2003: the data correspond to 69% of the total area prospected in 2006-2012.

Stratum	Survey									
	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
385	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00
387	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00
388	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00
389	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00
390	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00
391	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00
392	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00
729	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00
730	0.00	369.75	-	627.30	3312.88	4652.33	5262.97	3338.80	619.82	1706.80
731	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00
732	0.00	0.00	-	0.00	0.00	0.00	0.00	69.30	0.00	0.00
733	n.s.	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00
734	n.s.	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00
741	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00
742	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00
743	n.s.	31.90	-	0.00	0.00	0.00	93.59	0.00	0.00	0.00
744	n.s.	0.00	-	47.85	109.73	58.08	28.38	0.00	40.39	0.00
745	2.32	0.00	-	0.00	0.00	0.00	0.00	0.00	245.34	0.00
746	0.00	0.00	-	3541.07	3594.84	2407.60	1544.22	1888.13	2806.72	2353.63
747	n.s.	2944.27	-	2646.94	4354.53	4267.26	4816.77	4318.66	3767.42	3539.64
748	0.00	5879.82	-	2498.42	5694.85	12847.20	1946.16	13283.66	21612.87	4005.21
749	27688.50	2179.80	-	11481.75	28942.20	4461.66	16517.34	18738.09	14364.00	8899.76
750	n.s.	1556.80	-	3454.61	7772.42	6875.64	5085.02	471.21	951.50	2381.07
751	n.s.	n.s.	-	252.47	1008.75	865.62	1223.62	428.23	704.48	2186.95
TOTAL	27690.82	12962.34	-	24550.42	54790.18	36435.38	36518.07	42536.08	45112.55	25073.06
SD	6.19	2.07	-	3.78	8.45	5.62	5.63	6.56	6.95	3.87
	6.19	1.01	-	1.78	1.28	2.23	2.33	2.83	3.39	1.38

**TABLE 22.-** Survey estimates (by the swept area method) of **black dogfish** biomass (t.) by stratum and year and their SD on NAFO Div. 3L (R/V *Vizconde de Eza*). n.s. means stratum not surveyed. In 2003, the data correspond to 69% of the total area prospected in 2006-2012.


Stratum	Survey									
	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
385	0	0	-	0	0	0	0	0	0	0
387	0	0	-	0	0	0	0	0	0	0
388	0	0	-	0	0	0	0	0	0	0
389	0	0	-	0	0	0	0	0	0	0
390	0	0	-	0	0	0	0	0	0	0
391	0	0	-	0	0	0	0	0	0	0
392	0	0	-	0	0	0	0	0	0	0
729	0	0	-	0	0	0	0	0	0	0
730	0	33	-	58	294	433	468	300	56	152
731	0	0	-	0	0	0	0	0	0	0
732	0	0	-	0	0	0	0	6	0	0
733	n.s.	0	-	0	0	0	0	0	0	0
734	n.s.	0	-	0	0	0	0	0	0	0
741	0	0	-	0	0	0	0	0	0	0
742	0	0	-	0	0	0	0	0	0	0
743	n.s.	3	-	0	0	0	9	0	0	0
744	n.s.	0	-	4	10	5	3	0	4	0
745	0	0	-	0	0	0	0	0	22	0
746	0	0	-	315	325	227	139	167	248	209
747	n.s.	287	-	237	387	399	431	384	338	316
748	0	592	-	230	506	1181	170	1181	1954	356
749	2503	197	-	1004	2573	417	1468	1638	1298	804
750	n.s.	173	-	309	687	652	450	42	86	215
751	n.s.	n.s.	-	22	90	84	109	38	63	201
TOTAL	2503	1286	-	2179	4872	3399	3247	3756	4068	2253
SD	2546	695	-	994	721	1296	1340	1634	1964	819

**TABLE 23.- Black dogfish** length distribution per haul mean catches by sex and year. Number per stratified mean catches. Spanish Summer Survey on NAFO 3L: 2003-2012 (R/V *Vizconde de Eza*). Indet. means indeterminate.


Length (cm.)	2004				2006				2007				2008			
	M	F	I	T	M	F	I	T	M	F	I	T	M	F	I	T
16	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
18	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
20	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
22	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
24	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
26	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
28	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
30	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
32	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
34	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
36	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
38	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
40	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.01	0.00	0.00	0.01	0.01	0.00	0.00	0.01
42	0.01	0.00	0.00	0.01	0.01	0.01	0.00	0.02	0.00	0.03	0.00	0.03	0.02	0.00	0.00	0.02
44	0.00	0.00	0.00	0.00	0.02	0.03	0.00	0.05	0.02	0.00	0.00	0.02	0.01	0.04	0.00	0.05
46	0.00	0.00	0.00	0.00	0.02	0.02	0.00	0.04	0.01	0.04	0.00	0.04	0.04	0.06	0.00	0.09
48	0.01	0.02	0.00	0.04	0.05	0.02	0.00	0.06	0.01	0.02	0.00	0.03	0.03	0.01	0.00	0.04
50	0.03	0.06	0.00	0.08	0.00	0.03	0.00	0.03	0.03	0.09	0.00	0.12	0.07	0.03	0.00	0.10
52	0.03	0.06	0.00	0.10	0.03	0.06	0.00	0.10	0.05	0.06	0.00	0.11	0.09	0.08	0.00	0.17
54	0.06	0.05	0.00	0.11	0.04	0.06	0.00	0.09	0.11	0.18	0.00	0.28	0.18	0.10	0.00	0.28
56	0.02	0.12	0.00	0.14	0.04	0.06	0.00	0.11	0.11	0.14	0.00	0.25	0.19	0.12	0.00	0.30
58	0.13	0.06	0.00	0.19	0.08	0.12	0.00	0.20	0.28	0.36	0.00	0.64	0.28	0.15	0.00	0.43
60	0.11	0.19	0.00	0.30	0.15	0.15	0.00	0.29	0.45	0.22	0.00	0.68	0.55	0.16	0.00	0.71
62	0.08	0.15	0.00	0.23	0.11	0.23	0.00	0.35	0.65	0.45	0.00	1.10	0.63	0.12	0.00	0.75
64	0.09	0.00	0.00	0.10	0.17	0.19	0.00	0.35	0.38	0.39	0.00	0.77	0.58	0.13	0.00	0.72
66	0.13	0.10	0.00	0.23	0.14	0.18	0.00	0.32	0.23	0.29	0.00	0.51	0.17	0.17	0.00	0.34
68	0.01	0.07	0.00	0.08	0.07	0.14	0.00	0.21	0.13	0.25	0.00	0.38	0.08	0.10	0.00	0.18
70	0.04	0.07	0.00	0.11	0.01	0.15	0.00	0.16	0.05	0.24	0.00	0.29	0.01	0.12	0.00	0.13
72	0.01	0.00	0.00	0.01	0.01	0.15	0.00	0.16	0.00	0.24	0.00	0.24	0.02	0.02	0.00	0.04
74	0.00	0.03	0.00	0.03	0.00	0.11	0.00	0.11	0.00	0.21	0.00	0.21	0.00	0.08	0.00	0.08
76	0.00	0.00	0.00	0.00	0.00	0.03	0.00	0.03	0.00	0.10	0.00	0.10	0.00	0.07	0.00	0.07
78	0.00	0.00	0.00	0.00	0.00	0.02	0.00	0.02	0.00	0.06	0.00	0.06	0.00	0.00	0.00	0.00
80	0.00	0.00	0.00	0.00	0.00	0.02	0.00	0.02	0.00	0.04	0.00	0.04	0.00	0.00	0.00	0.00
82	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01
84	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01
86	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
88	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Total	0.77	0.99	0.00	1.77	0.94	1.77	0.00	2.71	2.51	3.41	0.00	5.92	2.95	1.59	0.00	4.53
Nº samples:				8				28				28				30
Nº Ind.:	58	55	0	113	99	184	0	283	179	245	0	424	269	152	0	421
Sampled catch:				127				397				593				526
Range:				17-75				41-84				41-81				41-85
Total catch:				132				397				593				526
Total valid hauls:				50				100				94				100

**TABLE 23 (cont).- Black dogfish** length distribution per haul mean catches by sex and year. Number per stratified mean catches. Spanish Summer Survey on NAFO 3L: 2003-2012 (R/V *Vizconde de Eza*). Indet. means indeterminate.


Length (cm.)	2009				2010				2011				2012			
	M	F	I	T	M	F	I	T	M	F	I	T	M	F	I	T
16	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
18	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
20	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
22	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
24	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
26	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
28	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
30	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
32	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
34	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
36	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.01	0.00	0.01	0.00	0.01	0.01	0.00	0.00	0.00
38	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.03	0.00	0.03	0.01	0.00	0.00	0.01
40	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	0.01	0.05	0.00	0.06	0.00	0.05	0.00	0.05
42	0.01	0.04	0.00	0.05	0.06	0.04	0.00	0.09	0.09	0.05	0.00	0.14	0.01	0.07	0.00	0.08
44	0.04	0.03	0.00	0.07	0.05	0.09	0.00	0.13	0.08	0.13	0.00	0.20	0.06	0.07	0.00	0.13
46	0.01	0.02	0.00	0.03	0.08	0.11	0.00	0.19	0.10	0.10	0.00	0.20	0.07	0.06	0.00	0.13
48	0.04	0.02	0.00	0.06	0.05	0.07	0.00	0.12	0.18	0.15	0.00	0.33	0.01	0.03	0.00	0.04
50	0.03	0.08	0.00	0.11	0.06	0.06	0.00	0.12	0.14	0.11	0.00	0.25	0.09	0.04	0.00	0.12
52	0.11	0.10	0.00	0.21	0.12	0.09	0.00	0.21	0.12	0.08	0.00	0.20	0.07	0.08	0.00	0.16
54	0.13	0.10	0.00	0.23	0.09	0.10	0.00	0.19	0.07	0.10	0.00	0.17	0.06	0.04	0.00	0.10
56	0.18	0.15	0.00	0.33	0.13	0.14	0.00	0.27	0.23	0.23	0.00	0.47	0.13	0.08	0.00	0.21
58	0.19	0.17	0.00	0.37	0.24	0.11	0.00	0.36	0.38	0.25	0.00	0.64	0.12	0.10	0.00	0.22
60	0.28	0.20	0.00	0.49	0.29	0.21	0.00	0.51	0.41	0.41	0.00	0.82	0.20	0.14	0.00	0.33
62	0.29	0.19	0.00	0.48	0.30	0.20	0.00	0.50	0.37	0.52	0.00	0.89	0.30	0.18	0.00	0.49
64	0.18	0.20	0.00	0.38	0.17	0.14	0.00	0.31	0.22	0.36	0.00	0.58	0.27	0.07	0.00	0.34
66	0.04	0.31	0.00	0.35	0.12	0.17	0.00	0.30	0.14	0.30	0.00	0.44	0.08	0.18	0.00	0.26
68	0.05	0.19	0.00	0.25	0.03	0.16	0.00	0.19	0.03	0.20	0.00	0.23	0.04	0.12	0.00	0.16
70	0.00	0.22	0.00	0.22	0.03	0.19	0.00	0.22	0.01	0.12	0.00	0.13	0.02	0.09	0.00	0.11
72	0.00	0.17	0.00	0.17	0.00	0.31	0.00	0.31	0.01	0.13	0.00	0.14	0.00	0.12	0.00	0.12
74	0.00	0.14	0.00	0.14	0.00	0.28	0.00	0.28	0.00	0.08	0.00	0.08	0.00	0.04	0.00	0.04
76	0.00	0.08	0.00	0.08	0.00	0.11	0.00	0.11	0.00	0.03	0.00	0.03	0.00	0.01	0.00	0.01
78	0.00	0.00	0.00	0.00	0.00	0.10	0.00	0.10	0.00	0.02	0.00	0.02	0.00	0.01	0.00	0.01
80	0.00	0.01	0.00	0.01	0.00	0.04	0.00	0.04	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01
82	0.00	0.00	0.00	0.00	0.00	0.02	0.00	0.02	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
84	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
86	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
88	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Total	1.58	2.46	0.00	4.04	1.84	2.73	0.00	4.57	2.58	3.47	0.00	6.05	1.55	1.62	0.00	3.16
Nº samples:				32				26				22				24
Nº Ind.:	157	234	0	391	172	275	0	447	214	301	0	515	150	137	0	287
Sampled catch:				554				624				612				360
Range:				41-89				37-87				36-78				39-80
Total catch:				554				624				612				360
Total valid hauls:				98				97				89				98




**FIGURE 1.-** Atlantic cod stratified mean catches in Kg and  $\pm$ SD by year. Spanish surveys in NAFO Division 3L: 2003 - 2012 (R/V “Vizconde de Eza”). In 2003, the data correspond to 69% of the total area prospected in 2006-2012.



**FIGURE 2.-** Atlantic cod abundance ('000), biomass in tonnes and  $\pm$ SD by year. Spanish surveys in NAFO Division 3L: 2003 - 2012 (R/V “Vizconde de Eza”). In 2003, the data correspond to 69% of the total area prospected in 2006-2012.



**FIGURE 3.-** Distribution of the catches per haul for **Atlantic cod**, **Roughhead grenadier**, **redfish**, **thorny skate** and **black dogfish** in 2012 Spanish 3L survey.



**FIGURE 4.-** Atlantic cod length distribution (cm) in NAFO 3L: 2003-2012. Number per stratified mean catches. In 2003, the data correspond to 69% of the total area prospected in 2006-2012.







FIGURE 5.- Atlantic cod, roughhead grenadier and redfish length distribution (cm) in NAFO 3L: 2003-2012.




**FIGURE 6.-** Roughhead grenadier stratified mean catches in Kg and  $\pm$ SD by year. Spanish surveys in NAFO Division 3L: 2003 - 2012 (R/V “Vizconde de Eza”). In 2003, the data correspond to 69% of the total area prospected in 2006-2012.




**FIGURE 7.-** Roughhead grenadier abundance ('000), biomass in tonnes and  $\pm$ SD by year. Spanish surveys in NAFO Division 3L: 2003 - 2012 (R/V “Vizconde de Eza”). In 2003, the data correspond to 69% of the total area prospected in 2006-2012.




**FIGURE 8.- Roughhead grenadier** length distribution (cm) in NAFO 3L: 2003-2012. Number per stratified mean catches. In 2003, the data correspond to 69% of the total area prospected in 2006-2012.




**FIGURE 9.-** Redfish stratified mean catches in Kg and  $\pm$ SD by year. Spanish surveys in NAFO Division 3L: 2003 - 2012 (R/V “Vizconde de Eza”). In 2003, the data correspond to 69% of the total area prospected in 2006-2012.




**FIGURE 10.-** Redfish abundance ('000), biomass in tonnes and  $\pm$ SD by year. Spanish surveys in NAFO Division 3L: 2003 - 2012 (R/V “Vizconde de Eza”). In 2003, the data correspond to 69% of the total area prospected in 2006-2012.




**FIGURE 11.- Redfish** length distribution (cm) in NAFO 3L: 2003-2012. Number per stratified mean catches. In 2003, the data correspond to 69% of the total area prospected in 2006-2012.



**FIGURE 12.-** Thorny skate stratified mean catches in Kg and  $\pm$ SD by year. Spanish surveys in NAFO Division 3L: 2003 - 2012 (R/V “*Vizconde de Eza*”). In 2003, the data correspond to 69% of the total area prospected in 2006-2012.



**FIGURE 13.-** Thorny skate abundance ('000), biomass in tonnes and  $\pm$ SD by year. Spanish surveys in NAFO Division 3L: 2003 - 2012 (R/V “*Vizconde de Eza*”). In 2003, the data correspond to 69% of the total area prospected in 2006-2012.



**FIGURE 14.-** Thorny skate length distribution (cm) in NAFO 3L: 2003-2012. Number per stratified mean catches. In 2003, the data correspond to 69% of the total area prospected in 2006-2012.







FIGURE 15.- Thorny skate and black dogfish length distribution (cm) in NAFO 3L: 2003-2012.



**FIGURE 16.- Black dogfish** stratified mean catches in Kg and  $\pm$ SD by year. Spanish surveys in NAFO Division 3L: 2003 - 2012 (R/V “Vizconde de Eza”). In 2003, the data correspond to 69% of the total area prospected in 2006-2012.



**FIGURE 17.- Black dogfish** abundance ('000), biomass in tonnes and  $\pm$ SD by year. Spanish surveys in NAFO Division 3L: 2003 - 2012 (R/V “Vizconde de Eza”). In 2003, the data correspond to 69% of the total area prospected in 2006-2012.



**FIGURE 18.-** Black dogfish length distribution (cm) in NAFO 3L: 2003-2012. Number per stratified mean catches. In 2003, the data correspond to 69% of the total area prospected in 2006-2012.