

NOT TO BE CITED WITHOUT PRIOR  
REFERENCE TO THE AUTHOR(S)

Northwest Atlantic



Fisheries Organization

Serial No. N6349

NAFO SCR Doc. 14/047

### SCIENTIFIC COUNCIL MEETING – SEPTEMBER 2014

Northern Shrimp (*Pandalus borealis*, Krøyer) from EU-Spain Bottom Trawl  
Survey 2014 in NAFO Div. 3LNO

by

Casas, J.M., E. Román, J. Teruel, and M. Álvarez  
Instituto Español de Oceanografía, Aptdo. 1552. 36200 - Vigo. Spain.

#### Abstract

The Spanish Institute of Oceanography carried out in 2014 two bottom trawl surveys in the NAFO Regulatory Area in Division 3NO and 3L during the months of June and August respectively. The results on Northern shrimp (*Pandalus borealis*) are presented and compared with those from previous surveys from the same series. As recent years in 2014 the shrimp catch (0.84 kg.) and estimated biomass (2.97 t.) in Divisions 3NO remain between the lowest of the series, confirming the decrease of shrimp importance from 2004. In a similar way, northern shrimp indices estimated from 2014 in 3L Division declined 40% with respect to 2013 and confirm the downward trend initiated in 2009. The biomass estimated in 2014 (10846 t.) was the second lowest in the historical series.

#### Introduction

Northern shrimp (*Pandalus borealis* Krøyer, 1883) is a protrandric, circumpolar species, discontinuously distributed in the North Atlantic and of considerable commercial importance. The greatest abundance is being in the Northwest Atlantic at latitudes above 46°N. The stock of this species in Div. 3LNO, NAFO is distributed along the entire edge of the grand banks, at depths generally ranging from 185 to 550 metres, although historically at least 92.7% of the 3LNO shrimp biomass had been found within Division 3L. The proportion of biomass in 3LNO within the NAFO Regulatory Area (NRA), over the period 1996 – 2012, accounted for between 6.3 and 32.6% (Orr and Sullivan, 2013).

Since 1995, Canadian multi-species stratified random surveys have been used to estimate northern shrimp biomass and abundance indices within NAFO Div. 3LNO. In this series of surveys, Div. 3N accounts for between 0.2 and 8.1% of the total 3LNO biomass. Between 0 and 100% of the 3N biomass was located outside the 200 Nmi limit. The biomass in Division 3O accounts for less than 1% of the biomass in Div. 3LNO and only a negligible amount of the biomass in Div. 3O is beyond the 200 mile limit (Orr and Sullivan, 2013).

The Oceanographic Spanish Institute (IEO) is conducting research cruises since 1995 in the NAFO Regulatory Area in Div. 3NO beyond Canada's EEZ. A stratified, random, bottom trawl, multi-species research sampling program was carried out to obtain abundance and biomass indices as well as other biological data for the most important commercial species present in the area. In the surveys conducted between 1995 and 2000, the catches of northern shrimp were insignificant. This could be explained by the low efficiency of the fishing gear "pedreira", with this species (Paz *et al.*, 1995), used in those years.

Since 2001, the survey was carried out on board R/V “*Vizconde de Eza*” using a Campelen 1800 net (Walsh *et al.*, 2001). Despite the improvements incorporated with the new vessel and the use of a Campelen 1800 net, which is highly efficient for this species (Vazquez, 2002), total catches in 2001 were poor, i.e., 29 kg. In the following years a significant increase of the catches of northern shrimp was noted in 3NO Division where catches were higher than 300 kg. Since 2007 the catches have declined to levels next to the lowest in the historical series.

Also, since 2003 a new research survey was conducted in Division 3L as an extension of the survey carried out in 3NO (Román *et al.*, 2008). The estimated biomass in 3L Division always was very superior to that estimated in 3NO. Since 2009 year the catches have declined to low levels staying in the last years between the lowest in the historical series.

This work presents data on the geographical distribution in the NAFO Regulatory Area (Div. 3LNO), on biomass, length frequencies, age structure and length-weight relationship of catches of northern shrimp on EU-Spanish bottom trawl surveys 2014.

### Materials and Methods

In 2014 the EU-Spanish bottom trawl surveys were carried out in June, from 1<sup>st</sup> to 21<sup>st</sup>, in 3NO and from 30<sup>th</sup> July to 19<sup>th</sup> August in 3L following set guidelines previously established for the series of Spanish research surveys (Walsh *et al.*, 2001). These surveys took place in Div. 3NO and 3L, with a total of 122 and 99 valid hauls respectively ranging depths between 40 and 1450 m approximately. This year all strata were surveyed.

Shrimp samples of approximately 1.5 kg were taken to determine length frequencies. In 2014 the residual catches in 3NO Divisions did not allow the length frequencies sampling. In 3L Division males and females were separated with reference to the endopod of the first pleopod (Rasmussen, 1953). Following this criterion, individuals that were in the middle of a sex change were considered as females. The females were differentiated into mature and immature, following the sternal spines criteria (McCray, 1971). Ovigerous females were considered as an independent group not included within the mature females.

Individuals were measured onboard by noting the distance from the base of the eye to the posterior mid dorsal point of the carapace -CL- (Shumway *et al.*, 1985). Such measurements were made to the lower half millimetre using electronic callipers.

Furthermore, in 2014 survey some samples were frozen onboard to determine the length-weight relationship in the laboratory. 4531 individuals were selected in 3L Divisions, dried and weighed with a precision of 0.01g to calculate the length-weight relationship in each Division.

### Results and Discussion

The Table 1 shows the catches, biomass and standard errors estimated by swept area method of northern shrimp from the EU-Spanish multi-species surveys, carried out by IEO Vigo from 1995-2014 in the NAFO Div. 3NO and from 2003-2014 in Division 3L. In the summer of 2005 the research survey could not be carried out in Division 3L. From the year 2002 an abrupt increase with respect to earlier years occurred in 3NO Division, both in terms of catch and biomass (Diaz *et al.*, 2002). These initial data were considered with caution due to the fact that, until 2001, the “Pedreira” gear used as a sampler (Paz *et al.*, 1995) was not efficient for catching shrimp. However, although in 2001, the gear “type Pedreira” was changed for a new type “Campelen 1800” (Walsh *et al.*, 2001) with high efficiency for catching this species (Vazquez, 2002), the catches and biomass estimated stayed at low levels.

From 2002 to 2006, the increase of shrimp catches in 3NO was confirmed, in terms of the period 1995-2001. After that, in the last eight years the catches and estimated biomasses of shrimp have decreased markedly and they are now at levels of the beginning of the series. The estimated biomass in 2014 was around 3 t. (Figure 1).

Unlike 3NO, the estimated biomass in 3L Division showed a general upward trend from 63647 t. in 2003 to 149265 t. in 2008. This trend changed in 2009 with the strong decline of the biomass estimated (74091 t, about 50% with respect to 2008) and since then the biomass decreased up to the historical minimum recorded in 2012 (10784 t.). In 2014 the biomass decreased again by 38% (10846 t.) compared to 2013 and remains between the lowest in the survey series (Figure 1).

The distribution of northern shrimp catches in the EU-Spanish trawl surveys 2014 is shown in Figure 2. As in previous years the catches in 3NO Division were residuals.

The Tables 2 and 3 show the shrimp biomass by depth strata from 1995 to 2014 surveys in 3NO Divisions and from 2003 to 2014 in 3L Division. Although it is considered that the shrimp in Div. 3LNO is distributed along the entire edge of the grand banks, at depths generally ranging from 51 to 300 fathoms (93-550 m.), the depth of the bulk of biomass in 3L Division was generally in depths lower than 200 ft (>95% of the biomass). This general pattern has changed in recent years and the percentage of the estimated biomass in depths lower than 200 ft has decreased up to 44% and 77% of the biomass in 2013 and 2014 respectively. In 3NO the percentage of the estimated biomass in depths lower than 200 ft. varied along the years, showing a deeper distribution in 2004, 2005 and 2011 where the percentage of biomass in depths bigger than 200 ft was higher than 43%.

The length distribution by sex estimated in 3L Division is presented in Table 4 and Figure 3. The main modes were around 20 mm. for males and 24 mm. for females. In 2014 the sex ratio showed a higher percentage of males (67%) in 3L Division. Due to the low catches in 3NO Division the length of shrimp could not be sampled and thus there was not estimated the length distribution.

The MIX modal size analysis programme was used with the length distribution by sex estimated in 3L Divisions (Table 5). From the cited analysis the males presented three modes at 13.17, 17.58 and 20.23 mm. corresponding with ages 2, 3 and 4 respectively. The females showed several modes at 19.92, 23.61 and 25.32 mm (ages 4, 5 and 6 respectively).

The Table 6 shows the length-weight relationship estimated in 2014 surveys by sex and maturity stage as well the parameters of the relationship, number of specimens sampled and determination coefficient  $R^2$ .

### References

- Díaz, P., T. Patrocinio, and X. Paz. 2002. Increased Catches of Northern Shrimp (*Pandalus borealis*, Krøyer) in a 2002 Spanish Bottom Trawl Survey in NAFO Division 3N. *NAFO SCR Doc.*, No. 143. Serial No. N4772, 11p.
- McCray, J.A. 1971. Sternal spines as a characteristic for differentiating between females of some Pandalidae. *J. Fish. Res. Bd. Can.*, **28**: 98-100.
- Orr, D. C. and D. J. Sullivan. 2013. The 2013 assessment of the Northern Shrimp (*Pandalus borealis*, Kroyer) resource in NAFO Divisions 3LNO. *NAFO SCR Doc.*, No. 064. Serial No. N6226, 78 p.
- Paz, X., J. Martínez, and E. De Cárdenas. 1995. Preliminary results from the 95 Spanish bottom trawl survey in the NAFO Regulatory Area for Divisions 3NO. *NAFO SCR Doc.*, No. 55. Serial No. N2568, 10 p.
- Rasmussen, B. 1953. On the geographical variation in growth and sexual development of the Deep Sea Prawn (*Pandalus borealis*, Kr.) . *Norweg. Fish. And Mar. invest. Rep.*, **10** (3):1-160.
- Román, E., C. González-Iglesias, Á. Armesto, and D. González-Troncoso. 2008. Results for the Spanish Survey in the NAFO Regulatory Area of Division 3L for the period 2003-2007. *NAFO SCR Doc.*, No. 20. Serial No. N5514, 25p.
- Shumway, S.E., H. C. Perkins, D. F. Schick, and A. P. Stikney. 1985. Synopsis of biological data on the Pink Shrimp (*Pandalus borealis*, Krøyer, 1838). *NOAA Techn. Rep. NMFS* **30**, 57 p.
- Vázquez, A. 2002. Catchability comparison between Lofoten and Campelen gears. *NAFO SCR. Doc.*, No.74. Serial No. N4688, 7p.
- Walsh, S. J., X. Paz, and P. Durán. 2001. A preliminary investigation of the efficiency of Canadian and Spanish survey bottom trawls on the southern Grand Bank. *NAFO SCR. Doc.*, No. 74. Serial No. N4453, 18 p.

**Table 1.-** Northern shrimp biomass estimated by swept area (t), standard error and catches (kg) from EU-Spanish bottom trawl surveys in NAFO Div. 3NO, 1995-2014 and 3L 2003-2014.

<b>3NO</b>			
Year	Biomass		Catch (kg)
	tons	Std. err.	
1995 <sup>1</sup>	14	13	5
1996 <sup>1</sup>	18	17	2
1997 <sup>1</sup>	1	1	0
1998 <sup>1</sup>	23	17	5
1999 <sup>1</sup>	81	36	13
2000 <sup>1</sup>	26	9	6
2001 <sup>2</sup>	178	72	29
2002 <sup>2</sup>	2043	814	408
2003 <sup>2</sup>	1618	716	325
2004 <sup>2</sup>	2654	1693	550
2005 <sup>2</sup>	1627	590	368
2006 <sup>2</sup>	1274	352	278
2007 <sup>2</sup>	401	285	71
2008 <sup>2</sup>	144	98	24
2009 <sup>2</sup>	140	111	33
2010 <sup>2</sup>	114	35	21
2011 <sup>2</sup>	37	24	9
2012 <sup>2</sup>	4	3	1
2013 <sup>2</sup>	38	15	9
2014 <sup>2</sup>	3	1	1

<b>3L</b>			
Year	Biomass		Catch (kg)
	tons	Std. err.	
2003 <sup>2</sup>	63647	20105	5836
2004 <sup>2</sup>	94270	40332	5093
2005	Not surveyed		
2006 <sup>2</sup>	125850	12690	17805
2007 <sup>2</sup>	113402	13445	18098
2008 <sup>2</sup>	149265	48490	23720
2009 <sup>2</sup>	74091	37999	12173
2010 <sup>2</sup>	37803	9836	6103
2011 <sup>2</sup>	24346	4449	4092
2012 <sup>2</sup>	10784	3724	1838
2013 <sup>2</sup>	17438	5363	3101
2014 <sup>2</sup>	10846	2764	1860

<sup>1</sup> Pedreira codend 35 mm. mesh size.

<sup>2</sup> Campelen codend 44 mm. mesh size. (inner codend 20mm)

**Table 2.-** Northern shrimp biomass (kg.) by strata from Spanish bottom trawl survey 1995-2014 in NAFO Div. 3NO.

Stratum	Area Mn <sup>2</sup>	Depth range	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
375	271	0-30	0	0		0	0	0	3453	0	25	0	0	1989	0	0	0	0	0	0	0	0
376	1334	0-30	0	0		0	0	0	1270	0	0	0	341	4203	0	0	0	0	34	0	0	0
353	269	31-50	0	0		0	0	0	79	0	48	0	0	0	126	0	16	0	0	0	0	0
360	2783	31-50	0	0		0	0	0	26423	1457	3470	24	0	0	445	0	110	1317	129	0	50	0
374	214	31-50	0	0		0	0	0	178	0	0	0	0	0	62	0	0	0	0	0	0	0
354	246	51-100	0	0		0	0	0	87612	0	292	6917	0	0	14	0	0	55	86	0	292	0
359	421	51-100	0	0		0	1389	0	6348	847	1309	43	41	22	98	42	0	543	47	0	30	28
377	100	51-100	0	0		0	208	44	0	2020	751	1471	3742	3704	83	60	40	0	0	0	0	0
382	343	51-100	0	0		0	213	206		112695	302	297	825	944	191	4131	0	0	0	0	0	0
355	74	101-150		0		0	0	0	15170	147	7635	6146	6183	9179	262	204	0	961	0	148	89	11
358	225	101-150	0	0		0	30129	0	717	3261	3900	10289	32548	258	2357	2902	0	17220	196	0	27	0
378	139	101-150	0	0		8968	10998	1196	17004	680353	11429	772	3985	10066	1357	481	73	192	0	0	0	0
381	144	101-150		0		63	11205	122		84984	20648	225280	1486	75176	303300	114294	466	25403	87	111	41	78
356	47	151-200		0		0	0	0	137	0	1337	12937	8046	2683	213	635	39	409	33	0	0	0
357	164	151-200	0	18097		0	0	0	606	16414	425145	163606	38796	114178	9307	1249	959	14877	29	0	0	144
379	106	151-200	0	0	720	0	135	0	12511	70342	254080	7709	329867	116970	12146	2238	5079	15709	19	28	897	175
380	96	151-200		0		1024	9346	10240		1000960	698502	258603	120866	607392	6488	11379	125767	26518	7269	3483	26188	1086
721	65	201-300		0		0	0	0	2889	3282	1112	852	256	3054	0	257	318	6	6339	11	315	569
723	155	201-300		0		0	16872	0	0	12667	92831	44044	3333	53799	14615	90	0	916	335	0	98	132
725	105	201-300	14315			0	0	0	271	527	91803	1814540	748369	206794	47133	578	239	7745	0	0	216	231
727	96	201-300		0		13213	0	11429		28660	2119	98477	326841	62635	1248	3172	179	632	22656	83	9350	512
722	84	301-400		0		0	37	734	2890	60	156	0	36	0	0	0	0	0	0	0	0	0
724	124	301-400	0	0		0	0	0	0	55	628	58	165	53	213	0	0	0	32	0	0	0
726	72	301-400	0	0		0	0	0	0	7	54	2048	0	406	170	0	5351	146	0	0	0	0
728	78	301-400		0		0	0	1671		7280	0	0	86	135	0	0	41	146	0	0	40	0
752	131	401-500		0		0	0	0		86	0	49	222	58	309	0	143	136	0	0	79	0
756	101	401-500		0		0	0	0	0	0	46	42	869	84	27	84	391	0	0	0	0	0
760	154	401-500		0		0	0	0	0	0	283	49	0	0	590	0	0	0	0	0	0	0
764	100	401-500		0		0	0	0	42	0	0	0	0	0	0	0	0	0	0	0	0	0
753	138	501-600		0		0	0	0		0	0	0	0	166	0	0	0	0	0	0	0	0
757	102	501-600		0		0	0	0		204	0	0	27	0	67	0	0	14	0	0	0	0
761	171	501-600		0		0	0	0	0	0	0	0	0	0	99	0	0	0	0	0	0	0
765	124	501-600		0		0	0	0	0	37	0	0	0	0	0	0	0	0	0	0	0	0
754	180	601-700		0		0	0	0		0	0	0	0	0	0	0	0	96	0	0	0	0
758	99	601-700		0		0	0	94		16302	0	19	88	0	0	0	0	0	0	0	0	0
762	212	601-700		0		0	0	0	0	85	0	0	0	0	0	0	0	0	0	0	0	0
766	144	601-700		0		0	0	0		19	58	0	0	0	0	0	0	32	0	0	0	0
755	385	701-800		0		0	0	89		0	174	0	68	0	0	1839	0	0	0	0	0	0
759	127	701-800		0		0	0	0		17	0	48	0	0	0	0	0	965	0	0	0	0
763	261	701-800		0		0	0	0		0	0	0	0	0	0	0	0	0	0	0	0	0
767	158	701-800		0		0	0	0		0	0	0	0	0	0	0	0	0	0	0	0	0
Biomass (t.)			14	18	1	23	81	26	178	2043	1618	2654	1627	1274	401	144	139	114	37	4	38	3
Std. Error (t.)			13	17	1	17	36	9	72	814	716	1693	590	352	285	98	111	35	24	3	15	1
Biomass % < 200fth			0	100	100	43	79	46	97	97	88	26	34	74	84	96	95	91	21	98	73	51

**Table 3.-** Northern shrimp biomass (kg.) by strata from Spanish bottom trawl survey 2003-2014 in NAFO Div. 3L.

Stratum	Area Mn <sup>2</sup>	Depth range fth.	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
385	118	51-100	420	175		2485867	2416545	8265541	140724	12046	975	4998	31	68
390	815	51-100	1014	3780		2577958	5404325	317330	37466118	145874	2020	49686	414	2340
389	509	101-150	14397492	41654297		53639329	49120205	74404070	25997291	21705956	979731	630153	149429	318135
391	282	101-150	1116135	1299793		3712072	12397477	24948041	28071	120096	11940	99221	3115	16223
387	256	151-200	17618619	21721973		29967360	11782827	14287154	6473372	7874303	15006844	6644446	5206921	3955026
388	357	151-200	25169595	24779540		32585066	26954928	21602795	2348269	5096163	8113071	2136050	1979045	3858773
392	145	151-200	2821419	1866379		193967	1199955	3675300	1564098	1608469	24550	118649	329956	155247
729	186	201-300	20371	1465049		88481	172095	16126	11533	95976	149	2618	11348	2331
731	216	201-300	2449416	1467221		177357	666240	1501056	54100	1083034	2647	799077	2191919	1644180
733	234	201-300		4077		390052	3281339	240647	6718	51397	194095	285343	7544711	833091
730	170	301-400	0	876		1485	76	32	20	581	92	0	36	907
732	231	301-400	34907	5643		14535	4723	1905	226	4266	1349	596	3229	34455
734	153	301-400		408		10554	136	2144	70	129	4910	1553	15628	16075
741	100	401-500	0	56		1379	22	486	0	0	662	189	402	1893
745	348	401-500	17642	0		1699	186	1950	0	2716	1911	250	1613	5068
748	159	401-500	292	696		366	499	66	0	49	108	0	21	83
742	64	501-600	0	0		462	0	0	0	1718	57	11202	9	0
746	392	501-600	0	0		134	0	74	70	225	381	0	395	1068
749	126	501-600	0	23		99	0	0	0	0	11	0	0	140
743	51	601-700		0		1020	0	23	0	0	2	20	0	18
747	724	601-700		0		147	0	41	201	51	32	0	116	753
750	556	601-700		0		58	0	132	295	0	308	0	37	178
744	66	701-800		0		185	0	0	0	0	0	0	0	9
751	229	701-800				0	0	0	0	0	0	0	21	21
Biomasa (t.)			63647	94270		125850	113402	149265	74091	37803	24346	10784	17478	10846
Std. Error (t)			20105	40332		12690	13445	48490	37999	9836	4449	3724	5363	2764
Biomass % < 200 fth			96	97		99	96	99	100	97	99	90	44	77

**Table 4.-** Northern shrimp size distribution ('000) by sex from Spanish bottom trawl survey 2014 in NAFO Div. 3L.

<b>CL (mm)</b>	<b>Males</b>	<b>Females</b>	<b>Total</b>
6	0	0	0
6.5	10	0	10
7	26	0	26
7.5	10	0	10
8	286	0	286
8.5	8	0	8
9	266	0	266
9.5	278	0	278
10	10	0	10
10.5	10	0	10
11	591	0	591
11.5	2706	0	2706
12	4682	266	4947
12.5	4860	0	4860
13	6085	683	6768
13.5	8331	0	8331
14	5917	532	6449
14.5	8138	0	8138
15	8585	0	8585
15.5	20065	328	20393
16	39187	0	39187
16.5	56884	0	56884
17	81176	1429	82605
17.5	93944	1613	95557
18	95112	2208	97320
18.5	96541	4628	101169
19	105067	3763	108830
19.5	118716	7479	126196
20	138481	13582	152062
20.5	105029	6909	111939
21	79110	12431	91542
21.5	65271	19442	84713
22	38070	24630	62700
22.5	23636	42805	66442
23	10562	67420	77981
23.5	6801	64433	71234
24	3665	80817	84481
24.5	0	78813	78813
25	0	58138	58138
25.5	0	47384	47384
26	0	27567	27567
26.5	0	16401	16401
27	0	7955	7955
27.5	0	8385	8385
28	0	1659	1659
29	0	1190	1190
29.5	0	0	0
30	0	0	0
30.5	0	0	0
<b>Total</b>	<b>1228115</b>	<b>602889</b>	<b>1831004</b>
	67%	33%	


**Table 5.** Results of the modal analysis (MIX) by sex and maturity stage Spanish bottom trawl survey 3L 2014.

3L				
Males			Females	
Age	Prop.	St. Dev.	Prop.	St. Dev.
1				
2	0.028	0.000		
3	0.325	0.001		
4	0.647	0.001	0.078	0.000
5			0.532	0.003
6			0.390	0.003
7				
Age	Mean CL	St. Dev.	Mean CL	St. Dev.
1				
2	13.17	0.007		
3	17.58	0.005		
4	20.23	0.004	19.92	0.006
5			23.61	0.005
6			25.32	0.006
7				
Age	Sigma	St. Dev.	Sigma	St. Dev.
1				
2	0.928	Const.CV.		
3	1.238	Const.CV.		
4	1.425	Const.CV.	0.897	Fixed C.V.
5			1.062	Fixed C.V.
6			1.139	Fixed C.V.
7				

**Table 6.** Northern shrimp length-weight relationship by sex, maturity stage and all combined from Spanish bottom trawl survey 2014 in NAFO Div. 3L.

	a	b	R <sup>2</sup>	N
Division 3L				
Males	0.00100	2.84414	0.95100	3394
Inmature females	0.00078	2.93113	0.86546	310
Mature females	0.00163	2.69444	0.80062	552
Ovigerous females	0.00291	2.51926	0.66458	275
All combined	0.00089	2.88513	0.96795	4531





**Figure 1.-** Northern shrimp biomass (tons) and catch (kg) from Spanish research surveys in NAFO Div. 3NO 2001-2014 and 3L 2003-2014.





Figure 2.- Geographic distribution of Northern shrimp catches from Spanish bottom trawls surveys 2014.



**Figure 3.-** Northern shrimp size distribution, by sex from Spanish bottom trawl survey (2014) in Div. 3L.