

NOT TO BE CITED WITHOUT PRIOR
REFERENCE TO THE AUTHOR(S)

Northwest Atlantic



Fisheries Organization

Serial No. N6437

NAFO SCR Doc. 15/016

SCIENTIFIC COUNCIL MEETING – JUNE 2015

Biomass and Abundance of Demersal Fish Stocks off West and East Greenland estimated
from the Greenland Institute of Natural resources Shrimp Fish Survey, 1988-2014.

By

Rasmus Nygaard
Greenland Institute of Natural Resources, Box 570, 3900 Nuuk, Greenland

And

Ole A. Jørgensen
DTU-Aqua, Technical University of Denmark
Charlottenlund Slot, 2920 Charlottenlund, Denmark

Abstract

Since 1988, the Greenland Institute of Natural Resources has annually conducted a bottom trawl survey off West Greenland. The survey was initially designed as a shrimp survey with the focus to evaluate the biomass and abundance of the Northern shrimp (*Pandalus borealis*). The survey has been continuously developed during the years particularly reflecting the needs of the shrimp assessments. Fish catches have systematically been recorded since 1992. The gear was changed prior to the 2005 survey. The East Greenland area has been surveyed since 2008. This paper contains; Biomass and abundance indices for West Greenland by NAFO subarea and East Greenland. The paper also contain length frequencies from West Greenland for Greenland halibut (*Reinhardtius hippoglossoides*), Atlantic cod (*Gadus morhua*), redfish (*Sebastes norvegicus* and *Sebastes mentella*), Atlantic wolffish (*Anarhichas lupus*), Spotted wolffish (*Anarhichas minor*), American plaice (*Hippoglossoides platessoides*) and Thorny skate (*Amblyraja radiata*). The paper also contains maps of tow biomass and abundance densities. Biomass and abundance estimates for Elasmobranchs, Teleosts, Cephalopods and crustaceans excl. Shrimp for the West-Greenland part of the GINR shrimp fish survey including the West-Greenlandic Shelf part of NAFO div 0A. In 2014, Greenland halibut abundance decreased to the second lowest level observed during the past decade mainly driven by poor recruitment of age 1 and biomass index decreased to the lowest level observed during the past decade. In West Greenland the cod abundance index was about the same level as in 2013, but the biomass index increased to the highest level observed in the time series, mainly caused a single large haul of cod in division 1C providing about 55% of the total abundance estimate and 45% of the total West Greenland biomass estimate. The offshore cod stock in East Greenland decreased compared to 2013, but is the second highest in the survey time series. In West Greenland the Golden redfish and deep-sea redfish biomass have increased in the past few years, but the abundance of redfish, mainly juvenile, has decreased about a factor 20 since the beginning of the time series and the 2014 abundance is the lowest estimate in the time series indicating continuous failing recruitment in the area. In East Greenland the biomass of deep-sea redfish was estimated at 68 Kt, which is the lowest observed since 2008 and the biomass of Golden redfish decreased to 138 Kt. Juvenile redfish less than 30 cm has virtually been absent since 2013 in this area. Spotted wolffish abundance and biomass indices have increased throughout the time series in West Greenland and the 2014 biomass estimate is among the highest observed.

Materials and Methods

The Greenland Shrimp and Fish survey in West Greenland (SFW survey)

The survey has throughout the time series been conducted with the 722 GRT stern trawler M/Tr 'Pâmiut'. The survey design, the area coverage and the trawl and its rigging has been unchanged since 2005. The years prior to 2005 experienced a number of survey developments that are detailed below. The survey period is June -July in West Greenland and August in East Greenland.

The Survey Gear and trawl: The survey initially used a Skjervoy 3000/20 trawl with steel bobbin gear and double bag. In 2005 the skjervoy trawl was replaced by a "Cosmos" trouser trawl (Wieland and Bergström, 2005). Until 2003, *Greenland Perfect* trawldoors were used (9.25 m^2 , 2.4 tons), but they were replaced in 2004 by Injector International trawl doors (7.5 m^2 , 2.8 tons) to facilitate the trawl change in the year after. Calibration experiments with the two trawls were conducted in the main shrimp areas in 2004 and 2005 and a formal analysis of conversion factors were established for shrimp (Rosing and Wieland, 2005). However the calibration factors for the different fish species were never finally evaluated. Preliminary conversion factors were derived as described by Rosing and Wieland (2005) and are given in table 3.

Survey area and stratification: The trawl survey initially covered the traditional offshore shrimp area, between 60° - 72° north, depth 150-600m. In 1991 the area was extended to include the Disko Bay. The area is delimited by a line 3nm off the base line and the 600 m depth curve. Areas shallower than 150 m was initially rather unsystematically covered but from 2004 two extra depth zones have been formally included (50-100m, and 100-150m).

The stratification is based on designated 'Shrimp Areas' that is divided into depth zones of: 151-200, 201-300, 301-400 and 401-600 m, as based on depth contour lines (figure. 1). The depth zones 0-100 m and 100-150m are delimited by the NAFO Subdivision boundaries. The "shrimp Areas" and their sizes are provided in table 1. The number of valid hauls by year and stratum is listed in table 2. If strata had no stations in a given year, the neighbouring strata in that year are geographically enhanced with area of the non-visited strata.

Fish species was prior to 2007 analysed using a re-stratification that followed the NAFO divisions. Re-stratification implies a potential bias and the survey information from 2005 and onwards has therefore been reanalysed in accordance with the shrimp strata actually used in the survey. Tow duration has over the years been gradually reduced from 60 min. (prior to 1997) to 30 and has been fixed to 15 min since 2005 (Wieland and Storr-Paulsen 2006). Towing speed has been about 2.5 knots throughout the years. Survey abundance and biomass is expressed per swept area: Wingspread*towed distance, where wingspread is inferred from Scanmar recordings and the towed distance is measured by GPS.

Allocation of trawl hauls. Trawl stations are allocated to strata with the objective to minimise the variances of the shrimp biomass. The allocation algorithm utilises the historically observed shrimp variances where highest weight is placed on the most recent information. Stations positions were initially selected at random but since 1999 station positions were chosen to secure a minimum distance between stations. Since 1998 about half of the haul positions were randomly selected from the previous year hauls; the rest of the hauls being selected at random.

The Greenland Shrimp and Fish survey in East Greenland (SFE survey)

The survey is carried out with the same gear and survey protocols as used in West Greenland after 2005. Stratification is based on the "Q-areas" used for the East Greenland survey for Greenland halibut. The areas are further depth stratified into 0-200 m, 200-400m and 400-600 m zones, the areas are shown in figure. 1 and the area sizes are given in table 1. The major difference between West and East Greenland is the bottom conditions that severely restrict the areas that can be trawled off East Greenland. Stations were randomly selected from historical known trawl-able sites.

Results

Greenland halibut (*Reinhardtius hippoglossoides*)

In West Greenland, Greenland halibut can be found in all divisions, but is most abundant in the important nursery areas in division 1A and 1BN and Disko Bay (1AX) (table 4 and5). The abundance index is mainly driven with the year to year variability in 1 year old recruits Greenland halibut abundance decreased to the

second lowest level observed during the past decade mainly driven by poor recruitment of age 1 and biomass index decreased to the lowest level observed during the past decade (figure 2). Clear modes can be found in the length distribution at 15 and 23 cm every year probably corresponding to year-classes 1 and 2 (figure 14). Distribution of survey catches in number pr. km² and kg pr. km² are given in figure 21.

In East-Greenland, recruits are rarely seen and both abundance and biomass indices are much lower on the shallow parts of the East Greenlandic shelf areas.

Greenland halibut recruitment.

A recruitment index was estimated for the entire survey area. By means of the Petersen-method ages 1, 2 and 3+ were separated in the survey catches. The number of one-year-old fish in the total survey area including Disko Bay increased gradually from 1996 to a peak of 500 million in 2001 (Fig. 9). During the 00' the recruitment was stable at around 300 mill. The number of one-year old peaked in 2011 with 530 million, which is the highest in the time series. The recruitment decreased in 2012 where the 2011 year-class was estimated to 175 mill., the lowest estimate since 1996 and at the level of the early 90's. The recruitment increased again in 2013 where the 2012 year-class was estimated at 444 mill. to decrease again in 2014 where the 2013 year-class was estimated to 180 mill. (Fig. 10a).

To allow comparison of abundance throughout the time series, the 2005 to 2014 catches were divided by a conversion factors to adjust the new Cosmos trawl catches to the old Skjervoy trawl catches. For Greenland halibut the conversion were length dependent and x in the equations is the individual fish length (Table 3).

The offshore recruitment has been rather stable between 2003 and 2010. The recruitment increased to the highest level in the time series in 2011 to decrease to lowest level seen since 1997 (1996 year-class) in 2012. The offshore recruitment (2012 year-class) increased again in 2013 to the second largest in the time series to decrease again to 130 mill. In 2014 78% of the one year old fish was found in the off shore areas (Fig. 11). The decrease in recruitment between 2013 and 2014 was seen in all divisions (Fig.12).

In Disko Bay the recruitment has been decreasing between 2003 and 2008 but increased since then to the highest level seen since 2001 in 2011. In 2012 the recruitment decreased to the lowest level seen since 2008 to increase again in 2013, but not as significantly as in the of shore areas. The recruitment decreased again in 2014 to the lowest estimate since 2008 (Fig. 12).

Generally there is a steep decline between abundance at age 1 and age 2 and 3+ which also was observed in the 2014 survey (Fig. 12). Further, it has been noted, that the year-classes estimated to be a very strong year-class at age 1 have not shown up as a particularly strong year-classes at age 5-8 in the fishery catches or in the 1CD survey for Greenland halibut.

Cod (*Gadus morhua*)

In West Greenland the cod abundance index was about the same level as in 2013, but the biomass index increased to the highest level observed in the time series, mainly caused a single large haul of cod in division 1C providing about 55% of the total abundance estimate and 45% of the total West Greenland biomass estimate. The survey was dominated by the 2010 YC, which was primarily caught in the large haul, followed by the 2009 YC which also dominated the survey in 2013, 2012 and 2011. The 2010 YC was the second most abundant YC in 2012 and 2013. The 2009 YC was mainly distributed in North-West Greenland (NAFO 1B) in 2010 and 2011, but moved further south (NAFO 1D-1E) in 2012 and was again mainly observed in this area in 2013 and 2014. The 2010 YC showed the same distribution pattern as the 2009 YC with main distribution area as 1 and 2 yr old in the northern part of the survey area, and at age 3 moving further to the south. Cod older than 6 years are almost absent in West Greenland. For further information on cod see the ICES Report of the North-Western Working Group (Anon., 2014)

Demersal Redfish (*Sebastes sp.*) combined.

Two species of redfish are common in the area, golden redfish, *Sebastes norvegicus* and deep-sea redfish *Sebastes mentella*. Due to difficulties in identification of species in some years redfish were classified as *Sebastes sp.* (prior to 2007). Redfish less than approximately 20 cm are classified as juvenile "*Sebastes sp.*". The distinction is however not straight forward and often small redfish are classified as *Sebastes Mentella*, which in West Greenland has also dominated individuals less than 30 cm in recent years.

Redfish are present in all divisions, but recruits are most abundant in division 1B (table 8, 9 and figure 16). Golden redfish and deep-sea redfish biomass has increased in West Greenland in the most recent years, but the abundance of redfish, mainly juvenile, has decreased about a factor 20 since the beginning of the time series and the 2014 abundance is the lowest estimate in the time series indicating continuous failing recruitment in the area (figure 4), mainly caused by the disappearance of juvenile redfish. Biomass of demersal species of redfish combined decreased to about a third the initial values throughout the 1990's, but stabilized during the 2000's and has been increasing since 2009 (table 9 and figure 4). Annual growth increments of 4 cm were indicated by repeatedly pronounced peaks in length compositions at 7-8 cm and 12 cm probably corresponding to age 1 and 2 (Nedreaas, 1990). Densities by haul in number pr. km² and kg pr. km² are given in figure 23 - 26. In East Greenland, *Sebastes mentella* is further divided into a demersal and two pelagic stocks of redfish (beeked redfish), a shallow and deep pelagic stock.

Classification of redfish by species.

A separation of redfish by species has been attempted since 2007 (table 10-15, figure 16). Particularly the separation of *Sebastes mentella* and juvenile redfish varies from year to year. The decrease in juvenile redfish is therefore partly cancelled by an increase in deep-sea redfish (table 10-13). Larger individuals are easier to separate, leading to increased credibility of the classification of golden redfish. Golden redfish biomass has increased since 2009 (table 15). The increase seen in 2013 and 2014 was seen particularly in division 1 E and 1F and is often caused by few hauls containing larger individuals (figure 26) giving more than half the total West Greenland biomass. The sudden increase in larger individuals of golden redfish could indicate a connection with the Icelandic and East Greenlandic stock of Golden redfish or stocks or the West Greenlandic fjords.

American plaice (*Hippoglossoides platessoides*)

In West Greenland, American plaice is common in all divisions, but the main areas are division 1A to 1E (table 16,17 and figure 27). The biomass and abundance indices decreased about 50% from 2013 to 2014 and the indices resemble the fluctuating Greenland halibut indices during the last decade (table 16, 17 and figure 5). Clear modes can be found at 5 and 15 cm indicating new incoming year-classes and individuals larger than 45 cm are rarely seen in Greenland (figure. 17). The highest concentrations are mainly found in West Greenland (figure 27).

Atlantic wolffish (*Anarhichas lupus*)

In West Greenland, Atlantic wolffish is common in all divisions, in both East and West Greenland (figure 28). Previously Atlantic wolffish was mainly found south of 68°N, but the distribution has shifted further north since the beginning of the time series (table 18, 19 and figure 6). In 2014 both abundance and biomass indices decreased to among the lower levels seen during the past decade. The length distribution reveals the dominance of fish < 35cm and although the proportion of larger individuals has increased in the past years, individuals above 65 cm are virtually absent (figure. 18).

Spotted wolffish (*Anarhichas minor*)

Spotted wolffish are common in all divisions in both East and West Greenland and inshore and offshore. Spotted wolffish abundance and biomass indices have increased throughout the time series and the 2014 biomass estimate is among the highest observed although less than 2013 (Table 20,21 and figure 7). The length distribution ranges from 10-120 cm and occasionally weak modes at 13 cm can be identified (figure 19). Recruits and juvenile individuals are mainly found in North-West Greenland (figure 19 and 29).

Thorny skate (*Amblyraja radiata*)

In West Greenland, thorny skate is common in all divisions but the majority of both the biomass and abundance is located in West Greenland (figure 30). The most significant change happened from 2004 to 2005 indicating a poor calibration factor between the new and old Gear. The general impression is a slightly increasing stock, but the 2014 indices decreased to the most recent decade average levels (table 22, 23 and figure 8). Although Lmax for thorny skates is reported to be more than 100 cm, it has been suggested that North American thorny skates grows to larger body size than East Atlantic individuals. Since individuals larger than 55 cm are rarely seen in Greenland waters and all individuals at about 50 cm seems fully mature it seems likely that thorny skates in Greenland resemble East Atlantic stocks. During the most recent years

thorny skate length distributions have revealed clear modes at 10-15 cm and 35-50 cm probably corresponding to recruits and overlapping year-classes of adult individual (figure. 20).

Other species

The Appendix contains biomass and abundance estimates for elasmobranchs, teleosts, cephalopods and crustaceans excl. Shrimp for the West-Greenland part of the GINR shrimp fish survey including the West-Greenlandic Shelf part of NAFO div 0A.

Discussion

Catch-ability is set at 1 for all species. However, since swept area is calculated for the trawl excluding doors and bridles, catchability may be higher than 1 for some species and below 1 for other species, implying that both biomass and abundance should be regarded as index values only, not absolute values.

Index changes from the end of the 1990's to 2001 for species related to shallow water and banks (50-150m) could reflect better coverage of these depths during the past decade. Changes from 2003 to 2004 could be influenced with the trawl door update from Greenland perfect to injector and may be species dependant, as no calibration experiments were made on this account. Index changes from 2004 to 2005 could be related to species specific data quality in the calibration experiments.

The main purpose of the survey is to evaluate the biomass of northern shrimp and the effort is concentrated in areas and depths where the commercial shrimp trawling is taking place, especially on the northern slopes of the bank Store Hellefiskebanke ($67^{\circ}50'N$ $55^{\circ}00'W$) and in the inshore area Disko Bay. As Store Hellefiskebanke and Disko Bay are important nursery areas for Greenland halibut and redfish, as well as other important species (Smidt, 1969; Tåning, 1949) it is likely, that the abundance estimates of the survey reflects the juvenile stock situation of these species.

The 2002 estimates in division 1AN may have been affected by low coverage in this division, since only 2 hauls was performed in this division in 2002 and therefore not all strata can have been covered. No record exists on whether any compensation for low coverage was made in subarea 1AN in 2002.

References

- Anon., 2015. Report of the North-Western Working Group (NWWG). *ICES CM 2015*.
- Cochran, W. G. 1977: Sampling Techniques, Third edition, Wiley & Sons.
- Kingsley, M.C.S., P. Kanneworff and D.M. Carlsson. 2004. Buffered random sampling: a sequential inhibited spatial point process applied to sampling in trawl survey for northern shrimp *Pandalus borealis* in West Greenland waters. *ICES J. Mar. Sci.* 61:12-24.
- Nedreaas, K. 1980: Age determination of Northeast Atlantic *Sebastes* species. *J. Cons. int.Mer.* 47: 47: 208-230.
- Rosing, M. & K. Wieland (2005): Preliminary results from shrimp trawl calibration experiments off West Greenland (2004, 2005) with notes on encountered design/analyses problems. NAFO SCR Doc. 05/92.
- Smidt, E.L.B., 1969: The Greenland Halibut *Reinhardtius hippoglossoides* (Walb.), Biology and Exploitation in the Greenland Waters. *Meddelelser fra Danmarks Fiskeri- og Havundersøgelser*, N.S.,6: 79-148.
- Sünksen, K. 2007: Discarded by-catch in shrimp fisheries in Greenlandic offshore waters 2006-2007. NAFO SCR Doc. 07/88
- Tåning, Å.V., 1949. On the breeding places and abundance of the redfish (*Sebastes*) in the North Atlantic. *Ibid. Journ. Cons.* Vol.16 No.1: 85-96.
- Wieland, K.,& M. Storr-Paulsen, 2006: Effect of tow duration on catch rates and mean length of Northern shrimp (*Pandalus borealis*) and Greenland halibut (*Reinhardtius hippoglossoides*) in the West Greenland Bottom Trawl Survey. *Fish. Res.* **78**: 276-285.

Table 1: The survey area (km²) in the Greenland Shrimp and Fish Survey.

West Greenland							
Area	Depthstrata						Total
	<100	100-150	150-200	200-300	300-400	400-600	
1A	3039	5220					8259
1B	11346	4966					16312
1C	4183	8169					12351
1D	4136	1538					5673
1E	494	2721					3215
1F	1497	5248					6745
C0			903	2179	1154		4236
I1		407	1963	2441	1499		6310
I2		419	815	1085	1338		3658
U1		2486	4633	4785	5129		17033
U2			6710	8481	7994		23185
U3		2012	3017	1675	2710		9413
W1		2873	6099	7520	816		17307
W2		1674	2612	1741	915		6941
W3		2122	4725	2085	2994		11926
W4		4119	1818	821	1961		8719
W5		3001	3648	1950	3021		11620
W6		1206	2006	1585	1234		6031
W7		2442	891	265	317		3915
W8		424	567	405	718		2114
W9		1711	938	516	430		3595
All strata	24695	27861	24895	41344	37534	32230	188559

East Greenland				
Area	Depthstrata			Total
	0001-0200	0201-0400	0401-0600	
Q1	217	35445	6975	42637
Q2	93	7657	1246	8996
Q3	3363	22547	9830	35740
Q4	1337	7770	2054	11161
Q5	469	2785	1819	5073
Q6	6307	6130	2063	14500
All strata				118107

Table 2. Numbers of valid hauls in West Greenland, 1988-2009. 1AX=Disko Bay. 0A= The West-Greenlandic shelf part in Canadian waters. (- area included in neighboring strata)

Year	West Greenland										Total
	0A	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	
1990		29	63	*	68	17	35	16	*	*	228
1991		18	39	41	44	18	11	16	*	*	187
1992		20	33	39	36	8	18	18	11	15	198
1993		16	22	31	39	10	21	15	12	13	179
1994		16	33	27	49	9	23	8	9	9	183
1995		17	33	33	48	13	29	13	14	11	211
1996		18	20	33	46	11	29	12	9	11	189
1997		17	33	34	47	9	32	12	12	19	215
1998		10	34	33	66	14	27	19	14	14	231
1999		10	40	34	63	18	33	16	14	17	245
2000		8	25	23	45	17	37	23	14	29	221
2001		9	28	23	59	16	36	24	15	26	236
2002		2	26	22	68	12	32	18	20	27	227
2003		11	21	19	51	12	30	18	15	22	199
2004		15	25	14	41	14	24	22	20	34	209
New survey gear introduced											
2005	6	20	30	16	45	10	26	19	23	23	212
2006	5	26	40	21	49	9	27	20	21	31	244
2007	8	18	38	18	47	9	27	27	31	39	254
2008	6	16	38	16	53	7	28	23	25	46	252
2009	8	21	31	24	60	13	28	22	24	48	271
2010	10	26	44	25	65	11	30	23	24	40	289
2011	-	17	31	26	54	9	24	18	12	25	216
2012	-	18	34	21	52	12	21	18	18	26	220
2013	4	19	37	17	44	8	20	13	21	28	211
2014	-	21	36	21	47	10	19	17	23	32	226

Note.

2005. Strata 1A 50-100 no stations - included in 1A 100-150.

2006. Strata 1C 50-100 no stations - included in 1C 100-150.

2007. Strata 1C 50-100 no stations - included in 1C 100-150.

2008. All strata surveyed.

2009. Strata 1A:50-100 no stations - included in 1A 100-150.

2010. Strata 1A:50-100, 1A:100-150 no stations - included in U1 150-200.

2011. Strata W1:400-600, C0:200-300, C0:300-400, C0:400-600 no stations - included in W1 300-400.

2012. Strata C0:200-300, C0:300-400, C0:400-600 no stations - included in W1 300-400.

2013. Strata C0:400-600 no stations - included in C0 300-400. Strata W9:400-600 no stations - included in W9 300-400.

2014. Strata C0:200-300, C0:300-400, C0:400-600 no stations - included in W1 equivalent depth strata.

Table 2 - continued. Numbers of valid hauls in East Greenland, 2008-2013.

East Greenland							
Year	Q1	Q2	Q3	Q4	Q5	Q6	Total
2007							35
2008	8	6	12	7	7	12	52
2009	21	12	26	19	6	13	97
2010	19	14	24	9	6	10	82
2011	20	11	21	12	7	14	85
2012	19	16	28	13	7	15	98
2013	25	12	22	14	5	14	92
2014	22	14	12	9	8	16	81

Note:

2007 Survey startup year. Only 35 hauls. No results presented.

2008. Strata Q1,Q2,Q4:0-200 no stations - included Strata Q1,Q2,Q3:200-400.

2009. Strata Q1:0-200 m no stations – included in Q1:200-400.

2010. Strata Q1:0-200 m no stations – included in Q1:200-400.

2011. Strata Q1:0-200 m no stations – included in Q1:200-400.

2012. Strata Q1:0-200 m no stations – included in Q1:200-400.

2013. Strata Q1:0-200 m no stations – included in Q1:200-400. Strata Q5:0-200 m no stations – included in Q4:0-200.

2014. strata Q1:0-200 m no stations – included in Q1:200-400. Strata Q3:0-200 included in Q4:0-200 (COD Q3:200-400.)

Table 3. Preliminary calibration factors to adjust the Cosmos trawl catches to the former Skjervoy trawl standard. For Greenland halibut and American plaice the conversion were length dependent and for those species x in the equations represents the individual fish length.

Fishspecies	Greenland halibut	Redfish	American plaice	Atlantic wolffish	Spotted wolffish	Cod	Thorny skate
Conversion factor	0.0404x+0.6527	2.4	-0.0825x + 5.3307	2.3	2.3	1.78	5.1
Trawl size factor	1/1.1516	1/1.1516	1/1.1516	1/1.1516	1/1.1516	1/1.1516	1/1.1516
Final factor	0.035x+0.567	2.0	-0.072x + 4.629	2.0	2.0	1.5	4.4

Table 4. Greenland halibut (*Reinhardtius hippoglossoides*). Abundance indices (Millions) for West Greenland with 95% confidence limits in percent of the stratified mean.

Year	0A	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	W-Grl	CI	E-Grl
1992		50.7	8.2	96.7	231.5	1.5	1.2	0.8	0.6	0.0	391.2	27	
1993		29.1	24.3	34.0	152.6	9.8	4.3	4.6	0.4	0.5	263.5	28	
1994		22.9	35.2	63.0	130.7	10.0	18.1	2.4	0.1	0.4	282.8	22	
1995		39.5	28.9	89.4	98.9	18.3	12.2	5.7	0.1	0.3	293.4	32	
1996		92.6	13.7	102.5	265.9	11.8	14.0	10.7	0.2	0.4	511.7	25	
1997		41.3	17.5	112.1	97.6	2.2	13.0	0.9	0.1	0.3	285.0	22	
1998		78.5	30.4	209.6	47.2	23.3	7.2	3.6	1.4	0.6	401.6	29	
1999		100.8	58.2	95.4	91.0	5.8	5.3	7.7	1.3	0.9	366.3	35	
2000		81.7	19.3	172.8	126.7	4.1	4.8	6.7	0.1	0.6	416.8	30	
2001		145.9	97.9	223.7	111.5	1.7	4.8	2.6	0.7	1.6	590.2	28	
2002		78.0	75.2	148.1	42.5	2.7	6.2	6.5	0.6	1.3	360.9	38	
2003		154.6	37.7	227.0	116.7	2.3	2.0	2.4	0.1	1.0	543.8	36	
2004		154.8	20.6	199.1	84.8	2.4	2.9	2.8	0.1	0.4	468.0	34	
New survey gear introduced											cv	cv	
2005	3.3	177.5	51.6	186.5	202	5.3	9.1	1.9	1.1	0.5	638.8	12	
2006	14.1	110.5	69.9	96.9	198.6	4.2	24.6	0.8	0.0	0.3	520.0	11	
2007	10.5	103.1	75.3	128.5	154.4	2.2	37.9	0.3	1.2	0.4	513.8	13	
2008	13.2	184.2	106.5	64.6	88.6	0.7	0.6	0.9	0.3	0.3	459.9	12	3.0 53
2009	8.0	132.8	75.3	72.9	119.6	6.9	0.7	0.9	0.3	0.1	417.5	8	4.1 12
													0
2010	9.1	154.5	117.7	123.3	115.9	4.5	0.7	1.2	0.1	0.1	527.1	9	3.0 43
2011	-	318.8	111.6	230.7	93.4	8.9	1.1	0.9	0	0.2	765.6	16	2.9 37
2012	-	80.2	48.8	105.9	52.3	5.0	0.4	0.1	0	0.1	292.8	8	1.7 46
2013	1.0	148.0	212.0	188.0	145.0	11.0	5.0	0.0	0.0	0.0	711.0	16	1.7 38
2014	-	79.1	117	103.7	32.6	1.7	1.6	0.3	0.6	0	336.1	15	0.8 25

Table 5. Greenland halibut (*Reinhardtius hippoglossoides*). Biomass indices (tons) for West Greenland with 95% confidence limits in percent of the stratified mean.

Year	0A	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	W-Grl	CI	E-Grl
1992		3516	785	4992	4203	402	206	97	48	0	14250	22	
1993		2483	1286	2507	4255	747	595	539	333	60	12804	27	
1994		2007	1697	3598	4748	1665	1458	91	10	25	15199	26	
1995		4367	1291	5786	2567	825	971	502	12	45	16365	51	
1996		3682	1294	8593	5496	439	1248	899	9	118	21778	22	
1997		4972	1746	6456	4929	421	1754	180	25	84	20567	29	
1998		7025	4976	11874	2821	1724	863	275	117	278	29955	35	
1999		10205	6025	8060	5224	555	778	261	48	318	31473	44	
2000		3411	1713	9537	3985	454	692	567	38	280	20676	30	
2001		8433	2478	10161	3802	278	1208	289	33	443	27126	32	
2002		6158	2067	9070	3108	779	737	670	39	402	23055	40	
2003		8297	3399	16556	5693	478	589	297	4	355	35668	28	
2004		15182	2079	28229	11755	1147	420	319	2	201	59332	36	
New survey gear introduced												cv	cv
2005	421	22894	7010	22580	17150	574	1129	347	263	412	72780	12	
2006	519	15179	4516	20246	13797	519	6693	93	0	206	61769	12	
2007	769	11603	5666	13137	6950	163	5920	3	82	246	44539	18	
2008	402	19559	4417	16422	7822	147	302	120	59	179	49429	13	4850
2009	229	21764	2634	19902	7047	478	324	164	16	25	52584	20	4454
2010	489	25880	7071	17559	8658	599	54	320	1	58	60688	11	5729
2011	-	18213	6778	23977	13945	640	364	49	0	14	64547	10	3825
2012	-	10331	4327	16168	9194	1475	225	7	4	81	41813	7	2439
2013	210	13661	12510	15103	11508	689	812	3	95	105	54695	9	2664
2014	-	8577	7388	11463	6014	942	829	45	111	9	35377	12	1611

Table 6. Cod (*Gadus morhua*). Abundance indices (million) for West Greenland with 95% confidence limits in percent of the stratified mean. () incomplete coverage of survey area.

Year	0A	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	W-Grl.	CI	E-Grl
1991	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.5			0.8	51	
1992	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.3	0.0	0.0	0.6	49	
1993	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.3	0.0	0.5	66	
1994	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.1	47	
1995	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.1	0.0	0.6	55	
1996	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.3	54	
1997	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	68	
1998	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	54	
1999	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.2	29	
2000	0.0	0.0	0.2	0.4	0.3	0.2	0.1	0.1	0.0	0.2	1.3	23	
2001	0.0	0.0	0.0	0.0	0.2	0.1	0.1	0.4	0.3	0.3	1.5	26	
2002	0.0	0.0	0.0	0.0	0.1	0.1	0.1	3.3	0.1	0.5	4.2	50	
2003	0.0	0.0	0.1	0.4	1.4	0.0	0.4	0.7	0.2	0.2	3.4	22	
2004	0.0	0.0	0.0	0.2	0.0	0.1	0.4	2.6	1.5	1.6	6.5	29	
New survey gear introduced											cv	cv	
2005	0.1	0.0	0.0	0.1	0.7	0.1	1.8	4.8	6.8	94.1	108.7	52	
2006	0.5	0.0	0.2	0.2	2.8	1.4	2.5	16.5	3.3	46.3	73.8	27	
2007	0.7	0.0	1.0	0.3	2.5	0.7	7.1	3.0	2.6	38.1	56.1	46	
2008	1.2	0.1	0.7	0.1	2.4	1.7	4.1	9.5	11.9	21.7	53.4	23	31.5
2009	0.9	0.0	0.8	0.1	3.2	1.0	3.2	2.8	1.4	1.7	15.1	11	54.7
2010	0.3	0.1	0.6	0.0	1.7	1.1	2.7	8.2	2.5	6.1	23.4	24	45.3
2011	-	0.0	3.4	3.7	35.8	7.8	2.1	19.5	1.0	7.4	80.9	17	42.7
2012	-	1.1	3.8	3.5	10.0	1.0	3.3	1.2	27.1	20.3	71.1	39	22.7
2013	4.7	1.0	4.0	3.8	11.1	1.6	6.1	7.5	30.0	55.5	125.2	36	67.2
2014	-	2.4	4.2	0.4	4.4	1.3	78.9	2.5	16.5	20.6	131.0	57	35.4

Table 7. Cod (*Gadus morhua*). Biomass indices (tons) for West Greenland with 95% confidence limits in percent of the stratified mean. () incomplete coverage of survey area.

Year	0A	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	W-Grl.	CI	E-Grl
1990		2	13	*	75	3	83	9005	*	*	(9180)	65	
1991		0	0	7	2	15	151	310	*	*	(485)	44	
1992		0	0	3	20	34	75	112	0	2	246	46	
1993		0	0	2	0	5	22	39	113	5	186	68	
1994		0	3	0	0	9	38	0	1	0	51	46	
1995		0	0	0	5	1	120	23	3	4	155	63	
1996		0	0	0	0	0	15	23	24	44	107	50	
1997		0	0	0	0	2	53	0	0	0	55	76	
1998		0	0	0	1	0	0	47	50	3	101	56	
1999		0	1	5	23	5	1	17	1	0	53	47	
2000		0	51	99	76	54	21	9	2	46	357	23	
2001		0	0	15	125	30	56	178	98	100	603	23	
2002		0	0	13	54	74	41	1489	42	150	1863	46	
2003		0	18	111	315	8	264	453	118	46	1332	26	
2004		0	0	496	46	7	176	680	685	305	2394	28	
New survey gear introduced													
											cv	cv	
2005	38	0	22	45	320	19	449	1077	1170	60546	63684	71	
2006	114	0	40	22	578	74	471	5513	536	19874	27221	32	
2007	247	13	317	56	711	121	1502	514	541	26843	30865	54	
2008	421	99	227	46	1218	794	923	1730	3321	19702	28481	37	47864 23
2009	212	0	184	42	1046	199	688	453	282	499	3604	13	58141 29
2010	183	38	215	7	821	144	573	2417	835	2899	8133	31	110656 53
2011	-	1	537	726	7468	1493	398	3963	196	3948	18730	16	81138 20
2012	-	194	1061	841	3000	313	1226	447	14104	15911	37098	39	64421 21
2013	2446	294	1026	1287	3443	447	1871	4361	19015	51622	85812	37	168771 28
2014	-	709	1644	286	1813	492	56061	2511	21714	27755	112984	50	153807 54

Table 8. Redfish species combined, all sizes (*Sebastes sp.*). Abundance indices (million) for West Greenland with 95% confidence limits in percent of the stratified mean.

Year	0A	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	W-Grl	CI	E-Grl
1992		1	146	9	1006	187	69	34	6	4	1462	32	
1993		4	210	17	361	22	157	182	97	520	1568	69	
1994		12	187	12	1573	225	273	85	10	84	2461	26	
1995		1	67	11	559	33	183	93	5	5	957	23	
1996		1	8	8	1688	59	124	63	11	54	2017	29	
1997		5	43	7	348	58	156	57	23	22	719	24	
1998		1	24	20	236	15	115	71	10	32	522	24	
1999		2	69	9	287	11	11	61	4	23	574	21	
2000		0	27	6	27	32	28	10	18	159	165	23	
2001		2	77	3	25	15	20	12	1	180	183	60	
2002		1	23	1	50	84	43	44	2	10	257	27	
2003		2	45	2	210	32	79	25	2	26	423	23	
2004		2	11	1	52	39	47	30	2	69	253	37	
New survey gear introduced											cv	cv	
2005	184	1	19	1	73	79	77	25	12	80	551		
2006	27	7	25	1	110	52	83	16	3	11	334		
2007	97	2	37	0	128	34	49	5	3	17	371		
2008	99	2	40	0	121	33	24	3	2	11	334	1847	
2009	37	2	29	0	115	26	12	1	2	9	234	1988	
2010	27	6	27	0	69	42	23	8	3	8	213	1566	
2011	-	10	23	1	121	18	31	13	1	9	225	2517	
2012	-	7	10	0	42	13	20	10	2	16	120	1018	
2013	12	2	10	0	29	4	30	6	44	7	145	1075	
2014	-	1	3	0	23	5	3	39	20	3	96	559	

Table 9. Redfish species combined, all sizes (*Sebastes sp.*). Biomass indices (tons) for West Greenland with 95% confidence limits in percent of the stratified mean.

Year	0A	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	W-Grl.	CI	E-Grl.
1992		69	18117	437	13423	2832	1576	1124	169	147	37894	43	
1993		195	4994	710	6420	300	1549	3835	1923	2138	22065	38	
1994		590	5076	538	16064	1986	3886	995	179	1272	30586	24	
1995		52	1585	775	5029	869	2963	1952	358	123	13705	22	
1996		18	117	782	12178	1694	2552	1980	304	1788	21413	28	
1997		599	1481	337	4913	1597	6766	1901	1099	1229	19922	31	
1998		39	1467	1423	6193	2130	3274	1953	606	1198	18283	22	
1999		164	4021	742	5596	999	2742	2976	207	1124	18671	25	
2000		0	1790	793	1045	2185	2337	463	2411	1214	12237	36	
2001		192	5380	536	1746	1460	2637	1069	60	2256	15337	50	
2002		55	1917	397	2536	2386	1676	2654	272	998	12891	28	
2003		279	2886	702	6357	2319	6185	1918	187	2476	23308	32	
2004		369	462	368	2210	2274	2996	1679	101	1026	11486	41	
New survey gear introduced											cv	cv	
2005	3491	134	1378	665	3370	6974	6212	2751	1388	2771	29132		
2006	591	1129	2196	759	4427	2717	6213	959	557	2350	21897		
2007	3367	248	2172	153	6886	1499	5166	358	282	1778	21908		
2008	2845	433	2221	210	7411	4007	2542	224	286	2585	22761	290666	
2009	1696	356	2472	193	5496	3688	1951	293	335	1339	17819	318517	
2010	1348	761	3363	910	4765	4193	3073	1043	369	1069	20895	336840	
2011	-	1917	1536	1486	8362	3538	5377	4026	226	2397	28865	511700	
2012	-	1382	1224	998	4380	2438	3560	1942	239	10341	26505	234650	
2013	965	429	1684	553	6063	1077	7327	1269	23081	3178	45626	454592	
2014		370	394	150	5241	1074	1092	11621	11017	2009	32969	206406	

Table 10. Juvenile redfish <20cm (*Sebastes sp.*). Abundance indices (million) for West Greenland with 95% confidence limits in percent of the stratified mean.

Year	0A	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	W-Grl	cv	E-Grl	cv
2006	27	6	20	0	107	51	79	15	2	7	314	16		
2007	96	0	28	0	120	33	44	4	2	15	342	15		
2008	97	0	36	0	118	0	9	2	1	6	269	26	452	53
2009	36	0	14	0	111	1	4	0	1	4	172	24	954	28
2010	24	0	9	0	33	0	3	4	2	5	80	21	507	31
2011	-	0	15	0	70	1	0	2	0	0	89	23	102	61
2012	-	0	2	0	16	0	1	0	1	0	21	23	253	56
2013	5	0	3	0	18	0	2	1	0	0	30	17	48	36
2014	0	1	0	8	0	0	0	0	0	0	9	31	21	33

Table 11. Juvenile redfish <20cm (*Sebastes sp.*). Biomass indices (tons) for West Greenland with 95% confidence limits in percent of the stratified mean. () incomplete coverage of survey area.

Year	0A	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	W-Grl	cv	E-Grl	cv
2006	489	650	1433	0	3270	2238	4623	746	111	1162	14722	18		
2007	2912	0	1475	0	3985	1142	4262	156	87	821	14839	16		
2008	2358	2	1678	0	5198	2	154	44	12	55	9503	29	9011	53
2009	1365	0	491	3	3571	36	160	9	24	90	5749	29	41341	63
2010	919	0	394	0	1524	0	105	161	66	154	3323	29	24135	31
2011	-	9	289	1	2579	47	19	88	25	18	3074	32	3781	56
2012	-	0	52	0	557	9	32	13	21	16	700	24	36567	79
2013	228	2	158	1	802	10	118	12	4	7	1340	21	1244	33
2014	0	22	0	227	7	4	3	4	2	268	28	550	26	

Table 12. Deep-sea (*Sebastes mentella*). Abundance indices (million) for West Greenland with 95% confidence limits in percent of the stratified mean.

Year	0A	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	W-Grl	cv	E-Grl	cv
2006	0	1	1	0	1	1	1	1	0	2	8	24		
2007	0	0	0	0	6	0	0	0	0	1	8	52		
2008	1	1	4	0	2	33	14	0	0	2	57	56	1286	49
2009	1	2	14	0	4	24	7	1	0	4	55	45	895	29
2010	2	5	16	0	35	41	19	4	0	3	127	36	995	36
2011	-	9	8	0	49	16	27	7	0	6	122	22	400	22
2012	-	6	7	0	24	12	16	8	1	4	79	10	305	27
2013	7	1	6	0	9	4	25	2	41	2	98	85	338	33
2014	0	2	0	13	5	2	37	19	2	80	25	172	25	

Table 13. Deep-sea (*Sebastes mentella*). Biomass indices (tons) for West Greenland with 95% confidence limits in percent of the stratified mean.

Year	0A	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	W-Grl	cv	E-Grl	cv
2006	11	168	65	77	252	274	543	64	34	544	2032	22		
2007	84	0	30	28	1443	0	8	28	11	277	1909	84		
2008	141	245	385	9	432	3989	2209	71	50	395	7926	49	247557	47
2009	79	239	1659	45	618	2989	1420	213	8	828	8098	38	236680	35
2010	141	606	1831	11	2281	4039	2322	608	56	621	12519	36	283924	36
2011	-	1300	998	4	3441	3156	4129	1509	71	1540	16149	24	122949	26
2012	-	1041	886	334	2084	2093	2340	1386	154	878	11195	9	100342	29
2013	655	124	958	217	2522	835	5974	454	20728	833	33301	59	162268	50
2014		71	264	8	1949	991	379	10466	9559	838	24525	22	67611	24

Table 14. Golden redfish (*Sebastes norvegicus*). Abundance indices (million) for West Greenland with 95% confidence limits in percent of the stratified mean.

Year	0A	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	W-Grl	cv	E-Grl	cv
2006	0	0	3	0	2	0	2	0	1	2	12	23		
2007	1	2	8	0	2	1	5	0	0	1	21	43		
2008	1	0	0	0	1	0	1	0	0	4	9	25	108	29
2009	1	0	1	0	1	1	1	0	1	2	6	15	139	27
2010	0	0	1	0	1	1	1	0	1	0	6	17	64	36
2011	-	1	0	1	2	1	3	3	1	2	14	12	2015	36
2012	-	0	1	0	2	0	2	2	0	12	20	54	460	36
2013	0	1	1	0	2	0	2	3	3	4	17	15	689	34
2014	0	0	0	2	0	1	1	1	1	1	7	17	366	40

Table 15. Golden redfish (*Sebastes norvegicus*). Biomass indices (tons) for West Greenland with 95% confidence limits in percent of the stratified mean.

Year	0A	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	W-Grl	cv	E-Grl.	cv
2006	92	311	698	682	905	204	1047	149	413	644	5144	16		
2007	371	248	667	126	1459	356	896	174	184	679	5160	27		
2008	345	185	157	201	1781	15	180	109	224	2135	5332	41	34098	32
2009	252	117	321	146	1308	663	370	70	303	422	3971	18	40496	27
2010	288	155	1137	899	960	154	646	275	246	293	5053	28	28781	32
2011	-	607	249	1482	2342	334	1229	2429	130	839	9642	21	384970	36
2012	-	341	286	664	1740	336	1188	543	65	9447	14609	61	97741	28
2013	82	303	568	335	2740	232	1234	803	2349	2338	10985	31	291080	57
2014	299	108	142	3066	77	709	1152	1454	1169	8176	19	138245	41	

Table 16. American plaice (*Hiploglossoides platessoides*). Abundance indices ('million) for West Greenland with 95% confidence limits in percent of the stratified mean. () incomplete coverage of survey area.

Year	0A	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	W-Grl.	CI	E-Grl
1992		1.4	1.0	2.8	1.4	0.6	1.7	1.4	0.5	0.1	10.8	22	
1993		1.3	1.7	1.2	3.0	0.8	2.0	1.7	0.6	0.9	13.2	24	
1994		2.2	3.7	3.3	14.9	7.0	9.5	0.7	0.3	0.5	42.0	32	
1995		1.0	0.6	1.8	6.3	0.9	2.7	3.0	0.3	0.5	17.2	29	
1996		1.6	3.4	7.3	4.6	1.7	4.2	3.1	0.1	0.7	26.6	18	
1997		6.6	2.0	2.7	15.1	1.0	10.4	2.0	0.3	0.7	40.8	47	
1998		1.6	1.9	2.4	3.6	1.2	1.5	6.4	0.9	5.6	25.2	27	
1999		0.5	1.7	2.0	6.8	1.2	5.3	1.9	1.0	0.6	20.9	18	
2000		1.8	4.8	6.7	14.8	1.9	3.5	3.8	0.5	0.5	38.5	23	
2001		1.3	1.3	2.2	13.6	1.5	3.5	2.3	0.6	0.6	26.8	31	
2002		0.0	3.8	4.7	8.8	1.8	5.1	31.8	1.5	1.8	59.5	49	
2003		2.2	5.2	5.5	25.7	1.6	13.7	15.0	0.8	2.0	71.7	22	
2004		0.7	1.4	5.3	11.9	2.8	10.2	8.9	0.6	1.6	43.4	27	
New survey gear introduced											cv	cv	
2005	3.5	2.1	13.7	5.8	59.9	17.3	74.5	19.5	6.1	5.6	208.0	9	
2006	3.9	1.3	15.0	10.9	40.3	10.0	38.1	21.6	4.4	4.5	150.0	10	
2007	6.5	0.6	12.3	6.3	46.8	9.4	31.2	11.9	2.6	2.7	130.8	12	
2008	7.1	3.7	10.8	4.0	29.5	2.2	13.5	11.4	1.3	1.3	84.8	8	17.2 22
2009	2.9	5.6	18.0	14.3	41.4	8.1	11.6	7.6	1.1	2.3	112.8	9	20.2 12
2010	9.5	2.8	40.7	14.3	50.7	9.2	34.4	20.3	1.2	2.1	185.2	8	17.4 19
2011	-	8.6	40.7	17.1	103.6	6.2	30.3	21.6	0.8	1.9	230.7	11	17.5 16
2012	-	2.3	18.3	15.3	51.3	8.3	19.4	11.4	0.9	1.1	129.4	9	10.4 22
2013	4.4	19.0	39.0	25.4	84.1	9.1	34.1	27.9	4.7	0.9	248.6	13	13.1 17
2014		7.0	22.0	11.0	36.0	6.0	15.0	10.0	2.0	1.0	110.0	7	10.0 19

Table 17. American plaice (*Hippoglossoides platessoides*). Biomass indices (tons) for West Greenland with 95% confidence limits in percent of the stratified mean. () incomplete coverage of survey area.

Year	0A	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	W-Grl.	CI	E-Grl
1992		57	54	213	78	51	137	128	55	6	779	23	
1993		56	72	87	90	28	107	141	69	43	694	26	
1994		112	293	277	487	308	284	60	22	64	1906	22	
1995		65	54	279	191	51	87	130	19	18	895	18	
1996		119	264	670	231	74	142	119	7	27	1654	18	
1997		323	150	287	398	87	367	135	31	25	1803	21	
1998		154	178	328	185	48	82	398	97	102	1573	20	
1999		81	136	170	287	43	202	145	65	44	1173	17	
2000		175	278	408	551	74	178	227	89	40	2021	18	
2001		169	79	140	403	65	162	153	38	67	1276	17	
2002		0	184	327	414	151	275	1061	92	67	2570	23	
2003		196	352	338	1013	125	680	1048	59	171	3980	20	
2004		138	143	192	537	128	715	747	38	150	2789	27	
New survey gear introduced												cv	cv
2005	246	346	944	722	3049	1136	4511	2196	470	569	14190	8	
2006	268	210	608	1148	2288	702	3534	2163	399	380	11699	13	
2007	356	112	544	731	2877	731	2418	810	230	308	9118	12	
2008	371	437	648	382	1889	212	1067	898	128	149	6181	9	2495
2009	183	556	649	643	1977	585	826	825	102	127	6473	7	2577
2010	355	235	1539	1214	2436	748	2128	1460	105	224	10442	9	3181
2011	-	463	1364	1459	5464	792	1391	2020	99	208	13256	7	4385
2012	-	236	756	730	3508	662	1350	777	121	178	8317	9	2702
2013	325	920	1472	1457	4066	521	2818	1425	568	121	13694	10	2269
2014		311	746	990	2245	337	1089	1027	203	130	7078	7	1203

Table 18. Atlantic wolffish (*Anarhichas lupus*). Abundance indices (millions) for West Greenland with 95% confidence limits in percent of the stratified mean.

Year	0A	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	W-Grl	CI	E-Grl
1992		0.0	0.0	0.0	0.0	0.1	0.3	0.1	0.1	0.1	0.8	33	
1993		0.0	0.0	0.0	0.0	0.0	0.2	0.2	0.3	0.3	1.2	41	
1994		0.0	0.1	0.0	0.3	0.2	1.9	0.4	0.4	1.6	4.9	36	
1995		0.0	0.0	0.0	0.1	0.1	0.4	0.1	0.4	0.2	1.3	34	
1996		0.0	0.1	0.0	0.1	0.1	0.4	0.2	0.6	0.6	2.0	50	
1997		0.0	0.0	0.0	0.2	0.0	0.8	0.2	0.3	0.1	1.7	28	
1998		0.0	0.3	0.0	0.3	0.1	0.7	0.4	0.5	0.7	3.0	30	
1999		0.0	0.2	0.1	0.1	0.1	0.4	0.2	0.4	0.4	2.0	44	
2000		0.0	0.5	0.1	0.3	0.4	0.4	0.6	1.7	0.0	4.0	69	
2001		0.0	0.1	0.2	0.4	0.1	0.2	0.4	0.2	0.1	1.5	40	
2002		0.0	0.4	0.1	0.5	0.0	0.9	0.6	1.1	0.5	4.1	33	
2003		0.0	0.1	0.6	2.3	0.2	1.5	0.5	2.3	1.1	8.8	27	
2004		0.0	0.1	0.3	1.6	0.9	0.2	0.3	0.7	0.3	4.4	39	
New survey gear introduced											cv	cv	
2005	0.1	0.0	0.5	0.0	1.5	0.6	6.9	2.4	5.5	1.5	19.1	16	
2006	0.0	0.1	0.8	0.1	0.9	0.6	2.4	1.2	1.0	2.8	9.8	14	
2007	0.0	0.0	0.7	0.0	1.3	0.6	0.7	0.3	0.4	0.7	4.8	15	
2008	0.2	0.0	1.0	0.1	0.7	0.5	1.0	0.5	0.4	0.3	4.6	15	4.7
2009	0.0	0.1	1.0	0.1	0.7	0.4	1.5	0.2	1.1	0.4	5.6	19	12.2
2010	0.1	0.4	2.6	0.1	2.3	0.6	3.1	0.8	0.9	0.6	11.4	8	5.3
2011	-	0.1	1.1	0.1	1.3	0.3	0.5	0.9	0.3	0.4	5.0	14	6.4
2012	-	0.0	2.2	0.2	1.6	0.3	0.7	1.2	0.3	0.6	7.0	18	3.8
2013	0.1	0.1	3.1	0.0	1.0	0.3	1.7	0.4	0.5	0.8	7.9	18	6.5
2014	-	0.4	2.3	0.3	0.7	0.8	0.3	0.1	0.2	0.3	5.5	16	2.6

Table 19. Atlantic wolffish (*Anarhichas lupus*). Biomass indices (tons) for West Greenland with 95% confidence limits in percent of the stratified mean.

Year	0A	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	W-Grl	CI	E-Grl
1992		0	7	0	8	21	47	22	28	31	163	33	
1993		0	5	6	1	2	26	35	29	188	292	64	
1994		0	12	9	40	39	198	30	65	249	644	38	
1995		0	0	0	22	9	38	24	90	36	219	40	
1996		0	1	3	17	23	41	35	103	101	324	53	
1997		0	3	0	21	1	115	16	58	15	228	30	
1998		0	3	2	13	9	60	34	104	133	358	38	
1999		0	4	13	21	12	8	6	202	62	329	79	
2000		0	20	3	52	31	55	54	396	15	626	90	
2001		0	1	3	11	1	16	21	42	23	117	40	
2002		0	9	2	77	5	73	78	216	118	579	37	
2003		0	2	41	267	64	361	60	205	148	1148	24	
2004		0	6	19	160	56	96	21	162	100	620	27	
New survey gear introduced											cv	cv	
2005	11	0	76	5	201	123	1564	348	890	640	3858	16	
2006	3	4	58	60	127	134	359	168	172	1241	2326	20	
2007	0	0	73	6	357	326	147	107	122	533	1670	21	
2008	15	0	76	31	124	55	348	88	136	226	1098	16	1091
2009	0	3	109	30	155	72	275	92	211	648	1595	31	2870
2010	10	12	509	51	350	144	668	134	104	345	2327	17	1295
2011	-	46	96	64	197	69	121	233	63	205	1094	17	1633
2012	-	0	239	82	506	44	79	252	64	444	1708	23	1172
2013	9	27	239	20	136	81	555	183	115	636	2000	25	1652
2014	-	9	79	69	158	62	107	78	100	458	1121	30	520

Table 20. Spotted wolffish (*Anarhichas minor*). Abundance indices (millions) for West Greenland with 95% confidence limits in percent of the stratified mean.

Year	0A	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	W-Grl	CI	E-Grl
1992		0.0	0.1	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.3	28	
1993		0.0	0.1	0.0	0.1	0.0	0.1	0.0	0.0	0.1	0.4	36	
1994		0.0	0.3	0.0	0.2	0.1	0.4	0.0	0.0	0.0	1.1	33	
1995		0.0	0.2	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.4	22	
1996		0.1	0.1	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.5	24	
1997		0.1	0.3	0.0	0.2	0.0	0.0	0.1	0.0	0.0	0.8	23	
1998		0.1	0.2	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.6	25	
1999		0.9	0.3	0.1	0.4	0.1	0.0	0.0	0.0	0.0	1.7	42	
2000		0.0	0.8	0.0	0.5	0.1	0.0	0.0	0.1	0.0	1.6	28	
2001		0.3	0.4	0.1	0.3	0.0	0.0	0.0	0.0	0.0	1.1	27	
2002		0.3	2.1	0.1	0.4	0.1	0.1	0.1	0.0	0.0	3.1	42	
2003		0.5	1.5	0.2	1.4	0.0	0.6	0.0	0.0	0.1	4.3	24	
2004		0.2	0.7	0.2	0.6	0.0	0.1	0.1	0.0	0.1	2.1	30	
New survey gear introduced										cv	cv		
2005	0.0	1.2	1.9	0.0	0.7	0.2	0.4	0.3	0.2	0.1	5.1	23	
2006	0.0	0.4	1.0	0.1	1.4	0.3	0.3	0.2	0.1	0.2	3.9	23	
2007	0.1	0.5	1.0	0.1	1.5	0.4	0.2	0.1	0.0	0.1	3.9	31	
2008	0.1	0.5	0.6	0.1	0.4	0.0	0.1	0.1	0.0	0.1	2.1	15	1.1
2009	0.0	0.4	1.7	0.1	0.6	0.1	0.0	0.0	0.0	0.1	3.1	26	2.3
2010	0.2	1.0	1.8	0.2	0.7	0.2	0.2	0.1	0.1	0.2	4.5	12	2.9
2011	-	0.9	1.8	0.2	0.9	0.1	0.2	0.1	0.0	0.1	4.4	14	2.3
2012	-	0.8	1.0	0.1	0.7	0.1	0.1	0.2	0.1	0.2	3.2	15	2.9
2013	0.1	1.3	2.2	0.2	1.7	0.1	0.3	0.2	0.4	0.4	6.9	20	2.5
2014		1.4	2.3	0.1	0.2	0	0.1	0.2	0.2	0.2	4.7	21	1.5

Table 21. Spotted wolffish (*Anarhichas minor*). Biomass indices (tons) for West Greenland with 95% confidence limits in percent of the stratified mean.

Year	0A	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	W-Grl	CI	E-Grl
1992		4	76	65	110	3	34	33	6	19	351	28	
1993		55	0	100	47	16	66	4	0	282	571	53	
1994		223	180	0	81	40	119	28	11	1	683	25	
1995		0	60	15	68	16	22	19	11	164	377	49	
1996		169	77	12	193	15	6	31	0	50	554	26	
1997		193	72	37	81	0	16	124	0	5	530	34	
1998		2	64	0	143	18	6	125	100	7	465	32	
1999		131	121	23	28	36	13	2	0	0	354	31	
2000		0	188	31	133	36	19	1	593	0	1000	114	
2001		523	30	25	310	80	4	0	0	10	982	52	
2002		135	194	20	169	81	74	233	71	126	1104	28	
2003		299	1416	195	978	22	741	107	0	226	3985	22	
2004		124	1270	623	567	2	78	603	352	545	4164	35	
New survey gear introduced												cv	cv
2005	150	764	1182	6	1058	155	741	2514	568	137	7275	26	
2006	0	472	1257	243	1345	1066	1336	716	350	1145	7930	19	
2007	14	543	705	196	1249	678	874	372	178	634	5442	17	
2008	63	1487	2050	74	730	24	347	995	425	372	6567	44	5262
2009	0	280	627	653	1453	154	35	129	189	160	3662	18	2890
2010	66	2363	1508	1195	1006	167	222	238	212	1715	8691	18	3877
2011	-	2537	2244	742	2460	1210	2294	479	218	769	12955	23	6133
2012	-	1227	683	464	3166	325	468	390	555	1104	8383	21	6871
2013	7	2026	2402	544	3135	632	1494	658	3163	4845	18906	22	4622
2014	-	779	3038	381	753	9	427	2740	1496	1916	11538	23	10468

Table 22. Thorny skate (*Amblyraja radiata*). Abundance indices in millions for West Greenland with 95% confidence limits in percent of the stratified mean.

Year	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	W-Grl	CI	E-Grl
1992	2.9	1.3	1.3	1.4	0.2	0.6	1.0	0.1	0.0	9.0	25	
1993	0.7	0.4	0.8	1.6	0.5	0.9	0.5	0.6	0.2	6.1	21	
1994	1.9	1.8	1.2	3.9	1.8	2.8	0.4	0.2	0.0	14.1	21	
1995	2.7	1.3	0.8	2.8	1.8	0.7	2.4	0.5	0.1	13.1	26	
1996	4.6	2.2	1.5	4.0	0.4	0.7	0.6	0.0	0.6	14.6	23	
1997	4.6	1.4	0.6	4.4	0.2	2.3	0.7	0.1	0.0	14.4	26	
1998	2.8	4.1	3.2	3.0	0.3	0.6	0.9	0.4	0.3	15.5	25	
1999	1.7	3.2	0.9	2.8	0.3	1.0	0.8	0.2	0.2	11.0	23	
2000	2.1	3.4	2.0	4.0	0.5	0.6	0.5	0.1	0.6	13.9	23	
2001	3.2	1.2	0.6	2.3	0.3	0.4	0.5	0.1	0.2	8.8	32	
2002	0.5	1.1	0.9	2.2	0.4	0.9	2.6	0.2	0.5	9.3	25	
2003	3.9	2.4	1.4	7.2	0.1	0.7	0.8	0.2	0.7	17.4	26	
2004	2.5	1.3	1.3	1.2	0.2	0.2	0.7	0.2	0.1	7.9	24	
New survey gear introduced										cv	cv	
2005	0.1	4.1	2.8	2.2	3.7	0.5	0.8	0.6	0.4	0.2	15.4	10
2006	0.2	3.8	2.2	3.0	2.8	0.2	2.3	2.8	0.5	1.1	18.8	10
2007	0.1	3.3	1.4	1.6	3.8	0.1	1.3	0.9	1.4	0.4	14.3	20
2008	0.2	5.4	1.4	1.0	1.9	0.1	0.5	1.1	0.1	0.2	11.9	16
2009	0.0	10.9	1.1	3.9	1.9	0.3	0.3	0.6	0.2	0.4	19.7	20
2010	0.1	5.6	4.0	4.1	3.3	0.8	2.7	1.2	0.1	0.0	21.8	12
2011	-	5.5	2.3	3.6	6.7	0.4	4.0	4.9	1.7	0.1	29.2	16
2012	-	2.7	2.2	2.4	4.7	0.5	1.3	2.1	0.1	0.0	16.1	15
2013	0.4	4.9	2.1	3.2	9.1	0.6	1.7	1.1	0.1	0.1	23.4	27
2014	2.4	2.4	1.6	1.9	0.5	0.7	0.2	0.1	0	9.8	13	0.6

Table 23. Thorny skate (*Amblyraja radiata*). Biomass (tons) for West Greenland with 95% confidence limits in percent of the stratified mean.




Fig 1: The Stratification areas used in the Greenland Shrimp and Fish survey. In West Greenland each strata is divided in depth strata of 150-200m, 200-300m, 300-400m and 400-600m. "Shallow" water strata of 0-100m and 100-150m delimited by the 3 nm line and the NAFO Div. Borders of the shallow water stratas are not shown. In East Greenland each strata is divided in depth strata of 200-400m and 400-600m. "Shallow" water strata of 0-200m is delimited by the 3 nm line.




Fig 2. Greenland halibut abundance (million) and biomass (Kt) for West Greenland. SK30 is the Skjaervoy trawl used from 1990 to 2004. CO26 is the Cosmos trawl used since 2005. Dashed lines are trawl calibrated indices.




Fig 3. Atlantic cod abundance (million) and biomass (Kt) for West Greenland. SK30 is the Skjaervoy trawl used from 1990 to 2004. CO26 is the Cosmos trawl used since 2005.




Fig 4. Redfish species (*S. mentella* and *S. norvegicus* combined) abundance (million) and biomass (Kt) for West Greenland. SK30 is the Skjaervoy trawl used from 1990 to 2004. CO26 is the Cosmos trawl used since 2005. Dashed lines are trawl calibrated indices.




Fig 5. American plaice abundance (million) and biomass (Kt) for West Greenland. SK30 is the Skjaervoy trawl used from 1990 to 2004. CO26 is the Cosmos trawl used since 2005. Dashed lines are trawl calibrated indices.




Fig 6. Atlantic wolffish abundance (million) and biomass (Kt) for West Greenland. SK30 is the Skjaervoy trawl used from 1990 to 2004. CO26 is the Cosmos trawl used since 2005. Dashed lines are trawl calibrated indices.




Fig 7. Spotted wolffish abundance (million) and biomass (Kt) for West Greenland. SK30 is the Skjaervoy trawl used from 1990 to 2004. CO26 is the Cosmos trawl used since 2005. Dashed lines are trawl calibrated indices.




Fig 8. Thorny skate abundance (million) and biomass (Kt) for West Greenland. SK30 is the Skjaervoy trawl used from 1990 to 2004. CO26 is the Cosmos trawl used since 2005. Dashed lines are trawl calibrated indices.




Fig 9. Total recruitment of age one including Disko Bay.




Fig 10. Distribution of one year old Greenland halibut in the 2012-2104 surveys.




Fig 11. Relative distribution of one-year old Greenland halibut between offshore areas and inshore Disko Bay.




Fig 12. Number of one-year of Greenland halibut by division and year.




Fig. 13. Distribution of age 2 and 3+ Greenland halibut in the 2104 survey.




Fig. 14. Greenland halibut (*Reinhardtius hippoglossoides*). Length frequencies for West Greenland 2009-2014 (left) and length frequencies per division in 2014 (right).




Fig. 15. Atlantic cod (*Gadus morhua*). Length frequencies for West Greenland 2009-2014 (left) and length frequencies per division in 2014 (right).




Fig. 16. Stacked Length frequencies for golden redfish (*Sebastes Norvegicus*). Deep-sea redfish (*Sebastes mentella*) and juvenile redfish (*Sebastes sp.*) for West Greenland 2009-2014 (left) and length frequencies per division in 2014 (right).




Fig. 17. American place (*Hippoglossoides platessoides*). Length frequencies for West Greenland 2009-2014 (left) and length frequencies per division in 2014 (right).




Fig. 18. Atlantic wolffish (*Anarhichas lupus*). Length frequencies for West Greenland by year (left) and length frequencies per division (right).




Fig. 19. Spotted wolffish (*Anarhichas lupus*). Length frequencies for West Greenland 2008-2013 (left) and length frequencies per division in 2013 (right).




Fig. 20. Thorny skate (*Amblyraja radiata*) length frequencies for West Greenland 2008-2013 (left) and length frequencies per division in 2013 (right).




Fig. 21. Greenland halibut survey biomass in kg/km² and abundance in numbers/km².




Fig. 22. Atlantic cod survey biomass in kg/km² and abundance in numbers/km².




Fig. 23. Juvenile redfish < 20 cm survey biomass in kg/km² and abundance in numbers/km².




Fig. 24. Deep-sea redfish survey biomass in kg/km² and abundance in numbers/km².




Fig. 25. Golden redfish survey biomass in kg/km² and abundance in numbers/km².




Fig. 26. Golden redfish and deep-sea redfish survey biomass in kg/km² and abundance in numbers/km².




Fig. 27. American plaice survey biomass in kg/km² and abundance in numbers/km².




Fig. 28. survey biomass in kg/km² and abundance in numbers/km².




Fig. 29. Spotted wolffish survey biomass in kg/km² and abundance in numbers/km².




Fig. 30. Thorny skate survey biomass in kg/km² and abundance in numbers/km².

Appendix I.

2014 biomass (in Kilotonnes) and abundance (in million individuals) indices for Elasmobranchs. Teleosts. Cephalopods and crustaceans excl. Shrimp species for the West and East-Greenland part of the GINR shrimp fish survey 0-600m including the West-Greenlandic Shelf part of NAFO div 0A (Canada).

NameLatin	West mill	East mill	Total mill	West Kt	East Kt	Total Kt
Gadus morhua	130,97	35,40	166,38	112,98	153,81	266,79
Sebastes norvegicus	6,72	366,33	373,05	8,18	138,25	146,42
Somniosus microcephalus	0,18	0,00	0,18	138,67	0,00	138,67
Sebastes mentella	79,74	171,56	251,29	24,53	67,61	92,14
Reinhardtius hippoglossoides	336,60	0,84	337,44	35,38	1,61	36,99
Mallotus villosus	2707,57	7,55	2715,12	28,95	0,12	29,07
Boreogadus saida	2789,63	66,83	2856,47	23,23	0,90	24,13
Anarhichas minor	4,71	1,53	6,24	11,54	10,47	22,01
Anarhichas denticulatus	0,99	0,91	1,90	4,76	8,49	13,25
Hippoglossoides platessoides	110,49	10,01	120,49	7,08	1,20	8,28
Ammodytes sp.	430,24	0,00	430,24	6,28	0,00	6,28
Myoxocephalus scorpius	18,75	0,00	18,75	4,64	0,00	4,64
Raja radiata	9,85	0,55	10,40	3,31	0,50	3,81
Triglops nybelini	442,81	1,02	443,82	3,67	0,01	3,67
Micromesistius poutassou	8,62	6,94	15,56	1,27	0,72	1,99
Hippoglossus hippoglossus	0,06	0,02	0,08	1,47	0,36	1,83
Anarhichas lupus	5,46	2,57	8,02	1,12	0,52	1,64
Argentina silus	6,71	8,29	14,99	0,27	1,33	1,61
Brosme brosme	0,05	0,73	0,78	0,16	1,05	1,21
Molva dipterygius	0,00	0,79	0,79	0,00	0,96	0,96
Leptoclinus maculatus	288,02	0,00	288,02	0,90	0,00	0,90
Raja fyllae	0,07	1,11	1,18	0,02	0,83	0,85
Sebastes sp.	9,46	20,89	30,35	0,27	0,55	0,82
Eumicrotremus spinosus	24,18	0,14	24,31	0,51	0,00	0,52
Benthosema glaciale	281,26	7,59	288,85	0,50	0,02	0,51
Coryphaenoides rupestris	4,88	0,00	4,88	0,46	0,00	0,46
Lycodes vahlii	7,79	0,23	8,02	0,44	0,00	0,44
Trisopterus esmarkii	0,03	11,14	11,17	0,00	0,44	0,44
Lycodes eudipleurostictus	3,37	0,49	3,86	0,28	0,03	0,31
Artediellus atlanticus	11,90	11,20	23,10	0,16	0,13	0,29
Gadus ogac	0,57	0,00	0,57	0,27	0,00	0,27
Lycodes seminudus	2,94	0,00	2,94	0,24	0,00	0,24
Macrourus berglax	0,53	0,05	0,58	0,07	0,17	0,24
Petromyzon norvegicus	0,00	0,56	0,56	0,00	0,22	0,22
Gonatus sp.	14,77	1,01	15,78	0,20	0,01	0,21
Molva molva	0,00	0,13	0,13	0,00	0,19	0,19
Triglops pingelii	27,22	0,00	27,22	0,17	0,00	0,17
Lycodes sp.	4,52	0,03	4,55	0,16	0,00	0,16
Lumpenus lampretaeformis	7,72	0,00	7,72	0,15	0,00	0,15
Triglops murrayi	2,57	2,36	4,93	0,05	0,08	0,13
Myctophidae	29,70	0,00	29,70	0,11	0,00	0,11
Dipterus linteus	0,03	0,00	0,03	0,11	0,00	0,11
Cyclopterus lumpus	0,11	0,03	0,13	0,08	0,02	0,10
Lycodes esmarkii	0,00	0,20	0,20	0,00	0,09	0,09

Hyas sp.	7,89	0,00	7,89	0,09	0,00	0,09
Notoscopelus kroeyeri	3,39	0,88	4,27	0,06	0,02	0,07
Gaidropsarus ensis	0,44	0,00	0,44	0,07	0,00	0,07
Gymnacanthus tricuspis	0,74	0,00	0,74	0,07	0,00	0,07
Eumesogrammus praecisus	1,96	0,00	1,96	0,07	0,00	0,07
Careproctus reinhardtii	1,55	0,43	1,98	0,05	0,01	0,06
Lycodes reticulatus	0,29	0,55	0,84	0,02	0,03	0,05
Myoxocephalus scorpioides	0,11	0,00	0,11	0,05	0,00	0,05
Arctozenius rissoii	1,25	0,09	1,33	0,04	0,00	0,04
Melanogrammus aeglefinus	0,04	0,01	0,05	0,03	0,01	0,03
Cottunculus microps	0,02	0,49	0,51	0,00	0,03	0,03
Gaidropsarus argentatus	0,09	0,69	0,77	0,01	0,02	0,03
Myctophum punctatum	4,55	0,00	4,55	0,03	0,00	0,03
Lampanyctus macdonaldi	3,61	0,00	3,61	0,02	0,00	0,02
Gonatus fabricii	1,10	0,00	1,10	0,02	0,00	0,02
Sebastes viviparus	0,01	0,19	0,21	0,00	0,02	0,02
Antimora rostrata	0,11	0,00	0,11	0,02	0,00	0,02
Raja hyperborea	0,12	0,00	0,12	0,02	0,00	0,02
Myxine glutinosa	0,27	0,14	0,41	0,01	0,01	0,01
Liparis gibbus	0,29	0,00	0,29	0,01	0,00	0,01
Liparis fabricii	0,74	0,02	0,77	0,01	0,00	0,01
Lycenchelys kolthoffi	0,18	0,86	1,05	0,00	0,01	0,01
Rossia sp.	0,91	0,25	1,16	0,01	0,00	0,01
Bathypolypus sp.	0,43	0,02	0,45	0,01	0,00	0,01
Leptagonus decagonus	1,35	0,16	1,51	0,01	0,00	0,01
Synapobranchus kaupi	0,44	0,00	0,44	0,01	0,00	0,01
Lithodes maja	0,07	0,00	0,07	0,01	0,00	0,01
Bathylagus euryops	0,86	0,03	0,89	0,01	0,00	0,01
Centroscyllium fabricii	0,19	0,00	0,19	0,01	0,00	0,01
Stomias boa	0,32	0,00	0,32	0,01	0,00	0,01
New Species No 1	0,12	0,03	0,15	0,00	0,00	0,00
Icelus sp.	0,83	0,00	0,83	0,00	0,00	0,00
Icelus bicornis	0,47	0,03	0,51	0,00	0,00	0,00
Lepidion eques	0,00	0,02	0,02	0,00	0,00	0,00
Lycodes paamiuti	0,00	0,08	0,08	0,00	0,00	0,00
Squid	0,00	0,27	0,27	0,00	0,00	0,00
Arctogadus glacialis	0,02	0,09	0,10	0,00	0,00	0,00
Gymnelus retrodorsalis	0,00	0,44	0,44	0,00	0,00	0,00
Gymnelus viridis	0,06	0,43	0,49	0,00	0,00	0,00
Octopodae	0,00	0,09	0,09	0,00	0,00	0,00
Rossia moelleri	0,03	0,13	0,16	0,00	0,00	0,00
Icelus spatula	0,33	0,03	0,36	0,00	0,00	0,00
Caulophryne jordani	0,00	0,03	0,03	0,00	0,00	0,00
Triglops sp.	0,12	0,00	0,12	0,00	0,00	0,00
Lycodes squamiventer	0,00	0,05	0,05	0,00	0,00	0,00
Lycenchelys sarsi	0,15	0,10	0,24	0,00	0,00	0,00
Serrivomer beani	0,04	0,03	0,06	0,00	0,00	0,00
Ammodytes dubius	0,11	0,00	0,11	0,00	0,00	0,00
Aspidophoroides monopterygius	0,36	0,00	0,36	0,00	0,00	0,00
Lampadena speculigera	0,03	0,00	0,03	0,00	0,00	0,00
Liparis tunicatus	0,04	0,00	0,04	0,00	0,00	0,00
Lampanyctus sp.	0,08	0,00	0,08	0,00	0,00	0,00

Melamphaidae	0,03	0,00	0,03	0,00	0,00	0,00
Cephalopoda	0,03	0,00	0,03	0,00	0,00	0,00
<i>Munida tenuimana</i>	0,00	0,12	0,12	0,00	0,00	0,00
<i>Bathypolypus arcticus</i>	0,00	0,00	0,00	0,00	0,00	0,00
<i>Microstomus kitt</i>	0,00	0,00	0,00	0,00	0,00	0,00
<i>Polyipnus polli</i>	0,00	0,02	0,02	0,00	0,00	0,00
<i>Argyropelecus hemigymnus</i>	0,00	0,02	0,02	0,00	0,00	0,00
<i>Artediellus uncinatus</i>	0,02	0,00	0,02	0,00	0,00	0,00
<i>Benthoctopus sp.</i>	0,08	0,00	0,08	0,00	0,00	0,00
<i>Anarhichas sp.</i>	0,08	0,00	0,08	0,00	0,00	0,00
<i>Cyclothona microdon</i>	0,02	0,00	0,02	0,00	0,00	0,00
<i>Gonatus sp.</i>	0,01	0,00	0,01	0,00	0,00	0,00
Liparidae	0,05	0,00	0,05	0,00	0,00	0,00
<i>Maurolicus muelleri</i>	0,02	0,00	0,02	0,00	0,00	0,00
<i>Melanolagus bericoides</i>	0,01	0,00	0,01	0,00	0,00	0,00
<i>Protomyctophum arcticum</i>	0,20	0,00	0,20	0,00	0,00	0,00
<i>Rouleina maderensis</i>	0,03	0,00	0,03	0,00	0,00	0,00