

NOT TO BE CITED WITHOUT PRIOR  
REFERENCE TO THE AUTHOR(S)

Northwest Atlantic



Fisheries Organization

Serial No. N6557

NAFO SCR Doc. 16/16

## SCIENTIFIC COUNCIL MEETING – JUNE 2016

Results for the Atlantic cod, roughhead grenadier, redfish, thorny skate and black dogfish of the Spanish Survey in the NAFO Div. 3L for the period 2003-2015

by

Esther Román, Ángeles Armesto, Diana González-Troncoso and Concepción González-Iglesias.

Instituto Español de Oceanografía  
P.O. Box 1552. Vigo, Spain  
e-mail: [esther.roman@vi.ieo.es](mailto:esther.roman@vi.ieo.es)

### Abstract

Since 2003, a stratified random spring bottom trawl survey was conducted by Spain in Division 3L of NAFO Regulatory Area (Flemish Pass). The surveys were carried out by the R/V "Vizconde de Eza" using bottom trawl net type *Campelen*. Entire series of mean catches, biomass and length distribution for Atlantic cod, roughhead grenadier, redfish, thorny skate and black dogfish are presented for the period 2003-2015.

KEYWORDS: Survey, Flemish Pass, Atlantic Cod, Roughhead grenadier, Redfish, Thorny skate, Black dogfish.

### Material and Methods

The Spanish surveys in Div. 3L of NAFO Regulatory Area (Flemish Pass) were initiated by Spain in 2003. The Research vessel "Vizconde de Eza" has carried out the entire surveys series following the same procedures and using the same bottom trawl gear *Campelen 1800*. In 2003 and 2004, the survey did not cover all strata adequately. In 2005, it was not possible to perform the survey due to problems with the winch of the ship; and in 2006, for the first time, an adequate prospecting survey was conducted in Division 3L with over 100 valid hauls. Table 1 shows the number of valid tows, the depth and number of covered strata and the dates of the survey series. To know more details about the technical specifications of the surveys, see Román *et al.*, 2016.

The catch from each haul was sorted out and weighted by species and a randomly selected sample of each species was taken in order to measure it and obtain the length distribution. In 2003 and 2004 the Atlantic cod samples were not sorted out by sex. There are two species of redfish in Division 3L (*Sebastes mentella* and *S. fasciatus*); the external characteristics of both species are very similar, which makes it difficult to distinguish between them and, as a result, they are treated together.

For Atlantic cod, redfish, thorny skate and black dogfish each individual of the sample was measured to the total length to the nearest lower cm and data are given in 2 cm intervals. However, roughhead grenadier individuals were measured from tip of snout to base of first anal-fin ray to the lower ½ cm., in 0.5 cm intervals, as adopted by NAFO in June 1980 (Atkinson, 1991) as a standard measurement for roundnose and roughhead grenadiers; length is presented as pre-anal-fin length (AFL) and data are given in 1 cm intervals.

It is presented the mean catch per haul, the stratified mean catch per haul and the biomass with their variance per year in the period 2003-2015. Length distribution in number per haul stratified mean catches per length, sex and year for these species are presented too. The following formula was used to obtain the biomass from length distribution: Weight=a(Length+0.5)<sup>b</sup> / Weight=a(Length+0.25)<sup>b</sup>.

## Results

### **Atlantic Cod (*Gadus morhua* Linnaeus, 1758)**

NAFO manages 3 cod stocks in Div. 3L, 3M (Flemish Cap) and 3NO (southern Gran Bank). After a dramatic decline of cod during the eighties and nineties, fishing bans were imposed in the 1990s. In recent assessment all stocks remain at a very low level although spawning biomass has increased in recent years. In 2010, after a decade long moratorium, a cod fishery on the Flemish Cap (Div. 3M) was re-opened but the moratoria (no directed fishery) continues for Div. 3NO and Div. 3L. (NAFO, 2015).

The cod fishery on Flemish Cap has traditionally been a directed fishery by Portuguese trawlers and gillnetters, Spanish pair-trawlers and Faroese longliners. Cod has also been taken as bycatch in the directed redfish fishery by Portuguese trawlers. In Div. 3L and 3NO, this stock has been under moratorium to directed fishing since February 1994. By-catch occurs primarily in the yellowtail flounder, skate and redfish fisheries.

#### **Mean catches and biomass**

Table 2 shows the swept area, the tow number, the mean catches and their variance per haul by stratum for Atlantic cod. Table 3 and Figure 1 present the stratified mean catches by stratum and year with their total variance. The entire time series (2003-2015) of biomass and their total variance for Atlantic cod are presented in Table 4 and Figure 2. Estimated parameters values of length-weight relationship are presented in Table 5 (2007-2015).

Figure 3 shows a map with the distribution of Atlantic cod catches per haul in 2015 Spanish 3L survey. Atlantic cod indices show a great variation, due to a few hauls in which the presence of cod was very high, however there is no clear trend along the whole period (2003-2015). Stratified mean catch and biomass decreased from 2003 to 2004; then, the values of these indices increased in 2006 and declined briefly again in 2007. A great increase is shown in 2008 but this was due to a single haul in which the presence of cod was very high (1298.5 kg). The great value of the variance in some years is produced by the tows with a large catch. In 2009 declined again and since then an increasing trend in the biomass can be seen. In 2011 the biomass reaches the highest value in the time series. The highest values in the estimated biomass have been observed in the shallow strata, in a range of depth from 93 to 274 meters. In 2012, the biomass decreases at the same level than in 2008, increased briefly in 2013 and 2014 declined again. In 2015, the biomass increases at the same level than in 2009.

#### **Length distribution**

Table 6 presents the length distribution of stratified mean catches per haul for this species, by sex and year (2008-2015), with the number of samples in which there were length measurements, the sampled catch, the total number of individuals measured in each sample and the range of lengths achieved, as well as the total catch of this species and the total hauls made in the survey. In Figures 4 and 5 the evolution throughout the period can be followed.

In this period, individuals between 12 and 25 cm can be seen although in 2004 there was no presence of individuals below 24 cm. In general all lengths presence is very low, even it is very difficult to follow the modal values. In 2008 we have a good presence of individuals between 26 and 33 cm, probably due to the haul with great catch of that year, 29 cm is the mode in the length distribution. In 2009 the dominant lengths were between 36 and 41 cm (mode = 37 cm). In 2010 the mode was 44 cm with the dominant length between 40 and 47 cm. In 2011 the mode observed was 51 cm and the dominant lengths were between 47 and 55 cm. and in 2012 the dominant lengths were between 34 and 56 cm (mode = 46 cm). In 2013 we have the best presence of individuals between 12-25 cm and there were two modes, one in 28 cm and another in 47cm with

the dominant length between 23-31 and 41-58 cm. In 2014 the mode is in 38 cm. and in 2015 the dominant lengths were between 22 and 44 cm (mode = 30-32 cm).

In last years no good recruitments were seen.

#### **Roughhead grenadier (*Macrourus berglax* Lacépède, 1802)**

The stock structure of this species in the North Atlantic remains unclear because there is little information on the number of different populations that may exist and their relationship. Roughhead grenadier is distributed throughout

NAFO Subareas 0 to 3 in depths between 300 and 2 000 m.. There is no directed fishery for this species and most catches are taken as by-catch in Greenland halibut fishery in Subareas 2 and 3. Most of the catches were taken in Divs. 3LMN by EU-Spain, EU-Estonia and EU-Portugal fleets.

Roughhead grenadier is taken mainly in Div. 3LMN of NAFO Regulatory Area. The highest level of observed catches was reached in 1998. Survey indices indicate a stable or declining stock in recent years. Fishing mortality indices have remained at low levels since 2005. Roughhead grenadier is not a regulated species (NAFO, 2015).

#### **Mean catches and biomass**

Roughhead grenadier haul mean catches by stratum are presented in Table 7; swept area, number of hauls and SD are also shown in this table. Stratified mean catches per tow by stratum and year and their variance are presented in Table 8. The entire time series (2003-2015) of biomass and their SD estimates of this species are shown in Table 9 and length-weight relationships are shown in Table 5 (2007-2015).

The indices of roughhead grenadier show no clear trend throughout the whole period, with an increasing in 2004 compared to 2003 and then remains stable (2006-2007). In 2008 the biomass increased, reaching the highest value of the series, but declining afterwards in the period 2009-2012. Biomass reaches the lowest value in the time series in the 2012. There was an increase in 2013, reaching the maximum value since 2010 but still lower than the 2009. In 2014 the values of these indices declined again.

In 2015 the biomass increased, reaching the second highest value of the series (Fig. 6 and 7). Figure 3 shows a map with the distribution of roughhead grenadier catches per haul in 2014 Spanish 3L survey.

#### **Length distribution**

Table 10 shows the stratified mean catches per haul length distribution, for roughhead grenadier, by sex and year (2008-2015), with the number of samples in which there was length measurements, the sampled catch, the total number of individuals measured in these samples and the range of lengths found. The total catch of this species and the total hauls made in the survey are shown too. In Figures 5 and 8 the evolution along the years can be followed. A slight recruitment can be seen in all period but it was quite good in 2013 (mode =16 cm). In 2015, the mode observed was 16.5 cm and the dominant lengths were between 5.5 and 24.5 cm.

Females attain larger lengths than males in all years.

#### **Redfish (*Sebastes* spp. Cuvier, 1829)**

There are two species of redfish that have been commercially fished in Div. 3LN, *Sebastes fasciatus* (Acadian redfish) and *S. mentella* (deepwater redfish). The external characteristics are very similar, making them difficult to distinguish, and as a consequence they are reported collectively as "redfish" in the commercial fishery. The redfish stocks in 3LN, 3M, 3O, as well as those in Subarea 2 and Div. 1F+3K are managed by NAFO. From 1998-2010 a moratorium was on 3LN stocks (no directed fishery) and the fishery was reopened in 2010 and have reached just over 6 300 t and 5 781 t in 2013 and 2014, the highest level recorded on 20 years. Catches from EU-Portugal, Russian and Canadian fleets justified most of the increase on the redfish catch observed on both Divisions 3L and 3N (NAFO, 2015).

#### **Mean catches and biomass**

Table 11 shows the swept area, the tow number, the mean catches per haul and year (2011-2015) and their variance for redfish. Table 12 and Figure 9 present the stratified mean catches per stratum with the total

variance per year. Figure 3 shows a map with the distribution of redfish catches per haul in 2015 Spanish 3L survey.

Table 13 and Figure 10 show the biomass estimate per swept area per stratum and their total variance by year and also the estimated abundance. Redfish shows a great annual variability probably due to its pelagic habitat. Redfish biomass indices decreased in 2004, 2007 and 2011 with a great decrease in 2013. In 2014 the biomass remains at the same value as the last year; and they increased in 2006, 2008 and 2009 with a sharp increase in 2010. In 2012, the redfish indices show the greater increasing reaching the highest value of the series (this was due to some hauls in which the presence of redfish was very high). Redfish biomass indices decreased in 2015 again. The length-weight relationships are presented in Table 5 (2007-2015).

#### Length distribution

Table 14 presents the length distribution of the stratified mean catches per haul for redfish, by sex and year (2008-2015), with the number of samples in which there was length measurements, the sampled catch, the total number of individuals measured in these samples and the range of lengths found. The total catch of this species and the total hauls made in the survey are also shown. In Figures 5 and 11 the evolution along the years can be followed. The highest proportions of small individuals in the catches (smaller than 20 cm) were found in the period 2010-2012. In 2015, the mode observed was 24 cm and the dominant lengths were between 18 and 34 cm.

#### **Thorny skate (*Amblyraja radiata* Donovan, 1808)**

Commercial catches of skates comprise a mix of skate species. However, thorny skate dominates, comprising about 95% of the skate species taken in the Canadian and EU-Spain catches. Thus, the skate fishery on the Grand Banks can be considered a fishery for thorny skate. In 2005, NAFO Fisheries Commission established a TAC of 13 500 t for thorny skate in Div. 3LNO. In 2010 and 2011, the TAC for Div. 3LNO has been reduced to 12 000 t. The TAC was further reduced to 8 500 t for 2012, and to 7 000 t for 2013-2015. Based on the continuous distribution and lack of physical barriers between Div. 3LNO and Subdiv. 3Ps, thorny skate in Div. 3LNOPs is considered to constitute a single stock. Div. 3LNO is managed by NAFO. The stock has been increasing very slowly from low levels since the mid-1990s (NAFO, 2015).

#### Mean catches and biomass

Table 15 shows the swept area, the tow number, the mean catches per haul and year (2011-2015) and their variance for thorny skate. Table 16 presents the length-weight relationships (2007-2015). Table 17 and Figure 12 present the stratified mean catches per stratum with the total variance per year. Table 18 and Figure 13 present the biomass per swept area by stratum and year, their total variance per year and the abundance index. The indices of the thorny skate decreased from 2003 to 2004, increased in 2006-2007 and decreased again in the period 2008-2011. In 2012 the indices of the thorny skate increased and they slightly decreased again in 2013. The thorny skate indices increased slightly since 2014.

Figure 3 shows a map with the distribution of thorny skate catches per haul in 2015 Spanish 3L survey.

#### Length distribution

Table 19 presents the stratified mean catches per haul length distribution for this species, by sex and year (2008-2015), with the number of samples in which there was length measurements, the sampled catch, the total number of individuals measured in these samples and the range of lengths achieved, as well as the total catch of this species and the total hauls made in the survey. In Figures 14 and 15, the evolution along the years can be followed. The highest proportion of small thorny skate in the catches was in 2007 and 2015. In 2015, the modes observed were 30 and 72 cm and the dominant lengths were between 22-36 and 50-80 cm.

#### **Black dogfish (*Centroscyllium fabricii* Reinhardt, 1825)**

Black dogfish is present in all Divisions, but is more abundant in Div. 3NO and in depths greater than 900 m. Black dogfish is not a regulated species and commercial catches of this species are mainly a by-catch of the Greenland halibut fishery in Div. 3LMNO (González-Costas *et al.*, 2006).

### Mean catches and biomass

Black dogfish haul mean catches by stratum are presented in Table 20, including swept area, number of hauls and SD. Stratified mean catches per tow by stratum and year and their variance are presented in Table 21. The entire time series (2003-2015) of biomass and their SD estimates of black dogfish are shown in Table 22. Length-weight relationships are presented in Table 16 (2007-2015).

The abundance and biomass present the same trend as mean catches. Biomass estimated from the 3L survey displays an increasing trend since 2004 until 2007 and decreased in 2008, 2009 and 2012, increasing again in last years. In 2003, the catches occurred only in two strata (745 and 749), in which the catches were much different, what explain why the variance in that year is so large. In 2015 the biomass increased, reaching the highest value of the series. (Fig. 16 and 17). Figure 3 shows a map with the distribution of black dogfish catches per haul in 2015 Spanish 3L survey.

### Length distribution

Table 23 presents the length distribution of the stratified mean catches per haul for black dogfish, by sex and year (2008-2015), with the number of samples in which there was length measurements, the sampled catch, the total number of individuals measured in these samples and the range of lengths met. The total catch of this species and the total hauls made in the survey are shown too. In Figures 15 and 18 the evolution throughout the years can be followed.

There is no presence of small individual (smaller 37 cm). Size compositions are mainly between 46 and 76 cm of length. In 2015 the observed mode was 62 cm and the dominant lengths were between 46 and 72 cm.

### Acknowledge

The data used in this paper have been funded by the EU through the European Maritime and Fisheries Fund (EMFF) within the National Program of collection, management and use of data in the fisheries sector and support for scientific advice regarding the Common Fisheries Policy.

### References

- ATKINSON, D. B. 1991. Relationships Between Pre-anal Fin Length and Total Length of Roughhead Grenadier (*Macrourus berglax* Lacépède) in the Northwest Atlantic. *J. Northw. Atlan. Fish. Sci.*, **11**: 7-9
- GONZÁLEZ-COSTAS, F., D. GONZÁLEZ-TRONCOSO, M. CASAS, and G. RAMILO. 2006. Spiny Dogfish (*Squalus acanthias*) and Black Dogfish (*Centroscyllium fabricii*) Spanish Data (Surveys and Fishery) in NAFO Divisions 3LMNO. *NAFO SCR Doc.*, No. 30, Serial No. N5250, 10 p.
- NAFO, 2015. Report of the Scientific Council Meeting, 29 May - 11 June 2015.
- ROMÁN, E., C. GONZÁLEZ-IGLESIAS, D. GONZÁLEZ-TRONCOSO, and M. ÁLVAREZ. 2016. Results for the Spanish Survey in the NAFO Regulatory Area of Division 3L for the period 2003-2015. *NAFO SCR Doc.*, No. XX, Serial No NXXXX.

Table 1. Spanish bottom trawl surveys in NAFO Division 3L for the period 2003-2015.

Year	Vessel	Valid tows	Depth strata covered (m)	Surveyed strata (no.)	Dates
2003	R/V "Vizconde de Eza"	39	118-1100	17	June 2 - June 6, June 29
2004	R/V "Vizconde de Eza"	50	141-1452	23	August 7 - August 15
2005	-	-	-	-	-
2006	R/V "Vizconde de Eza"	100	116-1449	24	July 31 - August 18
2007	R/V "Vizconde de Eza"	94	119-1449	24	July 23 - August 11
2008	R/V "Vizconde de Eza"	100	105-1455	24	July 24 - August 11
2009	R/V "Vizconde de Eza"	98	111-1458	24	July 25 - August 12
2010	R/V "Vizconde de Eza"	97	119-1462	24	July 25 - August 14
2011	R/V "Vizconde de Eza"	89	115-1419	24	August 10 - August 24
2012	R/V "Vizconde de Eza"	98	112-1478	24	July 30 - August 18
2013	R/V "Vizconde de Eza"	100	117-1420	24	July 30 - August 19
2014	R/V "Vizconde de Eza"	102	104-1411	24	July 30 - August 19
2015	R/V "Vizconde de Eza"	97	112-1458	24	July 28 - August 17

Table 2.- Swept area, number of hauls and Atlantic cod mean catch (Kg) and SD (\*\*) by stratum. Spanish Survey on NAFO Div. 3L in the period 2011-2015, on board R/V "Vizconde de Eza".

Stratum	2011			2012			2013			2014			2015		
	Swept Tow area No.	Mean catch	SD	Swept Tow area No.	Mean catch	SD	Swept Tow area No.	Mean catch	SD	Swept Tow area No.	Mean catch	SD	Swept Tow area No.	Mean catch	SD
<b>385</b>	0.0229 2	93.750	118.723	0.0225 2	4.820	2.871	0.0229 2	4.556	4.144	0.0225 2	8.360	8.712	0.0236 2	8.271	9.279
<b>387</b>	0.0450 4	36.505	32.228	0.0450 4	6.760	4.899	0.0450 4	92.938	97.705	0.0461 4	39.932	36.630	0.0458 4	17.995	19.617
<b>388</b>	0.0563 5	15.241	14.829	0.0570 5	162.020	264.788	0.0570 5	91.360	68.284	0.0585 5	28.395	23.211	0.0574 5	23.188	18.824
<b>389</b>	0.0675 6	26.796	42.096	0.0799 7	34.169	26.422	0.0791 7	74.413	71.762	0.0814 7	26.084	37.415	0.0814 7	70.861	40.597
<b>390</b>	0.1009 9	217.889	231.959	0.1354 12	43.245	27.872	0.1358 12	42.393	23.638	0.1369 12	20.592	24.738	0.1260 11	10.735	8.941
<b>391</b>	0.0458 4	150.275	91.993	0.0458 4	44.280	47.163	0.0450 4	14.288	19.423	0.0465 4	13.695	17.396	0.0465 4	24.148	22.449
<b>392</b>	0.0229 2	3.268	3.129	0.0225 2	13.470	4.992	0.0225 2	27.297	2.626	0.0225 2	1.485	0.092	0.0229 2	2.669	0.471
<b>729</b>	0.0338 3	0.000	0.000	0.0338 3	0.000	0.000	0.0341 3	0.759	1.314	0.0338 3	0.000	0.000	0.0345 3	0.000	0.000
<b>730</b>	0.0334 3	0.000	0.000	0.0338 3	0.000	0.000	0.0334 3	0.000	0.000	0.0345 3	0.000	0.000	0.0345 3	0.000	0.000
<b>731</b>	0.0334 3	0.000	0.000	0.0341 3	0.000	0.000	0.0334 3	0.173	0.300	0.0345 3	0.000	0.000	0.0345 3	1.540	2.667
<b>732</b>	0.0454 4	0.000	0.000	0.0454 4	0.000	0.000	0.0450 4	0.000	0.000	0.0454 4	0.000	0.000	0.0465 4	0.000	0.000
<b>733</b>	0.0454 4	0.545	0.642	0.0454 4	0.000	0.000	0.0450 4	5.008	7.845	0.0458 4	0.107	0.213	0.0454 4	0.349	0.492
<b>734</b>	0.0225 2	0.000	0.000	0.0233 2	0.000	0.000	0.0221 2	0.000	0.000	0.0225 2	0.085	0.120	0.0225 2	0.000	0.000
<b>741</b>	0.0218 2	0.000	0.000	0.0218 2	0.000	0.000	0.0221 2	0.000	0.000	0.0225 2	0.000	0.000	0.0236 2	0.000	0.000
<b>742</b>	0.0225 2	0.000	0.000	0.0206 2	0.000	0.000	0.0218 2	0.000	0.000	0.0221 2	0.000	0.000	0.0233 2	0.000	0.000
<b>743</b>	0.0221 2	0.000	0.000	0.0206 2	0.000	0.000	0.0218 2	0.000	0.000	0.0221 2	0.000	0.000	0.0233 2	0.000	0.000
<b>744</b>	0.0221 2	0.000	0.000	0.0221 2	0.000	0.000	0.0221 2	0.000	0.000	0.0225 2	0.000	0.000	0.0225 2	0.000	0.000
<b>745</b>	0.0446 4	0.000	0.000	0.0570 5	0.000	0.000	0.0559 5	0.000	0.000	0.0578 5	0.000	0.000	0.0578 5	0.000	0.000
<b>746</b>	0.0566 5	0.000	0.000	0.0675 6	0.000	0.000	0.0675 6	0.000	0.000	0.0683 6	0.000	0.000	0.0686 6	0.000	0.000
<b>747</b>	0.0893 8	0.000	0.000	0.1121 10	0.000	0.000	0.1125 10	0.000	0.000	0.1125 10	0.000	0.000	0.1028 9	0.000	0.000
<b>748</b>	0.0221 2	0.000	0.000	0.0225 2	0.000	0.000	0.0225 2	0.000	0.000	0.0229 2	0.000	0.000	0.0233 2	0.000	0.000
<b>749</b>	0.0221 2	0.000	0.000	0.0221 2	0.000	0.000	0.0225 2	0.000	0.000	0.0225 2	0.000	0.000	0.0225 2	0.000	0.000
<b>750</b>	0.0668 6	0.000	0.000	0.0885 8	0.000	0.000	0.0896 8	0.000	0.000	0.0904 8	0.000	0.000	0.0934 8	0.000	0.000
<b>751</b>	0.0334 3	0.000	0.000	0.0218 2	0.000	0.000	0.0446 4	0.000	0.000	0.0334 3	0.000	0.000	0.0341 3	0.000	0.000

$$(**) SD = \frac{\sum (x_i - \bar{x})}{n-1}$$

Table 3.- Stratified mean catches (Kg) of Atlantic cod by stratum and year (2003-2015) and SD. Research Vessel *Vizconde de Eza*. n.s. means stratum not surveyed. In 2003: the data correspond to 69% of the total area prospected in 2006-2015.

Stratum	Survey												
	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
385	7.26	53.10	-	210.34	98.53	713.96	623.63	91.45	11062.50	568.76	537.61	986.48	975.98
387	1123.84	482.56	-	101.12	509.82	1378.75	5940.16	878.72	9345.28	1730.43	23792.19	10222.59	4606.72
388	2809.59	468.74	-	2509.00	2653.87	6663.55	2646.51	22129.72	5441.04	57841.14	32615.52	10136.94	8278.04
389	429.34	259.59	-	5386.31	2118.59	15536.35	20804.94	76812.24	13639.08	17391.88	37876.07	13276.54	36068.25
390	0.00	0.00	-	65.94	1115.80	7076.10	18289.28	30271.32	177579.44	35245.01	34550.09	16782.48	8749.32
391	47.00	0.00	-	4043.18	3153.47	96519.44	18404.45	40629.15	42377.55	12486.96	4029.29	3862.06	6809.67
392	58.00	1916.68	-	296.53	2027.75	0.00	9.14	10248.60	473.79	1953.15	3958.07	215.33	387.01
729	234.36	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	141.11	0.00	0.00
730	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
731	4839.48	107.03	-	0.00	110.16	28.08	0.00	53.28	0.00	0.00	37.44	0.00	332.64
732	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
733	n.s.	0.00	-	0.00	99.84	0.00	0.00	0.00	127.59	0.00	1171.76	24.92	81.67
734	n.s.	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	13.01	0.00
741	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
742	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
743	n.s.	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
744	n.s.	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
745	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
746	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
747	n.s.	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
748	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
749	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
750	n.s.	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
751	n.s.	n.s.	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
TOTAL ( $\bar{y}$ )	9548.87	3287.70	-	12612.40	11887.83	127916.23	66718.10	181114.48	260046.27	127217.33	138709.14	55520.34	66289.29
SD	0.57	0.30	-	0.55	0.42	13.89	2.75	9.17	10.15	6.72	3.47	1.74	1.48

Table 4.- Survey estimates (by the swept area method) of Atlantic cod biomass (t.) by stratum and year and their SD on NAFO Div. 3L (R/V *Vizconde de Eza*). n.s. means stratum not surveyed. In 2003, the data correspond to 69% of the total area prospected in 2006-2015.

Stratum	Survey												
	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
385	1	5	-	18	9	62	55	8	967	51	47	88	83
387	98	45	-	9	45	127	542	77	831	154	2115	887	403
388	253	45	-	222	236	596	238	1941	484	5074	2861	866	721
389	38	23	-	474	188	1394	1815	6763	1212	1524	3351	1142	3103
390	0	0	-	6	99	609	1599	2667	15844	3124	3054	1471	764
391	4	0	-	359	280	8509	1609	3582	3705	1092	358	332	586
392	5	179	-	26	180	0	1	911	41	174	352	19	34
729	22	0	-	0	0	0	0	0	0	0	12	0	0
730	0	0	-	0	0	0	0	0	0	0	0	0	0
731	423	9	-	0	10	3	0	5	0	0	3	0	29
732	0	0	-	0	0	0	0	0	0	0	0	0	0
733	n.s.	0	-	0	9	0	0	0	11	0	104	2	7
734	n.s.	0	-	0	0	0	0	0	0	0	0	1	0
741	0	0	-	0	0	0	0	0	0	0	0	0	0
742	0	0	-	0	0	0	0	0	0	0	0	0	0
743	n.s.	0	-	0	0	0	0	0	0	0	0	0	0
744	n.s.	0	-	0	0	0	0	0	0	0	0	0	0
745	0	0	-	0	0	0	0	0	0	0	0	0	0
746	0	0	-	0	0	0	0	0	0	0	0	0	0
747	n.s.	0	-	0	0	0	0	0	0	0	0	0	0
748	0	0	-	0	0	0	0	0	0	0	0	0	0
749	0	0	-	0	0	0	0	0	0	0	0	0	0
750	n.s.	0	-	0	0	0	0	0	0	0	0	0	0
751	n.s.	n.s.	-	0	0	0	0	0	0	0	0	0	0
TOTAL	844	306	-	1114	1057	11300	5859	15953	23095	11192	12258	4809	5729
SD	222	180	-	315	245	7745	1556	5265	5833	3877	1984	1001	829

Table 5.- Length-weight relationships in the calculation of biomass, for Division 3L (out ZEE Canada), 2006-2015 for Atlantic cod, roughhead grenadier and redfish. The equation is Weight=a(Length+0.5)b or Weight=a(Length+0.25)b. To calculate the parameters for the indeterminate individuals, we used the total data (males+females+indeterminate individuals).

Atlantic cod							Roughhead grenadier							Redfish						
Year	Sex	L-W Equations	N	r <sup>2</sup>	Sex	L-W Equations	N	r <sup>2</sup>	Sex	L-W Equations	N	r <sup>2</sup>	All	L-W Equations	N	r <sup>2</sup>				
2007	All	$W = 0.0055 L^{3.1370}$	225	0.983		All	$W = 0.0885 L^{2.9691}$	1950	0.9895		All	$W = 0.0080 L^{3.1588}$	881	0.9842						
	Males	$W = 0.0061 L^{3.1114}$	107	0.991		Males	$W = 0.0946 L^{2.9435}$	754	0.9859		Males	$W = 0.0140 L^{2.9836}$	432	0.9858						
	Females	$W = 0.0047 L^{3.1750}$	118	0.9735		Females	$W = 0.0877 L^{2.9727}$	1165	0.9897		Females	$W = 0.0133 L^{3.0115}$	392	0.9868						
2008	All	$W = 0.0083 L^{3.0479}$	819	0.9856		All	$W = 0.1237 L^{2.8681}$	1773	0.9871		All	$W = 0.0142 L^{2.9849}$	699	0.9701						
	Males	$W = 0.0083 L^{3.0493}$	403	0.9855		Males	$W = 0.1174 L^{2.8868}$	754	0.9832		Males	$W = 0.0337 L^{2.7219}$	338	0.9343						
	Females	$W = 0.0084 L^{3.0467}$	416	0.9856		Females	$W = 0.1144 L^{2.8938}$	1024	0.988		Females	$W = 0.0314 L^{2.7511}$	340	0.9412						
2009	All	$W = 0.0084 L^{3.0256}$	684	0.9824		All	$W = 0.0903 L^{2.9583}$	1457	0.9911		All	$W = 0.0083 L^{3.1392}$	818	0.9854						
	Males	$W = 0.0089 L^{3.0085}$	296	0.9824		Males	$W = 0.0847 L^{2.9803}$	540	0.9871		Males	$W = 0.0135 L^{2.9882}$	354	0.9738						
	Females	$W = 0.0083 L^{3.0299}$	388	0.9821		Females	$W = 0.0927 L^{2.9505}$	899	0.9904		Females	$W = 0.0174 L^{2.9204}$	389	0.9763						
2010	All	$W = 0.0086 L^{3.0302}$	756	0.980		All	$W = 0.1006 L^{2.9369}$	1539	0.991		All	$W = 0.0110 L^{3.0593}$	808	0.9859						
	Males	$W = 0.0076 L^{3.0636}$	364	0.980		Males	$W = 0.0909 L^{2.9770}$	547	0.984		Males	$W = 0.0153 L^{2.9565}$	372	0.9754						
	Females	$W = 0.0095 L^{3.0027}$	392	0.979		Females	$W = 0.1071 L^{2.9152}$	947	0.990		Females	$W = 0.0161 L^{2.9484}$	397	0.9706						
2011	All	$W = 0.0090 L^{3.0101}$	1421	0.9874		All	$W = 0.0962 L^{2.9550}$	1545	0.9899		All	$W = 0.0105 L^{3.0803}$	1218	0.9882						
	Males	$W = 0.0102 L^{2.9790}$	682	0.9852		Males	$W = 0.1018 L^{2.9403}$	543	0.9796		Males	$W = 0.0129 L^{3.0158}$	529	0.9836						
	Females	$W = 0.0082 L^{3.0334}$	739	0.9892		Females	$W = 0.1169 L^{2.8873}$	913	0.9884		Females	$W = 0.0109 L^{3.0768}$	559	0.9855						
2012	All	$W = 0.0106 L^{2.9627}$	878	0.982		All	$W = 0.1070 L^{2.9148}$	1607	0.988		All	$W = 0.0126 L^{3.0228}$	978	0.9847						
	Males	$W = 0.0109 L^{2.9573}$	403	0.982		Males	$W = 0.1008 L^{2.9374}$	609	0.980		Males	$W = 0.0135 L^{2.9979}$	476	0.9856						
	Females	$W = 0.0123 L^{2.9243}$	474	0.980		Females	$W = 0.1081 L^{2.9117}$	934	0.988		Females	$W = 0.0157 L^{2.9616}$	491	0.9806						
2013	All	$W = 0.0072 L^{3.0592}$	1717	0.992		All	$W = 0.0979 L^{2.9309}$	1784	0.991		All	$W = 0.0080 L^{3.1741}$	1130	0.99						
	Males	$W = 0.0071 L^{3.0636}$	785	0.992		Males	$W = 0.0919 L^{2.9562}$	643	0.985		Males	$W = 0.0130 L^{3.0249}$	497	0.9803						
	Females	$W = 0.0073 L^{3.0554}$	932	0.993		Females	$W = 0.0995 L^{2.9248}$	1036	0.991		Females	$W = 0.0132 L^{3.0237}$	522	0.9822						
2014	All	$W = 0.0071 L^{3.0532}$	685	0.990		All	$W = 0.1003 L^{2.9350}$	1604	0.992		All	$W = 0.0094 L^{3.1208}$	925	0.9840						
	Males	$W = 0.0067 L^{3.0666}$	317	0.987		Males	$W = 0.0958 L^{2.9529}$	582	0.987		Males	$W = 0.0161 L^{2.9557}$	424	0.981						
	Females	$W = 0.0076 L^{3.0345}$	365	0.991		Females	$W = 0.1091 L^{2.9071}$	940	0.992		Females	$W = 0.0121 L^{3.0495}$	457	0.9624						
2015	All	$W = 0.0079 L^{3.0271}$	867	0.989		All	$W = 0.1107 L^{2.9089}$	1832	0.993		All	$W = 0.0088 L^{3.1436}$	1088	0.9909						
	Males	$W = 0.0080 L^{3.0280}$	393	0.989		Males	$W = 0.1127 L^{2.9084}$	662	0.987		Males	$W = 0.0148 L^{2.9886}$	500	0.9893						
	Females	$W = 0.0080 L^{3.0264}$	473	0.989		Females	$W = 0.1197 L^{2.8800}$	1097	0.992		Females	$W = 0.0104 L^{3.0946}$	554	0.9898						

Table 6.- Atlantic cod length distribution per haul mean catches by sex and year. Number per stratified mean catches. Spanish Summer Survey on NAFO 3L: 2008-2015 (R/V *Vizconde de Eza*). Indet. means indeterminate.

Length (cm.)	2008				2009				2010				2011				
	M	F	I	T	M	F	I	T	M	F	I	T	M	F	I	T	
<12	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
12	0.01	0.02	0.00	0.04	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
14	0.01	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	
16	0.08	0.03	0.00	0.11	0.01	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
18	0.19	0.15	0.00	0.34	0.00	0.00	0.00	0.00	0.03	0.00	0.00	0.03	0.00	0.01	0.00	0.01	
20	0.14	0.13	0.00	0.27	0.02	0.01	0.00	0.03	0.00	0.00	0.00	0.00	0.01	0.01	0.00	0.03	
22	0.12	0.19	0.00	0.31	0.06	0.06	0.00	0.12	0.01	0.00	0.00	0.01	0.04	0.05	0.00	0.08	
24	1.21	1.36	0.00	2.56	0.08	0.05	0.00	0.13	0.07	0.07	0.00	0.13	0.07	0.16	0.00	0.23	
26	5.14	6.23	0.00	11.37	0.12	0.12	0.00	0.24	0.21	0.24	0.00	0.45	0.37	0.31	0.00	0.68	
28	8.51	10.05	0.00	18.56	0.14	0.18	0.00	0.32	0.49	0.88	0.00	1.37	0.46	0.66	0.00	1.12	
30	6.60	7.42	0.00	14.02	0.20	0.15	0.00	0.36	0.99	1.06	0.00	2.05	0.58	0.71	0.00	1.29	
32	2.99	3.61	0.00	6.60	0.39	0.37	0.00	0.77	1.34	1.23	0.00	2.57	0.67	0.78	0.00	1.45	
34	1.94	0.81	0.00	2.74	0.66	1.04	0.00	1.70	0.87	1.07	0.00	1.95	0.81	0.72	0.00	1.53	
36	0.83	0.78	0.00	1.61	1.11	1.16	0.00	2.26	1.27	1.35	0.00	2.62	0.68	0.75	0.00	1.43	
38	0.32	0.35	0.00	0.67	1.09	1.42	0.00	2.51	1.31	1.44	0.00	2.75	0.71	0.75	0.00	1.46	
40	0.14	0.29	0.00	0.43	0.92	1.07	0.00	1.99	1.65	2.08	0.00	3.72	0.76	1.09	0.00	1.85	
42	0.06	0.37	0.00	0.43	0.49	0.76	0.00	1.25	1.91	2.12	0.00	4.02	0.95	0.86	0.00	1.82	
44	0.13	0.05	0.00	0.19	0.28	0.47	0.00	0.75	1.79	2.52	0.00	4.31	0.99	1.29	0.00	2.28	
46	0.09	0.29	0.00	0.37	0.15	0.37	0.00	0.52	1.60	2.24	0.00	3.85	1.18	1.61	0.00	2.79	
48	0.07	0.24	0.00	0.31	0.04	0.15	0.00	0.18	1.17	1.48	0.00	2.65	1.41	2.14	0.00	3.55	
50	0.06	0.09	0.00	0.16	0.08	0.14	0.00	0.22	0.51	0.95	0.00	1.46	2.26	2.42	0.00	4.68	
52	0.22	0.07	0.00	0.29	0.07	0.13	0.00	0.20	0.28	0.43	0.00	0.71	1.86	2.21	0.00	4.07	
54	0.04	0.06	0.00	0.10	0.07	0.08	0.00	0.15	0.18	0.31	0.00	0.49	1.34	2.00	0.00	3.34	
56	0.04	0.02	0.00	0.06	0.09	0.11	0.00	0.20	0.05	0.21	0.00	0.25	0.71	1.05	0.00	1.75	
58	0.19	0.03	0.00	0.22	0.01	0.13	0.00	0.14	0.12	0.13	0.00	0.26	0.49	0.62	0.00	1.11	
60	0.02	0.02	0.00	0.04	0.02	0.07	0.00	0.09	0.16	0.06	0.00	0.22	0.36	0.32	0.00	0.68	
62	0.05	0.03	0.00	0.09	0.03	0.04	0.00	0.07	0.05	0.07	0.00	0.12	0.08	0.22	0.00	0.30	
64	0.01	0.05	0.00	0.06	0.01	0.06	0.00	0.07	0.05	0.01	0.00	0.06	0.09	0.06	0.00	0.15	
66	0.04	0.00	0.00	0.04	0.01	0.03	0.00	0.04	0.02	0.05	0.00	0.07	0.07	0.05	0.00	0.12	
68	0.00	0.01	0.00	0.01	0.01	0.02	0.00	0.03	0.04	0.01	0.00	0.05	0.02	0.09	0.00	0.11	
70	0.01	0.00	0.00	0.01	0.01	0.03	0.00	0.04	0.01	0.00	0.00	0.01	0.00	0.05	0.00	0.05	
72	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	
74	0.00	0.01	0.00	0.01	0.01	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.01	0.01	0.00	0.02	
76	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	
78	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
80	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
82	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
84	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	
86	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
88	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
90	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	
92	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	
Total	29.27	32.78	0.00	62.05	6.18	8.25	0.00	14.44	16.19	20.07	0.00	36.25	17.01	21.02	0.00	38.03	
Nº samples:					34				32				36				34
Nº Ind.:	739	827	0	1566	580	781	0	1361	1014	1265	0	2279	1147	1440	0	2587	
Sampled catch:					1814				957				2509				3141
Range:					12-74				13-77				12-93				19-85
Total catch:					1814				957				2509				3141
Total valid hauls:					100				98				97				89

Table 6 (cont.).- Atlantic cod length distribution per haul mean catches by sex and year. Number per stratified mean catches. Spanish Summer Survey on NAFO 3L: 2008-2015 (R/V *Vizconde de Eza*). Indet. means indeterminate.

Length (cm.)	2012				2013				2014				2015				
	M	F	I	T	M	F	I	T	M	F	I	T	M	F	I	T	
<12	0.00	0.00	0.01	0.01	0.01	0.01	0.00	0.02	0.01	0.01	0.02	0.04	0.00	0.00	0.00	0.00	
12	0.00	0.00	0.00	0.00	0.12	0.06	0.00	0.18	0.00	0.00	0.01	0.01	0.00	0.00	0.00	0.00	
14	0.01	0.00	0.00	0.01	0.28	0.41	0.00	0.68	0.01	0.04	0.00	0.05	0.00	0.00	0.00	0.00	
16	0.02	0.02	0.00	0.04	0.54	0.41	0.00	0.95	0.02	0.01	0.00	0.03	0.01	0.00	0.00	0.01	
18	0.07	0.04	0.00	0.11	0.19	0.22	0.00	0.41	0.01	0.03	0.00	0.04	0.11	0.07	0.00	0.18	
20	0.01	0.00	0.00	0.01	0.22	0.29	0.00	0.51	0.02	0.07	0.00	0.08	0.25	0.12	0.00	0.37	
22	0.02	0.01	0.00	0.03	0.33	0.45	0.00	0.78	0.09	0.14	0.00	0.23	0.51	0.47	0.00	0.98	
24	0.05	0.03	0.00	0.08	0.66	0.63	0.00	1.30	0.12	0.11	0.00	0.23	0.60	0.69	0.00	1.28	
26	0.06	0.04	0.00	0.10	0.67	0.60	0.00	1.27	0.16	0.28	0.00	0.44	0.85	0.91	0.00	1.75	
28	0.15	0.03	0.00	0.18	0.71	0.65	0.00	1.36	0.19	0.17	0.00	0.36	0.79	0.86	0.00	1.65	
30	0.14	0.06	0.00	0.20	0.60	0.60	0.00	1.21	0.24	0.23	0.00	0.47	0.90	0.90	0.00	1.80	
32	0.15	0.07	0.00	0.22	0.33	0.43	0.00	0.76	0.28	0.23	0.00	0.52	1.08	0.72	0.00	1.80	
34	0.39	0.33	0.00	0.72	0.28	0.29	0.00	0.58	0.38	0.38	0.00	0.76	0.68	0.68	0.00	1.36	
36	0.44	0.58	0.00	1.03	0.41	0.36	0.00	0.78	0.63	0.39	0.00	1.02	0.66	0.62	0.00	1.28	
38	0.68	0.98	0.00	1.66	0.58	0.46	0.00	1.05	0.55	0.62	0.00	1.17	0.68	0.63	0.00	1.31	
40	0.73	0.82	0.00	1.55	0.50	0.35	0.00	0.86	0.43	0.41	0.00	0.84	0.49	0.42	0.00	0.90	
42	0.71	1.08	0.00	1.79	0.54	0.67	0.00	1.21	0.36	0.45	0.00	0.81	0.28	0.44	0.00	0.72	
44	0.75	0.85	0.00	1.60	0.73	0.98	0.00	1.71	0.18	0.34	0.00	0.52	0.37	0.50	0.00	0.88	
46	0.91	0.97	0.00	1.88	0.86	0.76	0.00	1.62	0.23	0.23	0.00	0.47	0.28	0.28	0.00	0.56	
48	0.64	0.97	0.00	1.61	0.75	0.80	0.00	1.54	0.24	0.27	0.00	0.51	0.24	0.34	0.00	0.58	
50	0.63	0.79	0.00	1.42	0.52	0.75	0.00	1.27	0.24	0.26	0.00	0.50	0.26	0.22	0.00	0.48	
52	0.48	0.62	0.00	1.10	0.50	0.62	0.00	1.11	0.15	0.27	0.00	0.42	0.10	0.11	0.00	0.22	
54	0.45	0.54	0.00	0.99	0.36	0.72	0.00	1.09	0.16	0.19	0.00	0.35	0.21	0.13	0.00	0.33	
56	0.55	0.48	0.00	1.03	0.42	0.44	0.00	0.86	0.09	0.18	0.00	0.27	0.13	0.18	0.00	0.31	
58	0.22	0.22	0.00	0.45	0.29	0.47	0.00	0.76	0.12	0.18	0.00	0.30	0.07	0.14	0.00	0.21	
60	0.16	0.33	0.00	0.48	0.17	0.31	0.00	0.49	0.06	0.09	0.00	0.15	0.04	0.10	0.00	0.13	
62	0.10	0.19	0.00	0.29	0.19	0.33	0.00	0.52	0.05	0.10	0.00	0.15	0.11	0.06	0.00	0.16	
64	0.05	0.17	0.00	0.22	0.12	0.17	0.00	0.28	0.02	0.10	0.00	0.12	0.05	0.02	0.00	0.07	
66	0.02	0.12	0.00	0.14	0.10	0.12	0.00	0.21	0.02	0.05	0.00	0.08	0.03	0.02	0.00	0.06	
68	0.04	0.04	0.00	0.08	0.10	0.09	0.00	0.19	0.02	0.02	0.00	0.04	0.00	0.01	0.00	0.01	
70	0.01	0.06	0.00	0.07	0.02	0.04	0.00	0.06	0.01	0.04	0.00	0.05	0.02	0.01	0.00	0.03	
72	0.00	0.00	0.00	0.00	0.04	0.09	0.00	0.13	0.00	0.04	0.00	0.04	0.01	0.02	0.00	0.04	
74	0.00	0.00	0.00	0.00	0.04	0.04	0.00	0.09	0.00	0.00	0.00	0.00	0.01	0.01	0.00	0.02	
76	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.01	0.00	0.02	0.00	0.02	0.00	0.00	0.00	0.00	
78	0.00	0.01	0.00	0.01	0.01	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	
80	0.00	0.00	0.00	0.00	0.01	0.01	0.00	0.02	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
82	0.00	0.03	0.00	0.03	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
84	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.01	0.00	0.00	0.00	0.00	
86	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
88	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
90	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
92	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
Total	8.65	10.47	0.01	19.14	12.24	13.65	0.00	25.89	5.12	5.97	0.03	11.12	9.83	9.68	0.00	19.51	
Nº samples:					35				41				38				39
Nº Ind.:	603	693	1	1297	1085	1200	0	2285	463	546	3	1012	848	840	0	1688	
Sampled catch:					1809				2002				806				927
Range:					5-82				11-87				9-84				17-79
Total catch:					1809				2002				806				927
Total valid hauls:					98				100				99				97

Table 7.- Swept area, number of hauls and roughhead grenadier mean catch (Kg) and SD (\*\*) by stratum. Spanish Survey on NAFO Div. 3L in the period 2011-2015, on board R/V "Vizconde de Eza".

<b>Stratum</b>	2011			2012			2013			2014			2015		
	Swept Tow area No.	Mean catch	SD	Swept Tow area No.	Mean catch	SD	Swept Tow area No.	Mean catch	SD	Swept Tow area No.	Mean catch	SD	Swept Tow area No.	Mean catch	SD
<b>385</b>	0.0229 2	1.010	1.428	0.0225 2	0.000	0.000	0.0229 2	0.000	0.000	0.0225 2	0.000	0.000	0.0236 2	0.000	0.000
<b>387</b>	0.0450 4	11.304	9.250	0.0450 4	16.012	11.119	0.0450 4	48.039	29.999	0.0461 4	34.291	36.472	0.0458 4	56.143	42.743
<b>388</b>	0.0563 5	5.022	4.969	0.0570 5	14.019	22.081	0.0570 5	11.737	9.670	0.0585 5	19.183	19.378	0.0574 5	35.214	22.023
<b>389</b>	0.0675 6	4.711	3.126	0.0799 7	11.893	9.022	0.0791 7	7.694	11.153	0.0814 7	4.613	7.433	0.0814 7	14.689	18.122
<b>390</b>	0.1009 9	2.856	7.168	0.1354 12	0.000	0.000	0.1358 12	0.418	0.995	0.1369 12	0.203	0.530	0.1260 11	0.173	0.573
<b>391</b>	0.0458 4	153.179	92.811	0.0458 4	21.670	8.743	0.0450 4	6.940	6.438	0.0465 4	18.675	19.226	0.0465 4	118.535	108.870
<b>392</b>	0.0229 2	83.417	29.674	0.0225 2	73.339	76.293	0.0225 2	462.715	55.388	0.0225 2	165.300	98.005	0.0229 2	92.963	28.656
<b>729</b>	0.0338 3	3.398	2.102	0.0338 3	23.722	12.954	0.0341 3	13.044	2.954	0.0338 3	20.597	10.873	0.0345 3	21.267	8.882
<b>730</b>	0.0334 3	66.456	55.464	0.0338 3	27.264	5.665	0.0334 3	16.433	3.745	0.0345 3	24.237	12.193	0.0345 3	43.188	29.351
<b>731</b>	0.0334 3	2.002	1.506	0.0341 3	5.244	2.400	0.0334 3	5.861	7.211	0.0345 3	11.131	11.131	0.0345 3	12.921	8.486
<b>732</b>	0.0454 4	2.393	2.786	0.0454 4	3.022	2.324	0.0450 4	9.399	5.783	0.0454 4	20.145	14.299	0.0465 4	18.716	4.826
<b>733</b>	0.0454 4	6.622	8.721	0.0454 4	9.322	10.885	0.0450 4	25.366	26.819	0.0458 4	48.449	47.653	0.0454 4	22.976	35.302
<b>734</b>	0.0225 2	8.413	1.874	0.0233 2	20.968	0.803	0.0221 2	51.715	2.849	0.0225 2	52.870	32.286	0.0225 2	57.250	48.154
<b>741</b>	0.0218 2	7.707	9.880	0.0218 2	5.764	2.452	0.0221 2	26.100	18.526	0.0225 2	9.559	5.316	0.0236 2	26.240	14.199
<b>742</b>	0.0225 2	14.545	14.221	0.0206 2	6.851	3.796	0.0218 2	4.829	4.554	0.0221 2	39.490	39.330	0.0233 2	8.550	3.323
<b>743</b>	0.0221 2	18.488	1.660	0.0206 2	5.421	7.609	0.0218 2	23.750	18.314	0.0221 2	14.015	16.567	0.0233 2	12.869	6.178
<b>744</b>	0.0221 2	6.254	3.743	0.0221 2	8.725	9.086	0.0221 2	27.217	13.266	0.0225 2	9.081	3.064	0.0225 2	3.869	1.951
<b>745</b>	0.0446 4	2.802	4.240	0.0570 5	1.932	1.671	0.0559 5	7.092	4.649	0.0578 5	14.445	16.588	0.0578 5	14.563	7.820
<b>746</b>	0.0566 5	8.981	7.193	0.0675 6	14.447	14.048	0.0675 6	19.411	13.114	0.0683 6	18.434	11.243	0.0686 6	16.779	6.548
<b>747</b>	0.0893 8	22.273	17.958	0.1121 10	19.457	7.563	0.1125 10	22.433	9.574	0.1125 10	20.426	14.337	0.1028 9	35.466	21.325
<b>748</b>	0.0221 2	25.955	33.074	0.0225 2	106.350	134.562	0.0225 2	50.520	62.607	0.0229 2	72.050	46.457	0.0233 2	105.925	34.189
<b>749</b>	0.0221 2	27.713	30.670	0.0221 2	9.800	8.061	0.0225 2	16.950	0.495	0.0225 2	15.900	4.384	0.0225 2	56.302	40.590
<b>750</b>	0.0668 6	9.292	4.047	0.0885 8	18.823	14.451	0.0896 8	6.988	4.947	0.0904 8	10.760	11.655	0.0934 8	23.339	13.464
<b>751</b>	0.0334 3	14.880	6.137	0.0218 2	34.850	33.022	0.0446 4	9.238	3.941	0.0334 3	9.612	6.745	0.0341 3	56.233	55.211

$$(**) SD = \frac{\sum (x_i - \bar{x})}{n-1}$$

Table 8.- Stratified mean catches (Kg) of roughhead grenadier by stratum and year (2003-2015) and SD. Research Vessel *Vizconde de Eza*. n.s. means stratum not surveyed. In 2003: the data correspond to 69% of the total area prospected in 2006-2015.

Stratum	Survey												
	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
385	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	119.18	0.00	0.00	0.00	0.00
387	0.00	15356.54	-	8906.24	11773.44	5201.92	7691.52	3686.21	2893.70	4099.14	12297.98	8778.56	14372.48
388	0.00	15458.10	-	9426.94	13445.76	5374.85	9862.70	6131.05	1792.71	5004.78	4190.04	6848.33	12571.40
389	0.00	954.38	-	725.69	1565.18	9674.64	15832.37	4189.80	2397.73	6053.54	3916.39	2347.80	7476.48
390	456.40	5.43	-	0.00	0.00	472.70	3787.71	872.79	2327.28	0.00	340.94	165.72	140.77
391	4.70	4.94	-	50230.55	24400.05	70203.05	20551.46	47806.05	43196.41	6110.94	1957.08	5266.35	33426.87
392	565.50	29094.25	-	17113.63	18842.75	8435.38	8835.43	5082.25	12095.47	10634.08	67093.68	23968.50	13479.56
729	7021.50	5482.35	-	4680.44	4927.20	3709.46	1858.39	2011.90	632.09	4412.23	2426.25	3830.98	3955.60
730	17178.50	5731.55	-	9055.90	13834.26	5970.29	12827.07	4488.00	11297.58	4634.82	2793.67	4120.23	7341.96
731	758.16	2257.20	-	2270.52	3095.93	3095.93	1075.61	2269.73	432.36	1132.78	1266.05	2404.37	2791.01
732	7946.40	9122.19	-	5119.88	2575.96	4976.90	1914.82	3709.74	552.67	698.08	2171.17	4653.55	4323.40
733	n.s	3639.48	-	5487.30	4470.26	5601.67	4471.16	2055.69	1549.49	2181.41	5935.70	11337.07	5376.33
734	n.s	10075.05	-	6015.20	3580.20	4678.66	4402.88	10040.63	1287.19	3208.03	7912.32	8089.11	8759.25
741	870.00	105.53	-	1755.70	465.00	1035.90	1133.40	1435.00	770.65	576.35	2610.00	955.90	2624.00
742	1561.60	300.80	-	1339.68	927.55	1079.10	219.20	247.68	930.85	438.46	309.02	2527.33	547.20
743	n.s	1338.50	-	539.27	1512.97	1300.93	677.18	1577.79	942.89	276.45	1211.25	714.74	656.29
744	n.s	168.30	-	1014.09	2241.69	3872.22	541.70	879.05	412.73	575.85	1796.29	599.31	255.32
745	6106.24	2018.40	-	2866.88	1261.09	4970.83	1317.95	2769.59	975.10	672.20	2468.16	5026.86	5068.06
746	25009.60	10272.36	-	16372.53	13565.94	12042.24	9201.61	5107.56	3520.47	5663.35	7609.05	7226.19	6577.37
747	n.s	31585.71	-	30630.47	45257.17	20791.04	24022.61	26632.56	16125.29	14086.51	16241.27	14788.28	25677.71
748	8900.82	3579.89	-	10799.28	5331.80	34557.06	14680.47	8005.65	4126.85	16909.65	8032.68	11455.95	16842.08
749	18295.20	5783.40	-	3267.18	3616.20	5978.95	1726.20	2580.67	3491.84	1234.80	2135.70	2003.40	7093.99
750	n.s	31553.00	-	9377.25	10850.99	6636.90	9393.86	7096.23	5166.44	10465.52	3885.26	5982.56	12976.62
751	n.s	n.s	-	973.82	5597.91	2069.59	20196.12	5072.35	3407.52	7980.65	2115.39	2201.22	12877.43
TOTAL $(\bar{y})$	94674.62	183887.34	-	197968.44	193139.30	221730.20	176221.39	153747.96	120444.46	107049.61	160715.33	135292.31	205211.18
SD	3.38	5.27	-	7.41	4.86	6.12	4.97	1.71	2.51	2.92	1.75	2.44	3.31

Table 9.- Survey estimates (by the swept area method) of roughhead grenadier biomass (t.) by stratum and year and their SD on NAFO Div. 3L (R/V *Vizconde de Eza*). n.s. means stratum not surveyed. In 2003, the data correspond to 69% of the total area prospected in 2006-2015.

Stratum	Survey												
	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
385	0	0	-	0	0	0	0	0	10	0	0	0	0
387	0	1437	-	792	1047	478	701	322	257	364	1093	761	1257
388	0	1472	-	832	1195	481	889	538	159	439	368	585	1096
389	0	85	-	64	139	868	1381	369	213	531	346	202	643
390	41	0	-	0	0	41	331	77	208	0	30	15	12
391	0	0	-	4465	2169	6189	1797	4214	3777	534	174	453	2875
392	49	2722	-	1496	1675	763	772	452	1058	945	5964	2131	1179
729	669	496	-	416	438	330	163	179	56	392	213	341	344
730	1553	518	-	833	1230	555	1140	403	1016	412	251	358	638
731	66	194	-	200	275	281	95	202	39	100	114	209	243
732	706	869	-	460	229	446	170	330	49	62	193	410	372
733	n.s	331	-	484	397	520	397	183	137	192	528	991	474
734	n.s	995	-	535	318	423	405	893	114	276	715	719	779
741	77	10	-	161	41	99	102	128	71	53	236	85	222
742	134	25	-	117	82	103	21	22	83	43	28	228	47
743	n.s	143	-	48	134	128	67	140	85	27	111	65	56
744	n.s	17	-	89	206	350	52	77	37	52	162	53	23
745	537	190	-	251	112	448	118	246	87	59	221	435	439
746	2242	913	-	1455	1226	1133	827	451	311	503	676	635	575
747	n.s	3082	-	2739	4023	1945	2150	2367	1445	1256	1444	1315	2249
748	818	360	-	993	474	3178	1284	712	373	1503	714	1002	1449
749	1654	523	-	286	321	559	153	226	316	112	190	178	631
750	n.s	3506	-	840	959	629	831	631	464	946	347	530	1112
751	n.s	n.s	-	86	498	201	1795	451	306	734	190	198	1132
TOTAL	8546	17887	-	17641	17190	20148	15641	13612	10672	9535	14308	11898	17846
SD	1340	3240	-	4271	2799	3534	2844	972	1466	1676	1010	1393	1864

Table 10.- Roughhead grenadier length distribution per haul mean catches by sex and year. Number per stratified mean catches. Spanish Summer Survey on NAFO 3L: 2008-2015 (R/V *Vizconde de Eza*). Indet. means indeterminate.

Length (cm.)	2008				2009				2010				2011				
	M	F	I	T	M	F	I	T	M	F	I	T	M	F	I	T	
1.5	0.00	0.00	0.02	0.02	0.00	0.00	0.01	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
2.5	0.00	0.03	0.09	0.13	0.01	0.00	0.13	0.15	0.03	0.00	0.26	0.29	0.02	0.00	0.17	0.19	
3.5	0.28	0.08	1.42	1.78	0.27	0.16	1.01	1.44	0.07	0.05	0.33	0.46	0.00	0.01	1.41	1.42	
4.5	0.11	0.01	0.03	0.15	0.07	0.00	0.05	0.12	0.04	0.09	0.01	0.15	0.03	0.03	0.07	0.14	
5.5	0.10	0.13	0.01	0.24	0.12	0.13	0.00	0.25	0.29	0.20	0.00	0.48	0.06	0.16	0.01	0.24	
6.5	0.69	0.64	0.03	1.36	0.38	0.45	0.00	0.83	0.58	0.59	0.00	1.17	0.24	0.30	0.02	0.56	
7.5	0.24	0.38	0.00	0.62	0.11	0.23	0.00	0.35	0.26	0.22	0.00	0.47	0.22	0.18	0.00	0.40	
8.5	0.39	0.46	0.00	0.85	0.25	0.30	0.00	0.54	0.28	0.36	0.00	0.64	0.44	0.34	0.00	0.78	
9.5	0.74	0.58	0.00	1.31	0.38	0.51	0.00	0.89	0.54	0.43	0.00	0.97	0.29	0.46	0.00	0.75	
10.5	0.87	0.77	0.00	1.63	0.56	0.52	0.00	1.08	0.76	0.66	0.00	1.42	0.31	0.42	0.00	0.72	
11.5	1.19	1.32	0.00	2.51	0.56	0.99	0.00	1.55	0.95	0.89	0.00	1.83	0.50	0.29	0.00	0.79	
12.5	1.07	1.20	0.00	2.26	1.24	0.91	0.00	2.15	1.26	1.10	0.00	2.37	0.62	0.63	0.00	1.25	
13.5	1.58	1.36	0.00	2.93	1.33	1.44	0.00	2.77	1.84	1.74	0.00	3.59	0.81	0.79	0.00	1.61	
14.5	2.16	1.77	0.00	3.94	1.58	1.53	0.00	3.11	2.46	2.38	0.00	4.85	1.48	1.13	0.00	2.61	
15.5	2.61	2.21	0.00	4.82	1.92	1.90	0.00	3.81	2.29	2.10	0.00	4.40	2.22	1.37	0.00	3.59	
16.5	2.60	2.67	0.00	5.26	1.96	1.80	0.00	3.76	2.32	2.49	0.00	4.80	2.24	1.41	0.00	3.65	
17.5	1.92	1.97	0.00	3.89	1.71	1.96	0.00	3.67	1.89	2.35	0.00	4.24	1.35	1.79	0.00	3.14	
18.5	1.60	1.74	0.00	3.34	1.31	1.52	0.00	2.83	1.35	2.30	0.00	3.65	1.31	1.99	0.00	3.30	
19.5	1.36	1.77	0.00	3.13	0.97	1.24	0.00	2.22	0.75	1.78	0.00	2.52	0.58	1.78	0.00	2.36	
20.5	0.82	1.89	0.00	2.71	0.59	1.22	0.00	1.81	0.36	1.26	0.00	1.62	0.16	1.26	0.00	1.42	
21.5	0.37	1.71	0.00	2.09	0.30	1.23	0.00	1.53	0.16	1.20	0.00	1.36	0.06	0.85	0.00	0.91	
22.5	0.10	1.82	0.00	1.91	0.15	1.21	0.00	1.37	0.04	0.85	0.00	0.89	0.06	0.66	0.00	0.72	
23.5	0.03	1.83	0.00	1.86	0.01	1.33	0.00	1.35	0.04	0.93	0.00	0.96	0.00	0.58	0.00	0.58	
24.5	0.00	2.28	0.00	2.29	0.00	1.25	0.00	1.25	0.00	0.56	0.00	0.56	0.01	0.73	0.00	0.74	
25.5	0.00	1.87	0.00	1.87	0.01	1.18	0.00	1.19	0.00	0.80	0.00	0.80	0.00	0.58	0.00	0.58	
26.5	0.00	1.53	0.00	1.53	0.00	1.19	0.00	1.19	0.00	0.56	0.00	0.56	0.00	0.63	0.00	0.63	
27.5	0.00	0.88	0.00	0.88	0.00	0.82	0.00	0.82	0.00	0.44	0.00	0.44	0.00	0.50	0.00	0.50	
28.5	0.00	0.62	0.00	0.62	0.00	0.52	0.00	0.52	0.00	0.38	0.00	0.38	0.00	0.37	0.00	0.37	
29.5	0.00	0.58	0.00	0.58	0.00	0.46	0.00	0.46	0.00	0.23	0.00	0.23	0.00	0.17	0.00	0.17	
30.5	0.00	0.15	0.00	0.15	0.00	0.27	0.00	0.27	0.00	0.11	0.00	0.11	0.00	0.10	0.00	0.10	
31.5	0.00	0.11	0.00	0.11	0.00	0.23	0.00	0.23	0.00	0.09	0.00	0.09	0.00	0.03	0.00	0.03	
32.5	0.00	0.07	0.00	0.07	0.00	0.14	0.00	0.14	0.00	0.06	0.00	0.06	0.00	0.04	0.00	0.04	
33.5	0.00	0.03	0.00	0.03	0.00	0.09	0.00	0.09	0.00	0.06	0.00	0.06	0.00	0.03	0.00	0.03	
34.5	0.00	0.00	0.00	0.00	0.00	0.08	0.00	0.08	0.00	0.06	0.00	0.06	0.00	0.01	0.00	0.01	
35.5	0.00	0.01	0.00	0.01	0.00	0.03	0.00	0.03	0.00	0.02	0.00	0.02	0.00	0.05	0.00	0.05	
36.5	0.00	0.02	0.00	0.02	0.00	0.03	0.00	0.03	0.00	0.05	0.00	0.05	0.00	0.01	0.00	0.01	
37.5	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.04	0.00	0.04	0.00	0.02	0.00	0.02	
38.5	0.00	0.02	0.00	0.02	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.03	0.00	0.03	
39.5	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
40.5	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	
41.5	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.02	0.00	0.02	0.00	0.00	0.00	0.00	
Total	20.84	34.48	1.59	56.91	15.78	26.93	1.21	43.93	18.58	27.44	0.61	46.63	13.01	19.73	1.68	34.43	
Nº samples:									87				84				83
Nº Ind.:	2022	3019	176	5217	1409	2319	105	3833	1486	1997	65	3548	1037	1506	140	2683	
Sampled catch:									3287				2541				1710
Range:				1.5-42.5				2.0-41.5				2.5-42				2.5-39	
Total catch:				3287				2543				2234				1710	
Total valid hauls:				100				98				97				89	

Table 10 (cont).- Roughhead grenadier length distribution per haul mean catches by sex and year. Number per stratified mean catches. Spanish Summer Survey on NAFO 3L: 2008-2015 (R/V *Vizconde de Eza*). Indet. means indeterminate.

Length (cm.)	2012				2013				2014				2015				
	M	F	I	T	M	F	I	T	M	F	I	T	M	F	I	T	
1.5	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
2.5	0.00	0.00	0.24	0.24	0.01	0.03	1.12	1.15	0.00	0.05	0.24	0.29	0.00	0.00	1.16	1.16	
3.5	0.00	0.02	1.01	1.04	0.24	0.09	4.47	4.80	0.04	0.01	1.38	1.43	0.10	0.00	3.19	3.30	
4.5	0.10	0.03	0.03	0.16	0.17	0.20	0.24	0.60	0.09	0.01	0.07	0.17	0.06	0.04	0.04	0.14	
5.5	0.22	0.23	0.00	0.44	1.23	1.16	0.00	2.38	0.56	0.51	0.01	1.08	0.67	1.14	0.02	1.83	
6.5	0.92	0.88	0.00	1.80	1.66	2.08	0.00	3.74	0.85	1.04	0.00	1.88	2.00	2.51	0.00	4.51	
7.5	0.38	0.35	0.00	0.74	0.39	0.45	0.00	0.84	0.17	0.27	0.00	0.43	0.73	0.77	0.00	1.50	
8.5	0.38	0.28	0.00	0.66	1.13	1.52	0.00	2.65	0.27	0.38	0.00	0.65	2.08	2.32	0.00	4.39	
9.5	0.44	0.53	0.00	0.98	1.23	3.14	0.00	4.37	0.46	0.36	0.00	0.81	1.38	1.64	0.00	3.02	
10.5	0.57	0.42	0.00	0.99	0.63	1.16	0.00	1.78	0.61	0.47	0.00	1.08	0.73	0.87	0.00	1.59	
11.5	0.68	0.60	0.00	1.28	1.10	2.29	0.00	3.39	0.70	0.62	0.00	1.32	0.94	1.08	0.00	2.01	
12.5	0.65	0.59	0.00	1.24	1.52	2.64	0.00	4.16	0.75	0.69	0.00	1.43	1.54	1.23	0.00	2.77	
13.5	0.79	0.74	0.00	1.53	2.42	3.03	0.00	5.46	1.33	1.03	0.00	2.36	1.57	1.29	0.00	2.86	
14.5	1.26	0.91	0.00	2.17	1.77	2.40	0.00	4.17	1.24	1.14	0.00	2.38	1.94	1.38	0.00	3.31	
15.5	1.52	1.13	0.00	2.65	2.04	2.84	0.00	4.88	1.46	1.15	0.00	2.61	2.76	2.02	0.00	4.78	
16.5	1.63	1.02	0.00	2.65	2.18	2.17	0.00	4.35	1.84	1.26	0.00	3.10	3.18	1.87	0.00	5.05	
17.5	1.54	1.46	0.00	2.99	1.98	2.97	0.00	4.95	1.49	1.74	0.00	3.23	2.76	2.25	0.00	5.01	
18.5	1.06	1.38	0.00	2.45	1.51	2.30	0.00	3.81	0.91	1.71	0.00	2.62	2.67	2.08	0.00	4.75	
19.5	0.64	1.19	0.00	1.83	0.65	2.34	0.00	2.99	0.51	1.64	0.00	2.15	1.05	2.55	0.00	3.60	
20.5	0.29	1.25	0.00	1.55	0.33	1.70	0.00	2.03	0.40	1.84	0.00	2.24	0.47	2.62	0.00	3.09	
21.5	0.09	0.96	0.00	1.05	0.16	1.40	0.01	1.57	0.19	1.76	0.00	1.95	0.17	2.34	0.00	2.51	
22.5	0.01	0.98	0.00	0.99	0.01	1.44	0.00	1.45	0.04	1.36	0.00	1.40	0.11	1.70	0.00	1.81	
23.5	0.01	0.61	0.00	0.63	0.00	1.16	0.00	1.16	0.04	1.22	0.00	1.26	0.02	1.44	0.00	1.47	
24.5	0.00	0.70	0.00	0.70	0.00	0.70	0.00	0.70	0.00	1.02	0.00	1.02	0.00	1.26	0.00	1.26	
25.5	0.00	0.49	0.00	0.49	0.00	0.63	0.00	0.63	0.00	0.67	0.00	0.67	0.00	0.80	0.00	0.80	
26.5	0.00	0.45	0.00	0.45	0.00	0.47	0.00	0.47	0.00	0.51	0.00	0.51	0.01	0.76	0.00	0.77	
27.5	0.00	0.44	0.00	0.44	0.01	0.29	0.00	0.30	0.00	0.45	0.00	0.45	0.00	0.45	0.00	0.45	
28.5	0.00	0.23	0.00	0.23	0.00	0.36	0.00	0.36	0.00	0.29	0.00	0.29	0.00	0.44	0.00	0.44	
29.5	0.00	0.10	0.00	0.10	0.00	0.18	0.00	0.18	0.00	0.21	0.00	0.21	0.00	0.34	0.00	0.34	
30.5	0.00	0.08	0.00	0.08	0.00	0.18	0.00	0.18	0.00	0.15	0.00	0.15	0.00	0.17	0.00	0.17	
31.5	0.00	0.16	0.00	0.16	0.00	0.08	0.00	0.08	0.00	0.05	0.00	0.05	0.00	0.20	0.00	0.20	
32.5	0.00	0.02	0.00	0.02	0.00	0.06	0.00	0.06	0.00	0.04	0.00	0.04	0.00	0.13	0.00	0.13	
33.5	0.00	0.02	0.00	0.02	0.00	0.03	0.00	0.03	0.00	0.04	0.00	0.04	0.00	0.09	0.00	0.09	
34.5	0.00	0.02	0.00	0.02	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.08	0.00	0.08	
35.5	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.03	0.00	0.03	0.00	0.05	0.00	0.05	
36.5	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.03	0.00	0.03	0.00	0.02	0.00	0.02	
37.5	0.00	0.02	0.00	0.02	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.01	
38.5	0.00	0.03	0.00	0.03	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	
39.5	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
40.5	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
41.5	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
Total	13.19	18.35	1.29	32.83	22.36	41.53	5.84	69.73	13.94	23.75	1.70	39.39	26.93	37.94	4.41	69.28	
Nº samples:					82				83				83				82
Nº Ind.:	1077	1413	113	2603	0	1986	427	3731	1126	1892	154	3172	2276	3199	444	5919	
Sampled catch:					1508				2379				2043				2954
Range:					2.5-38.5				2.5-39				2.5-39				2.5-38
Total catch:					1508				2379				2043				2954
Total valid hauls:					98				100				99				97

Table 11.- Swept area, number of hauls and redfish mean catch (Kg) and SD (\*\*) by stratum. Spanish Survey on NAFO Div. 3L in the period 2011-2015, on board R/V "Vizconde de Eza".

Stratum	2011			2012			2013			2014			2015			
	SweptTow area No.	Mean catch	SD	SweptTow area No.	Mean catch	SD	SweptTow area No.	Mean catch	SD	SweptTow area No.	Mean catch	SD	Swept Tow area No.	Mean catch	SD	
<b>385</b>	0.0229	2	0.205	0.290	0.0225	2	0.000	0.000	0.0229	2	0.114	0.161	0.0225	2	0.000	0.000
<b>387</b>	0.0450	4	471.900	592.192	0.0450	4	456.188	146.956	0.0450	4	903.875	221.080	0.0461	4	692.755	574.493
<b>388</b>	0.0563	5	400.680	561.867	0.0570	5	3649.824	2735.118	0.0570	5	2614.156	2779.770	0.0585	5	2063.600	2163.327
<b>389</b>	0.0675	6	314.072	337.845	0.0799	7	5366.45013039.715	0.0791	7	1522.331	2830.529	0.0814	7	672.973	1713.444	
<b>390</b>	0.1009	9	0.298	0.893	0.1354	12	0.307	0.723	0.1358	12	0.250	0.567	0.1369	12	0.096	0.316
<b>391</b>	0.0458	4	270.078	524.098	0.0458	4	1317.264	848.814	0.0450	4	9.546	9.721	0.0465	4	39.913	51.137
<b>392</b>	0.0229	2	7489.781	7767.171	0.0225	2	4138.815	2411.128	0.0225	2	1336.512	1473.062	0.0225	2	2692.510	923.665
<b>729</b>	0.0338	3	1405.563	2154.649	0.0338	3	1491.733	2440.054	0.0341	3	1933.319	1952.744	0.0338	3	1061.297	884.322
<b>730</b>	0.0334	3	98.992	73.752	0.0338	3	214.100	203.592	0.0334	3	143.300	121.829	0.0345	3	92.793	111.735
<b>731</b>	0.0334	3	45.227	32.987	0.0341	3	37.000	4.590	0.0334	3	82.897	60.702	0.0345	3	110.933	80.154
<b>732</b>	0.0454	4	12.480	9.605	0.0454	4	7.236	4.921	0.0450	4	5.558	2.888	0.0454	4	39.853	27.312
<b>733</b>	0.0454	4	255.160	236.623	0.0454	4	129.800	140.677	0.0450	4	418.230	374.577	0.0458	4	2467.588	3626.885
<b>734</b>	0.0225	2	7.888	0.972	0.0233	2	9.015	1.393	0.0221	2	168.600	170.554	0.0225	2	42.250	1.909
<b>741</b>	0.0218	2	0.500	0.707	0.0218	2	0.700	0.990	0.0221	2	2.003	2.833	0.0225	2	0.360	0.509
<b>742</b>	0.0225	2	0.208	0.294	0.0206	2	0.000	0.000	0.0218	2	0.000	0.000	0.0221	2	0.000	0.000
<b>743</b>	0.0221	2	0.000	0.000	0.0206	2	0.000	0.000	0.0218	2	0.000	0.000	0.0221	2	0.000	0.000
<b>744</b>	0.0221	2	0.858	1.213	0.0221	2	0.000	0.000	0.0221	2	0.000	0.000	0.0225	2	0.000	0.000
<b>745</b>	0.0446	4	0.745	1.007	0.0570	5	0.348	0.506	0.0559	5	0.490	0.565	0.0578	5	1.204	1.597
<b>746</b>	0.0566	5	0.000	0.000	0.0675	6	0.000	0.000	0.0675	6	0.000	0.000	0.0683	6	0.009	0.022
<b>747</b>	0.0893	8	0.379	1.071	0.1121	10	0.000	0.000	0.1125	10	0.000	0.000	0.1125	10	0.000	0.000
<b>748</b>	0.0221	2	0.595	0.134	0.0225	2	0.000	0.000	0.0225	2	7.045	8.846	0.0229	2	0.000	0.000
<b>749</b>	0.0221	2	0.000	0.000	0.0221	2	0.000	0.000	0.0225	2	0.000	0.000	0.0225	2	0.000	0.000
<b>750</b>	0.0668	6	0.242	0.592	0.0885	8	0.039	0.110	0.0896	8	0.000	0.000	0.0904	8	0.000	0.000
<b>751</b>	0.0334	3	0.000	0.000	0.0218	2	0.000	0.000	0.0446	4	0.000	0.000	0.0334	3	0.000	0.000

$$(**) SD = \frac{\sum(x_i - \bar{x})}{n-1}$$

Table 12.- Stratified mean catches (Kg) of redfish by stratum and year (2003-2015) and SD. Research Vessel *Vizconde de Eza*. n.s. means stratum not surveyed. In 2003: the data correspond to 69% of the total area prospected in 2006-2015.

Stratum	Survey												
	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
385	0.12	0.59	-	0.00	4.84	58.35	32.45	0.00	24.19	0.00	13.45	0.00	42.78
387	439.04	14336.00	-	29103.36	20582.40	47392.00	145517.18	71328.00	120806.40	116784.00	231392.00	177345.28	187906.56
388	2303.84	4212.60	-	23576.28	57861.85	75951.75	602000.18	329247.18	143042.76	1302987.17	933253.69	736705.20	312767.70
389	407.58	16822.45	-	23418.22	5458.01	196133.55	163604.53	1755783.21	159862.48	2731523.05	774866.41	342543.18	354032.77
390	472.70	0.00	-	153.59	141.00	751.23	70.36	3.93	242.69	250.00	203.95	78.10	465.14
391	24.44	404.67	-	2012.07	1695.53	308262.66	68687.02	659127.27	76161.86	371468.38	2691.97	11255.33	84684.81
392	6713.50	177236.40	-	633242.55	139149.25	30326.75	115644.17	69614.50	1086018.17	600128.18	193794.24	390413.95	202241.22
729	16516.80	57706.50	-	37603.00	23973.29	115034.80	9454.32	52966.60	261434.78	277462.40	359597.27	197401.18	42352.20
730	39283.60	9443.50	-	24806.97	62515.29	5064.30	28492.00	25065.93	16828.70	36397.00	24361.00	15774.87	40800.85
731	8502.84	17182.80	-	4115.52	8013.60	28720.80	7992.00	19231.20	9768.96	7992.00	17905.68	23961.60	107211.60
732	16678.20	9707.78	-	1302.46	2798.49	2766.23	1919.90	3849.62	2882.88	1671.40	1283.78	9205.93	2691.15
733	n.s	26130.00	-	16988.40	27066.00	31028.40	13975.65	40802.00	59707.44	30373.20	97865.82	577415.48	151614.45
734	n.s	823.65	-	1886.11	3783.31	3440.21	2481.66	909.59	1206.79	1379.30	25795.80	6464.25	11559.15
741	224000.00	25.50	-	0.00	0.00	55.50	90.25	0.00	50.00	70.00	200.30	36.00	138.70
742	0.00	21.18	-	0.00	19.20	0.00	0.00	0.00	13.31	0.00	0.00	0.00	28.74
743	n.s	106.59	-	0.00	0.00	0.00	284.33	0.00	0.00	0.00	0.00	0.00	0.00
744	n.s	0.00	-	0.00	31.58	0.00	0.00	8.78	56.63	0.00	0.00	0.00	0.00
745	610078.80	0.00	-	41.47	132.24	126.74	0.00	151.73	259.26	121.10	170.59	418.92	97.72
746	0.00	0.00	-	46.39	0.00	0.00	16.99	20.91	0.00	0.00	0.00	3.59	117.99
747	n.s	144.80	-	0.00	0.00	8.98	0.00	0.00	274.22	0.00	0.00	0.00	15.69
748	429.30	69.96	-	20.67	131.97	682.11	250.50	0.00	94.61	0.00	1120.16	0.00	488.93
749	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	24.19	0.00
750	n.s	0.00	-	0.00	0.00	0.00	127.88	102.17	134.37	21.68	0.00	0.00	82.22
751	n.s	n.s	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	63.36
TOTAL $(\bar{y})$	925850.76	334374.97		798317.04	353357.83	845804.35	1160641.36	3028212.59	1938870.48	5478628.86	2664516.12	2489047.05	1499403.71
SD	206.94	53.43		123.06	54.47	130.38	178.92	466.81	298.89	844.56	410.75	383.70	231.14
	136.03	28.87		90.99	11.94	36.35	69.07	285.47	130.15	396.90	115.72	101.29	56.47

Table 13.- Survey estimates (by the swept area method) of redfish biomass (t.) by stratum and year and their SD on NAFO Div. 3L (R/V *Vizconde de Eza*). n.s. means stratum not surveyed. In 2003, the data correspond to 69% of the total area prospected in 2006-2015.

Stratum	Survey												
	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
385	0	0	-	0	0	5	3	0	2	0	1	0	4
387	38	1341	-	2587	1830	4358	13267	6236	10738	10381	20568	15380	16429
388	207	401	-	2082	5143	6797	54234	28881	12715	114297	81864	62966	27256
389	36	1495	-	2062	485	17602	14271	154597	14210	239382	68551	29466	30454
390	42	0	-	14	13	65	6	0	22	22	18	7	41
391	2	37	-	179	151	27175	6005	58105	6659	32478	239	968	7285
392	578	16584	-	55365	12369	2741	10111	6188	94952	53345	17226	34703	17682
729	1573	5216	-	3342	2131	10225	831	4708	23239	24663	31613	17547	3683
730	3551	854	-	2281	5557	471	2533	2253	1513	3235	2190	1372	3548
731	743	1478	-	362	712	2611	703	1709	878	703	1609	2084	9323
732	1483	925	-	117	249	248	171	342	254	147	114	812	231
733	n.s.	2375	-	1498	2406	2878	1242	3627	5263	2678	8699	50484	13365
734	n.s.	81	-	168	336	311	228	81	107	119	2332	575	1027
741	19911	2	-	0	0	5	8	0	5	6	18	3	12
742	0	2	-	0	2	0	0	0	1	0	0	0	2
743	n.s.	11	-	0	0	0	28	0	0	0	0	0	0
744	n.s.	0	-	0	3	0	0	1	5	0	0	0	0
745	53633	0	-	4	12	11	0	13	23	11	15	36	8
746	0	0	-	4	0	0	2	2	0	0	0	0	10
747	n.s.	14	-	0	0	1	0	0	25	0	0	0	1
748	39	7	-	2	12	63	22	0	9	0	100	0	42
749	0	0	-	0	0	0	0	0	0	0	0	2	0
750	n.s.	0	-	0	0	0	11	9	12	2	0	0	7
751	n.s.	n.s.	-	0	0	0	0	0	0	0	0	0	6
TOTAL	81837	30825		70066	31410	75567	103675	266754	170632	481469	235158	216405	130418
SD	50717	17163		50718	6885	20435	40871	164597	72507	229026	66637	57523	31673

Table 14.- Redfish length distribution per haul mean catches by sex and year. Number per stratified mean catches. Spanish Summer Survey on NAFO 3L: 2008-2015 (R/V *Vizconde de Eza*). Indet. means indeterminate.

Length (cm.)	2008				2009				2010				2011			
	M	F	I	T	M	F	I	T	M	F	I	T	M	F	I	T
4	0.00	0.00	0.16	0.16	0.00	0.00	0.01	0.01	0.00	0.00	0.05	0.05	0.00	0.00	0.20	0.20
6	0.00	0.00	8.19	8.19	0.00	0.00	1.44	1.44	0.00	0.00	3.06	3.06	0.00	0.00	5.36	5.36
8	0.00	0.00	17.35	17.35	0.00	0.00	7.73	7.73	0.00	0.00	5.23	5.23	0.00	0.00	6.74	6.74
10	0.81	0.21	57.74	58.76	0.12	0.14	6.53	6.79	0.20	0.00	4.23	4.43	0.14	0.08	5.23	5.45
12	3.70	2.13	17.78	23.62	0.78	0.36	8.74	9.87	0.21	0.12	3.91	4.24	1.55	1.30	2.51	5.36
14	8.31	3.62	0.11	12.04	3.23	2.04	5.53	10.80	2.31	8.76	2.81	13.87	2.58	2.02	1.26	5.86
16	19.39	18.88	0.00	38.27	46.42	22.66	0.79	69.87	52.93	20.23	0.33	73.50	13.55	6.58	0.00	20.13
18	66.37	46.99	0.05	113.41	133.26	137.85	0.00	271.11	362.56	228.57	0.00	591.13	54.39	33.52	0.00	87.90
20	96.85	63.72	0.00	160.57	115.15	92.22	0.08	207.45	557.56	698.41	0.00	1255.97	141.06	124.18	0.00	265.25
22	81.51	63.44	0.00	144.94	117.95	120.09	0.00	238.03	260.01	387.04	0.00	647.05	115.55	123.27	0.00	238.82
24	49.16	50.05	0.00	99.21	67.44	106.44	0.00	173.88	91.63	122.89	0.00	214.51	165.60	80.38	0.00	245.98
26	25.59	33.03	0.00	58.62	15.72	82.79	0.00	98.51	53.99	95.89	0.00	149.88	110.11	66.27	0.00	176.37
28	22.11	21.05	0.00	43.16	9.27	17.36	0.00	26.62	21.46	66.19	0.00	87.65	33.80	104.64	0.00	138.43
30	10.25	9.73	0.00	19.99	2.75	10.77	0.00	13.52	8.10	14.77	0.00	22.87	5.54	79.03	0.00	84.57
32	3.50	4.98	0.00	8.48	2.46	4.50	0.00	6.96	4.85	10.51	0.00	15.36	2.92	27.91	0.00	30.82
34	1.11	2.86	0.00	3.96	2.23	2.06	0.00	4.29	2.69	4.84	0.00	7.54	1.12	17.35	0.00	18.48
36	0.49	0.68	0.00	1.18	0.60	1.49	0.00	2.10	1.25	2.39	0.00	3.64	1.18	5.13	0.00	6.31
38	0.06	0.29	0.00	0.35	0.15	0.03	0.00	0.19	0.60	1.72	0.00	2.31	0.21	0.67	0.00	0.88
40	0.01	0.12	0.00	0.13	0.32	0.37	0.00	0.70	0.06	0.95	0.00	1.01	0.01	0.05	0.00	0.06
42	0.01	0.11	0.00	0.12	0.00	0.04	0.00	0.04	0.06	1.79	0.00	1.85	0.02	0.00	0.00	0.02
44	0.00	0.00	0.00	0.00	0.00	0.31	0.00	0.31	0.00	0.11	0.00	0.11	0.00	0.00	0.00	0.00
46	0.00	0.13	0.00	0.13	0.00	0.00	0.00	0.00	0.00	0.05	0.00	0.05	0.00	0.05	0.00	0.05
48	0.00	0.00	0.00	0.00	0.00	0.30	0.00	0.30	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
50	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
52	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.04	0.00	0.04
54	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.03	0.00	0.03	0.00	0.00	0.00	0.00
56	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
58	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
60	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Total	389.23	322.03	101.39	812.65	517.84	601.84	30.85	1150.53	1420.47	1665.26	19.63	3105.35	649.33	672.46	21.31	1343.10
Nº samples:				52				51				48				51
Nº Ind.:	3957	3147	1372	8476	3016	2723	558	6297	3216	3082	1178	7476	3017	3572	443	7032
Sampled catch:				12283				16615				42525				27586
Range:				5-47				5-49				5-55				5-52
Total catch:				12283				16615				42526				27586
Total valid hauls:				100				98				97				89

Table 14 (cont).- Redfish length distribution per haul mean catches by sex and year. Number per stratified mean catches. Spanish Summer Survey on NAFO 3L: 2008-2015 (R/V *Vizconde de Eza*). Indet. means indeterminate.

Length (cm.)	2012				2013				2014				2015				
	M	F	I	T	M	F	I	T	M	F	I	T	M	F	I	T	
4	0.00	0.00	0.02	0.02	0.00	0.00	0.09	0.09	0.00	0.00	0.38	0.38	0.00	0.00	0.18	0.18	
6	0.00	0.00	11.79	11.79	0.00	0.00	5.15	5.15	0.00	0.00	4.76	4.76	0.00	0.00	66.62	66.62	
8	0.06	0.82	16.26	17.13	0.01	0.00	9.49	9.50	0.00	0.05	6.74	6.79	0.00	0.00	31.23	31.23	
10	3.18	4.43	12.65	20.26	0.06	0.33	10.90	11.30	0.50	0.15	9.67	10.33	0.11	0.53	6.73	7.36	
12	9.66	4.04	4.21	17.91	4.92	2.02	3.30	10.23	2.37	2.20	3.86	8.42	3.45	1.38	7.14	11.97	
14	5.06	2.67	0.94	8.68	9.57	4.45	0.04	14.06	2.69	2.45	0.57	5.71	7.85	7.33	0.61	15.79	
16	12.20	5.77	0.00	17.97	10.71	8.41	0.00	19.12	7.88	4.42	0.00	12.30	5.87	6.21	0.00	12.08	
18	134.16	83.98	0.00	218.14	21.03	10.38	0.00	31.41	20.55	13.12	0.00	33.67	9.82	9.49	0.00	19.31	
20	635.81	404.59	0.00	1040.41	172.77	104.22	0.00	276.99	64.32	49.96	0.00	114.27	44.36	26.09	0.00	70.45	
22	783.26	916.84	0.00	1700.11	247.48	310.10	0.00	557.58	200.92	140.84	0.00	341.76	130.55	68.48	0.00	199.03	
24	279.36	676.30	0.00	955.66	166.92	323.66	0.00	490.58	173.58	217.21	0.00	390.78	116.13	122.70	0.00	238.83	
26	118.77	229.31	0.00	348.08	89.86	137.72	0.00	227.57	127.00	173.62	0.00	300.62	64.97	85.12	0.00	150.09	
28	23.11	113.92	0.00	137.02	27.74	80.12	0.00	107.86	68.06	94.45	0.00	162.51	35.75	54.34	0.00	90.09	
30	6.96	74.74	0.00	81.70	13.10	58.07	0.00	71.17	27.14	57.35	0.00	84.49	10.82	36.53	0.00	47.36	
32	3.54	30.04	0.00	33.58	4.06	22.19	0.00	26.25	8.34	32.35	0.00	40.68	6.52	23.12	0.00	29.64	
34	3.37	6.71	0.00	10.08	3.59	10.79	0.00	14.37	5.01	12.67	0.00	17.68	3.37	15.38	0.00	18.75	
36	1.21	2.74	0.00	3.96	1.19	4.65	0.00	5.84	4.13	4.05	0.00	8.18	1.69	7.33	0.00	9.01	
38	1.21	1.64	0.00	2.85	0.06	2.10	0.00	2.16	2.02	1.93	0.00	3.95	0.43	2.55	0.00	2.98	
40	0.06	0.46	0.00	0.53	0.13	0.10	0.00	0.23	0.13	0.37	0.00	0.50	0.03	0.09	0.00	0.12	
42	0.00	0.01	0.00	0.01	0.00	0.94	0.00	0.94	0.01	0.10	0.00	0.11	0.00	0.01	0.00	0.01	
44	0.00	0.01	0.00	0.01	0.00	1.94	0.00	1.94	0.00	0.08	0.00	0.08	0.00	0.05	0.00	0.05	
46	0.09	0.06	0.00	0.15	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
48	0.00	0.11	0.00	0.11	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	
50	0.00	0.00	0.00	0.00	0.01	0.02	0.00	0.03	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.01	
52	0.03	0.00	0.00	0.03	0.00	0.02	0.00	0.02	0.00	0.01	0.00	0.01	0.00	0.03	0.00	0.03	
54	0.05	0.03	0.00	0.08	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
56	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.01	0.00	0.00	0.00	0.00	
58	0.02	0.00	0.00	0.02	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	
60	0.02	0.00	0.00	0.02	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.01	
Total	2021.20	2559.23	45.88	4626.30	773.21	1082.22	28.97	1884.40	717.47	810.54	26.08	1554.09	441.75	466.78	112.50	1021.03	
Nº samples:					49				52				50				56
Nº Ind.:	3715	3954	502	8171	3635	4233	866	8734	3205	3251	1162	7618	3604	3365	1350	8319	
Sampled catch:					76987				38588				37262				21880
Range:					5-61				5-53				5-56				5-62
Total catch:					76988				38588				37262				21880
Total valid hauls:					98				100				99				97

Table 15.- Swept area, number of hauls and thorny skate mean catch (Kg) and SD (\*\*) by stratum. Spanish Survey on NAFO Div. 3L in the period 2011-2015, on board R/V "Vizconde de Eza".

<b>Stratum</b>	2011				2012				2013				2014				2015			
	Swept Tow area No.	Mean catch	SD	Swept Tow area No.	Mean catch	SD	Swept Tow area No.	Mean catch	SD	Swept Tow area No.	Mean catch	SD	Swept Tow area No.	Mean catch	SD	Swept Tow area No.	Mean catch	SD	Swept Tow area No.	
<b>385</b>	0.0229	2	40.870	7.722	0.0225	2	38.670	8.358	0.0229	2	18.500	15.570	0.0225	2	8.912	1.912	0.0236	2	18.864	4.574
<b>387</b>	0.0450	4	5.241	5.174	0.0450	4	7.559	6.290	0.0450	4	23.395	7.473	0.0461	4	62.785	26.835	0.0458	4	64.881	63.854
<b>388</b>	0.0563	5	9.356	7.705	0.0570	5	42.734	32.557	0.0570	5	32.704	9.754	0.0585	5	70.966	52.957	0.0574	5	97.970	58.681
<b>389</b>	0.0675	6	11.893	10.892	0.0799	7	14.376	12.301	0.0791	7	21.343	11.010	0.0814	7	32.745	32.251	0.0814	7	23.159	12.273
<b>390</b>	0.1009	9	20.264	12.350	0.1354	12	18.599	15.739	0.1358	12	14.574	21.619	0.1369	12	15.477	15.734	0.1260	11	7.690	7.359
<b>391</b>	0.0458	4	32.718	28.277	0.0458	4	38.843	29.385	0.0450	4	37.358	37.052	0.0465	4	36.052	35.657	0.0465	4	52.499	19.346
<b>392</b>	0.0229	2	40.537	19.861	0.0225	2	178.990	196.916	0.0225	2	56.130	25.725	0.0225	2	53.836	58.357	0.0229	2	152.976	185.437
<b>729</b>	0.0338	3	4.906	5.481	0.0338	3	35.344	8.527	0.0341	3	28.835	4.548	0.0338	3	42.980	19.122	0.0345	3	22.367	16.344
<b>730</b>	0.0334	3	1.467	2.540	0.0338	3	3.670	6.357	0.0334	3	11.360	7.412	0.0345	3	22.237	11.856	0.0345	3	6.492	6.242
<b>731</b>	0.0334	3	4.470	5.812	0.0341	3	3.263	2.986	0.0334	3	14.460	9.648	0.0345	3	21.310	15.539	0.0345	3	21.632	12.445
<b>732</b>	0.0454	4	0.000	0.000	0.0454	4	0.000	0.000	0.0450	4	0.848	1.695	0.0454	4	1.980	3.960	0.0465	4	3.333	2.617
<b>733</b>	0.0454	4	2.899	3.869	0.0454	4	5.995	4.874	0.0450	4	18.918	20.706	0.0458	4	32.181	22.484	0.0454	4	6.778	7.155
<b>734</b>	0.0225	2	0.000	0.000	0.0233	2	0.010	0.014	0.0221	2	0.000	0.000	0.0225	2	0.000	0.000	0.0225	2	0.000	0.000
<b>741</b>	0.0218	2	0.000	0.000	0.0218	2	0.000	0.000	0.0221	2	0.000	0.000	0.0225	2	0.000	0.000	0.0236	2	0.000	0.000
<b>742</b>	0.0225	2	0.000	0.000	0.0206	2	0.000	0.000	0.0218	2	0.000	0.000	0.0221	2	0.000	0.000	0.0233	2	0.000	0.000
<b>743</b>	0.0221	2	0.000	0.000	0.0206	2	0.000	0.000	0.0218	2	0.000	0.000	0.0221	2	0.000	0.000	0.0233	2	0.000	0.000
<b>744</b>	0.0221	2	0.000	0.000	0.0221	2	0.000	0.000	0.0221	2	0.000	0.000	0.0225	2	0.000	0.000	0.0225	2	0.000	0.000
<b>745</b>	0.0446	4	0.000	0.000	0.0570	5	0.004	0.008	0.0559	5	0.000	0.000	0.0578	5	0.000	0.000	0.0578	5	0.000	0.000
<b>746</b>	0.0566	5	0.000	0.000	0.0675	6	0.000	0.000	0.0675	6	0.000	0.000	0.0683	6	0.000	0.000	0.0686	6	0.000	0.000
<b>747</b>	0.0893	8	0.424	1.199	0.1121	10	0.000	0.000	0.1125	10	0.000	0.000	0.1125	10	0.559	1.227	0.1028	9	0.698	2.095
<b>748</b>	0.0221	2	0.000	0.000	0.0225	2	0.000	0.000	0.0225	2	0.000	0.000	0.0229	2	1.530	2.164	0.0233	2	0.000	0.000
<b>749</b>	0.0221	2	0.000	0.000	0.0221	2	0.000	0.000	0.0225	2	0.000	0.000	0.0225	2	0.000	0.000	0.0225	2	0.000	0.000
<b>750</b>	0.0668	6	0.000	0.000	0.0885	8	0.000	0.000	0.0896	8	0.493	1.393	0.0904	8	0.000	0.000	0.0934	8	0.000	0.000
<b>751</b>	0.0334	3	0.000	0.000	0.0218	2	0.000	0.000	0.0446	4	0.154	0.308	0.0334	3	0.000	0.000	0.0341	3	0.000	0.000

$$(**) SD = \frac{\sum (x_i - \bar{x})}{n-1}$$

Table 16.- Length-weight relationships in the calculation of biomass, for Division 3L (out ZEE Canada), 2007-2015 for thorny skate and black dogfish. The equation is Weight=a(Length+0.5)b. To calculate the parameters for the indeterminate individuals, we used the total data (males+females+indeterminate individuals).

Thorny skate				Black dogfish								
Year	Sex	L-W Equations	N	r <sup>2</sup>	Sex	L-W Equations	N	r <sup>2</sup>	Sex	L-W Equations	N	r <sup>2</sup>
2007	All	$W = 0.0080 L^{3.0609}$	539	0.9848		$W = 0.0008 L^{3.4421}$	362	0.9155				
	Males	$W = 0.0091 L^{3.0242}$	255	0.9868		$W = 0.0099 L^{2.8281}$	147	0.9029				
	Females	$W = 0.0072 L^{3.0929}$	284	0.9839		$W = 0.0006 L^{3.5445}$	215	0.9373				
2008	All	$W = 0.0071 L^{3.0883}$	598	0.9884		$W = 0.0014 L^{3.3183}$	279	0.9006				
	Males	$W = 0.0077 L^{3.0618}$	282	0.9903		$W = 0.0087 L^{2.8575}$	160	0.8956				
	Females	$W = 0.0064 L^{3.1175}$	316	0.9867		$W = 0.0008 L^{3.4541}$	119	0.9283				
2009	All	$W = 0.0072 L^{3.0862}$	283	0.9864		$W = 0.0007 L^{3.4922}$	236	0.9246				
	Males	$W = 0.0093 L^{3.0231}$	171	0.9848		$W = 0.0132 L^{2.7605}$	75	0.8865				
	Females	$W = 0.0057 L^{3.1507}$	112	0.9881		$W = 0.0007 L^{3.5184}$	161	0.9465				
2010	All	$W = 0.0060 L^{3.1361}$	290	0.9906		$W = 0.0019 L^{3.2510}$	299	0.9506				
	Males	$W = 0.0060 L^{3.1285}$	149	0.9892		$W = 0.0137 L^{2.7559}$	130	0.9408				
	Females	$W = 0.0056 L^{3.1630}$	141	0.9927		$W = 0.0012 L^{3.3617}$	169	0.9637				
2011	All	$W = 0.0031 L^{3.2899}$	218	0.9937		$W = 0.0020 L^{3.2316}$	455	0.9518				
	Males	$W = 0.0036 L^{3.2468}$	136	0.9941		$W = 0.0059 L^{2.9580}$	171	0.9493				
	Females	$W = 0.0024 L^{3.3657}$	82	0.9941		$W = 0.0014 L^{3.3220}$	284	0.9568				
2012	All	$W = 0.0065 L^{3.1140}$	352	0.9918		$W = 0.0019 L^{3.2460}$	242	0.9531				
	Males	$W = 0.0085 L^{3.0429}$	219	0.9925		$W = 0.0107 L^{2.8100}$	116	0.9571				
	Females	$W = 0.0040 L^{3.2467}$	133	0.9933		$W = 0.0010 L^{3.4151}$	126	0.9718				
2013	All	$W = 0.0057 L^{3.1365}$	336	0.9926		$W = 0.0007 L^{3.4877}$	352	0.9275				
	Males	$W = 0.0054 L^{3.1470}$	218	0.9914		$W = 0.0084 L^{2.8679}$	81	0.8884				
	Females	$W = 0.0054 L^{3.1631}$	118	0.9955		$W = 0.007 L^{3.4843}$	271	0.9385				
2014	All	$W = 0.0066 L^{3.1037}$	577	0.9836		$W = 0.0010 L^{3.3969}$	259	0.9283				
	Males	$W = 0.0077 L^{3.0639}$	402	0.9764		$W = 0.0067 L^{2.9222}$	77	0.9222				
	Females	$W = 0.0049 L^{3.1865}$	175	0.994		$W = 0.009 L^{3.4286}$	182	0.9338				
2015	All	$W = 0.0064 L^{3.1098}$	532	0.9944		$W = 0.0013 L^{3.3416}$	578	0.9544				
	Males	$W = 0.0075 L^{3.0685}$	337	0.9945		$W = 0.0056 L^{2.9683}$	178	0.959				
	Females	$W = 0.0050 L^{3.1760}$	195	0.9941		$W = 0.0011 L^{3.4038}$	400	0.9604				

Table 17.- Stratified mean catches (Kg) of thorny skate by stratum and year (2003-2015) and SD. Research Vessel *Vizconde de Eza*. n.s. means stratum not surveyed. In 2003: the data correspond to 69% of the total area prospected in 2006-2015.

Stratum	Survey												
	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
385	0.00	831.90	-	713.19	3570.68	4437.69	2696.89	499.14	4822.66	4563.06	2183.00	1051.62	2225.95
387	1355.52	2739.20	-	4208.00	8316.16	6726.59	5271.04	5721.60	1341.76	1935.04	5989.12	16072.83	16609.41
388	4738.58	5961.90	-	15774.40	11101.27	13261.69	11952.50	12470.58	3340.16	15256.04	11675.33	25334.72	34975.36
389	3045.60	5548.10	-	16786.09	13163.25	16830.16	6593.66	13829.31	6053.28	7317.60	10863.51	16667.21	11788.00
390	154.85	1627.28	-	4506.21	6003.36	4110.66	11444.98	10513.50	16515.07	15158.46	11878.15	12613.48	6267.28
391	485.98	18118.50	-	42606.68	28385.42	53804.19	8995.45	6779.63	9226.41	10953.66	10534.89	10166.52	14804.72
392	1457.25	9033.50	-	21677.50	47864.50	23090.82	5991.69	5325.49	5877.79	25953.48	8138.85	7806.15	22181.52
729	10221.63	26109.75	-	9162.48	30645.36	6373.35	7084.74	1200.20	912.52	6573.92	5363.25	7994.28	4160.32
730	12138.00	0.00	-	739.22	0.00	0.00	0.00	2.04	249.33	623.90	1931.20	3780.23	1103.58
731	8360.28	3998.16	-	10099.44	12408.84	1974.24	4934.88	2400.70	965.52	704.74	3123.36	4602.96	4672.51
732	17602.20	0.00	-	465.47	0.00	167.94	1640.10	0.00	0.00	0.00	195.77	457.38	769.98
733	n.s.	2191.02	-	3410.14	1503.84	3438.05	1009.71	1304.02	678.31	1402.83	4426.70	7530.41	1585.94
734	n.s.	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	1.53	0.00	0.00	0.00
741	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
742	0.00	0.00	-	0.00	0.00	0.00	0.00	0.70	0.00	0.00	0.00	0.00	0.00
743	n.s.	0.00	-	0.00	0.00	0.00	71.15	0.00	0.00	0.00	0.00	0.00	0.00
744	n.s.	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
745	7682.68	0.00	-	0.00	0.00	0.00	0.00	226.20	0.00	1.32	0.00	0.00	0.00
746	908.46	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
747	n.s.	0.00	-	0.00	0.00	0.00	0.00	0.00	306.80	0.00	0.00	404.72	505.67
748	10369.98	0.00	-	133.03	0.00	0.00	0.00	0.00	0.00	0.00	0.00	243.27	0.00
749	1015.56	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
750	n.s.	764.50	-	218.69	0.00	0.00	0.00	0.00	0.00	0.00	273.83	0.00	0.00
751	n.s.	n.s.	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	35.21	0.00	0.00
TOTAL	79536.57	76923.81	-	130500.54	162962.67	134215.36	67686.78	60273.11	50289.61	90445.57	76612.16	114725.78	121650.25
( $\bar{y}$ )	17.78	12.29	-	20.12	25.12	20.69	10.43	9.29	7.75	13.94	11.81	17.69	18.75
SD	2.41	4.54	-	3.27	5.19	1.92	1.44	1.30	0.98	3.36	1.36	2.25	3.58

Table 18.- Survey estimates (by the swept area method) of thorny skate biomass (t.) by stratum and year and their SD on NAFO Div. 3L (R/V *Vizconde de Eza*). n.s. means stratum not surveyed. In 2003, the data correspond to 69% of the total area prospected in 2006-2015.

Stratum	Survey												
	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
385	0	73	-	62	317	388	240	44	422	406	191	93	188
387	119	256	-	374	739	619	481	500	119	172	532	1394	1452
388	426	568	-	1393	987	1187	1077	1094	297	1338	1024	2165	3048
389	268	493	-	1478	1170	1510	575	1218	538	641	961	1434	1014
390	14	142	-	397	534	354	1001	926	1473	1344	1050	1106	547
391	43	1666	-	3787	2523	4743	786	598	807	958	936	875	1274
392	125	845	-	1895	4255	2087	524	473	514	2307	723	694	1939
729	973	2360	-	814	2724	567	623	107	81	584	471	711	362
730	1097	0	-	68	0	0	0	0	22	55	174	329	96
731	731	344	-	888	1103	179	434	213	87	62	281	400	406
732	1565	0	-	42	0	15	146	0	0	0	17	40	66
733	n.s.	199	-	301	134	319	90	116	60	124	393	658	140
734	n.s.	0	-	0	0	0	0	0	0	0	0	0	0
741	0	0	-	0	0	0	0	0	0	0	0	0	0
742	0	0	-	0	0	0	0	0	0	0	0	0	0
743	n.s.	0	-	0	0	0	7	0	0	0	0	0	0
744	n.s.	0	-	0	0	0	0	0	0	0	0	0	0
745	675	0	-	0	0	0	0	20	0	0	0	0	0
746	81	0	-	0	0	0	0	0	0	0	0	0	0
747	n.s.	0	-	0	0	0	0	0	28	0	0	36	44
748	954	0	-	12	0	0	0	0	0	0	0	21	0
749	92	0	-	0	0	0	0	0	0	0	0	0	0
750	n.s.	85	-	20	0	0	0	0	0	0	0	24	0
751	n.s.	n.s.	-	0	0	0	0	0	0	0	3	0	0
TOTAL	7164	7031	-	11531	14486	11968	5982	5310	4448	7991	6783	9956	10577
SD	942	2642	-	1887	2993	1124	808	740	560	2008	779	1263	1981

Table 19.- Thorny skate length distribution per haul mean catches by sex and year. Number per stratified mean catches. Spanish Summer Survey on NAFO 3L: 2008-2015 (R/V *Vizconde de Eza*). Indet. means indeterminate.

Length (cm.)	2008				2009				2010				2011				
	M	F	I	T	M	F	I	T	M	F	I	T	M	F	I	T	
10	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	
12	0.09	0.11	0.00	0.20	0.11	0.05	0.00	0.16	0.05	0.05	0.00	0.10	0.09	0.03	0.00	0.11	
14	0.12	0.09	0.00	0.20	0.06	0.07	0.00	0.13	0.08	0.07	0.00	0.15	0.06	0.08	0.00	0.14	
16	0.03	0.03	0.00	0.06	0.02	0.02	0.00	0.04	0.00	0.03	0.00	0.03	0.01	0.00	0.00	0.01	
18	0.04	0.01	0.00	0.05	0.00	0.05	0.00	0.05	0.01	0.02	0.00	0.03	0.00	0.01	0.00	0.01	
20	0.09	0.01	0.00	0.10	0.01	0.03	0.00	0.04	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	
22	0.02	0.01	0.00	0.03	0.00	0.00	0.00	0.00	0.00	0.02	0.00	0.02	0.00	0.00	0.00	0.00	
24	0.02	0.02	0.00	0.04	0.02	0.02	0.00	0.05	0.01	0.00	0.00	0.01	0.00	0.00	0.00	0.00	
26	0.08	0.07	0.00	0.14	0.01	0.00	0.00	0.01	0.01	0.02	0.00	0.03	0.00	0.00	0.00	0.00	
28	0.02	0.05	0.00	0.08	0.02	0.01	0.00	0.03	0.00	0.01	0.00	0.01	0.00	0.03	0.00	0.03	
30	0.04	0.05	0.00	0.10	0.02	0.03	0.00	0.06	0.04	0.02	0.00	0.07	0.02	0.01	0.00	0.03	
32	0.07	0.05	0.00	0.12	0.00	0.01	0.00	0.01	0.00	0.03	0.00	0.03	0.01	0.00	0.00	0.01	
34	0.05	0.04	0.00	0.10	0.01	0.01	0.00	0.02	0.01	0.04	0.00	0.05	0.00	0.01	0.00	0.01	
36	0.03	0.05	0.00	0.08	0.00	0.00	0.00	0.00	0.04	0.02	0.00	0.07	0.02	0.01	0.00	0.03	
38	0.01	0.03	0.00	0.04	0.02	0.01	0.00	0.03	0.02	0.01	0.00	0.03	0.00	0.02	0.00	0.02	
40	0.05	0.01	0.00	0.06	0.02	0.00	0.00	0.02	0.00	0.01	0.00	0.01	0.01	0.01	0.00	0.03	
42	0.02	0.05	0.00	0.07	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	
44	0.01	0.02	0.00	0.03	0.01	0.04	0.00	0.05	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	
46	0.03	0.06	0.00	0.09	0.00	0.01	0.00	0.01	0.01	0.00	0.00	0.01	0.00	0.00	0.00	0.00	
48	0.02	0.01	0.00	0.03	0.01	0.02	0.00	0.03	0.01	0.00	0.00	0.01	0.00	0.00	0.00	0.00	
50	0.06	0.03	0.00	0.09	0.05	0.01	0.00	0.06	0.00	0.01	0.00	0.01	0.01	0.00	0.00	0.01	
52	0.07	0.08	0.00	0.15	0.02	0.02	0.00	0.04	0.01	0.02	0.00	0.03	0.00	0.00	0.00	0.00	
54	0.08	0.09	0.00	0.17	0.05	0.05	0.00	0.09	0.00	0.02	0.00	0.02	0.00	0.01	0.00	0.01	
56	0.03	0.13	0.00	0.16	0.02	0.15	0.00	0.17	0.02	0.04	0.00	0.07	0.00	0.01	0.00	0.01	
58	0.12	0.22	0.00	0.34	0.13	0.09	0.00	0.22	0.08	0.09	0.00	0.17	0.02	0.05	0.00	0.07	
60	0.22	0.28	0.00	0.50	0.16	0.08	0.00	0.24	0.03	0.12	0.00	0.16	0.06	0.09	0.00	0.15	
62	0.29	0.35	0.00	0.65	0.23	0.24	0.00	0.47	0.08	0.10	0.00	0.18	0.03	0.08	0.00	0.10	
64	0.35	0.45	0.00	0.81	0.23	0.14	0.00	0.36	0.12	0.16	0.00	0.28	0.15	0.06	0.00	0.20	
66	0.39	0.45	0.00	0.84	0.25	0.18	0.00	0.43	0.21	0.18	0.00	0.38	0.13	0.09	0.00	0.23	
68	0.32	0.44	0.00	0.76	0.28	0.18	0.00	0.47	0.19	0.23	0.00	0.42	0.19	0.12	0.00	0.31	
70	0.25	0.37	0.00	0.62	0.19	0.07	0.00	0.26	0.21	0.07	0.00	0.28	0.17	0.15	0.00	0.32	
72	0.19	0.15	0.00	0.34	0.17	0.09	0.00	0.25	0.13	0.08	0.00	0.21	0.18	0.03	0.00	0.22	
74	0.26	0.16	0.00	0.42	0.19	0.01	0.00	0.20	0.11	0.05	0.00	0.16	0.16	0.05	0.00	0.21	
76	0.10	0.13	0.00	0.23	0.02	0.03	0.00	0.06	0.09	0.03	0.00	0.12	0.11	0.00	0.00	0.11	
78	0.09	0.03	0.00	0.12	0.04	0.03	0.00	0.07	0.09	0.01	0.00	0.10	0.05	0.00	0.00	0.05	
80	0.07	0.00	0.00	0.07	0.01	0.00	0.00	0.01	0.03	0.00	0.00	0.03	0.04	0.00	0.00	0.04	
82	0.05	0.02	0.00	0.07	0.01	0.00	0.00	0.01	0.02	0.00	0.00	0.02	0.01	0.00	0.00	0.01	
84	0.03	0.00	0.00	0.03	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.01	
86	0.02	0.00	0.00	0.02	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
88	0.01	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.01	0.04	0.00	0.00	0.04	
Total	5.10	6.71	0.00	11.81	2.41	1.76	0.00	4.17	1.73	1.62	0.00	3.35	1.59	0.96	0.00	2.55	
Nº samples:					43				44				46				39
Nº Ind.:	457	621	0	1078	211	156	0	367	159	145	0	304	136	82	0	218	
Sampled catch:					2325				996.2				853				663
Range:					12-82				12-82				12-88				11-88
Total catch:					2325				996.2				853				663
Total valid hauls:					94				98				97				89

Table 19.- Thorny skate length distribution per haul mean catches by sex and year. Number per stratified mean catches. Spanish Summer Survey on NAFO 3L: 2008-2015 (R/V *Vizconde de Eza*). Indet. means indeterminate.

Length (cm.)	2012				2013				2014				2015			
	M	F	I	T	M	F	I	T	M	F	I	T	M	F	I	T
10	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
12	0.09	0.05	0.00	0.14	0.01	0.06	0.00	0.07	0.01	0.02	0.00	0.03	0.02	0.04	0.00	0.06
14	0.07	0.05	0.00	0.11	0.10	0.03	0.00	0.13	0.12	0.12	0.00	0.24	0.03	0.11	0.00	0.14
16	0.01	0.00	0.00	0.01	0.03	0.06	0.00	0.10	0.02	0.04	0.00	0.06	0.02	0.03	0.00	0.05
18	0.03	0.00	0.00	0.03	0.02	0.01	0.00	0.03	0.06	0.01	0.00	0.07	0.03	0.11	0.00	0.14
20	0.04	0.01	0.00	0.05	0.00	0.03	0.00	0.03	0.05	0.09	0.00	0.14	0.12	0.09	0.00	0.21
22	0.03	0.03	0.00	0.06	0.00	0.00	0.00	0.00	0.08	0.06	0.00	0.14	0.23	0.08	0.00	0.31
24	0.05	0.06	0.00	0.11	0.00	0.00	0.00	0.00	0.08	0.04	0.00	0.12	0.21	0.15	0.00	0.36
26	0.04	0.03	0.00	0.07	0.00	0.01	0.00	0.01	0.10	0.09	0.00	0.19	0.30	0.28	0.00	0.58
28	0.03	0.11	0.00	0.14	0.02	0.00	0.00	0.02	0.03	0.11	0.00	0.14	0.33	0.27	0.00	0.60
30	0.08	0.01	0.00	0.09	0.00	0.00	0.00	0.00	0.17	0.11	0.00	0.27	0.39	0.40	0.00	0.79
32	0.04	0.04	0.00	0.08	0.00	0.00	0.00	0.00	0.13	0.08	0.00	0.20	0.38	0.27	0.00	0.65
34	0.04	0.04	0.00	0.08	0.01	0.02	0.00	0.03	0.07	0.06	0.00	0.12	0.27	0.24	0.00	0.51
36	0.06	0.06	0.00	0.12	0.00	0.03	0.00	0.03	0.06	0.08	0.00	0.14	0.24	0.19	0.00	0.43
38	0.06	0.04	0.00	0.10	0.00	0.02	0.00	0.02	0.10	0.10	0.00	0.20	0.13	0.10	0.00	0.23
40	0.07	0.05	0.00	0.12	0.02	0.04	0.00	0.06	0.11	0.04	0.00	0.15	0.03	0.08	0.00	0.11
42	0.06	0.01	0.00	0.07	0.02	0.04	0.00	0.06	0.12	0.04	0.00	0.17	0.13	0.10	0.00	0.23
44	0.02	0.02	0.00	0.05	0.06	0.03	0.00	0.10	0.15	0.10	0.00	0.25	0.12	0.10	0.00	0.22
46	0.02	0.03	0.00	0.05	0.06	0.05	0.00	0.11	0.20	0.08	0.00	0.29	0.11	0.10	0.00	0.21
48	0.01	0.00	0.00	0.01	0.05	0.01	0.00	0.06	0.19	0.10	0.00	0.29	0.08	0.10	0.00	0.17
50	0.02	0.05	0.00	0.07	0.05	0.00	0.00	0.05	0.14	0.03	0.00	0.17	0.14	0.21	0.00	0.34
52	0.05	0.03	0.00	0.08	0.06	0.00	0.00	0.06	0.18	0.09	0.00	0.26	0.12	0.09	0.00	0.21
54	0.02	0.01	0.00	0.03	0.02	0.03	0.00	0.06	0.12	0.02	0.00	0.14	0.12	0.09	0.00	0.21
56	0.01	0.00	0.00	0.01	0.02	0.03	0.00	0.05	0.13	0.06	0.00	0.19	0.12	0.04	0.00	0.16
58	0.03	0.04	0.00	0.07	0.04	0.02	0.00	0.06	0.06	0.04	0.00	0.11	0.05	0.10	0.00	0.16
60	0.03	0.05	0.00	0.08	0.04	0.03	0.00	0.07	0.09	0.09	0.00	0.17	0.12	0.09	0.00	0.21
62	0.07	0.10	0.00	0.18	0.10	0.09	0.00	0.18	0.11	0.02	0.00	0.13	0.11	0.12	0.00	0.23
64	0.10	0.11	0.00	0.21	0.07	0.14	0.00	0.21	0.08	0.06	0.00	0.14	0.16	0.07	0.00	0.24
66	0.10	0.14	0.00	0.24	0.11	0.14	0.00	0.25	0.24	0.19	0.00	0.42	0.19	0.18	0.00	0.37
68	0.28	0.22	0.00	0.50	0.25	0.17	0.00	0.42	0.27	0.10	0.00	0.37	0.32	0.09	0.00	0.41
70	0.30	0.09	0.00	0.38	0.23	0.17	0.00	0.39	0.31	0.12	0.00	0.44	0.32	0.10	0.00	0.42
72	0.37	0.12	0.00	0.49	0.23	0.06	0.00	0.30	0.36	0.15	0.00	0.51	0.40	0.08	0.00	0.48
74	0.23	0.03	0.00	0.26	0.24	0.02	0.00	0.26	0.31	0.07	0.00	0.39	0.31	0.04	0.00	0.35
76	0.19	0.07	0.00	0.26	0.24	0.06	0.00	0.30	0.33	0.04	0.00	0.36	0.32	0.03	0.00	0.35
78	0.21	0.03	0.00	0.24	0.17	0.01	0.00	0.18	0.19	0.04	0.00	0.23	0.27	0.02	0.00	0.29
80	0.14	0.01	0.00	0.15	0.14	0.00	0.00	0.14	0.22	0.01	0.00	0.23	0.19	0.04	0.00	0.23
82	0.09	0.01	0.00	0.10	0.07	0.00	0.00	0.07	0.13	0.00	0.00	0.13	0.03	0.00	0.00	0.03
84	0.07	0.00	0.00	0.07	0.02	0.00	0.00	0.02	0.06	0.00	0.00	0.06	0.08	0.00	0.00	0.08
86	0.02	0.00	0.00	0.02	0.01	0.00	0.00	0.01	0.05	0.00	0.00	0.05	0.03	0.00	0.00	0.03
88	0.01	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.01	0.01	0.00	0.00	0.01
Total	3.19	1.74	0.00	4.93	2.53	1.44	0.00	3.97	5.23	2.51	0.00	7.74	6.59	4.24	0.00	10.83
Nº samples:					44			49				50				49
Nº Ind.:	266	151	0	417	225	117	0	342	474	217	0	691	607	390	0	997
Sampled catch:					1309			1128				1695				1748
Range:					12-88			13-86				13-89				12-91
Total catch:					1309			1128				1695				1748
Total valid hauls:					98			100				99				97

Table20.- Swept area, number of hauls and black dogfish mean catch (Kg) and SD (\*\*) by stratum. Spanish Survey on NAFO Div. 3L in the period 2011-2015, on board R/V "Vizconde de Eza".

Stratum	2011				2012				2013				2014				2015			
	Swept area	Tow No.	Mean catch	SD	Swept area	Tow No.	Mean catch	SD	Swept area	Tow No.	Mean catch	SD	Swept area	Tow No.	Mean catch	SD	Swept area	Tow No.	Mean catch	SD
385	0.0229	2	0.000	0.000	0.0225	2	0.000	0.000	0.0229	2	0.000	0.000	0.0225	2	0.000	0.000	0.0236	2	0.000	0.000
387	0.0450	4	0.000	0.000	0.0450	4	0.000	0.000	0.0450	4	0.000	0.000	0.0461	4	0.000	0.000	0.0458	4	0.000	0.000
388	0.0563	5	0.000	0.000	0.0570	5	0.000	0.000	0.0570	5	0.000	0.000	0.0585	5	0.000	0.000	0.0574	5	0.000	0.000
389	0.0675	6	0.000	0.000	0.0799	7	0.000	0.000	0.0791	7	0.000	0.000	0.0814	7	0.000	0.000	0.0814	7	0.000	0.000
390	0.1009	9	0.000	0.000	0.1354	12	0.000	0.000	0.1358	12	0.000	0.000	0.1369	12	0.000	0.000	0.1260	11	0.000	0.000
391	0.0458	4	0.000	0.000	0.0458	4	0.000	0.000	0.0450	4	0.000	0.000	0.0465	4	0.000	0.000	0.0465	4	0.000	0.000
392	0.0229	2	0.000	0.000	0.0225	2	0.000	0.000	0.0225	2	0.000	0.000	0.0225	2	0.000	0.000	0.0229	2	0.000	0.000
729	0.0338	3	0.000	0.000	0.0338	3	0.000	0.000	0.0341	3	0.000	0.000	0.0338	3	0.000	0.000	0.0345	3	0.000	0.000
730	0.0334	3	3.646	6.315	0.0338	3	10.040	17.053	0.0334	3	0.000	0.000	0.0345	3	0.000	0.000	0.0345	3	16.964	28.977
731	0.0334	3	0.000	0.000	0.0341	3	0.000	0.000	0.0334	3	0.000	0.000	0.0345	3	0.000	0.000	0.0345	3	0.000	0.000
732	0.0454	4	0.000	0.000	0.0454	4	0.000	0.000	0.0450	4	0.000	0.000	0.0454	4	0.000	0.000	0.0465	4	0.000	0.000
733	0.0454	4	0.000	0.000	0.0454	4	0.000	0.000	0.0450	4	0.000	0.000	0.0458	4	0.000	0.000	0.0454	4	0.000	0.000
734	0.0225	2	0.000	0.000	0.0233	2	0.000	0.000	0.0221	2	0.000	0.000	0.0225	2	0.000	0.000	0.0225	2	0.000	0.000
741	0.0218	2	0.000	0.000	0.0218	2	0.000	0.000	0.0221	2	0.000	0.000	0.0225	2	0.000	0.000	0.0236	2	0.000	0.000
742	0.0225	2	0.000	0.000	0.0206	2	0.000	0.000	0.0218	2	0.000	0.000	0.0221	2	0.598	0.845	0.0233	2	0.523	0.740
743	0.0221	2	0.000	0.000	0.0206	2	0.000	0.000	0.0218	2	0.945	1.336	0.0221	2	2.505	3.543	0.0233	2	5.060	2.206
744	0.0221	2	0.612	0.865	0.0221	2	0.000	0.000	0.0221	2	3.550	5.020	0.0225	2	0.000	0.000	0.0225	2	0.506	0.716
745	0.0446	4	0.705	1.410	0.0570	5	0.000	0.000	0.0559	5	0.620	1.386	0.0578	5	0.000	0.000	0.0578	5	51.731115.643	
746	0.0566	5	7.160	9.335	0.0675	6	6.004	4.804	0.0675	6	26.233	40.751	0.0683	6	10.215	14.886	0.0686	6	29.042	14.767
747	0.0893	8	5.204	3.122	0.1121	10	4.889	4.861	0.1125	10	11.874	6.025	0.1125	10	11.466	4.719	0.1028	9	7.979	7.512
748	0.0221	2	135.930	187.058	0.0225	2	25.190	35.624	0.0225	2	25.780	36.458	0.0229	2	63.850	2.758	0.0233	2	100.365117.401	
749	0.0221	2	114.000	69.141	0.0221	2	70.633	84.905	0.0225	2	42.515	34.104	0.0225	2	66.725	41.260	0.0225	2	107.620	22.316
750	0.0668	6	1.711	2.351	0.0885	8	4.283	6.729	0.0896	8	7.622	10.816	0.0904	8	12.006	13.261	0.0934	8	11.718	17.339
751	0.0334	3	3.076	2.976	0.0218	2	9.550	5.388	0.0446	4	7.797	3.881	0.0334	3	3.267	3.348	0.0341	3	15.593	0.655

$$(**) SD = \frac{\sum (x_i - \bar{x})}{n-1}$$

Table 21.- Stratified mean catches (Kg) of black dogfish by stratum and year (2003-2015) and SD. Research Vessel *Vizconde de Eza*. n.s. means stratum not surveyed. In 2003: the data correspond to 69% of the total area prospected in 2006-2015.

Stratum	Survey												
	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
385	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
387	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
388	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
389	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
390	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
391	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
392	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
729	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
730	0.00	369.75	-	627.30	3312.88	4652.33	5262.97	3338.80	619.82	1706.80	0.00	0.00	2883.94
731	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
732	0.00	0.00	-	0.00	0.00	0.00	0.00	69.30	0.00	0.00	0.00	0.00	0.00
733	n.s	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
734	n.s	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
741	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
742	0.00	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	38.24	33.47
743	n.s	31.90	-	0.00	0.00	0.00	93.59	0.00	0.00	0.00	48.20	127.76	258.06
744	n.s	0.00	-	47.85	109.73	58.08	28.38	0.00	40.39	0.00	234.30	0.00	33.40
745	2.32	0.00	-	0.00	0.00	0.00	0.00	0.00	245.34	0.00	215.76	0.00	18002.53
746	0.00	0.00	-	3541.07	3594.84	2407.60	1544.22	1888.13	2806.72	2353.63	10283.47	4004.35	11384.59
747	n.s	2944.27	-	2646.94	4354.53	4267.26	4816.77	4318.66	3767.42	3539.64	8596.56	8301.38	5776.96
748	0.00	5879.82	-	2498.42	5694.85	12847.20	1946.16	13283.66	21612.87	4005.21	4099.02	10152.15	15958.04
749	27688.50	2179.80	-	11481.75	28942.20	4461.66	16517.34	18738.09	14364.00	8899.76	5356.89	8407.35	13560.12
750	n.s	1556.80	-	3454.61	7772.42	6875.64	5085.02	471.21	951.50	2381.07	4237.97	6675.48	6514.93
751	n.s	n.s	-	252.47	1008.75	865.62	1223.62	428.23	704.48	2186.95	1785.40	748.07	3570.87
TOTAL ( $\bar{y}$ )	27690.82	12962.34	-	24550.42	54790.18	36435.38	36518.07	42536.08	45112.55	25073.06	34857.56	38454.77	77976.90
SD	6.19	2.07	-	3.78	8.45	5.62	5.63	6.56	6.95	3.87	5.37	5.93	12.02

Table 22.- Survey estimates (by the swept area method) of black dogfish biomass (t.) by stratum and year and their SD on NAFO Div. 3L (R/V *Vizconde de Eza*).  
 n.s. means stratum not surveyed. In 2003, the data correspond to 69% of the total area prospected in 2006-2015.


Stratum	Survey												
	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
385	0	0	-	0	0	0	0	0	0	0	0	0	0
387	0	0	-	0	0	0	0	0	0	0	0	0	0
388	0	0	-	0	0	0	0	0	0	0	0	0	0
389	0	0	-	0	0	0	0	0	0	0	0	0	0
390	0	0	-	0	0	0	0	0	0	0	0	0	0
391	0	0	-	0	0	0	0	0	0	0	0	0	0
392	0	0	-	0	0	0	0	0	0	0	0	0	0
729	0	0	-	0	0	0	0	0	0	0	0	0	0
730	0	33	-	58	294	433	468	300	56	152	0	0	251
731	0	0	-	0	0	0	0	0	0	0	0	0	0
732	0	0	-	0	0	0	0	6	0	0	0	0	0
733	n.s.	0	-	0	0	0	0	0	0	0	0	0	0
734	n.s.	0	-	0	0	0	0	0	0	0	0	0	0
741	0	0	-	0	0	0	0	0	0	0	0	0	0
742	0	0	-	0	0	0	0	0	0	0	0	3	3
743	n.s.	3	-	0	0	0	9	0	0	0	4	12	22
744	n.s.	0	-	4	10	5	3	0	4	0	21	0	3
745	0	0	-	0	0	0	0	0	22	0	19	0	1559
746	0	0	-	315	325	227	139	167	248	209	914	352	995
747	n.s.	287	-	237	387	399	431	384	338	316	764	738	506
748	0	592	-	230	506	1181	170	1181	1954	356	364	888	1373
749	2503	197	-	1004	2573	417	1468	1638	1298	804	476	747	1205
750	n.s.	173	-	309	687	652	450	42	86	215	378	591	558
751	n.s.	n.s.	-	22	90	84	109	38	63	201	160	67	314
TOTAL	2503	1286	-	2176	4872	3399	3247	3756	4068	2253	3102	3398	6789
SD	2546	695	-	994	721	1296	1340	1634	1964	819	773	466	2012

Table 23.- Black dogfish length distribution per haul mean catches by sex and year. Number per stratified mean catches. Spanish Summer Survey on NAFO 3L: 2008-2015 (R/V *Vizconde de Eza*). Indet. means indeterminate.


Length (cm.)	2008				2009				2010				2011			
	M	F	I	T	M	F	I	T	M	F	I	T	M	F	I	T
16	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
18	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
20	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
22	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
24	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
26	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
28	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
30	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
32	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
34	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
36	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.01	0.00	0.01	0.00	0.01
38	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.00	0.03	0.00
40	0.01	0.00	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	0.01	0.05	0.00	0.06
42	0.02	0.00	0.00	0.02	0.01	0.04	0.00	0.05	0.06	0.04	0.00	0.09	0.09	0.05	0.00	0.14
44	0.01	0.04	0.00	0.05	0.04	0.03	0.00	0.07	0.05	0.09	0.00	0.13	0.08	0.13	0.00	0.20
46	0.04	0.06	0.00	0.09	0.01	0.02	0.00	0.03	0.08	0.11	0.00	0.19	0.10	0.10	0.00	0.20
48	0.03	0.01	0.00	0.04	0.04	0.02	0.00	0.06	0.05	0.07	0.00	0.12	0.18	0.15	0.00	0.33
50	0.07	0.03	0.00	0.10	0.03	0.08	0.00	0.11	0.06	0.06	0.00	0.12	0.14	0.11	0.00	0.25
52	0.09	0.08	0.00	0.17	0.11	0.10	0.00	0.21	0.12	0.09	0.00	0.21	0.12	0.08	0.00	0.20
54	0.18	0.10	0.00	0.28	0.13	0.10	0.00	0.23	0.09	0.10	0.00	0.19	0.07	0.10	0.00	0.17
56	0.19	0.12	0.00	0.30	0.18	0.15	0.00	0.33	0.13	0.14	0.00	0.27	0.23	0.23	0.00	0.47
58	0.28	0.15	0.00	0.43	0.19	0.17	0.00	0.37	0.24	0.11	0.00	0.36	0.38	0.25	0.00	0.64
60	0.55	0.16	0.00	0.71	0.28	0.20	0.00	0.49	0.29	0.21	0.00	0.51	0.41	0.41	0.00	0.82
62	0.63	0.12	0.00	0.75	0.29	0.19	0.00	0.48	0.30	0.20	0.00	0.50	0.37	0.52	0.00	0.89
64	0.58	0.13	0.00	0.72	0.18	0.20	0.00	0.38	0.17	0.14	0.00	0.31	0.22	0.36	0.00	0.58
66	0.17	0.17	0.00	0.34	0.04	0.31	0.00	0.35	0.12	0.17	0.00	0.30	0.14	0.30	0.00	0.44
68	0.08	0.10	0.00	0.18	0.05	0.19	0.00	0.25	0.03	0.16	0.00	0.19	0.03	0.20	0.00	0.23
70	0.01	0.12	0.00	0.13	0.00	0.22	0.00	0.22	0.03	0.19	0.00	0.22	0.01	0.12	0.00	0.13
72	0.02	0.02	0.00	0.04	0.00	0.17	0.00	0.17	0.00	0.31	0.00	0.31	0.01	0.13	0.00	0.14
74	0.00	0.08	0.00	0.08	0.00	0.14	0.00	0.14	0.00	0.28	0.00	0.28	0.00	0.08	0.00	0.08
76	0.00	0.07	0.00	0.07	0.00	0.08	0.00	0.08	0.00	0.11	0.00	0.11	0.00	0.03	0.00	0.03
78	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.10	0.00	0.10	0.00	0.02	0.00	0.02
80	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.04	0.00	0.04	0.00	0.00	0.00	0.00
82	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.02	0.00	0.02	0.00	0.00	0.00	0.00
84	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
86	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
88	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Total	2.95	1.59	0.00	4.53	1.58	2.46	0.00	4.04	1.84	2.73	0.00	4.57	2.58	3.47	0.00	6.05
Nº samples:				30				32				26				22
Nº Ind.:	269	152	0	421	157	234	0	391	172	275	0	447	214	301	0	515
Sampled catch:				526				554				624				612
Range:				41-85				41-89				37-87				36-78
Total catch:				526				554				624				612
Total valid hauls:				100				98				97				89

Table 23 (cont).- Black dogfish length distribution per haul mean catches by sex and year. Number per stratified mean catches. Spanish Summer Survey on NAFO 3L: 2008-2015 (R/V *Vizconde de Eza*). Indet. means indeterminate.


Length (cm.)	2012				2013				2014				2015				
	M	F	I	T	M	F	I	T	M	F	I	T	M	F	I	T	
16	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
18	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
20	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
22	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.01	
24	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
26	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
28	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
30	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
32	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
34	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
36	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	
38	0.01	0.00	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
40	0.00	0.05	0.00	0.05	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	0.01	0.02	0.00	0.03	
42	0.01	0.07	0.00	0.08	0.00	0.04	0.00	0.04	0.03	0.07	0.00	0.10	0.03	0.13	0.00	0.16	
44	0.06	0.07	0.00	0.13	0.00	0.06	0.00	0.06	0.04	0.04	0.00	0.07	0.05	0.13	0.00	0.19	
46	0.07	0.06	0.00	0.13	0.04	0.03	0.00	0.07	0.04	0.12	0.00	0.15	0.23	0.29	0.00	0.53	
48	0.01	0.03	0.00	0.04	0.01	0.04	0.00	0.05	0.05	0.11	0.00	0.17	0.23	0.32	0.00	0.55	
50	0.09	0.04	0.00	0.12	0.02	0.06	0.00	0.08	0.03	0.09	0.00	0.12	0.22	0.28	0.00	0.50	
52	0.07	0.08	0.00	0.16	0.01	0.06	0.00	0.07	0.15	0.07	0.00	0.22	0.21	0.31	0.00	0.52	
54	0.06	0.04	0.00	0.10	0.05	0.10	0.00	0.15	0.09	0.10	0.00	0.19	0.16	0.30	0.00	0.46	
56	0.13	0.08	0.00	0.21	0.04	0.12	0.00	0.16	0.08	0.24	0.00	0.32	0.13	0.32	0.00	0.46	
58	0.12	0.10	0.00	0.22	0.12	0.11	0.00	0.22	0.13	0.27	0.00	0.40	0.21	0.41	0.00	0.63	
60	0.20	0.14	0.00	0.33	0.26	0.21	0.00	0.47	0.21	0.21	0.00	0.43	0.31	0.37	0.00	0.68	
62	0.30	0.18	0.00	0.49	0.13	0.25	0.00	0.38	0.28	0.34	0.00	0.62	0.42	0.61	0.00	1.02	
64	0.27	0.07	0.00	0.34	0.15	0.37	0.00	0.52	0.16	0.26	0.00	0.42	0.31	0.57	0.00	0.88	
66	0.08	0.18	0.00	0.26	0.08	0.35	0.00	0.42	0.06	0.30	0.00	0.36	0.16	0.58	0.00	0.74	
68	0.04	0.12	0.00	0.16	0.00	0.36	0.00	0.36	0.05	0.25	0.00	0.29	0.09	0.63	0.00	0.72	
70	0.02	0.09	0.00	0.11	0.00	0.21	0.00	0.21	0.01	0.17	0.00	0.18	0.02	0.60	0.00	0.63	
72	0.00	0.12	0.00	0.12	0.00	0.16	0.00	0.16	0.00	0.15	0.00	0.15	0.00	0.38	0.00	0.38	
74	0.00	0.04	0.00	0.04	0.00	0.13	0.00	0.13	0.00	0.13	0.00	0.13	0.00	0.18	0.00	0.18	
76	0.00	0.01	0.00	0.01	0.00	0.05	0.00	0.05	0.00	0.07	0.00	0.07	0.00	0.05	0.00	0.05	
78	0.00	0.01	0.00	0.01	0.00	0.03	0.00	0.03	0.00	0.05	0.00	0.05	0.00	0.04	0.00	0.04	
80	0.00	0.01	0.00	0.01	0.00	0.02	0.00	0.02	0.00	0.02	0.00	0.02	0.00	0.03	0.00	0.03	
82	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.00	
84	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
86	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
88	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
Total	1.55	1.62	0.00	3.16	0.91	2.77	0.00	3.68	1.41	3.08	0.00	4.50	2.81	6.56	0.00	9.37	
Nº samples:					24				31				27				35
Nº Ind.:	150	137	0	287	85	264	0	349	125	282	0	407	260	594	0	854	
Sampled catch:					360				517				549				1124
Range:					39-80				39-81				37-82				22-81
Total catch:					360				517				549				1124
Total valid hauls:					98				100				99				97




**Fig 1.-** Atlantic cod stratified mean catches in Kg and  $\pm$ SD by year. Spanish surveys in NAFO Division 3L: 2003 - 2015 (R/V "Vizconde de Eza"). In 2003, the data correspond to 69% of the total area prospected in 2006-2015.



**Fig. 2.-** Atlantic cod abundance ('000), biomass in tonnes and  $\pm$ SD by year. Spanish surveys in NAFO Division 3L: 2003 - 2015 (R/V "Vizconde de Eza"). In 2003, the data correspond to 69% of the total area prospected in 2006-2015.



**Fig. 3.-** Distribution of the catches per haul for **Atlantic cod**, **Roughhead grenadier**, **redfish**, **thorny skate** and **black dogfish** in 2015 Spanish 3L survey.



**Fig. 4..- Atlantic cod length distribution (cm) in NAFO 3L: 2003-2015.** Number per stratified mean catches. In 2003, the data correspond to 69% of the total area prospected in 2006-2015.







Fig. 5.- Atlantic cod, roughhead grenadier and redfish length distribution (cm) in NAFO 3L: 2003-2015.




**Fig. 6.-** Roughhead grenadier stratified mean catches in Kg and  $\pm$ SD by year. Spanish surveys in NAFO Division 3L: 2003 - 2015 (R/V "Vizconde de Eza"). In 2003, the data correspond to 69% of the total area prospected in 2006-2015.




**Fig. 7. .-** Roughhead grenadier abundance ('000), biomass in tonnes and  $\pm$ SD by year. Spanish surveys in NAFO Division 3L: 2003 - 2015 (R/V "Vizconde de Eza"). In 2003, the data correspond to 69% of the total area prospected in 2006-2015.




**Fig. 8.- Roughhead grenadier** length distribution (cm) in NAFO 3L: 2003-2015. Number per stratified mean catches. In 2003, the data correspond to 69% of the total area prospected in 2006-2015.



**Fig. 9.- Redfish** stratified mean catches in Kg and  $\pm$ SD by year. Spanish surveys in NAFO Division 3L: 2003 - 2015 (R/V "Vizconde de Eza"). In 2003, the data correspond to 69% of the total area prospected in 2006-2015.




**Fig. 10..- Redfish** abundance ('000), biomass in tonnes and  $\pm$ SD by year. Spanish surveys in NAFO Division 3L: 2003 - 2015 (R/V "Vizconde de Eza"). In 2003, the data correspond to 69% of the total area prospected in 2006-2015.




**Fig. 11.- Redfish length distribution (cm) in NAFO 3L: 2003-2015.** Number per stratified mean catches. In 2003, the data correspond to 69% of the total area prospected in 2006-2015.

### Thorny skate stratified mean catches




**Fig. 12.** **Thorny skate** stratified mean catches in Kg and  $\pm$ SD by year. Spanish surveys in NAFO Division 3L: 2003 - 2015 (R/V "Vizconde de Eza"). In 2003, the data correspond to 69% of the total area prospected in 2006-2015.


### Thorny skate biomass and abundance




**Fig. 13.-** **Thorny skate** abundance ('000), biomass in tonnes and  $\pm$ SD by year. Spanish surveys in NAFO Division 3L: 2003 - 2015 (R/V "Vizconde de Eza"). In 2003, the data correspond to 69% of the total area prospected in 2006-2015




**Fig. 14.** Thorny skate length distribution (cm) in NAFO 3L: 2003-2015. Number per stratified mean catches. In 2003, the data correspond to 69% of the total area prospected in 2006-2015.




**Fig. 15. Thorny skate and black length distribution (cm) in NAFO 3L: 2003-2015.**



**Fig. 16.** Black dogfish stratified mean catches in Kg and  $\pm$ SD by year. Spanish surveys in NAFO Division 3L: 2003 - 2015 (R/V "Vizconde de Eza"). In 2003, the data correspond to 69% of the total area prospected in 2006-2015.



**Fig. 17.** Black dogfish abundance ('000), biomass in tonnes and  $\pm$ SD by year. Spanish surveys in NAFO Division 3L: 2003 - 2015 (R/V "Vizconde de Eza"). In 2003, the data correspond to 69% of the total area prospected in 2006-2015.



**Fig. 18.- Black dogfish** length distribution (cm) in NAFO 3L: 2003-2015. Number per stratified mean catches. In 2003, the data correspond to 69% of the total area prospected in 2006-2015.