

Northwest Atlantic



Fisheries Organization

Serial No. N6590

NAFO SCR Doc. 16/041

**NAFO/ICES *PANDALUS* ASSESSMENT GROUP—September 2016**

The West Greenland trawl survey for *Pandalus borealis*, 2016, with reference to earlier results.

by

AnnDorte Burmeister and Michael C.S. Kingsley

Greenland Institute of Natural Resources  
Box 570, 3900 Nuuk, Greenland

**Abstract**

A trawl survey is carried in NAFO Subarea 1 and NAFO Division 0A east of 60°30'W, as a contribution to the assessment of the stock of the Northern Shrimp (*Pandalus borealis*). In 2015, 252 stations were fished in 42 fishing days; 156 provided data to the shrimp survey in all strata.

The 2003 peak in total survey and fishable biomasses has been followed by continuous decline, reaching in 2014 the lowest levels since 1997. Total survey biomass increased by 58% over 2014, but were followed by a decrease of 25.2% over 2015. Offshore survey biomass were 28.3% less in 2016 than in 2015, about 62% of its previous maximum in 2010, above the past 5-year mean, but below the 20-year lower quantile. In Disko Bay and Vaigat the survey biomass is 17% less than in 2015, 74% of its past 5-year mean, and below the 20-year lower quantile. Offshore regions comprise 69% of the total survey biomass, and 31% is inshore in Disko Bay and Vaigat.

Surveyed regions showed decrease in survey biomass of about 20% in North (U1-U3) and 54% in the West (W1-W7) compared with 2015 results. Biomass in the southern part area, W8-W9, amounted only 1% of the total estimated survey biomass in 2016.

The Fishable proportion of the survey biomass is a little above the mean of the foregoing 20 years. While female biomass is close to its 20-year median and proportion of the fishable biomass is high and above the 20-year upper quantiles, in both regions, the biomass of fishable males is low, and the stock appearing deficient in these length classes.

That the spawning stock compose a high proportion of the total survey biomass makes the stock sensitive to fishing pressure and short-term recruitment to the spawning stock is likely to be low.

Offshore the index of age-2 shrimps is well below its 20-year mean when considered relative to survey biomass, but above its 20-year lower quantiles, while index of age-2 shrimps inshore is slightly above 20-year mean.

Since the late 1990s the stock is found in shallower water than before, this trend have continued since 2011, but in 2016 the survey biomass depth index increased at 21m from its 2015 value.

Area-weighted mean bottom temperature in the survey area increased at the beginning of the 1990s, by about 1.6°C, but despite a decrease from 2015 the relatively warm period continued in 2016.

## Introduction

Since 1988, the Greenland Institute of Natural Resources has carried out annual trawl surveys on the West Greenland shelf between June and August to assess the biomass and recruitment of the stock of *Pandalus borealis* and to obtain information on the size and sex composition of the stock as well as on the environmental conditions. This document presents the results of the 2014 survey, and compares them with revised series from previous surveys.

## Material and Methods

### *Survey design and area coverage*

The offshore survey area for the Northern shrimp, *Pandalus borealis*, covers waters on the West Greenland continental shelf from Kap Farvel in the south to latitude 72°30'N, comprising NAFO Sub area 1 and, where the shelf bulges into the Canadian EEZ, a small area in the eastern part of NAFO Div. 0A. In the late 1980s when the survey was initiated, *P. borealis* was fished in waters between about 150 m and 550 m deep, and the shrimp survey has always been restricted to depths between 150 m and 600 m. Since 1991 the survey has included the inshore areas of Disko Bay and Vaigat in NAFO Div. 1A but, along most of the coast, the survey does not cover areas closer to shore than 3 miles offshore of the fishery baseline. In some coastal areas fishable concentrations of *P. borealis* exist closer to shore than this, including areas that extend into some fjords (see e.g. Fig. 4a of Hammeken and Kingsley 2010).

The survey area is divided into primary and secondary strata. The survey primary strata correspond to geographical areas identified on the basis of logbook information on the distribution of the fishery (Carlsson *et al.* 2000). They are subdivided into four secondary (depth) strata at 150–200 m, 200–300 m, 300–400 m, and 400–600 m. When the survey was initiated, bathymetric information in Disko Bay, as well as offshore north of 69°30'N, did not support this depth stratification, and these regions were therefore originally subdivided into geographical substrata not based on depth. Depth data logged by the survey and other investigations eventually allowed these waters to be stratified on depth and a new geographical stratification with depth sub-strata was introduced in 2004 (Wieland and Kannevorff, 2004). At the same time, the geographical strata in the Canadian zone, formerly two, were merged into one.

From 1988 through 1999, trawl stations were allocated to strata in proportion to stratum area, but since 2000 more stations have been allocated to strata where biomass variances have been high in previous years in order to improve the precision of the overall biomass estimate (Kingsley *et al.*, 1999). An exponential smoothing of previous years' stratum variances was applied in the allocation procedure, giving higher influence to the more recent years. Past variance data for Atlantic cod and Greenland halibut is also made available to the allocation procedure, which is now set to minimize a weighted combination of the expected survey precision for the three species.

Generally the station layout is based on a division of the survey area into elements about 2 nautical miles square. Since 1999 survey stations have been positioned using 'buffered random' sampling, in which stations are placed randomly with the constraint that a minimum distance between them, which depends on station density within the stratum, must be observed (Kingsley *et al.*, 2004).

From 1988 through 1998, survey designs were independent from year to year, stations being placed anew in the strata. Since 1999 about 50% of the stations included in the preceding year's design have been repeated as fixed stations in the following year, the others being placed, as before, using the buffered sampling rules.

Catches are correlated from year to year by position, and fixing stations improves the ability of the survey to indicate year-to-year changes in stock size by inducing serial correlation in survey error (Kingsley, 2001a).

In 2016, 256 stations were planned at depths between 150 and 600 m in the survey area, with 40 'extra' stations mapped and available to be included if time permitted. 252 of the 256 planned stations incl. 14 'extra' stations were fished. 8 stations were reported as having been moved more than 2 n.mi. from the planned position, with a mean of 9.2 n.mi.; 31 stations were discarded either of untrawlable bottom owing to general trawl difficulties.

Of the 252 stations fished, 156 provided usable data to the shrimp survey. In the course of the shrimp survey, 39 CTD casts were made along standard transects offshore and in Disko Bay and Vaigat; 13 stations were fished to cover deep water Greenland halibut and at 92 extra station were fished with an beam trawl for benthos monitoring.

#### *Survey period and daily sampling period*

The trawl survey has been carried out every year between mid-June and the end of August to minimize the effect of seasonal cycles in the biology of the species. Trawling is carried out between 0800 and 2000 UTC; it appears that the daily vertical migration of the Northern shrimp is quite abrupt at sunrise and sunset and a shorter trawling day is not necessary.

#### *Tow duration*

From 1988 to 1997 all tows in the shrimp survey lasted 60 min. However, shorter tows give just as accurate results (Carlsson *et al.*, 2000; Kingsley, 2001b; Kingsley *et al.*, 2002; Wieland and Storr-Paulsen, 2006; Ziemer and Siegstad, 2009) and since 2005 the survey has been operated with 15-minute tows alone.


#### *Fishing practices*

The surveys have been conducted with the research trawler *Paamiut* (722 GRT) since 1991; similar vessels were used in 1988–1990. Initially, a 3000/20-mesh *Skjervøy* bottom trawl with a twin cod-end, and equipped with a heavy steel-bobbin footrope, was used. A 20-mm (stretched-mesh) liner was added to the 44-mm-mesh cod-end in 1993. From 1988 to 1991 estimates of trawl geometry—door spread and height of head-rope—were based on results from tank experiments at the Danish Institute for Fisheries Technology and Aquaculture. Since 1991 *Scanmar* acoustic sensors have been mounted on the trawl doors, and a *Furuno* trawl-eye on the head rope. Door-spread readings are recorded during the tow, and the reading from the trawl-eye is used to judge when the trawl has settled and the tow can be deemed started.

From 1988 through 2003 *Greenland Perfect* trawl doors were used, measuring 9.25 m<sup>2</sup> and weighing 2420 kg. They were replaced in 2004 by *Injector International* 7.5 m<sup>2</sup> trawl doors weighing 2800 kg to facilitate a change of survey trawl in 2005. In 2005 the *Skjervøy* 3000 trawl was replaced by a *Cosmos* 2000 trawl with 'rock-hopper' ground gear comprising steel bobbins and rubber disks. Towing speeds have been about 2.5 knots in all surveys.

#### *Swept area calculation*

Nominal swept area for each tow was calculated as the straight-line distance between its GPS start and end positions multiplied by the wingspread. The distance between the trawl doors should be recorded 3 or 5 times during each tow; provided it was recorded at least once, wingspread for a tow was calculated from the mean door spread and the geometry of the trawl. For both trawls the wingspread (i.e. the width of the swept area)  $V$  has been calculated as follows. The trawl and the trawl plus bridles are assumed to form two similar triangles, bridles and wings making a straight line:



and the lengths of the bridles ( $s$ ) and the trawl wings ( $t$ ) are known. The wingspread  $V$  is then calculated as:

$$V = (t_t * L) / (t_t + s_t)$$

where  $L$  is the distance between the doors (doorspread).

The length of the *Skjervøy* trawl is 67.15 m and the length of the *Cosmos* trawl is 71.8 m, both measures excluding the cod-ends. Since 2004 the bridle length, i.e. the total length of lines, chains and shackles between the trawl doors and the tip of the trawl wing, has been 54 m for either trawl; other bridle lengths were used in earlier years (Table 1). In the case of the *Skjervøy* trawl, 0.7 m has been added to the calculated wingspread because the *Skjervøy* trawl is a three-winged trawl and the lower wings (directly attached to the ground-rope) were estimated to spread 0.35 meters wider than the middle wings on each side in tank experiments at the Danish Institute for Fisheries Technology and Aquaculture (Per Kanneworff, pers. com.).

If doorspread was not recorded in the course of a tow, the unweighted mean of the calculated wingspreads for the year was used, without regard to fishing depth or wire length. If the speed of the tow calculated from its duration and its start and end positions lay outside a range of 3.0 to 6.5 km/hr, its length was recalculated from the survey average towing speed and the tow duration.

#### *Biomass estimation*

Each haul's catch was divided by its estimated swept area calculated from wingspread and track length to estimate a biomass density. Unweighted mean stratum densities were multiplied by the stratum area to compute stratum biomass, and a corresponding error variance for the stratum biomass estimate was also calculated for strata with two or more accepted hauls. For strata with only one accepted haul, an error coefficient of variation of 0.95 was assigned. Stratum biomasses and their error variances were summed to get regional and overall estimates.

#### *Indices of distribution and location of shrimp biomass*

Indices to summarize how widely the survey biomass is distributed and a measure of its central latitude were calculated (Kingsley 2008). Data from surveys executed in 1994–2012 was used: there was no survey before 1994 in the southernmost areas and before 1991 in Disko Bay or Vaigat, but since 1994 the series has been consistent. Biomass estimates from the annual survey are customarily presented (e.g Ziemer 2008) for 7 divisions of the survey area:

- a northern division, formerly stratified as N1–N9, and re-stratified according to depth information (Wieland and Kanneworff 2004) as U1–U3 with depth strata;
- Disko Bay and Vaigat, formerly stratified as D1–D9, restratified as I1 and I2 with depth strata;
- Canadian Exclusive Economic Zone, once 2 divisions, now 1;

- subdivisions of the west coast, from the mouth of Disko Bay and adjacent shelf waters to Paamiut;
- an extreme southerly division, comprising Julianehåb Bay and adjacent waters.

These divisions were given southerly ranks: Julianehåb Bay and the adjacent shelf waters were 1; the subdivisions up the coast were given ranks 2, 3 and 4. The small Canadian sub-division was split, half being included with the subdivisions W1 and W2, and the other half included with survey subdivisions W3 and W4. Disko Bay and Vaigat were given rank 5, and the northernmost subdivision of the survey area was ranked 6.

Then a 'lat. index' was calculated as a mean rank for the survey, weighting by estimated total survey biomass. This index summarises how far north a (weighted) centre of gravity of the stock biomass lies.

A 'spread index' was calculated as a Simpson diversity index (Simpson 1949) of the distribution of the biomass:

$$SpreadIndex = \left( \sum_{Subdivisions} Biomass_{Subdiv.} \right)^2 / \sum_{Subdivisions} (Biomass_{Subdiv.})^2$$

This index summarizes how widely the survey stock biomass is distributed among survey subdivisions.

#### *Depth distribution of biomass.*

The overall depth distribution of the estimated survey biomass was calculated according to available depth information. Up to 2003, such information was only available for the west-coast area W and the Canadian EEZ, and the depth distribution of the biomass was analyzed only for those areas. Since 2004, the northern area and the inshore areas in Disko Bay and Vaigat have been sub-stratified depth and the depth analysis extended to those areas.

A single depth index, in meters, was calculated by assuming that the entire survey biomass in any depth stratum was concentrated at the stratum's midpoint depth, except for the deepest stratum: the fishery takes little from water deeper than 520 m, so the central depth for the 400–600 m stratum was set at 460 m.

#### *Sampling, weighting and "area expansion"*

The composition of the stock by size and sex is based on a two-stage analysis of lengths and weights. From catch samples, 1000–2000 individual shrimps are both weighed and measured, and these measurement pairs are used to estimate a weight-length relationship. From each catch a sample of about 0.5 to 3 kg was taken and sorted to species. All specimens of Northern shrimp were classified: juveniles and males composed one class, primiparous and multiparous females two others (Allen 1959, McCrary 1971). The oblique carapace length (CL) of each shrimp in the sample was measured to the nearest 0.1 mm. These length measurements are then supplemented with weight estimates based on the fitted weight-length curve. Aggregated, and then averaged over the stations in a stratum, these observations of sex and measurements of length and their associated estimates of weight are used to estimate the distribution of the stratum biomass between sex and length classes as well as the numbers of shrimps in the stratum in the various sex and length classes.

The descriptions of calculation methods that follow consider only one stratum. Survey strata are considered here as independent sampling problems. Survey totals are got by summing stratum results. 'Length class' can be generalized to include sex or sex-length class. From the catch and swept area at a station, the single-station estimate of stratum biomass is

$$\hat{B}_{ts} = \frac{Strat.Area_t \cdot Catch_{ts}}{Sw.Area_{ts}}$$

These single-station estimates are averaged. An unweighted analysis gives

$$\hat{B}_t = \frac{Strat.Area_t}{K_t} \sum_s \frac{Catch_{ts}}{Sw.Area_{ts}}$$

where  $t$  is the stratum and  $s$  is the station, of which there are  $K_t$  in stratum  $t$ . Shrimp density does not vary much within a haul's distance (Kingsley et al. 2002) and so it is statistically preferable to use the unweighted mean estimate of stratum biomass. This is what is done in the West Greenland survey.

If the number of shrimps in class  $l$  in the Length Sample, of weight  $Samp.Wt_s$  from station  $s$  in stratum  $t$  is  $n_{lts}$ , the corresponding single-station estimate of the number of shrimps in the class in the stratum is

$$\hat{N}_{lts} = Strat.Area_t \frac{n_{lts}}{Samp.Wt_{ts}} \cdot \frac{Catch_{ts}}{Sw.Area_{ts}}$$

The stratum estimate from many stations is:

$$\hat{N}_{lt} = \frac{Strat.Area_t}{K_t} \sum_s \left( \frac{n_{lts}}{Samp.Wt_{ts}} \cdot \frac{Catch_{ts}}{Sw.Area_{ts}} \right)$$

where the divisor  $K_t$ , the number of stations, includes those with no catch. The coefficient

$$\frac{Catch_{ts}}{Samp.Wt_{ts} \cdot Sw.Area_{ts}}$$

common to all length (and sex) classes counted in the Length Sample for a station can be called its 'raising factor'. Given these estimates of numbers, and if estimates  $w(l)$  of individual weight at length are available, length-class biomass is given by

$$\hat{W}_{lt} = w(l) \frac{Strat.Area_t}{K_t} \sum_s \left( \frac{n_{lts}}{Samp.Wt_{ts}} \cdot \frac{Catch_{ts}}{Sw.Area_{ts}} \right)$$

Provided that for all Length Samples  $\sum_l n_{lts} w(l) = Samp.Wt_{ts}$ , i.e. the weight of every Length Sample

answers exactly to the number and size of the shrimps that compose it, the sum of length-class biomasses equals the stratum total biomass calculated from catches and swept areas. This is ensured by using, as the weight of the Length Sample, the total weight of the shrimps it comprises, calculated from the weight-length function, instead of using its weighed weight. In effect, the length-measurement data is used only to partition the stock biomass between sex and length classes.

Where catches were, for one reason or another, not sampled for lengths, the mean of the length frequency distributions from the available samples in that stratum, each raised by its catch and swept area, was applied to the entire stratum biomass. If it occurred that there were no length samples from the catches in an entire stratum, the length distribution estimated for the entire survey from data for the strata that did have length samples would be applied to its estimated biomass.

Results from these calculations were subsequently used to construct area-specific length frequency distributions (LFD). LFD results were used to calculate indices of numbers by sex and length, biomass by sex,

an index of fishable biomass (comprising shrimps at least 17 mm CPL), and numbers of small pre-recruits ('age 2 shrimps'), which are expected to enter the fishery in coming years.

For data since 2007 the following weight-length relationship has been fitted:

$$w(l) = w_0 + a \cdot (l - l_0)^z$$

with the parameter  $l_0$  constrained to be less than 4.5 mm. A weighted fit was used with constant coefficient of variation about the fitted line.

As there is no reliable method of aging even young shrimps, indices of numbers at age are obtained by modal analysis of length frequencies for juveniles and males, attempting to fit overlapping age-specific normal distributions to the aggregated length distributions.

Modal analysis is carried out using Partiel Mix (Kingsley, 2014, MCSKMIX 2014 © Greenland Institute of Natural Resources), implemented in an Excel spread sheet. Partiel MIX fits numbers of shrimps at age 2 years and is estimated by decomposing the length distributions, fitting Normally distributed components.. To prevent a skewness to the plus age 3 group, analysis was only done on shrimps below or equal to 19 mm CPL. While using this new method in 2014, all age-2 results from 2005 to 2014 was re-calculated. A more comprehensive description of the method is found in Kingsley (2014).

As growth is probably affected by temperature, the survey area is divided into 6 regions, defined from gradients in bottom temperature (Wieland, 2004), and estimated numbers in strata are pooled over regions. Length-frequency histograms are not smoothed before being analyzed. The analyses of sex and length distribution in the stock, and the modal analyses, were re-done in 2014 for data from 2005–2014. To simplify the modal analysis, and because the main objective is to estimate the numbers at age 2, only the numbers for the 'Juveniles and Males' sex class were put through it. For this re-analysis, strata were grouped: U1 to U3, I1 and I2, W1 to W4, W5 and W6, and W7 to W9. Within each group, the estimated stock numbers in each stratum were aggregated over depth substrata, and the stratum aggregate numbers presented to Partial MIX as a haul.

### *Bottom temperature*

Bottom temperature was measured with a *Starmon* sensor mounted on one of the trawl doors. It records at intervals of 30 s with a resolution of 0.01°C. The average temperature for each haul was calculated after retrieval of the sensor. All measurements taken at greater depths than 150 m were used to calculate a mean bottom temperature weighted for the areas of the survey strata between 150 and 600 m depth.

## **Results and Discussion**

### *Survey conduct and progress.*

The survey in 2016 started from Nuuk on 3 June. The first cruise occupied stations from Nuuk north to Disko Bay having occupied 83 shrimp and fish stations and 17 CTD cast in 14 days. The second cruise occupied 76 shrimp and fish stations and 22 CTD cast in 22 days in Vaigat, north to the survey limit, and west of Disko on its way back south, the northern area being less densely sampled. The third cruise started from Nuuk on 09 July to occupy 93 stations south to the southern limit of the survey at 13 days until July the 21. No CTD cast were sampled at the third trip.

All cruise was accompanied by a crew engaged in taking photographs of the sea bottom, who used the ship at night when the survey was suspended. From all planned shrimp and fish stations, at the three cruises, by-catch of benthic faunal invertebrates were sorted from each stations at all, identified to the lowest possible taxonomic level, and some species were preserved in 10% buffered formalin solution for later identification,

and both projects aims to investigate the long-term effects of trawling on bottom ecosystems. Same procedure were done for benthic species sampled with a beam-trawl. 27 beam-trawl stations were done at the first cruise, 42 beam-trawl stations at the second cruise and 23 beam-trawl stations at the third cruise.

Of the 58 planned survey strata, 1 (the Canadian zone C0) had no stations trawled caused by the ice cover; owing to general trawl difficulties Substrata (W2-4) were not fished and only one stations were fished in each of substrata W6-4 and W7-4. 33 stations had no catch of *P. borealis*; 6 strata, with 12 occupied stations between them, had no catches and only strata U1 – U3, I1, I2, W2 and W4 had catches of *P. borealis* at all stations. Length samples were measured from 125 stations with catches even very small ones, and length & weight samples were measured at 21 stations.

There were no exceptionally large catches and only at one station in I2-1 catches was closed to 1 ton (977 kg). There were 14 catches over 200 kg, of which 7 were made in Disko Bay or Vaigat, 1 stations north from 69°30 N, 4 in strata W1-W4 and 1 in W6 and 1 in W8-4. In southerly areas, strata W7-W9 yielded an average catch of 9 kg, and 242 of 34 stations had no catch of *P. borealis*. A catch of 305 kg were observed in substrata W8-4, whereas the average catch were 0.01 kg at the other 33 stations in the southern regions.

#### *Overall Biomass and Area Distribution.*

For all strata biomass estimates have been calculated (Table 2) on the basis of the nominal swept area. The biomass estimates (in tons) for the five main regions and the entire survey area in 2014 were:

Region-2014	Biomass estimate (t)	Number of stations	ECV (%)
North (U1-U3)	29 381	32	26.8
Canadian zone (C0)*	-	0	-
West (W1-W9)	57 015	131	17.2
Disko Bay & Vaigat (I1, I2)	93 461	26	11.6
Total	179 857	189	9.2

- Area C0 was not surveyed in 2014, owing to time trouble.

and in 2015:

Region-2016	Biomass estimate (t)	Number of stations	ECV (%)
North (U1-U3)	75 394	32	26.8
Canadian zone (C0)*	-	0	-
West (W1-W9)	130 004	132	29.1
Disko Bay & Vaigat (I1, I2)	78 215	22	10.7
Total	212 031	156	14.7

- Area C0 was not surveyed in 2015, owing to ice cover

and in 2016:

Region-2016	Biomass estimate (t)	Number of stations	ECV (%)
North (U1-U3)	60 548	26	12.0
Canadian zone (C0)*	-	0	-
West (W1-W9)	86 689	112	15.1
Disko Bay & Vaigat (I1, I2)	67 795	18	18.4
Total	212 031	156	9.0

- Area C0 was not surveyed in 2016, owing to ice cover and W1-4 due to bottom trouble.



Survey biomass decreased over 2010 and the declining trend in survey biomass continued until 2014, where biomass estimate for Disko Bay and Vaigat was little changed from 2011, but were further large decline in all offshore regions. In 2015, perhaps caused by an over optimistic survey (Burmeister and Kingsley, 2015a; Burmeister and Kingsley, 2015; NIPAG, 2015), the total survey biomass increased over 2014 and was 57.6% higher in 2015 (Table 2, Figure 2a and 2b). All offshore surveyed regions showed increases in survey biomass compared with the 2014 results, especially in North (U1-U3) recorded its biggest survey biomass since 2009, but an increase were also observed in the total West regions. Surveyed biomass declined about 16% in Disko Bay and Vaigat, but remained above their 1988–2015 mean by about 15%. The Canadian area was not surveyed, but usually contributes only a few percent of the survey biomass.

The 2016 total survey biomass (212.03 Kt) had been decreasing and is 25.2% less than in 2015, is 92% of its past 5 year mean (2011–2015), but below its serial mean of 292.5 Kt. All the surveyed region showed decreases in survey biomass compared with the 2015 result. Survey biomass in Disko Bay and Vaigat was 17% less and 74% (87.5 Kt) of its past 5 year mean (2011–2015). However, due to general trawl difficulties, the southern part below 69°N in Disko Bay was not surveyed in 2016. In average, over the past five year, 42% of the biomass has been located north of 69°N in stratum I1, but since 2013 the majority of the total biomass in stratum I1 has been found south of 69°N. It is therefore reasonably to assume, that 2016 biomass in Disco Bay might be underestimated due to the lack of sampling in the southern part of the Bay.

The Canadian area was not surveyed in 2016, but usually contributes only a few percent of the biomass. In all offshore regions, except the most southern strata W8-W9, survey biomass (147.2 Kt) decreased 28.3% over 2015 (Table 2, Figure 2a), remain close to its past five year mean (142 Kt), but was only 65.5% of its serial mean (1988–2016). There were large decreases in biomass in W3-W4, of 57% and in W5-W7 of 43%, whereas the decline amounted to 6.2% in more northern regions W1-W2 and 19.8% further north in U1-U3 (Table 3).

Survey biomass in W8-W9, increased by 37% over 2015, nevertheless, biomass estimates for the southernmost area only amounted only 1 % of the total estimated survey biomass in 2016.

Densities of *P. borealis* in Disko Bay have been almost stable from 2010 to 2014, followed by a continuous decline since 2015, and is in 2016 both below its past 5 years mean and the 1988 – 2015 mean. However, the inshore region, had far higher densities (6.50 t/km<sup>2</sup>) than other areas (Fig 3a), over three times as high as the second highest, strata W1 and W2 combined (1.98 t/km<sup>2</sup>). Lowest densities were found in the Southern areas W5-W7 (0.48 t/km<sup>2</sup>) and W8-W9 (0.49t/km<sup>2</sup>) (Table 5). Over all mean density decrease by 21.4% over 2015, and was estimated to 1.69 t/km<sup>2</sup> in 2016, below the 1988–2015 means (2.30 t/km<sup>2</sup>), but equal to its 2011–2015 mean (Fig. 3b).

The spread index, of how widely the survey thinks the stock biomass is distributed, showed an increase in 2016 indicating that the stock is a bit more widely distributed than compared to the past 5 years. However, the north index remained close to its maximum value (Fig. 4).

#### *Depth distribution of the shrimp biomass.*

During the early and mid-1990s the depth distribution of the survey biomass was stable, with a significant contribution from the 400–600 m stratum, but most of the biomass found in 300–400 m of water (Fig. 5a). From 1990 to 1998 the average distribution was 1.9% in 150–200 m of water, 25.2% in 200–300m, 55.9% in 300–400 m, and 17.1% deeper than 400 m; the survey biomass depth index was stable near 350 m (Fig 5.b). In the late 1990s this situation started to change. The proportion in water 300–600 m deep decreased and that in the 200–300-m stratum increased greatly. In the late 1990s, the stock biomass started the sustained increase that peaked in 2003.

Somewhat later, starting in the early 2000s, the shallowest stratum, which is only 50 m from shallowest to deepest, also started to contribute a little more to the biomass than it had done.

The 200–300 m stratum started showing biomass increases as early as the late-1990s, and at that time the deepest stratum also started its decline—hence the start of the change in the mean-depth index (Fig. 5.c.). The 150–200 and 300–400 m strata increased later, and less. After 2004 and until 2012, the biomass has declined in all strata, but proportionally by more in the deepest two strata.

Compared with this shift in the depth distribution of the survey biomass, a shift in the depth distribution of the commercial catches appears to have started earlier. In 1991–1994 the median catch depth was 347 m but in 1995–98 it had already decreased by about 30 m, and by 2003–2006 it was 100 m less at 247 m (Kingsley 2011). This change has stayed in place in the case of the inshore fleet, which in 2007–2010 was still taking its median catch at 255 m, but the offshore catches show slight signs of moving back to deeper water with a median catch depth of 277 m in 2007–2010. The fraction of catches taken in water shallower than the lower limit of the survey, i.e. 150 m, is however still only a few percent, so this is probably not the reason for the recent decrease in survey biomass; besides, the great increase in survey biomass between about 1997 and 2003 coincided with the first years of this shift of the biomass, and the fishery, into shallower water.

In 2012 16.5% of the survey biomass was estimated to be in 150–200 m of water, 57.5% in 200–300 m, 22.3% in 300–400 m and 3.6% deeper than 400 m. In fact, the distribution was reversed: 17% deeper than 400 m in the early years was replaced by 17% shallower than 150 m now, and 55% in 300–400 m of water by 55% in 200–300 m.

Depth distribution of survey biomass changed in 2016, while only 3% of the total surveyed biomass was observed in the shallow water (150 – 200m). The 2016 value is way below the 2011–2015 mean for this depth stratum (17%), and change was caused by a significant decrease of biomass in shallow water in Disko Bay & Vaigat and in region W1, W3, W4 and W5. In fact - there was no shrimp at the shallow water at W3. At the same time proportion of biomass in 200 – 300 m was 10% higher in 2016 (61%) compared to the past 5 years. Only a slightly increase in biomass was observed in 300-400m (31.%) and in the deep water 400-600 (Fig 5a and Fig 5b). The depth distribution of the biomass in region U1-U3 were comparable with the previous years.

In 2001–2016 the survey biomass depth index has ranged between 265 and 300 m, with an average near 281 m compared to an average of 329 m in the period from 1988 to 2000 (Fig 5.c). In 2016 the depth index was estimated to approx. 288 m, a decrease of 21 m compared with 267 m in 2015.

#### *Bottom temperature and biomass*

The overall mean bottom temperature in the shrimp survey area was stable near 0.9°C in the early 1990s. Between the mid- and the late 1990s it underwent a step increase and since the late 1990s it has been stable in the neighborhood of 1.6°C. The increase has affected all depth strata and all areas. The depth distribution of the shrimp survey biomass appears to have shifted after, and not concurrently with, the temperature shift (Figure 6 and figure 14). The latitude distribution of the stock has also shifted, but also *after* the temperature shift: it is *since* the late 1990s that the survey 'North index' has continually increased, and the mean catch latitude in the fishery reached its decided minimum, at 64°41'N, in 1998.

In summary: bottom temperature increased in a stepwise fashion in the mid- to late 90s, and after that the survey biomass moved north and into shallower water, and the fishery catches also moved north; but fishery catches started moving into shallower water when the temperature started to increase.

### **Demography and recruitment**

#### *Length-weight relationships*

In 2016, only 1125 shrimps were individually weighed and measured (Table below). A single weight-length relationship was fitted to all sex classes. The same relationship was retrospectively fitted to data from 2005–2010 to provide a consistent basis for estimating numbers of pre-recruits for those years; we have compared

estimated weight at length from the weight-length relationships used in different years. Parameter values estimated in 2016 were little changed from the past years.

Table: *Pandalus borealis* in West Greenland: parameters of a relationship  $w(l) = w_0 + a \cdot (l - l_0)^z$  fitted to weight-length data for individual shrimps.

Year	$w_0$ (g)	coefficient $a$ (mg)	$l_0$ (mm)	exponent $z$	sample	Scatter c.v. (%)
1988–2000	0	0.669	0	2.96		
2001–02	0	0.483	0	3.0576		
2003	0	0.752	0	2.9177		
2004	0	0.765	0	2.9092		
2005	0.03103	1.726	1.91	2.7188	1616	6.79
2006	0.05771	1.426	1.591	2.761204	1907	7.89
2007	0.7700	1.789	4.5*	2.78216	487	6.42
2008	-0.03285	1.4162	0.797398	2.7501	2147	6.67
2007–08	-0.121034	0.4031	-1.66043	3.052731	2634	6.88
2009	0.01823	2.774	2.19026	2.58902	1768	6.86
2010	0.1533	8.155	4.5*	2.32036	1096	6.66
2011	0.12993	9.753	4.5*	2.25051	2569	6.64
2012	0.08185	8.928	4.229	2.27317	2300	7.31
2013	0.12644	9.541	4.5*	2.26021	2353	6.66
2014	0.10582	9.554	4.5*	2.27123	2371	6.75
2015	0.07269	4.165	3.1	2.49521	1088	6.79
2016	0.18877	8.655	4.5*	2.30198	1125	5.68

\*  $L_0$  must be no greater than 4.5 mm.

The scatter about the fitted weight-length relationship is consistent from year to year, except that the 2012 data, when the sample is relatively large, and has a slightly larger scatter, than the other years. In 2016 also the scatter was lower than most other years.

Table: *Pandalus borealis* in West Greenland: weights at length predicted from fitted weight-length relationships.

Year	Weight (g) at length (mm):			
	10	15	20	25
1988–2000	0.61	2.03	4.75	9.19
2001–2002	0.55	1.91	4.59	9.08
2003	0.62	2.03	4.70	9.02
2004	0.62	2.02	4.66	8.92
2005	0.54	1.91	4.56	8.82
2006	0.57	1.91	4.50	8.67
2007	0.98	2.01	4.44	8.75
2008	0.60	2.06	4.76	9.02
2007–2008	0.61	2.04	4.70	8.96
2009	0.59	2.06	4.82	9.12
2010	0.58	2.06	4.87	9.17
2011	0.58	2.07	4.79	8.86

Year	Weight (g) at length (mm):			
	10	15	20	25
2012	0.56	2.06	4.80	8.90
2013	0.58	2.07	4.80	8.92
2014	0.56	2.10	4.93	9.21
2015	0.59	2.08	4.90	9.28
2016	0.63	2.13	4.95	9.24

Estimated weights at length have been consistent over years, and there is very little change from 2010–2015. Though 2016 has rather heavy shrimps at the small medium size, 10, 15 and 20 mm.

The mean weight of both sexes has decreased steadily over the 27-year history of the survey (Fig. 8), by, on average, 54 mg/yr. for females and 23 mg/yr. for males. We do not know the reason for this; it might be due to a progressive reduction in the size at sex change, which would affect the mean size of both sexes. There is serial correlation in the residuals about the trend lines, 0.56\* for males and 0.84\*\* for females, indicating that there is some underlying mechanism.

#### *Recruitment and mean length at age 2*

Length-frequency plots—e.g. regions I1 – I2 and W1–W4 in 2014 (Fig. 9)—show a first insignificant component with mean CPL at 7–8 mm. These shrimps had probably hatched in the spring of the year before. Catches of this first component are small owing to the mesh size used in the survey; even the second component is not fully retained by the survey gear (Wieland, 2002b).

Regional differences in length at age 2 are obvious: they tend to be larger in more southerly areas and in deeper water. Year-to-year changes in the mean length at age 2 have been related to shifts in bottom temperature and changes in stock density of shrimp (Wieland, 2005). For the most recent six years, however, estimates of the mean size at age 2 were fairly consistent (Table 9).

In 1993–1995 estimates of numbers at age 2 were low, particularly in Disko Bay and Vaigat (Table 10, Fig. 11); in 1996 they were exceptionally high, especially offshore, but dropped again in 1997. After 1997, age-2 numbers increased steadily to peak in 2001, followed by a steady decline to 2005. A series minimum observed in 1995 was reached again in 2007 and 2012.

Since 2013 numbers of age-2 shrimps increased and more than double to 10.85 bn in 2015, considerably above the series 1988–2015 mean of 6.27 bn (Fig. 11 and table 10). The high level of age-2 shrimps was not maintained and numbers of age-2 shrimp was about 61% less in 2016 (4.26 bn), but comparable to 2013 and 2014 values and close to its past 5 years mean (2011–2015) at 4.85 bn. The decline in numbers of 2-age shrimp is caused by a significant drop of age-2 shrimps in area W1W4 of 81%, 52% in North (U1–U3) and 31% in Disko Bay & Vaigat, while there were almost no age-2 shrimps found in W5–W9.

Disko Bay and Vaigat, which include only about 7% of the survey area, contributed 28–72% of the numbers at age 2 in 1997–2005, and more than 40% in the period from 2006–2010 (Table 10). Whereas the contribution from Disko Bay and Vaigat was about 50 and 68% in 2013 and 2014 respectively, it was only 33% in 2015 but 58.9% in 2016 (Table 10, Fig 11a). However, 2016 number of age-2 shrimps was above the 1988–2016 mean and 99% of the mean for the past 5 years.

In the north most region U1– U3 numbers of age 2 shrimps decreased significantly over 2015, but is still above the serial mean and far above the mean or the 5 past year (Table 10).

Age-2 shrimps have always been few in the southernmost region W7 to W9, and since 2004 they have also been very few in W5 and W6. In the most recent years contributions of age-2 shrimps have been low in

region W1-W4/C0 (8 – 12%), but were its constitute about 50% of the total numbers in 2015 is was only 24% in 2016 (Table 10).

The age-2 index is correlated with the fishable biomass two, three and four years later (Fig. 12); lags of two to four years in such a correlation might be reasonable considering that the main contribution to the fishable biomass comes from sizes corresponding to ages 4 to 6 years. The recent low numbers of small shrimps presage a decrease of the fishable biomass in the coming years to below the average of the late 1990s. However, these estimates are uncertain: the series both of age-2 numbers and of fishable biomass have strong serial correlation, which invalidates the nominal levels of statistical significance of the simple correlations.

#### *Numbers, spawning stock biomass and fishable biomass*

Given that the survey biomass is down to a low level compared to 2002-2006, survey biomass and fishable biomass increased considerable over 2014 to a 2015 level comparable with 2009 and 2010 (Table 2 and table 8). The progress in fishable biomass were not maintained and was 21.2% less in 2016 (195.6 Kt). Spawning stock biomass—i.e. of females—accounted for more than 40% of the total survey biomass from 2010 to 2014, reached its highest level at 47.8% in 2014, but dropped over the past year to 35% in 2015 (Table 7). In 2016 females biomass constitute 46.4% of the total biomass, despite the biomass of females, estimated at 98.4 Kt in 2016, was almost unchanged from 2015 and below the 1988 – 2015 mean (106.6 Kt) (Table 6). The increase of female proportion of the total survey biomass was caused by a decrease of male biomass.

Compared with the length distributions in past twelve years, there are a greater proportion of the very smallest shrimps, those below about 15 mm CPL caught by 2015 survey in Disko Bay and Vaigat as well as in the offshore regions (Fig10). The proportion of those small shrimps was in 2016 comparable with observations from 2013 and 201. The proportion of the largest females above 23 mm CPL were comparable with the 2012 - 2015 level in Disko Bay and Vaigat, and with the 2015 level in the offshore regions (Fig. 7 and Fig. 10). Female biomass in offshore regions were low in 2014, increased significantly in 2015 remain on a comparable level in 2016, but is still below the 20-yr mean, while female biomass in Disko Bay declined over 2014, remain stable in 2016 and is below its 20-yr mean.

Number of pre-recruits (14-16.5 mm CL) were considerably lower 57.9% offshore, but close to the 2014 value, while a 37% decrease was observed in Disko Bay. A fraction of the high proportion of pre-recruits observed in 2015 were expected to enter the fishery in 2016, but both inshore and offshore regions reveal no increase in male biomass (shrimps from 17- 23 mm). Nor the high proportions of fishable males observed in 2015, resulted in increased female biomass in 2016. In both regions length distribution is relatively deficient in the intermediate lengths. In spite of this, the fishable biomass is estimated at 92% of the total survey biomass, a proportion that is equal to the average, probably because of the high proportion of females. The high proportion of large shrimps is expected to make the stock in its present state sensitive to fishing pressure, as the fishery prefers, in general, to catch the largest shrimps; while a deficiency in intermediate sizes bodes ill for short-term recruitment to the fishable and to the spawning stock.

#### *Bottom temperature*

Area weighted bottom temperatures is given in Fig. 6. Bottom temperature have been stable at somewhat 3.1°C from 2008 to 2012, but starting a decrease in 2015 (2.98°C) which continuing in 2016 to 2.66 °C.

Regionally the temperature in the Northern regions (U1-U3 and I1-I2) at 2.6°C is compared to previous years. In W1-W4 temperature increased about to 3.9 °C in 2014, followed by a significant decrease in over 2015 to 2.22°C. In the more southern areas (W5-W9) temperature has decrease little since 2014 but is above the overall mean temperature, and bottom temperature remain warm in the southernmost regions W7-W9 at 4.6°C. In shallow water in depth between 150-200 m the area weighted average bottom temperature is continuously rising from 2007 to 2014, but minor decrease have been observed since 2015. In conclusion the overall temperature has decreased by 0.87°C since 2014.

## Conclusions

### *Stock size*

In aggregate, the stock has continued the decline it has maintained, interrupted by small increase in 2008 - 2010 and again in 2013 and 2015, since it last peak in 2003. In fact at short term, there has been fluctuations in the stock with ups and downs in biomass with no clear trends since 2010. The overall survey biomass decreased by 25.2% over 2015, mainly caused by a significant decrease of 53.8% in the offshore regions W3-W7, and a minor decline of 17% in the survey biomass in Disko Bay & Vaigat. The offshore regions is now constitute 69% of the total surveyed biomass, and below the 1988-2016 mean (77%) but above the past 5 year mean (60%).

The offshore and inshore components have had very different trajectories since 2008. Inshore, in Disko Bay and Vaigat, the 2008–10 increase constituted almost a doubling of the survey biomass, which in the succeeding two years stayed close to the level reached in 2010. Surveyed biomass in Disko Bay decreased by 12% over 2012, but increase with 14.9% over 2013. This turns to a further continuing decline over 2014–2015, and is now below the 1988-2016 mean and 73.9% its past 5-year mean. Offshore—and in aggregate, the subdivisions having had somewhat different trajectories—there was almost no increase from 2008–10; and since 2010 the rapid decline offshore continued until 2014. Offshore survey biomass more than double over 2014, but turned to a decrease in the following year and is now in 2016 at 65% of its 28-year mean, but 3.7% larger than its past 5-year mean.

Fishable biomass decreased over 2015, again caused by a significantly decrease in offshore regions, as well as a 16% decline fishable biomass in Disko Bay. In 2016 fishable biomass constitute 92% of the total surveyed biomass. In both regions fishable biomass is at 65.6% of its 28-year mean offshore and 96.8% of its 28-year mean inshore (Disko Bay & Vaigat).

### *Stock distribution*

The area over which the stock is distributed has increased a little more since 2013, and it remains concentrated in Disko Bay, Vaigat and the Northern part of its range. Despite a decline over 2015, mean densities are high in Disko Bay and Vaigat, but is in 2016 a little below the 28-year mean. In almost all strata in the Offshore regions, densities decreased over 2015; was about 1.2 times lower the Northern regions, about 2.3 times lower in W3-W4 north, south, and west of Store Hellefiskebanke, 1.7 times lower in W5-W7 and is at 64% of its 27-year mean. Densities are practically zero on the continental slope west of the banks all the way from Store Hellefiskebanke to Kap Farvel.

The latitude index of the survey has declined from 2010 – 2015, interrupted by a increase in 2014 and 2016. The North index remain stable owing to high proportion of biomass in Disko Bay 34.8 – 52% from 2012 – 2016, and a low proportion of biomass in the W – regions. Survey biomass from the northern (above 66°N) and inshore regions constitute almost 94% of the total offshore survey biomass.

Since the late 1990s, the stock appears to be found in shallower water than before. Some large catches were made in shallow water in 2011–15 but the 150–200m stratum yielded its highest proportion of biomass in 2015 (21.7%). Only 3% of the survey biomass were observed in the shallow water in 2016. Proportions, of the survey biomass slightly decreased over 2007 - 2011 in the central depths from 201 to 400m, but the depth index for the survey biomass increased about 21 meter to 288m over 2015.

### *Stock composition*

By numbers both, the stock is heavily weighted toward males, but in biomass toward the females, which composed a high proportion of the total fishable and surveyed biomass. But in the same time, it seems deficient in large males. As a result the fishable biomass is a bit more than the average as a proportion of the total survey biomass, but compose to an exceptionally high degree of females.

Numbers at age-2 are overall, below the 20-year mean and in their lower quantile. The number of pre-recruits (14-16,5 mm CL) in total, at a very low level and even relative to the survey biomass below their lower quantile both in offshore as well as inshore regions.

### References

- ALLEN, J.A. 1959. On the biology of *Pandalus borealis* Kroger, with reference to a population off the Northumberland coast. *J. Mar. Biol. Assoc. U.K.*, **38**: 189-220.
- AITCHISON, J. 1955. On the distribution of a positive random variable having a discrete probability mass at the origin. *J. Am. Stat. Assoc.* 50: 901-908
- BURMEISTER, AD. AND KINGSLEY, M.C.S. 2015. The west Greenland trawl survey for *Pandalus borealis*, 2015, with reference to earlier results. *NAFO SCR Doc.*, No. 15/043, Serial No. N6478.
- BURMEISTER, AD. AND KINGSLEY, M.C.S. 2015. A provisional Assessment of the Shrimp Stock off West Greenland in 2015. *NAFO SCR Doc.*, No. 15/044, Serial No. N6479.
- CARLSSON, D.M. and P. KANNEWORFF. 2000. Stratified-random trawl survey for shrimp (*Pandalus borealis*) in NAFO Sub area 0+1, in 2000. *NAFO SCR Doc.*, No. 78, Serial No. N4335.
- CARLSSON, D., O. FOLMER, P. KANNEWORFF, M.C.S. KINGSLEY and M. PENNINGTON. 2000. Improving the West Greenland Trawl Survey for shrimp (*Pandalus borealis*). *J. North. Atl. Fish. Sci.*, **27**: 151-160.
- KINGSLEY, M.C.S. 2014. Numbers of Age-2 Shrimps in West Greenland—again. *NAFO SCR Doc.*, No. 14/058, Serial No. N6360.
- KINGSLEY, M.C.S. 2001a. Effects in 2001 of recent modifications to the design of the West Greenland Shrimp Survey. *NAFO SCR Doc.*, No. 176, Serial No. N4565.
- KINGSLEY, M.C.S. 2001b. Studies in 2001 on the end effect of the Skjervøy 3000 trawl in the West Greenland shrimp survey. *NAFO SCR Doc.*, No. 177, Serial No. N4566.
- KINGSLEY, M.C.S., P. KANNEWORFF and D.M. CARLSSON. 1999. Modifications to the design of the trawl survey for *Pandalus borealis* in West Greenland waters: effects on bias and precision. *NAFO SCR Doc.*, No. 105, Serial No. N4184.
- KINGSLEY, M.C.S., D.M. CARLSSON, P. KANNEWORFF and M. PENNINGTON. 2002. Spatial structure of the resource of *Pandalus borealis* and some implications for trawl survey. *Fish. Res.* **58**: 171-183.
- KINGSLEY, M.C.S., P. KANNEWORFF and D.M. CARLSSON. 2004. Buffered random sampling: a sequential inhibited spatial point process applied to sampling in a trawl survey for Northern shrimp *Pandalus borealis* in West Greenland waters. *ICES J. Mar. Sci.*, **61**: 12-24.
- KINGSLEY, M.C.S. 2008. Indices of distribution and location of shrimp biomass for the West Greenland research trawl survey. *NAFO SCR Doc.*, No 78, Serial No. N5610.
- MCCRARY, J.A. 1971. Sternal spines as a characteristic for differentiating between females of some Pandalidae. *J. Fish. Res. Board Can.*, **28**: 98-100.
- NAFO/ICES 2015. NAFO/ICES *Pandalus* Assessment Group Meeting, 9-16 September 2015, Northwest Atlantic Fisheries Centre, St. John's, Newfoundland, Canada. *NAFO SCS Doc.* 15/13 Serial No. N6497, ICES CM 2015/ACOM:14
- PENNINGTON, M. 1983. Efficient estimators of abundance for fish and plankton surveys. *Biometrics* 39: 281-286
- WIELAND, K. 2002a. A new length-weight relationship for Northern shrimp (*Pandalus borealis*) off West Greenland (NAFO Subareas 0+1). *NAFO SCR Doc.*, No. 144, Serial No. N4773.
- WIELAND, K. 2002b. The use of fine-meshed bags for sampling juvenile Northern shrimp (*Pandalus borealis*) in the West Greenland Bottom Trawl Survey. *NAFO SCR Doc.*, No. 145, Serial No. N4774.

- WIELAND, K. 2004. Abundance, mean size at age and growth of Northern shrimp (*Pandalus borealis*) juveniles and males off West Greenland in 1993–2004. *NAFO SCR Doc.*, No. 04/73, Serial No. N5043.
- WIELAND, K. 2005. Changes in recruitment, growth and stock size of Northern shrimp (*Pandalus borealis*) at West Greenland: temperature and density-dependent effects at released predation pressure. *ICES J. Mar. Sci.* **62**: 1454-1462.
- WIELAND, K. and D.M. CARLSSON. 2001. Geographical distribution and mean size of different life stages of Northern shrimp (*Pandalus borealis*) off West Greenland. *NAFO SCR Doc.*, No. 01/178, Serial No. N4567.
- WIELAND, K. and P. KANNEWORFF. 2004. Revision of depth contours and stratification of the West Greenland Bottom Trawl Survey for Northern shrimp. Technical report No. 56, Greenland Institute of Natural Resources ([http://www.natur.gl/publikationer/tekniske\\_rapporter](http://www.natur.gl/publikationer/tekniske_rapporter)).
- WIELAND, K. and M. STORR-PAULSEN. 2006. Effect of tow duration on catch rates and mean length of Northern shrimp (*Pandalus borealis*) and Greenland halibut (*Reinhardtius hippoglossoides*) in the West Greenland Bottom Trawl Survey. *Fish. Res.* **78**: 276-285.



Table 1. *Pandalus borealis* in West Greenland: vessels, trawl types and rigging parameters used in the West Greenland Bottom Trawl Survey for shrimp and fish, 1988–2015

	Vessel	Trawl	Bridle length (m)	Wing-spread (m)	
1988	Elias Kleist	Skjervøy	59.9	23.1	*
1989	Sisimiut	"	81.1	17.9	*
1990	Maniitsoq	"	59.9	23.1	*
1991	Paamiut	"	75.1	28.3	**
1992–2003	"	"	60.1	20.1–25.2	**
2004	"	"	54.0	25.7	**
2005–2015	"	Cosmos	54.0	27.4–28.8	**

(\*: from tank experiments (Per Kannevorff, pers.com.), \*\*: average for all valid tows calculated from measures of door spread and approximate geometry of the trawl).

Table 2. *Pandalus borealis* in West Greenland: survey estimates of total biomass 2016.

Stratum	Area (km <sup>2</sup> )	Stations	Biomass density (t/km <sup>2</sup> )	Biomass (Kt)	Biomass error variance	Error coefft of variation (%)
C0-2	903	0	0.000	0.000	0.0000	0.0
C0-3	2179	0	0.000	0.000	0.0000	0.0
C0-4	1154	0				
Overall C0	4236	0	0.000	0.000	0.0000	0.0
I1-1	407	2	2.942	1.196	0.0012	2.8
I1-2	1963	3	9.097	17.859	51.9404	40.4
I1-3	2441	2	8.149	19.895	14.2569	19.0
I1-4	1499	2	0.599	0.897	0.0023	5.4
I2-1	419	2	5.664	2.375	0.5006	29.8
I2-2	815	2	22.349	18.214	70.2419	46.0
I2-3	1085	3	3.064	3.324	3.5610	56.8
I2-4	1338	2	0.774	1.035	0.9767	95.5
Overall I	9967	18	6.501	64.795	141.4809	18.4
U1-1	2486	2	0.017	0.042	0.0013	84.5
U1-2	4633	2	0.949	4.399	5.6481	54.0
U1-3	4785	3	0.772	3.695	0.8505	25.0
U1-4	5129	2	0.018	0.092	0.0066	88.4
U2-2	6710	3	0.949	6.367	2.2635	23.6
U2-3	8481	3	1.334	11.318	26.2007	45.2
U2-4	7994	3	0.015	0.117	0.0046	57.6
U3-1	2012	2	0.366	0.736	0.0816	38.8
U3-2	3017	2	8.008	24.157	11.4742	14.0
U3-3	1675	2	4.728	7.918	3.7589	24.5
U3-4	2710	2	0.630	1.706	2.1863	86.7
Overall U	49631	26	1.220	60.548	52.4762	12.0
W1-1	2873	2	0.186	0.535	0.1789	79.0
W1-2	6099	6	1.658	10.112	3.4669	18.4

W1-3	7520	3	0.922	6.930	2.3601	22.2
W1-4*	816	2	0.000	0.000	0.0000	100.0
W2-1	1674	2	0.595	0.995	0.4094	64.3
W2-2	2612	3	8.244	21.530	61.4245	36.4
W2-3	1741	2	4.525	7.878	0.0291	2.2
W2-4	915	0	0.000	0.000	0.0000	0.0
W3-1	2122	2	0.000	0.000	0.0000	0.0
W3-2	4725	12	3.579	16.909	59.3612	45.6
W3-3	2085	2	0.974	2.031	0.2811	26.1
W3-4	2994	3	0.752	2.251	1.5760	55.8
W4-1	4119	2	0.074	0.305	0.0922	99.5
W4-2	1818	3	1.375	2.501	1.7303	52.6
W4-3	821	2	1.795	1.474	2.1182	98.7
W4-4	1961	2	0.038	0.074	0.0043	89.6
W5-1	3001	3	0.001	0.004	0.0000	58.4
W5-2	3648	13	0.379	1.382	0.6924	60.2
W5-3	1950	2	0.826	1.611	2.5951	100.0
W5-4	3021	2	0.001	0.003	0.0000	80.9
W6-1	1206	3	0.039	0.047	0.0006	50.8
W6-2	2006	3	3.126	6.271	27.1274	83.1
W6-3	1585	3	0.649	1.028	1.0573	100.0
W6-4	1234	1	0.000	0.000	0.0000	0.0
W7-1	2442	3	0.000	0.000	0.0000	0.0
W7-2	891	5	0.000	0.000	0.0000	52.7
W7-3	265	2	0.000	0.000	0.0000	0.0
W7-4	317	1	0.000	0.000	0.0000	0.0
W8-1	424	2	0.000	0.000	0.0000	100.0
W8-2	567	2	0.002	0.001	0.0000	100.0
W8-3	405	2	0.004	0.001	0.0000	100.0
W8-4	718	2	3.919	2.814	7.9106	100.0
W9-1	1711	8	0.000	0.000	0.0000	100.0
W9-2	938	3	0.000	0.000	0.0000	0.0
W9-3	516	2	0.001	0.000	0.0000	52.4
W9-4	430	2	0.000	0.000	0.0000	100.0
Overall W	72169	112	1.201	86.689	172.4157	15.1
Survey totals	136003	156	1.559	212.031	366.3728	9.0

† strata with 1 trawled station have been assigned an error coefficient of variation of 95%

\* area C0 was not surveyed in 2016 owing to ice cover and substrata W2-4 due to general trawl difficulties..

Table 3. *Pandalus borealis* in West Greenland: biomass estimates (Kt) for survey subdivisions and standard errors for the entire survey in 1988–2016.

Year	N1-9/ U1-3 <sup>1,5</sup>	C1-3/C0 <sup>1,6</sup>	W1-2	W3-4	W5-7 <sup>2</sup>	S/W8-9 <sup>1</sup>	D1-9/I1-2 <sup>3,1</sup>	Total	SE <sup>4</sup>
1988	22.6	9.5	55.1	85.5	17.7		39.2	229.7	24.7
1989	11.1	3.7	50.0	82.7	39.0		39.2	225.7	32.3
1990	11.0	9.1	78.6	53.9	23.5		39.2	215.3	32.6
1991	5.1	4.2	26.8	47.4	23.3		43.1	149.9	23.0
1992	18.1	22.2	46.2	30.6	45.8		41.4	204.4	32.5
1993	6.9	2.9	93.8	36.7	62.2		28.3	230.8	30.9
1994	6.6	6.0	95.0	44.5	32.6	16.7	34.0	235.4	51.7
1995	6.8	3.9	39.0	52.4	48.7	1.6	39.1	191.4	30.6
1996	8.8	1.5	46.4	31.5	80.0	3.3	44.3	215.9	40.4
1997	5.7	0.2	34.7	13.1	57.9	21.8	44.3	177.7	31.1
1998	7.0	0.4	37.8	100.6	45.1	18.6	51.8	261.2	57.6
1999	17.6	10.5	50.1	23.2	50.5	56.0	52.6	260.6	42.1
2000	8.4	10.7	62.1	69.8	71.0	21.8	73.0	316.9	40.3
2001	34.1	3.7	74.3	47.6	58.5	36.3	72.1	326.7	44.2
2002	17.4	5.4	114.0	62.1	94.9	40.5	85.8	420.2	60.0
2003	109.3	5.9	148.6	93.3	98.0	35.0	107.7	597.8	77.0
2004	111.2	3.5	152.8	96.5	102.6	15.4	81.4	563.4	103.7
2005	100.5	9.3	159.9	87.2	53.4	1.9	139.6	551.9	88.4
2006	54.7	45.8	108.9	60.6	92.2	12.5	110.7	484.0	65.1
2007	61.2	1.7	128.1	64.0	21.3	1.2	79.1	356.6	44.3
2008	91.7	16.7	61.3	40.0	20.9	0.7	50.8	282.1	28.3
2009	91.7	4.3	62.9	30.1	18.4	1.0	70.1	278.4	27.1
2010	73.1	3.0	89.6	65.3	13.5	0.9	99.3	344.7	44.6
2011	55.5	-( <sup>1</sup> )	69.2	6.1	34.3	2.6	92.9	260.6	36.1
2012	33.5	-( <sup>1</sup> )	33.8	7.0	23.1	0.4	92.5	190.3	20.6
2013	54.1	0.4	51.6	37.4	8.6	0.2	81.4	233.8	23.2
2014	29.4	-( <sup>1</sup> )	34.5	12.1	10.4	0.0	93.5	179.9	16.6
2015	75.4	-( <sup>1</sup> )	51.5	59.9	18.0	0.6	78.2	283.6	41.7
2016	60.5	-( <sup>1</sup> )	48.0	25.5	10.3	2.8	64.8	212.0	19.1

<sup>1</sup>: New stratification introduced in 2003 (regions N and S) and in 2004 (regions U, C and D)

<sup>2</sup>: Areas W6 and W7 were sampled from 1990 and 1993, respectively

<sup>3</sup>: D1-D9 1988-90 not sampled, but set to mean of 1991-1997.

<sup>4</sup>: Standard error calculated excluding D1-D9 in 1988-1990

<sup>5</sup>: Probably underestimated due to poor coverage of the northern part of the area N

<sup>6</sup>: Canada(C) in 2011, 2012, 2015 and 2016 was not sampled due to ice condition

Table 4. *Pandalus borealis* in West Greenland: error coefficients of variation (%) for the biomass estimates of five main survey regions and the entire survey area in 1988–2016.

Year	N1-9/ U1-3	C1-3/C0	W1-2	W3-4	W5-7	S/W8-9	D1-9/I1-2	Total suvey	Number of hauls
1988	31.4	40.0	17.9	26.4	42.5	-	-	14.41	131
1989	22.2	42.8	23.9	33.2	35.0	-	-	18.60	130
1990	43.5	39.9	22.7	39.5	42.0	-	-	18.30	109
1991	40.2	27.1	21.2	30.4	30.6	-	22.9	13.37	194
1992	16.9	68.9	15.5	17.9	42.0	-	15.7	13.84	167
1993	51.6	53.3	14.6	28.0	28.5	-	19.4	11.66	157
1994	48.7	18.3	27.2	16.8	45.9	99.2	26.0	19.11	157
1995	47.1	44.7	20.0	22.6	36.3	74.0	17.7	13.93	163
1996	52.6	91.0	23.9	19.8	36.2	95.0	10.6	16.31	148
1997	37.9	61.9	15.8	20.2	43.1	14.6	14.5	15.26	167
1998	40.4	44.0	33.2	44.5	30.6	58.8	18.4	19.19	209
1999	51.1	80.0	13.1	14.8	27.7	52.1	14.2	14.08	227
2000	36.1	7.8	21.5	32.7	18.5	56.8	12.9	11.08	198
2001	26.5	44.5	24.9	45.0	25.6	22.8	18.6	11.77	224
2002	56.0	45.4	16.0	28.9	36.5	55.0	18.7	12.44	216
2003	26.8	44.4	17.5	16.0	34.9	49.9	17.5	11.21	172
2004	24.9	22.6	22.8	27.1	24.1	71.4	11.6	16.03	187
2005	17.4	97.3	22.8	38.0	50.4	48.7	34.6	16.02	194
2006	20.5	79.0	19.2	20.5	41.3	79.4	12.9	13.41	223
2007	21.9	45.1	19.8	32.6	8.8	56.8	20.8	12.43	223
2008	17.3	80.7	15.6	33.1	43.5	44.8	11.5	10.02	204
2009	22.1	59.6	12.8	18.7	39.0	53.0	15.9	9.72	247
2010	11.1	29.7	27.0	61.0	66.9	98.7	10.0	12.94	270
2011	9.8	-(1)	19.1	26.2	89.0	80.5	12.4	13.85	192
2012	16.0	-(1)	12.4	23.9	45.8	84.6	17.4	10.83	193
2013	16.1	76.8	21.1	41.8	55.6	78.6	11.6	9.93	181
2014	26.8	-(1)	19.6	43.2	34.7	65.0	11.6	9.23	189
2015	20.3	-(1)	33.2	58.3	45.5	51.8	10.7	14.69	186
2016	12.0	-(1)	15.7	31.6	54.2	99.9	18.4	9.03	156
Mean 1994-2016								13.15	

1: C (Canada) in 2011 – 2016, except 2014 was not sampled

Table 5. *Pandalus borealis* in West Greenland: estimated mean densities (t/km<sup>2</sup>) for survey subdivisions in 1988–2016.

Year	N1-9/ U1- 3	C1-3/C0	W1-2	W3-4	W5-7	S/W8-9	D1-9/I1-2
1988	0.54	2.77	2.34	3.94	1.76	-	-
1989	0.25	1.08	2.76	3.81	3.88	-	-
1990	0.25	2.65	3.33	2.48	1.59	-	-
1991	0.12	1.23	1.14	2.18	1.57	-	4.60
1992	0.44	6.46	1.96	1.41	3.09	-	4.42
1993	0.17	0.85	3.55	1.68	3.32	-	3.02
1994	0.17	1.76	3.59	2.03	1.74	3.22	3.63
1995	0.18	1.15	1.47	2.39	2.60	0.24	4.17
1996	0.23	0.44	1.75	1.44	4.27	0.51	4.73
1997	0.15	0.06	1.31	0.60	3.09	3.35	4.73
1998	0.18	0.11	1.43	4.59	2.41	2.85	5.54
1999	0.46	3.06	1.89	1.10	2.70	8.59	5.62
2000	0.22	3.10	2.35	3.18	3.79	3.35	7.80
2001	0.89	1.08	2.81	2.17	3.12	5.57	7.70
2002	0.45	1.57	4.31	4.46	5.07	6.21	9.16
2003	2.22	1.39	6.11	6.25	5.23	5.80	11.49
2004	2.20	0.82	6.25	4.71	4.76	2.65	8.37
2005	1.99	2.20	6.54	4.25	2.48	0.34	14.19
2006	1.08	10.81	4.46	2.96	4.28	2.20	11.26
2007	1.21	0.40	5.24	3.12	0.99	0.21	8.04
2008	1.85	3.94	2.53	1.94	0.97	0.13	5.09
2009	1.85	1.01	2.59	1.46	0.85	0.17	7.03
2010	1.47	0.70	3.70	3.16	0.63	0.16	9.96
2011	1.12	-	2.95	0.30	1.59	0.46	9.32
2012	0.67		1.40	0.34	1.07	0.07	9.28
2013	1.09	0.13	2.13	1.81	0.40	0.05	8.16
2014	0.59	0.00	1.42	0.59	0.48	0.00	9.38
2015	1.52	-	2.12	2.90	0.84	0.10	7.85
2016	1.22	-	1.98	1.24	0.48	0.49	6.50

Table 6. *Pandalus borealis* in West Greenland: estimated numbers (billions) by sex in 1988–2016.

Year	Males	Females	Total	Males %	Females %
1988 <sup>1</sup>	26.8	9.3	36.1	74.3	25.7
1989 <sup>1</sup>	39.0	6.9	45.9	85.0	15.0
1990 <sup>1</sup>	29.3	8.9	38.1	76.8	23.2
1991	19.6	5.1	24.7	79.3	20.7
1992	29.4	6.5	35.9	81.9	18.1
1993	34.8	8.3	43.1	80.7	19.3
1994	32.0	8.9	40.9	78.3	21.7
1995	27.7	6.5	34.2	80.9	19.1
1996	38.2	6.6	44.8	85.2	14.8
1997	27.2	6.3	33.5	81.2	18.8
1998	41.0	9.9	50.9	80.5	19.5
1999	42.5	9.9	52.3	81.1	18.9
2000	62.4	11.1	73.4	84.9	15.1
2001	56.6	11.8	68.4	82.7	17.3
2002	85.3	14.9	100.1	85.1	14.9
2003	99.4	24.9	124.4	80.0	20.0
2004	89.4	26.3	115.8	77.3	22.7
2005	94.5	25.1	119.6	79.0	21.0
2006	78.3	24.6	102.9	76.1	23.9
2007	55.1	16.0	71.1	77.4	22.6
2008	42.4	11.8	54.2	78.2	21.8
2009	41.3	12.0	53.3	77.4	22.6
2010	52.9	13.5	66.3	79.7	20.3
2011 <sup>2</sup>	36.8	13.0	49.8	74.0	26.0
2012 <sup>2</sup>	28.8	8.7	37.4	77.0	23.2
2013	31.1	12.0	43.1	72.2	27.8
2014	22.8	9.2	32	71.3	28.8
2015	51.3	10.6	61.9	82.9	17.1
2016	27.6	10.5	38.1	72.4	27.6
Average	47.0	12.1	58.4	79.1	20.9

<sup>1</sup> mean values for Disko Bay and Vaigat in 1991–1997 have been inserted for 1988–1990, and included in the calculation of the average.

<sup>2</sup> area C0 was not surveyed in 2011 – 2016, except 2013 owing to sea ice; and in 2014 due to time trouble; no correction has been made.

<sup>3</sup>Substrata W1-4 in 2011 and 2015, W9-4 in 2015 and W9-4 were not surveyed due to trawl difficulties and poor bottom conditions.

Table 7. *Pandalus borealis* in West Greenland: biomass estimates ('000 t) by sex based on length-weight distributions from the West Greenland shrimp survey in 1988–2016

Year	Males	Females	Total	Males %	Females %
1988	134.7	94.8	229.5	58.7	41.3
1989	157.1	68.6	225.7	69.6	30.4
1990	129.4	85.4	214.9	60.2	39.8
1991	100.5	49.4	149.9	67.0	33.0
1992	141.3	63.1	204.4	69.1	30.9
1993	149.2	81.9	231.1	64.6	35.4
1994	146.5	88.9	235.4	62.2	37.8
1995	124.5	66.9	191.4	65.0	35.0
1996	147.9	68.0	215.9	68.5	31.5
1997	114.7	62.9	177.7	64.6	35.4
1998	170.4	90.9	261.3	65.2	34.8
1999	166.7	93.9	260.6	64.0	36.0
2000	213.8	100.2	314.0	68.1	31.9
2001	199.1	108.3	307.4	64.8	35.2
2002	293.6	126.6	420.2	69.9	30.1
2003	389.2	208.6	597.8	65.1	34.9
2004	353.1	210.3	563.4	62.7	37.3
2005	355.2	196.7	551.9	64.4	35.6
2006	297.4	188.0	485.4	61.3	38.7
2007	227.8	128.7	356.6	63.9	36.1
2008	182.6	99.5	282.1	64.7	35.3
2009	173.5	105.0	278.4	62.3	37.7
2010	222.3	122.4	344.7	64.5	35.5
2011	148.5	112.0	260.6	57.0	43.0
2012	115.4	74.9	190.3	60.7	39.3
2013	129.9	103.9	233.8	55.6	44.4
2014	93.8	86.0	179.8	52.2	47.8
2015	184.6	99.0	283.6	65.1	34.9
2016	113.6	98.4	212.0	53.6	46.4
Average	185.4	106.3	291.7	63.3	36.7

<sup>1</sup> 1991–1997 mean values for Disko Bay and Vaigat have been used for 1988–1990, and included in the calculation of the average.

<sup>2</sup> data for 2005–2010 was re-analyzed in 2011.

Table 8. *Pandalus borealis* in West Greenland: estimates of fishable biomass ( $\geq 17$ mm CL; '000 t) from the West Greenland shrimp survey in 1988–2016

Year	Offshore fishable		Disko fishable		Overall fishable	
	biomass	%	biomass	%	biomass	%
1988	186.2		37.0		223.2	
1989	171.9		37.0		209.0	
1990	170.0		37.0		207.0	
1991	104.7	98%	41.3	96%	146.0	97%
1992	154.8	95%	39.4	95%	194.2	95%
1993	189.4	94%	27.1	96%	216.5	94%
1994	191.0	95%	32.1	94%	223.1	95%
1995	144.9	95%	38.3	98%	183.2	96%
1996	150.6	88%	41.5	94%	192.1	89%
1997	127.7	96%	39.4	89%	167.1	94%
1998	197.2	94%	47.1	91%	244.3	94%
1999	195.0	94%	42.3	80%	237.3	91%
2000	219.8	90%	60.6	83%	280.3	88%
2001	216.8	85%	63.7	88%	280.5	86%
2002	302.2	90%	67.2	78%	369.5	88%
2003	454.0	93%	94.3	88%	548.3	92%
2004	457.5	95%	70.8	87%	528.3	94%
2005	381.8	93%	112.3	80%	494.2	90%
2006	358.6	96%	92.4	83%	451.0	93%
2007	264.7	95%	71.3	90%	336.1	94%
2008	216.8	94%	45.8	90%	262.6	93%
2009	192.2	92%	62.8	90%	255.1	92%
2010	229.8	94%	88.9	90%	318.7	92%
2011	155.9	93%	83.1	89%	239.0	92%
2012	89.5	91%	83.9	91%	173.3	91%
2013	143.6	94%	74.6	92%	218.1	93%
2014	81.7	95%	85.5	91%	167.2	93%
2015	178.8	87%	69.3	89%	248.1	87%
2016	137.3	70%	58.3	90%	195.6	92%
Average	209.1	93%	60.2	89%	269.3	92%

<sup>1</sup> 1991–1997 mean values for Disko Bay and Vaigat have been used for 1988–1990, and included in the calculation of the average.

<sup>2</sup> data for 2005–2010 was re-analysed in 2011.

<sup>3</sup> area C0 was not surveyed in 2011–2016, except 2013 nor substratum W1–4 in 2011 and 2015, owing to sea ice; and W9–4 in 2015 and W2–4 due to trawl difficulties.



Table 9. *Pandalus borealis* in West Greenland: mean carapace length (mm) at age 2 in 2005–2016, with standard deviations and coefficients of variation (- : not present, ( ) : fixed in the final MIX run, na: no data).

Mean:

Year	Region / Strata					
	U1-U3 150-600 m	I1+I2 150-600 m	C0 200-600 m	W1-W4 150-600 m	W5+W6 150-600 m	W7-W9 150-600 m
2005	11.1	11.5	-	12.5	-	-
2006	11.5	11.4	-	12.3	12.5	-
2007	-	11.5	-	-	12.5	-
2008	13.4	12.6	-	12.4	11.5	-
2009	12.2	11.8	-	12.3	12.2	-
2010	10.8	11.9	-	13.1	-	-
2011	-	11.2	na	11.3	-	-
2012	11.7	11.0	na	-	11.7	-
2013	11.2	13.1	-	13.5	13.0	-
2014	12.3	13.5	na	13.1	-	-
2015	11.5	13.4	na	12.1	12.0	-
2016	12.6	12.0	na	12.1	13.7	-

Standard deviation:

Year	Region / Strata					
	U1-U3 150-600 m	I1+I2 150-600 m	C0 200-600 m	W1-W4 150-600 m	W5+W6 150-600 m	W7-W9 150-600 m
2005	0.9	1.5	-	1.3	-	-
2006	1	1.1	-	1.36	1.26	-
2007	-	1.37	-	-	1.32	-
2008	1.65	1.87	-	1.28	0.83	-
2009	1.36	1.39	-	1.12	1.26	-
2010	1.28	1.20	-	1.27	-	-
2011	-	1.00	na	1.17	-	-
2012	1.53	1.09	na	-	0.40	-
2013	0.96	1.32	-	1.08	1.4	-
2014	0.99	1.38	na	1.41	-	-
2015	0.88	1.51	na	1.13	0.96	-
2016	1.25	1.53	na	1.89	0.94	-

Coefficient of  
variation:

Year	Region / Strata					
	U1-U3 150-600 m	I1+I2 150-600 m	C0 200-600 m	W1-W4 150-600 m	W5+W6 150-600 m	W7-W9 150-600 m
2005	0.08	0.13	-	0.1	-	-
2006	0.09	0.09	-	0.11	0.1	-
2007	-	0.12	-	-	0.11	-
2008	0.12	0.15	-	0.10	0.07	-
2009	0.11	0.12	-	0.09	0.11	-
2010	0.12	0.10	-	0.10	-	-
2011	-	0.09	na	0.10	-	-
2012	0.13	0.10	na	-	0.03	-
2013	0.09	0.10	-	0.08	0.11	-
2014	0.08	0.1	na	0.11	-	-
2015	0.08	0.11	na	0.09	0.08	-
2016	0.08	0.1	na	0.1	0.07	-

Table 10. *Pandalus borealis* in West Greenland: survey estimate of numbers (billions) at age 2 years, 1993–2016.

Year	U1 to U3	I1 and I2	C0 and W1 to W4		W5 and W6	W7 to W9	Total
1993	0.06	0.08	2.60		1.54	0.00	4.28
1994	0.01	0.21	1.51		1.20	0.00	2.92
1995	0.02	0.11	0.82		1.37	0.00	2.32
1996	0.11	1.25	2.45		6.20	0.00	10.01
1997	0.05	1.37	0.52		1.27	0.00	3.22
1998	0.04	1.79	2.01		2.60	0.00	6.44
1999	0.42	5.06	2.66		3.22	0.00	11.36
2000	0.33	5.54	4.92		3.50	0.01	14.29
2001	1.66	5.44	7.79		1.01	0.01	15.90
2002	0.02	3.98	3.41		2.97	0.04	10.42
2003	0.76	1.11	1.70		2.88	0.03	6.48
2004	0.64	3.39	2.24		0.47	0.01	6.75
2005	0.32	3.47	0.23		0.05	0.00	4.07
2006	0.33	1.86	2.12		0.78	0.00	5.09
2007	0.00	0.96	0.00	0.00	0.38	0.00	1.34
2008	2.96	1.48	0.00	0.86	0.36	0.00	5.66
2009	1.95	2.05	0.00	0.91	0.14	0.00	5.05
2010	0.95	2.23	0.00	1.13	0.00	0.00	4.31
2011	0.00	2.52	-	0.44	0.00	0.00	2.96
2012	0.30	1.20	-	0.00	0.07	0.00	1.57
2013	0.40	2.03	0.00	1.63	0.00	0.00	4.06
2014	0.17	3.30	-	1.36	-	0.00	4.83
2015	1.38	3.62	-	5.35	0.50	0.00	10.85
2016	0.66	2.51	-	1.04	0.05	0.00	4.26
Average:	0.56	2.36	2.03		1.33	0.00	6.19

<sup>1</sup> data for 1993 to 2004 has been converted from Skervøy to Cosmos trawl;

<sup>2</sup> for area C0 – numbers of age 2 shrimps is in general very low. Estimating the components in the mixture distribution to the data using maximum-likelihood estimation for the grouped data, resulting is very poor fits.

\* Numbers of age-2 shrimps from 2005 – 2014 have been recalculated in 2014.




Fig.1. *Pandalus borealis* in West Greenland: density distribution from 156 trawl-survey stations in 2016.




Fig.2a. *Pandalus borealis* in West Greenland: estimated survey biomass, 1988–2016. Area C0-4 was not surveyed in 2011 – 2016, except 2013 because of ice cover, in 2014 W9-4 due to poor bottom conditions, area C0 in 2014 owing to time trouble and W2-4 due to trawl difficulties and poor bottom conditions.




Fig.2b. *Pandalus borealis* in West Greenland: distribution of survey biomass between major survey regions, 1991 – 2016. Area C0-4 was not surveyed in 2011 – 2016, except 2013 because of ice cover, in 2014 W9-4 due to poor bottom conditions, area C0 in 2014 owing to time trouble and W2-4 due to trawl difficulties and poor bottom conditions.




Fig.3 *Pandalus borealis* in West Greenland: mean survey density of Northern shrimp in a) ●- Disko Bay and Vaigat (since 1991 7–8% of the survey area), ▲-offshore (92–93%) and b) total surveyed regions.




Fig. 4. *Pandalus borealis* in West Greenland: indices of distribution and location of shrimp biomass in the West Greenland trawl survey 1994–2016 (3-point moving averages). ○- North index (3-pt moving average) and ●- Spread index (3-pt moving average).




Fig. 5a. *Pandalus borealis* in West Greenland: survey biomass estimates by depth stratum, 1988–2016. 3-point moving averages; Until 2003, only areas C and W were substratified by depth. -●-: 150 – 200m, -○-: 200–300m, -▼-: 300–400 and -△-400–600m.




Fig. 5b. *Pandalus borealis* in West Greenland: distribution of survey biomass between 150 and 600 m by depth, 1988–2016. Until 2003, only areas C and W were substratified by depth. -●-: 150 – 200 m, -○-: 200–300 m, -▼-: 300–400 m and -△-400–600 m




Fig. 5c. *Pandalus borealis* in West Greenland: depth index for survey biomass, 1988–2016. Until 2003, only areas C and W were substratified by depth.




Fig. 6. *Pandalus borealis* in West Greenland: depth index of survey biomass vs. area-weighted mean bottom temperature from survey trawl-door measurements, 1990–2016.






Fig. 7. *Pandalus borealis* in West Greenland: distribution of lengths from survey length analyses in 2012 (black line), 2013 (pink line), 2014 (grøn line), 2015 (blue line) and 2016 (rød lined).




Fig. 8. *Pandalus borealis* in West Greenland: mean weight, by sex, from survey data, 1988–2016.




Fig. 9. *Pandalus borealis* in West Greenland: regional length frequencies in 2016.




Fig. 10. *Pandalus borealis* in West Greenland: length frequencies offshore, in Disko Bay and Vaigat, and overall, 2007–2016.




Fig. 11a. *Pandalus borealis* in West Greenland: index of numbers at age 2, 1993–2016.




Fig. 11b. *Pandalus borealis* in West Greenland: index of numbers at age 2, 1993–2016.




Fig. 12. *Pandalus borealis* in West Greenland: survey estimates of numbers at age 2 in 1993–2016 against female biomass 3 years earlier (labels denote years in which age-2 numbers were estimated).




Fig.13. *Pandalus borealis* in West Greenland: lagged fishable biomass vs. survey estimates of numbers at age 2 from 1993 to 2016 (autocorrelated regressions with 95% confidence and prediction intervals).




Fig.14. *Pandalus borealis* in West Greenland: area-weighted mean bottom temperature for survey regions (see Fig.1 for locations), depth strata in offshore areas C and W1–W7, and the entire survey area in 1990–2016.