

NOT TO BE CITED WITHOUT PRIOR  
REFERENCE TO THE AUTHOR(S)

Northwest Atlantic



Fisheries Organization

**Serial No. N6709**

**NAFO SCR Doc. 17-049**

**SCIENTIFIC COUNCIL MEETING – JUNE 2017**

An assessment of the witch flounder resource in NAFO Divisions 3NO

by

E. Lee, J. Morgan, R. M. Rideout, D. Ings, L. Wheeland  
Fisheries and Oceans Canada  
Northwest Atlantic Fisheries Center  
P.O. Box 5667 St. John's,  
Canada, A1C 5X1

**Abstract**

This stock underwent full assessment in 2014 based on survey indices and in 2015 utilizing a surplus production model in a Bayesian framework. An interim monitoring report was provided in 2016. Witch flounder in Divs. 3NO was under moratorium to directed fishing from 1995 to 2014. The 2015 assessment indicated that the stock steadily increased since 1999 and was at 81%  $B_{msy}$ . In 2015 the risk of the stock being below  $B_{lim}$  or above  $F_{lim}$  was concluded to be less than 1%. Based upon this information, the NAFO Fisheries Commission set a total allowable catch (TAC) for 2015, 2016 and 2017 at 1,000 tonnes (t), 2,172 t and 2,225 t respectively. Despite the 1,000 t quota available, the catch reported for 2015 (389 t) was consistent with the bycatch range (300-400 t) reported since 2010. The reported catch for 2016 was 1,062 t of the available TAC of 2,172 t.

Indices of abundance and biomass for witch flounder from Canadian spring surveys in Divs. 3NO although variable, had shown a general decreasing trend from 1985 to 1998, a general increasing trend from 1998 to 2003, and a general decreasing trend from 2003 to 2010. From 2010 to 2013 the index increased to values near the series high from 1987. Biomass values declined substantially from a high in 2013 to a value 49% of the time series average in 2015. Biomass values increased slightly in 2016.

Biomass and abundance indices from the Canadian fall survey series for NAFO Divs. 3NO were less variable, showing a general increasing trend from 1996 to 2009 but have declined since to 57% of the time series average in 2016.

Indices from Spanish surveys in the NAFO Regulatory Area of Divs. 3NO increased from 2002 to 2004, decreased from 2004 to 2009, were variable from 2010 to 2012, declined sharply from 2012 to 2014 and increased from 2014 to 2016.

Recruitment (defined as fish less than 21cm) in both the spring and fall Canadian surveys although somewhat variable has generally been low since 2003. Recruitment in spring and fall surveys in 2016 approached the lowest of the time series.



The surplus production model results indicate that stock size decreased from the late 1960s to the late 1990s and then increased from 1999 to 2010 but has since decreased. The model suggests that a maximum sustainable yield ( $MSY$ ) of 3 641 (2 263-5 689) t can be produced by total stock biomass of 50 000 (35 559-69 581) t ( $B_{msy}$ ) at a fishing mortality rate ( $F_{msy}$ ) of 0.07 (0.04-0.13).

### **Fisheries and Management**

As noted in previous reports (Lee et.al. 2014 and Brodie et.al. 2011), species-specific catch statistics for flatfish prior to 1973 were largely developed from breakdowns of unspecified flounders and therefore should be considered with caution. Catches in the 1960s peaked at 11,000-12,000 tons in 1967-68 and remained relatively high during the next several years (Table 1; Fig. 1). Catch peaked at a time series high of 15,000 tons in 1971 and subsequently declined over the next decade to levels between 2000 and 4000 tons in the early 1980s (Table1; Fig 1).

The first total allowable catch (TAC) for witch flounder was introduced by ICNAF in 1974 at a level of 10,000 tons, largely based on average historical catches (Table 1; Fig. 1). This remained in effect until 1979 when it was reduced to 7,000 tons in consideration of declining commercial catch rates. It was further reduced to 5,000 tons in 1981 and remained at that level until 1993. The Scientific Council advised that for 1994, catches from this stock should not exceed 3,000 tons. A TAC of 3,000 tons was agreed by the NAFO Fisheries Commission, however, it was also agreed that no directed fishery would be conducted for witch flounder in 1994 to permit rebuilding due to the poor state of the stock. The NAFO Fisheries Commission introduced a complete moratorium for directed fishing in 1995, which was continued through 2014. There was no directed fishing on this stock from 1994 to 2014. A 1,000 t TAC was adopted for 3NO Witch Flounder beginning in 2015 with a TAC increase to 2,172 t and 2,225 t in 2016 and 2017 respectively. The 2016 Canadian catch was 799 t from a Canadian TAC of 1,303 t. Non-Canadian catch in 2016 was 263 t for a total estimated catch of 1,015 t of an available 2,172 t quota. The NAFO Fisheries Commission set a TAC for 2015, 2016 and 2017 at 1,000 t, 2,172 t and 2,225 t respectively. Despite the 1,000 ton quota available, the catch reported for 2015 (359 t) was consistent with the bycatch range (300-400 t) reported since 2010. The reported catch for 2016 was 1,015 t of the available TAC of 2,172 t (Table 1; Fig. 1). Not all Contracting Parties with quota have resumed directed fishing for witch flounder. In 2016 the catch (a combination of directed and bycatch fisheries) was estimated to be 1,062 t.

Annual catches rose rapidly to around 9,000 tons in 1985 and 1986 as a result of an increase in fishing effort in the NAFO Regulatory Area, primarily on the "tail" of the Grand Bank in Division 3N. Catches remained relatively high in 1987 and 1988 at around 7,500 tons. During 1990-93 estimated catches were in the range of 4,200-5,000 tons. The estimated catch for 1994 was in the order of 1,100 tons. A moratorium was introduced for this stock in 1995 (Table 1; Fig. 1). The catch dropped to 300 tons in 1995 likely as a result of a substantial reduction in fishing effort for Greenland halibut where witch flounder comprises a bycatch. Bycatch then increased steadily and by 1999 was about 800 tons, although it declined again to an estimated 450 tons in 2002. In 2003 several sources of catch data were available and a single source could not be considered as the most valid. As a result, catches were estimated to be 1544 t in 2003 (midpoint of a range of estimates) which declined to about 200 t in 2007, increased to 421 t in 2010 then declined slightly to about 335 t in 2014 (Table 1; Fig. 1).

Historically, the fishery was conducted primarily by Canada and the former Soviet Union. Canadian catches fluctuated from between 1,200 and 3,000 tons from 1985-91 but increased to about 4,300 tons in 1992 and 1993 (Table 1; Fig. 1). Canadian catches since the 1995 moratorium have averaged 31 t per year. Catches by

the USSR/Russian vessels declined from between 1,000 and 2,000 tons in the period 1982-88 and have averaged 29 t per year since the 1995 moratorium. Combined catch from other countries since 1995 has been in the range of 170 (2007) to 1500 t (2003) with an average annual catch of about 330 t (Table 1; Fig. 1).

#### *Data from commercial fisheries*

Length frequencies were available from observer data for Canadian witch flounder directed fisheries in NAFO Div. 30 in 2016. Canadian data indicated the catch ranged between 35 and 50 cm with a mean length of 42 cm (Fig. 2). Length frequencies were available from bycatches in directed fisheries for yellowtail flounder, redfish, Greenland halibut, and skate by Spain, Portugal, and Russia in 2016 (Fig. 2). The Spanish data (SCS 17/04) from Divs. 3NO indicated most of the witch flounder bycatch was between 28 and 46 cm in length. In the Portuguese data (SCS 17/05) for Div. 30 the witch flounder bycatch was dominated by lengths between 26 cm and 34 cm, with a mode at 32 cm and mean length of 31.7 cm (Fig. 2). In the Russian data (SCS 17/11) length frequencies of witch flounder bycatch in Div. 30 indicated the length ranged from 38 to 54 cm with a mean of 44.1 cm (Fig. 2).

### **Research Vessel Surveys**

#### *Canadian RV surveys*

#### *Spring Surveys*

Stratified-random research vessel surveys have been carried out by Canada on the Grand Banks in NAFO Divs. 3NO during spring since 1971, although during the early period coverage was limited and, in fact, for most years up to 1990, only surveyed depths to 366 meters (Tables 2 and 3). However, since 1991, depth coverage was extended to 731 meters. In 1993 only, spring surveys were completed to a depth of 914 m. During the course of the 2006 Canadian spring survey, operational difficulties lead to incomplete coverage of the survey in Divs. 3NO (Tables 2 and 3). Otherwise, spring surveys in Div. 3N were completed for most strata in all years from 1991 to 2016 to a depth of 731 m except for 1997, 2008, and 2012 which were each missing one stratum (Tables 2 and 3). Spring surveys in Div. 30 were completed for most strata in all years from 1991 to 2016 to a depth of 731 m except for 2011 which was missing one stratum and 2016 which was missing two strata (Tables 2 and 3).

#### *Fall Surveys*

In addition to spring surveys, a time series of fall surveys was begun in 1990 to investigate seasonal variation in stock distribution and abundance of various groundfish species (Tables 4 and 5). Note that due to operational difficulties there was no fall survey of NAFO Divs. 3NO in 2014. From fall 1998 the survey depth range in Div. 3N was further extended occasionally from the previous maximum depth range of 731 m to 1463 m (Table 4). Only four fall surveys have covered the Div. 3N deeper strata completely (2000, 2001, 2002, and 2007) or partially (2 missing in 1998, 4 missing in 2005, 8 missing in 2009, and 11 missing in 2010) (Table 4). Fall surveys in Div. 3N were limited to 366 m in 1990, and limited to 731 m from 1991 to 1997, in 1999, 2003, 2004, and 2008, and from 2010 to 2013 (Table 4). From fall 2000 the survey depth range in Div. 30 was extended occasionally from the previous maximum depth range of 1097 m to 1463 m (Table 5). Only six fall surveys since then have covered the Div. 30 deeper strata completely (2000 to 2002, 2005, 2007, and 2009) or partially (8 missing in 2003) (Table 5). Except for 1990 (549 m) and 1998 (1097 m) Div. 30 fall surveys have

primarily been limited to a depth range of 731 m (1991 to 1996, 1999, 2003, 2004, 2006, 2008, and 2010 to 2016) (Table 5).

Beginning with the fall survey in 1995, the survey gear was changed from an *Engel 145* groundfish trawl with steel bobbin footgear to a *Campelen 1800* shrimp trawl with rockhopper footgear. The data from the earlier Engel surveys have been converted to Campelen 1800 trawl catch equivalents. Only the converted survey data are presented but some caution should be used in comparing converted Engel data with data from the Campelen trawl series.

#### *Survey Stock Indices 3N and 30*

Biomass (Tables 6-9) and abundance (Tables 10-13) estimates by stratum are presented for the spring and fall surveys in NAFO Divs. 3N and 30 respectively. Mean numbers (Tables 14-17) and weights (Tables 18-21) per tow are also presented by stratum and division for the spring and fall surveys. Graphical plots to better illustrate the comparative trends in stock biomass, abundance, and mean numbers/weights per tow by season/year are presented for NAFO Divs. 3N and 30 separately and combined, in Figures 3-8. The time series from 1984 to 2016 indicates that the majority of the stock resides in NAFO Div. 30, with biomass and abundance estimates up to 10 times the estimates for NAFO Div. 3N during spring surveys and approximately 1.5 to 2.5 times the estimates for NAFO Div. 3N during fall surveys (Figs. 3 and 5).

Spring stock indices in NAFO Div. 3N indicate a high degree of variability over the time series with an upward trend from 2008 to 2011 followed by a sharp decline to 2013 and an increase from 2013 to 2014 to the highest biomass levels of the entire time series (Fig. 3). Indices declined in 2015 and increased slightly from 2015 to 2016. (Fig 3). Spring stock indices in NAFO Div. 30, which are also highly variable, indicate a downward trend from 2004 to 2010, a sharp increase from 2010 to 2013 equivalent to the highest levels of the time series. This is followed by a sharp decline in spring stock indices from 2013 to 2015 to levels at or below the mean levels for the time series and a small increase from 2015 to 2016 (Fig. 3).

Due to operational difficulties there was no fall survey of NAFO Divs. 3NO in 2014. Fall stock indices in NAFO Div. 3N were for the most part consistently low from 1990 to 2006 followed by a sharp increase from 2007 to 2009, a steep decrease from 2010 to 2011, a steep increase in 2012, and a sharp decrease from 2012 to 2016 (Fig. 5). Fall stock indices in NAFO Div. 30 were quite variable from 1990 to 2004 followed by a moderate decline from 2005 to 2007, a sharp increase to 2009, and a fairly consistent sharp decline from 2009 to 2015 and a slight increase from 2015 to 2016 (Fig 5).

#### *Spring Surveys 3N and 30*

Indices of biomass and mean weight per tow derived from spring surveys (which are the longer time series) in NAFO Div. 3N were at fairly low levels from 1988 to 2004 but have generally been higher (except for a sharp decrease in 2008) since then (Figs. 3 and 4; Tables 6 and 18). This was followed by a sharp decline from 2011 to 2013 to estimates which, although reduced, were still 2 times the range of the 1989-2004 “low” period and were slightly above the previous highest levels reported in 1984 and 1988 (Figs. 3 and 4; Tables 6 and 18). This was followed by a sharp increase from 2013 to 2014 to the highest estimates of the time series for biomass (2500 t) and mean weight per tow (1.0 kg) (Figs. 3 and 4; Tables 6 and 18). Indices declined from 2014 to 2015 prior to increasing slightly from 2015 to 2016 (biomass of 1500 to 1900 t and mean weight per tow of 0.6 to 0.8 kg) (Figs. 3 and 4; Tables 6 and 18).

The estimates for abundance and mean number per tow in NAFO Div. 3N were highly variable over the time series but show an increasing trend from 2008 to 2011 followed by a marked decline from 2011 to 2013 (Figs. 3 and 4; Tables 10 and 14). This was followed by a sharp increase from 2013 to 2014 to values approaching the highest estimates of the time series observed in 2011. Estimates for 2011 were an abundance of 6 million fish and mean number per tow of 2.5. Indices declined from 2014 to 2015 prior to increasing slightly from 2015 to 2016 (abundance of 2.7 to 3.6 million fish and mean number per tow of 1.1 to 1.5) (Figs. 3 and 4; Tables 10 and 14).

In most years (1989-2004) of the time series the spring stock indices in NAFO Div. 3N were estimated to be less than 1000 tons or 2 million fish, (Fig. 3; Tables 6 and 10). However, from 2005 onward, 7 of the 8 survey years in NAFO Div. 3N have produced higher than average index values for biomass and abundance, and the 2009, 2010 and 2011 values for both 3N indices are more than double the time series averages, however, this was followed by a decline in 2012 and 2013 back to levels that are still 1.5-2 times the mean of the time series. Indices increased sharply from 2013 to 2014 to the highest values of the time series for biomass and approaching the highest values of the time series for abundance (Fig. 3; Tables 6 and 10). Indices declined from 2014 to 2015 prior to increasing slightly from 2015 to 2016 (Fig. 3; Tables 6 and 10).

For NAFO Div. 30, where the majority of the stock resides, the indices from the spring surveys are more variable over the time series than 3N. Indices of stock size showed considerable annual fluctuations particularly in the 1980s and 1990's where biomass estimates ranged from about 24000 t in 1985 to about 2000 t in 1998 while abundance estimates ranged from about 43 million fish in 1994 to about 6 million fish in 1998 (Fig. 3; Tables 7 and 11). From 2003-2010, there was an overall decline in the spring indices for NAFO Div. 30. The biomass index in 2010 (5,000 t) is about half the long term mean and the 2010 abundance index (15 million) is about 25% below the long term mean (Fig. 3; Tables 7 and 11). From 2010 to 2013 indices of stock size have indicated a substantial upward trend to levels approaching the previous time series high for biomass and mean weight (kg) per tow in 1985 and 1988 and exceeding the previous time series high for abundance and mean number per tow in 1994 and 2003 (Figs. 3 and 4; Tables 11 and 15). In 2013 the highest abundance and mean number per tow estimates of the time series (about 66 million and 26 fish respectively) were encountered (Figs. 3 and 4; Tables 11 and 15). In 2013 the second highest estimates of the time series (in comparison to 1984-1985) of biomass (24,000 t) and mean weight per tow (7 kg) were encountered (Figs. 3 and 4; Tables 7 and 19). From 2013 to 2015 there were sharp decreases in indices of biomass to values near the time series mean and abundance to approximately 50% above the time series mean. Indices increased slightly from 2015 to 2016 (biomass of 3500 to 5000 t, mean weight per tow of 1.3 to 2.1 kg, abundance of 10 to 13 million, and mean number per tow of 4 to 5 (Figs. 3 and 4; Tables 15 and 19).

#### *Fall RV Surveys 3N and 30*

Due to operational difficulties there was no fall survey of NAFO Divs. 3NO in 2014. Stock indices means of biomass (3,000 t), mean weight per tow (1 kg), abundance (3 million fish), and mean number per tow (# fish) derived from the fall surveys in NAFO Div. 3N were consistently low from 1990 to 2007 (Figs. 5 and 6; Tables 8, 12, 16, and 20). From 2007 to 2009 there was an increase in stock indices of biomass (24000 t), mean weight per tow (5 kg), abundance (25 million fish), and mean number per tow (10 fish) to levels 5 to 10 times the mean of the previous time series. In 2010 and 2011, the stock indices in NAFO Div. 3N declined sharply, but were still about 2 times the 1990 to 2007 time series mean. In 2012 stock indices again increased to the highest levels of the time series which were comparable to the levels (5 to 10 times the time series mean) observed from 2007 to 2009. Stock indices in NAFO Div. 3N declined sharply from 2012 to 2016 (biomass 13000 to 1000 t, mean weight per tow 5.5 to 0.4 kg, abundance 25 to 2 million, and mean number per tow 10.5 to 0.8) (Fig.

5 and 6; Tables 8, 12, 16, and 20).

Stock indices for the fall surveys in NAFO Div. 30 from 1990 to 2007 exhibited a higher degree of variability over the time series in comparison to fall surveys in NAFO Div. 3N (Figs. 5 and 6; Tables 9, 13, 17, ands 21). Stock indices from 1990 to 2007 ranged about 2 times above and below the time series means for biomass (10000 t), abundance (23 million fish), mean number per tow (7 fish), and mean weight per tow (3 kg). Stock indices for the fall surveys in NAFO Div. 30 increased sharply from 2007 to 2009 to about 2 to 3 times the highest levels of the time series from previous years for biomass (25000 t), abundance (60 million fish), mean number per tow (24 fish), and mean weight per tow (7 kg). From 2010 to 2015 stock indices indicated a decreasing trend approaching the lowest levels of the 1990 to 2007 time series (Figs. 5 and 6; Tables 9, 13, 17, ands 21). From 2015 to 2016 indices increased marginally (biomass of 6000 to 6800 t, mean weight per tow of 2.5 to 2.7 kg, abundance of 16.2 to 16.5 million, and mean number per tow of 6.4 to 6.5) (Figs. 5 and 6; Tables 9, 13, 17, ands 21).

#### *3NO Combined*

For spring surveys in NAFO Divs. 3NO the stock indices trends are primarily driven by the significantly higher (approximately 10 times) overall numbers estimated for NAFO Div. 30. The NAFO Divs. 3NO-combined indices for spring show a slow decline in biomass and abundance from 1984 to the late-1990s (Fig. 7) and although fluctuations continue to occur, some minor improvement in the estimates had occurred from 1998 to 2003 until a decline in levels from 2003 to 2005. Values from 2007-2010 have fluctuated around the long-term mean, however from 2010 to 2013 estimates of both biomass (7000 to 24000 t) and abundance (20 to 70 million fish) increased substantially, with the time series highest values in 2013 peaking at about 3 times the long term mean (Fig. 7). This increase from 2010 to 2013 was followed by a sharp decline in both biomass and abundance from 2013 to 2015 to levels approaching the time series mean for biomass and to levels approximately 1.5 times the time series means for abundance. Spring survey indices for NAFO Divs 3NO increased slightly from 2015 to 2016 (biomass of 5000 to 7000 t, mean weight per tow of 1.0 to 1.5, abundance 13 to 17 million fish, and mean number per tow of 2.7 to 3.4) (Figs. 7 and 8)

Due to operational difficulties there was no fall survey of NAFO Divs. 3NO in 2014. The fall survey series for Divisions 3NO combined is less variable with a generally increasing trend in biomass and abundance from about 1997 until 2004 (Fig. 7). Variability increases substantially from 2006 to 2013. There was a decline in 2006 and 2007, and a large increase in the 2008-2010 estimates, to levels between 1.7 and 2.8 times the mean. This peak (the highest in the time series) is followed by a decrease in 2011, an increase in 2012 and a decrease in 2013. Although the values from 2008 to 2013 exhibit substantial variability the overall mean from this period is still 1.5 to 2 times greater than the mean for the time series (Fig 7). The fall survey indices for NASFO Divs. 3NO declined sharply from 2012 to 2016 to values approaching the lowest of the time series encountered in the 1990's (biomass of 27000 to 8000 t, mean weight per tow of 5.5 to 1.5 kg, abundance of 68 to 18 million fish, and mean number per tow of 14 to 4) (Figs. 7 and 8).

#### *Depth distribution*

Witch flounder have been described as a relatively deep water species, having been captured at depths of up to 1500

m. However, in the Newfoundland-Labrador area, they are thought to prefer depths of 184-366 m (Bowering and Brodie 1991). Because it was previously thought that witch flounder may not be adequately covered by

the survey depths, the issue was examined by analyzing the Canadian survey data (Dwyer 2008). It was concluded that the preferred depth of Divs. 3NO witch flounder differs by division and by time of year. A higher percentage of the biomass in 3N is found in deeper strata, but there is still a large percentage found in depths of less than 100m, especially in the fall. In Div. 30 where the main component of the stock is distributed, a large proportion of the biomass is found in depths less than 183 m in either spring or fall. This is despite the fact that in a number of years, the survey covered depths of up to 1500 m in the fall. The percent abundance by depth showed similar patterns.

Depths covered by the surveys have changed over the years as stated above. In the spring series, only 1994 was surveyed to 914 m, but only 1.4% of the Divs. 3NO biomass index value was found in these strata (Tables 6 and 7), although it was 17% for Div. 3N alone. For the fall surveys, in years and divisions where coverage was complete in depths 731 to 1462 m, between 15 and 25% of biomass estimates in Div. 3N were contained in these depths (Table 6). However, in Div. 30, there were very few fish found in this depth range, generally less than 5% (Table 7). Because Div. 30 contains the majority of the biomass estimate in the fall surveys (83% on average), the percentage of the total Div. 3NO biomass in the deeper strata is similarly low.

As discussed in Dwyer (2008), distribution plots indicated more witch flounder are distributed on the shallower, shelf area of the Grand Banks in some years, especially in Div. 30 and especially in the fall. Therefore, it seems likely that the RV survey coverage does adequately cover the depth distribution of witch flounder, particularly in the fall. The variation in the survey indices may be due to the movement of flounder onto and off of the shelf areas depending on water temperatures and spawning aggregations. Bowering and Orr (1996) suggested that the movement of witch flounder onto the shallow parts of the bank in large strata cause the high variability in annual stock size estimates. It is also likely that some witch flounder may be distributed outside the survey area, particularly in the spring, following spawning in deeper waters, and this may also contribute to variability in survey estimates.

A NAFO scientific council working paper prepared by Lee in 2015 examined the biomass distributions of witch flounder in NAFO Divs 3N and 30 in both spring and fall with respect to depth strata. The analysis indicated that for 3N, both spring and fall biomass proportions were fairly evenly distributed within depths ranging from 57 to 914 m. Although biomass proportions were low in both spring and fall in depths less than 56 m. For 30 spring and fall biomass proportions were highest in depths ranging from 57 to 183 m and uniformly low at depths ranging from 184 to 914 m. Both 3N and 30 fall distributions indicated low biomass proportions in depths ranging from 915 to 1463 m.

#### *Distribution Plots*

Geographic distributions of witch flounder from 1996-2007 spring and fall surveys (mean weight per tow) were plotted in Dwyer (2008), distributions from 2008 to 2012 were plotted in Lee et al. (2014). For recent years 2013-2016 (no fall survey in 2014), the spring and fall distribution plots are presented in Figures 9 and 10. The witch flounder stock for Div. 3NO is mainly distributed in Div. 30 along the southwestern slope of the Grand Bank. In most years the distribution is concentrated along this slope but in certain years (particularly in the fall) it is distributed in shallower parts of the bank in the larger strata. It is this variation in distribution from smaller to larger strata that is often responsible, in part, for the high variability in the annual biomass and abundance indices (Bowering and Orr 1996).

### *Length frequencies*

Canadian rv survey length frequency data for individual years from 1997 to 2016 are presented in Figure 11. Canadian and Spanish rv survey abundance at length is compiled across years and presented in Figure 12. Length frequencies from Canadian rv surveys are fairly flat and evenly distributed from 1997 to 2002, with few fish > 45 cm (Fig. 11). Length frequencies of 30-50 cm fish increase from 2002 to 2005, decrease to pre-2002 levels from 2006 to 2007, and are then consistently higher from 2008 to 2014 (note there was no survey data collected in the fall of 2014) with a mode generally within the mode of 40 cm (Fig. 11). The increase in 30-50 cm fish is generally more pronounced in the fall survey data as opposed to the flatter distributions of the spring surveys.

There have been a few identifiable peaks in the time series (presumably year classes) that could be followed in successive years (e.g. peak at 9 cm in 1997, 11 cm in 1998, and 20 cm in 1999; peak at 13 cm in 2011, 20 cm 2013), in 2002 a peak at 12 cm was not observed subsequently (Fig. 11). There have been less distinctive peaks, usually in the 10-20 cm range, observed in 2007, 2011, and 2015 although they were not identified in subsequent years (Fig. 11). It should be noted that no ageing information for this stock has been available from Canadian rv surveys since the mid 1990's, making the tracking of cohorts from length frequency data all but impossible given the relatively slow growth of witch flounder.

Abundance at length in the Canadian spring rv surveys appears to be fairly consistent since 2000 with few fish greater than 50 cm, and a mode generally around 38-40 cm (Fig. 12). However, since 2007 there has been an increase in the number of larger fish in the 40-45 cm range except for an anomalous 30-35 cm range encountered in 2014 (Fig. 12). Abundance at length in the Spanish spring rv surveys was fairly consistent at 33-35 cm from 2001 to 2007 (a smaller range than the Canadian surveys during the same time period). From 2008 to 2016 the size range has generally increased with more fish in the 38-40 cm range. In 2016 the mode was 42 cm which was higher than the rest of the time series (Fig. 12).

There were a small number of distinctive peaks in the 5-15 cm range (recruitment year classes) in both surveys that were evident and could be followed through successive years. This included the periods from 2007 to 2009 in the Canadian series and from 2002-2003, and 2005 -2006 in the Spanish series (Fig.12).

### *Recruitment*

Figure 13 shows the abundance index for fish less than 21 cm (a recruitment proxy) for NAFO Divs. 3NO combined, as measured in the spring and fall Canadian rv surveys. Highest spring levels were in 1997, highest fall levels were in 1998 and 1999. Values since 2002 for the fall have been consistently below the mean of the time series. Spring values were occasionally above the time series mean in 2003, 2005, 2009, and 2013. Recruitment in spring and fall surveys in 2016 approached the lowest values of the time series (Fig. 13).

The distributions of juvenile (< 21 cm) witch flounder over the spring and fall Canadian surveys indicate a marginal pattern of fish being more widely distributed over the shallower depths in the larger strata during the fall (Figs. 9 and 10). It is also possible that the weak pattern may be related to the distributions previously presented for the entire population which indicated a movement of fish to the shallower, larger strata during the fall.

### Spanish Div. 3NO surveys

Since 1995, Spain has carried out a stratified random spring bottom trawl survey in Div. 3NO of the NAFO Regulatory Area. In 2001, the trawl vessel (*C/V Playa de Menduiña*) and gear (*Pedreira*) were replaced by the R/V *Vizconde de Eza* using a Campelen trawl. Biomass estimates from 1995 to 2016 are presented in Table 22. Mean weight and numbers per tow series from 2002 are provided in Figure 14 and biomass estimates are provided in Figure 15 for the latter series, as data prior to 2000 have not yet been converted. Index values peaked in 2004 and 2010, declined in 2011, increased in 2012 and then declined sharply from 2012 to 2014 to the lowest values reported over the time series. Indices have steadily increased from 2014 to 2016.

Length frequency comparisons between Canadian and Spanish spring research vessel surveys from 2013 to 2016 are presented in Figure 16. In the years 2013, 2015, and 2016 there is a high degree of overlap between Canadian and Spanish length frequencies in each survey year. Length frequencies range from approximately 25-50 cm with modal frequencies around 40 cm and low numbers of fish greater than 50 cm or below 25 cm. Although Canadian length frequencies were relatively stable in 2014, Spanish length frequencies shifted slightly to a larger range. In 2014 Spanish length frequencies ranged from about 20 cm to 47 cm with a modal frequency of approximately 33 cm (Figure 16).

### Surplus production model

A surplus production model in a Bayesian framework was used for the assessment of this stock. This model was developed and accepted by the Scientific Council in 2015 (Morgan et al 2015). The updated surplus production model used in this assessment is fully described and diagnostics provided in Morgan and Lee (2017). The input data were catch from 1960-2016, Canadian spring survey series from 1984-1990, Canadian spring survey series from 1991-2016 (no 2006) and the Canadian autumn survey series from 1990-2016 (no 2014).

The priors used in the model were:

Initial population size	$\text{Pin} \sim \text{dunif}(0.5, 1)$	uniform(0.5 to 1)
Intrinsic rate of natural increase	$r \sim \text{dlnorm}(-1.763, 3.252)$	lognormal (mean, precision)
Carrying capacity	$K \sim \text{dlnorm}(4.562, 11.6)$	lognormal (mean, precision)
Survey catchability	$pq \sim \text{dgamma}(1, 1)$ $q = 1/pq$	gamma(shape, rate)
Process error	$\sigma \sim \text{dunif}(0, 10)$ $\text{isigma2} = \text{pow}(\sigma, -2)$	uniform(0 to 10)
Observation error	$\tau \sim \text{dgamma}(1, 1)$ $\text{itaue2} = 1/\tau$	gamma(shape, rate)

## Resource Status

Recruitment (defined as fish less than 21cm) in both the spring and fall Canadian surveys although somewhat variable has generally been low since 2003. Recruitment in spring and fall surveys in 2016 approached the lowest of the time series.

The surplus production model results indicate that stock size decreased from the late 1960s to the late 1990s and then increased from 1999 to 2010 but has since decreased (Fig. 17). The model suggests that a maximum sustainable yield (MSY) of 3 641 (2 263-5 689) t can be produced by total stock biomass of 50 000 (35 559-69 581) t (B<sub>msy</sub>) at a fishing mortality rate (F<sub>msy</sub>) of 0.07 (0.04-0.13) (Fig. 18). The intrinsic rate of natural increase is estimated to be 0.14 and carrying capacity 100 000 t.

The analysis showed that relative population size (median B/B<sub>msy</sub>) was near  $B_{lim}=30\% BMSY$  from 1993-1998. Biomass has since increased to a level of 52% BMSY in 2016 (Fig. 17). The probability of being below  $B_{lim}$  in 2016 is 0.15. Relative fishing mortality rate (median F/F<sub>msy</sub>) was mostly above 1.0 from the late 1960s to the mid-1990s (Fig. 18). F has been below F<sub>msy</sub> since the moratorium implemented in 1995. Median F was estimated to be 59% of F<sub>msy</sub> with a probability of 0.19 of being above F<sub>msy</sub> in 2016.

The posterior distributions (13500 samples) for r, K, sigma, and biomass and the production model equation were used to project the population to 2020. All projections assumed that the catch in 2017 was equal to the TAC of 2 225 t (which produces F2017). This was followed by constant fishing mortality for 2018 and 2019 at several levels of F (F2016, 75% F2016, 125% F2016, 2/3 FMSY, 75% FMSY, and 85% FMSY).

The probability that F >  $F_{lim}$  in 2017 is 57% at a catch of 2 225 t (Tables 23 and 24). The probability of F> $F_{lim}$  ranged from 15 to 42% for the catch scenarios tested. The population is projected to grow under all scenarios and the probability that the biomass in 2020 is greater than the biomass in 2016 is greater than 50% in all scenarios. The population is projected to remain below BMSY for all levels of F examined with a probability of greater than 70%. The probability of projected biomass being below  $B_{lim}$  was about 20% in all catch scenarios examined and was 14% by 2020 in the F=0 scenario (Tables 23 and 24).

The model used is best at forecasting trends in stock development and less precise in predicting year-to-year changes. Although the stock is estimated to be above  $B_{lim}$ , recent declining trends in survey indices and low recruitment will be monitored in future years. Uncertainty around parameter estimates has increased compared to the 2015 assessment.

Because of the uncertainty of the estimates and proximity to the limit reference points, the next full assessment of this stock is re-scheduled for 2018.

## References

- Bowering, W. R. and D. Orr. 1996. Distribution and trends in stock size of witch flounder in NAFO Divisions 3NO. NAFO SCR Doc. 96/70.
- Brodie, W., Parsons, D., Murphy, E., and Dwyer, K. 2011. An assessment of the witch flounder resource in NAFO Divisions 3NO. NAFO SCR Doc. 11/029.
- Brodie, W. and Stansbury, D. 2007. A brief description of Canadian Multispecies surveys in SA2 + Divisions 3KLMNO from 1995-2006. NAFO SCR Doc. 07/18.
- Dwyer, K. 2008. An assessment of witch flounder in NAFO Divisions 3NO. NAFO SCR Doc. 08/39., Ser. No. N5540.
- Fomin, K. and Pochtar M. 2017. Russian research report for 2016. NAFO SCS Doc. 17/11, Serial No. N6686.
- Fomin, K. , Khlevnov, V., Mishin T., and Zavoloka P. 2015. Russian research report for 2014. NAFO SCS Doc. 15/07, Serial No. N6433.
- Gonzalez-Costas, F., Ramilo, G., Roman, E., Gago, A. Sacau, M. Gonzalez-Troncoso, D., Casas, M., and Lorenzo, J. 2017. Spanish Research Report for 2016. NAFO SCS Doc. 17/04, Ser. No. N6656.
- Gonzalez-Costas, F., Ramilo, G., Roman, E., Gonzalez-Troncoso, D., Casas, M., Sacua, E., and Lorenzo, J. 2015. Spanish Research Report for 2014. NAFO SCS Doc. 15/05, Ser. No. N6423.
- Lee E., Regular, P., Brodie B., Rideout R. M., Dwyer K., Ings D., and Morgan J. 2014. An assessment of the witch flounder resource in NAFO Divisions 3NO. NAFO SCR Doc.14/029, Ser. No. N6325.
- Morgan, J. Lee, E. 2017. Surplus production model in a Bayesian framework applied to witch flounder in NAFO Divs. 3NO. NAFO SCR 17/047, Ser. No. N6707
- Morgan, M.J., C. Hvingel and M. Koen-Alonso. 2015. Surplus production models in a Bayesian framework applied to witch flounder in NAFO Div. 3NO. NAFO SCR 15/37.
- NAFO, 2013. Report of Scientific Council Meeting, 7-20 June 2013.
- NAFO, 2012. Canadian Request for Scientific Advice on management in 2013 of certain stocks in subareas 0 to 4. NAFO SCS Doc. 12/04., Ser. No. N6014.
- Power, D. and Richards, D. 2017. Canadian research report for 2016 Newfoundland and Labrador Region 2016. NAFO SCS Doc. 17/13, Ser. No. N6704
- Torra, T., Sirps. 2015. Estonian Research Report for 2015. NAFO SCS Doc. 15/04, Ser. No. N6420
- Vargas, J., R. Alpoim, E. Santos and A. M. Ávila de Melo. 2015. Portuguese Research Report for 2014. NAFO SCS Doc. 15/06, Ser. No. N6426.
- Vargas, J., R. Alpoim, E. Santos and A. M. Ávila de Melo. 2017. Portuguese Research Report for 2016. NAFO SCS Doc. 17/05, Ser. No. N6658.

Table 1. Catches and TACs (t) of Witch flounder in Div. 3NO from 1960 to 2016.

Year	Canada	USSR (Russia)	Other	Total	TAC
1959	-	-	-	5799	0
1960	-	-	-	4627	0
1961	-	-	-	1228	0
1962	-	-	-	2183	0
1963	895	485	803	1066	0
1964	1055	-	11	2177	0
1965	1324	849	4	7522	0
1966	3644	3828	50	11503	0
1967	2863	8565	75	10599	0
1968	1503	9078	18	4700	0
1969	479	4215	6	6763	0
1970	723	6039	1	14965	0
1971	178	14774	13	9177	0
1972	3419	5738	20	6691	0
1973	4943	1714	34	8045	10000
1974	2807	5235	3	6168	10000
1975	1137	5019	12	6035	10000
1976	3044	2991	-	5759	10000
1977	3013	2742	4	3616	10000
1978	1165	2275	33	3077	7000
1979	1193	1868	16	2420	7000
1980	425	1994	1	2425	5000
1981	381	2044	-	3732	5000
1982	1760	1969	3	3616	5000
1983	1674	1942	-	2802	5000
1984	834	1955	13	4117	5000
1985	2746	1908	4117	8771	5000
1986	2937	1724	4470	9131	5000
1987	2829	1425	3342	7596	5000
1988	1927	1037	4361	7325	5000
1989	1241	81	2366	3688	5000
1990	2654	9	1516	4179	5000
1991	2624	-	2223	4847	5000
1992	4328	-	632	4960	5000
1993	4337	3	250	4414	5000
1994	2	-	1117	1119	3000
1995	-	-	300	300	0
1996	64	-	294	358	0
1997	19	-	493	512	0
1998	2	5	605	612	0
1999	6	86	671	763	0
2000	12	50	483	545	0
2001	13	34	647	694	0
2002	26	112	312	450	0
2003*	62	59	1423	1544	0
2004	58	60	509	627	0
2005	49	8	200	257	0
2006	94	2	385	481	0
2007	21	27	174	222	0
2008	46	17	201	264	0
2009	41	22	313	376	0
2010	39	28	354	421	0
2011	11	2	337	350	0
2012	2	10	303	315	0
2013	62	54	212	328	0
2014	11	57	267	335	0
2015	221	36	102	359	1000
2016**	799	26	237	1062	2172
					2225

Although a TAC of 3000 tons was agreed by the Fisheries Commission (FC), it was also agreed that no directed fishing on witch flounder in NAFO Divs. 3NO take place during 1994 due to the poor state of the stock.

\*Estimated catch in 2003 is the mean of a range of catch from several sources.

3NO Witch flounder was under moratorium from 1995 to 2014

\*\*Non-Canadian Catch for 2016 was derived by the NAFO Catch Data Advisory Group (CDAG) methodology (NAFO scwp17-021 and NAFO scwp 17-022). A 2,225 t quota for witch flounder in 3NO was adopted by the FC for 2017.



Table 2. Canadian spring research vessel surveys in NAFO Division 3N, 1984-2016.

Depth Range (m)	Old Stratum Area	New Stratum Area	Stratum	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
				84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
<=56	1593 1499	1593 1499	375 376																																	
57-92	2992 1853 2520 2520 931 674	2992 1853 2520 2520 931 674	360 361 362 373 374 383																																	
93-183	421 100 647	421 100 647	359 377 382																																	
184-274	225 139 182	225 139 182	358 378 381																																	
275-366	164 106 116	164 106 116	357 379 380																																	
367-549	155 105 160	155 105 160	723 725 727																																	
550-731	124 72 156	124 72 156	724 726 728																																	
732-914		134 106 154	752 756 760																																	

Note dark grey indicates that a stratum was sampled. White indicates that a stratum was not sampled.

Table 3. Canadian spring research vessel surveys in NAFO Division 30, 1984-2016.

Depth Range (m)	Old Stratum Area	New Stratum Area	Stratum	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
57-92	2089	2089	330																																	
	456	456	331																																	
	1898	1898	338																																	
	1716	1716	340																																	
	2520	2520	351																																	
	2580	2580	352																																	
	1282	1282	353																																	
93-183	1721	1721	329																																	
	1047	1047	332																																	
	948	948	337																																	
	585	585	339																																	
	474	474	354																																	
184-274	151	147	333																																	
	121	121	336																																	
	103	103	355																																	
275-366	92	92	334																																	
	58	58	335																																	
	61	61	356																																	
367-549	93	166	717																																	
	76	76	719																																	
	76	76	721																																	
550-731	111	134	718																																	
	105	105	720																																	
	93	93	722																																	
732-914	.	105	764																																	
	.	135	772																																	

Note dark grey indicates that a stratum was sampled. White indicates that a stratum was not sampled.

Table 4. Canadian fall research vessel surveys in NAFO Division 3N, 1990-2016.

Depth Range (m)	Old Stratum Area	New Stratum Area	Stratum	90	91	92	93	94	95	96	97	98	99	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
<=56	1593 1499	1593 1499	375 376																											
57-92	2992 1853 2520 2520 931 674	2992 1853 2520 2520 931 674	360 361 362 373 374 383																											
93-183	421 100 647	421 100 647	359 377 382																											
184-274	225 139 182	225 139 182	358 378 381																											
275-366	164 106 116	164 106 116	357 379 380																											
367-549	155 105 160	155 105 160	723 725 727																											
550-731	124 72 156	124 72 156	724 726 728																											
732-914	.	134 106 154	752 756 760																											
915-1097	.	138 102 171	753 757 761																											
1098-1280	.	180 99 212	754 758 762																											
1281-1463	.	385 127 261	755 759 763																											

Note dark grey indicates that a stratum was sampled. White indicates that a stratum was not sampled.

Table 5. Canadian fall research vessel surveys in NAFO Division 30, 1990-2016.

Depth Range (m)	Old Stratum Area	New Stratum Area	Stratum	90	91	92	93	94	95	96	97	98	99	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
57-92	2089	2089	330																											
	456	456	331																											
	4898	4898	338																											
	1716	1716	340																											
	2520	2520	351																											
	2580	2580	352																											
	1282	1282	353																											
93-183	1721	1721	329																											
	1047	1047	332																											
	948	948	337																											
	585	585	339																											
	474	474	354																											
184-274	151	151	333																											
	121	121	336																											
	103	103	355																											
275-366	92	96	334																											
	58	58	335																											
	61	61	356																											
367-549	93	166	717																											
	76	76	719																											
	76	76	721																											
550-731	111	134	718																											
	105	105	720																											
	93	93	722																											
732-914	.	105	764																											
	.	99	768																											
	.	135	772																											
915-1097	.	124	765																											
	.	138	769																											
	.	128	773																											
1098-1280	.	144	766																											
	.	128	770																											
	.	135	774																											
1281-1463	.	158	767																											
	.	175	771																											
	.	155	775																											

Note dark grey indicates that a stratum was sampled. White indicates that a stratum was not sampled.



Table 6. Estimated biomass (tonnes) of Witch flounder (M+F) in each stratum from surveys in Div. 3N during spring of 1984-2016. (Engel 145 data converted to Campelen Units for 1984-95).

Depth Range (m)	Old Stratum Area	New Stratum Area	Stratum	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16
<=56	1593	1593	375	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	41	35	0	21	0	0	0	0	0			
	1499	1499	376	0	0	0	19	0	0	0	0	0	0	0	0	0	8	18	0	0	0	0	0	0	0	89	0	0	0	0	0	0				
57 - 92	2992	2992	360	1715	89	629	461	1519	175	0	0	29	165	0	0	0	115	33	120	266	0	0	19	97	983	264	543	85	0	395	156	72	188	135	0	0
	1853	1853	361	119	0	0	39	50	0	20	0	0	0	0	39	0	0	0	0	242	45	0	0	0	35	139	0	18	72	0	131	0	92	75	0	0
	2520	2520	362	0	82	23	18	147	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	133	0	0	0	0	0	0		
	2520	2520	373	0	0	43	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
	931	931	374	0	0	0	0	0	0	0	0	0	18	34	0	0	0	0	0	0	30	0	0	0	0	0	0	0	0	0	0	0	0	0		
	674	674	383	0	57	0	37	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
93 - 183	421	421	359	231	47	99	43	306	121	0	0	0	19	0	0	0	0	67	149	58	13	0	0	0	334	52	0	593	719	1365	299	83	835	612	117	
	100	100	377	8	0	0	72	3	32	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	38	0	0	0	0	0	0
	647	647	382	0	0	0	12	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	42	0	
184 - 274	225	225	358	40	308	42	137	20	29	57	0	44	132	106	7	51	49	134	6	9	154	14	168	0	42	316	68	237	156	241	86	189	135	24	884	
	139	139	378	22	19	32	155	31	42	0	0	29	0	0	0	0	3	0	0	0	5	8	1	0	0	0	0	0	0	0	14	55	0	0	6	0
	182	182	381	21	7	32	101	69	0	28	0	0	0	0	0	0	0	0	0	0	7	0	0	0	0	53	13	18	0	0	30	0	23	267	0	
275 - 366	164	164	357	8	87	154	4	60	21	0	31	49	81	20	36	12	159	21	75	17	26	65	42	0	19	0	4	31	83	134	25	42	94	56		
	106	106	379	36	12	23	173	44	20	35	3	18	0	4	0	0	9	2	26	4	4	0	4	0	6	0	0	7	12	23	101	88	237	5	0	
	116	116	380	6	53	0	134	24	7	4	0	0	0	0	0	0	0	6	0	0	0	3	0	0	0	5	0	0	0	22	5	12	4	0		
367 - 549	155	155	723																																	
	105	105	725																																	
	160	160	727																																	
550 - 731	124	124	724																																	
	72	72	726																																	
	156	156	728																																	
732 - 914	.	134	752																																	
	.	106	756																																	
	.	154	760																																	
Grand Total				2205	761	1078	1401	2218	485	164	655	484	862	510	308	170	443	566	525	1042	632	380	532	346	1807	577	1442	502	1936	1818	2395	1135	1188	2489	1470	1922
Biomass >366 m				0	0	0	0	0	0	0	652	333	480	284	242	84	255	230	262	296	343	289	272	207	366	0	335	313	984	505	328	278	517	953	458	865
Percent >366 m				0.0	0.0	0.0	0.0	0.0	0.0	0.0	99.5	68.8	55.7	55.7	78.6	49.2	57.6	40.6	49.9	28.4	54.2	76.0	51.0	59.9	20.3	0.0	23.2	62.4	50.8	27.8	13.7	24.5	43.5	38.3	31.2	45.0



Table 7. Estimated biomass (tonnes) of Witch flounder (M+F) in each stratum from surveys in Div. 30 during spring of 1984-2016. (Engel 145 data converted to Campelen Units for 1984-95).

Depth Range (m)	Old Stratum Area	New Stratum Area	Stratum	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16		
57 - 92	2089	2089	330	0	0	0	0	22	0	0	0	0	0	0	0	0	0	21	121	111	0	0	0	117	129	569	0	278	0	0	875	55	36	294	0	0		
	456	456	331	1912	302	36	18	444	0	0	0	0	0	0	0	74	0	36	537	28	375	102	0	0	292	1301	425	1124	17	212	81	10	352	20	0			
	1898	1898	338	134	7806	1108	1184	3075	1827	434	0	109	295	0	228	870	0	357	780	183	1354	121	320	1171	646	1675	1016	450	990	769	948	2569	2641	455	804	119		
	1716	1716	340	40	146	0	21	0	0	15	0	147	0	0	0	0	0	0	0	83	0	0	0	0	26	90	0	0	182	0	0	0	4	45	0	0		
	2520	2520	351	688	211	385	222	978	217	109	0	0	0	0	0	0	0	21	22	0	0	0	0	0	0	0	65	0	0	21	0	0	0	0	0			
	2580	2580	352	82	951	225	1275	1330	664	1426	40	105	60	40	63	59	100	53	1196	130	53	693	27	628	551	1199	733	555	102	562	791	1754	298	85	30	0		
	1282	1282	353	4519	1122	1067	1609	7208	2486	1637	0	243	209	0	42	23	2	272	2209	1300	469	688	470	572	430	3390	576	529	172	299	1078	2982	1265	1264	413	0		
93 - 183	1721	1721	329	0	0	0	0	789	48	27	494	0	0	5071	193	0	11	51	240	26	0	0	2209	0	147	559	215	983	559	752	1117	7541	66	495	0			
	1047	1047	332	3779	8589	2485	3367	6829	1485	4599	2426	2182	359	58	1791	1180	235	460	981	407	3025	2458	10236	7945	1075	641	3188	2005	1669	1270	911	9766	4888	629	2120			
	948	948	337	50	4129	1415	1506	1061	1543	1627	1581	580	675	50	654	330	163	321	879	936	1823	752	715	233	655	333	1211	563	630	198	1958	1007	140	453	1704			
	585	585	339	335	0	16	223	136	0	0	0	0	0	0	1	0	0	1	0	5	2	0	0	189	825	4	37	284	2	58	0	14	56	0	0			
	474	474	354	495	105	1231	233	345	47	240	144	149	841	0	0	36	0	226	1062	826	914	553	163	496	640	393	1148	430	147	968	164	378	429	478	56			
184 - 274	151	147	333	10	48	10	0	67	16	129	498	79	80	5196	162	7	109	25	27	30	122	375	63	36	39	27	9	32	20	6	9	42	0	2	155			
	121	121	336	12	7	43	25	63	0	53	492	1374	100	1057	62	180	293	23	47	27	163	598	211	61	51	44	61	16	16	26	10	38	18	15	74			
	103	103	355	45	181	38	71	0	97	126	136	16	34	129	43	86	48	50	18	14	87	193	340	117	12	27	34	67	44	12	26	14	3	24	797			
275 - 366	92	96	334	0	42	42	18	22	23	26	20	108	20	860	15	150	362	4	7	11	2	143	133	29	3	11	5	14	6	6	1	10	4	2	92			
	58	58	335	0	98	18	2	51	22	92	42	1107	65	103	43	78	109	2	62	128	8	8	53	10	11	2	1	4	3	3	3	17	12	8	0	3		
	61	61	356	5	83	17	23	18	29	55	39	129	77	75	62	40	11	29	23	14	34	38	49	13	18	3	6	6	5	0	4	29	2	9	73			
367 - 549	93	166	717																																			
	76	76	719																																			
	76	76	721																																			
550 - 731	111	134	718																																			
	105	105	720																																			
	93	93	722																																			
732 - 914	.	105	764																																			
	.	135	772																																			
Grand Total				12108	23820	8136	9799	22438	8503	10594	6415	7734	3364	15769	3748	3915	6691	2121	8411	4448	8786	7182	15323	11479	5057	7747	5746	8323	7243	4821	7349	11727	23208	8212	3457	5211		
Biomass >366 m				0	0	0	0	0	0	0	504	1405	550	3128	390	800	5247	192	201	172	354	459	336	51	144	0	75	107	268	75	124	70	104	102	82	18		
Percent >366 m											7.9	18.2	16.4	19.8	10.4	20.4	78.4	9.1	2.4	3.9	4.0	6.4	2.2	0.4	2.9	0.0	1.3	1.3	3.7	1.6	1.7	0.6	0.4	1.3	2.4	0.4		



Table 8. Estimated biomass (tonnes) of Witch flounder (M+F) in each stratum from surveys in Div. 3N during fall of 1990-2016 (Engel 145 data converted to Campelen units for 1990-94).

Depth Range (m)	Old Stratum Area	New Stratum Area	Stratum	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	
<=56	1593	1593	375	0	73	0	0	0	0	0	0	0	0	0	0	35	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	1499	1499	376	0	0	0	0	0	14	0	22	0	0	0	0	38	28	0	0	0	0	0	67	0	0	59	202	23	0		
57 - 92	2992	2992	360	265	171	1297	173	75	888	23	427	431	177	535	326	520	586	836	2364	100	0	4788	10335	1627	1311	11991	7294	736	566		
	1853	1853	361	28	467	463	0	32	0	0	14	0	268	28	170	148	99	0	168	38	584	25	0	410	190	188	78	0	28		
	2520	2520	362	400	221	87	0	0	0	0	0	0	32	0	0	0	136	0	0	40	0	0	46	192	55	70	90	0	31		
	2520	2520	373	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
	931	931	374	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
	674	674	383	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	25	0	0	0	0	0	27	23	0	
93 - 183	421	421	359	0	0	278	0	0	22	0	0	1213	1	0	121	42	110	139	43	151	192	442	1080	288	398	190	156	523	42		
	100	100	377	0	0	0	8	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	39	31	10	94	0	0	0	0	
	647	647	382	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
184 - 274	225	225	358	0	20	66	24	0	74	0	11	30	19	40	45	0	145	22	107	144	28	141	86	83	104	374	98	0	28		
	139	139	378	0	41	15	0	0	0	0	1	0	0	0	3	5	0	0	0	93	4	7	4	22	56	191	446	24	0		
	182	182	381	0	0	0	0	0	1	0	0	0	7	0	0	0	0	0	0	0	0	0	3	0	0	0	0	0	0		
275 - 366	164	164	357	0	234	9	187	43	85	0	27	0	52	18	21	41	27	37	103	59	90	17	39	5	93	31	166	7			
	106	106	379	4	4	0	0	0	1	7	0	0	2	111	33	8	867	0	3	0	156	13	29	662	18	4	40	6	0		
	116	116	380	0	0	0	0	0	0	1	2	5	0	0	0	9	11	0	0	0	0	0	0	0	0	0	0	12	0		
367 - 549	155	155	723	41	163	180	57	15	28	74	27	28	66	16	123	20	98	38	17	98	93	27	62	37	38	1278	4	0			
	105	105	725	15	376	46	19	0	135	10	33	19	7	5	10	7	7	11	21	40	12	12	71	83	17	0	0	0	0		
	160	160	727	0	38	0	0	29	7	4	0	10	0	0	7	21	0	0	143	82	21	22	32	17	70	0	0	0			
550 - 731	124	124	724	172	414	180	104	60	197	72	181	87	70	90	70	95	206	127	455	204	117	143	72	79	10	40	0	0			
	72	72	726	310	54	48	40	21	38	34	16	22	59	52	32	19	49	45	42	105	6	17	23	4	57	53	0	0	0		
	156	156	728	153	35	21	76	78	106	153	103	286	178	93	19	122	191	269	404	434	51	125	213	108	145	0	0	0	0		
732 - 914	.	134	752								120		23	0	1								6								
	.	106	756								124		51	83	9								67								
	.	154	760								88		41	78	173								221								
915 - 1097	.	138	753								0		0	0	3								0								
	.	102	757								0		0	37	7								0								
	.	171	761								46		147	42	10								118	7		102					
1098 - 1280	.	180	754								0		0	0	0								0								
	.	99	758								0		0	0	0								0								
	.	212	762								0		109	0									15	28		40					
1281 - 1463	.	385	755								0		0	0	0								0								
	.	127	759								0		0	2	0								0								
	.	261	763								19		5	10									0	0		3					
Grand Total				696	1441	2235	1647	808	1346	160	993	2333	884	1244	1435	1511	1516	2122	3221	1093	1475	6703	12986	3306	3064	13432	8590	3575	991		
Biomass >366 m				0	213	15	1263	651	263	137	485	657	385	582	634	669	363	222	491	423	609	1029	1294	606	307	300	420	1606	259		
Percent >366 m				0.0	14.8	0.7	76.7	80.5	19.5	85.6	48.8	28.2	43.5	46.8	44.2	44.3	23.9	10.5	15.2	38.7	41.3	15.4	10.0	18.3	10.0	11.4	4.9	44.9	26.1		



Table 9. Estimated biomass (tonness) of witch flounder (M+F) in each stratum from surveys in Div. 3O during fall of 1990-2016 (Engel 145 data converted to Campelen units for 1990-94.

Depth Range (m)	Old Stratum Area	New Stratum Area	Stratum	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16
57 - 92	2089	2089	330	122	67	79	0	0	247	0	72	168	208	48	284	342	438	74	312	383	362	508	1087	344	708	49	837	984	431	
	456	456	331	22	315	134	0	0	108	0	0	256	946	243	468	775	306	14	394	108	144	114	564	1219	793	75	688	83.4	48	
	1898	1898	338	2226	438	837	3966	2193	4684	503	1329	483	2736	375	943	976	2666	3899	1931	604	543	1407	2044	5483	2554	643	1222	884	231	
	1716	1716	340	173	280	63	0	0	204	0	22	0	415	104	172	123	57	28	116	654	1	494	116	81	142	575	959	132	154	
	2520	2520	351	1690	284	72	0	0	0	0	37	205	0	172	0	25	35	54	369	158	165	28	75	65	234	0	33.7	89		
	2580	2580	352	1415	896	1352	946	228	379	273	573	374	1491	920	430	789	964	3377	1663	1109	558	1409	5915	2305	2597	1335	1635	476	63	
	1282	1282	353	2405	343	477	0	732	538	789	168	1066	2996	2379	1360	1490	1204	2657	3710	1587	1121	1431	8037	8234	3098	4323	1446	1204	3689	
93 - 183	1721	1721	329	99	85	0	18	0	417	0	173	305	0	0	282	732	97	484	250	2974	0	4484	1977	171	1616	1518	1096	0	465	
	1047	1047	332	2102	155	1724	813	321	1114	4569	190	245	1664	544	343	1155	807	1512	2061	3887	708	2453	500	1393	284	3372	283	485	963	
	948	948	337	1333	188	954	563	2132	421	492	322	479	978	344	67	211	352	114	1721	190	576	1592	352	989	158	328	150	222	100	
	585	585	339	1132	224	651	119	742	1911	0	481	261	344	338	1927	457	3755	1854	1070	1060	1147	2405	2693	2359	882	320	1273	489		
	474	474	354	1291	23	316	75	210	191	4647	215	201	103	766	258	470	967	438	316	505	694	306	1320	544	312	78	294	531	65	
184 - 274	151	147	333	221	11	22	30	92	26	4	6	33	4	20	17	48	0	3	24	3	2	5	6	14	0	3	1.31	6		
	121	121	336	82	151	76	298	13	35	32	19	19	67	31	37	23	10	5	35	3	53	142	22	18	8	13	17.5			
	103	103	355	497	93	120	25	16	343	6	14	110	35	5	6	6	21	2	5	17	72	23	20	15	41	3	1.86	8		
275 - 366	92	96	334	24	16	0	9	17	4	5	1	7	5	14	9	8	0	16	0	0	0	10	2	4	4	8	0.2	12		
	58	58	335	194	25	25	30	18	1	23	0	1	23	8	3	9	1	5	3	3	1	6	0	0	0	7	0	0.82	1	
	61	61	356	11	7	430	98	7	60	3	4	32	22	7	3	6	2	7	0	0	0	10	1	8	4	3	0	0		
367 - 549	93	166	717	30	0	32	37	12	42	260	0	13	11	54	9	2	14	9	102	40	14	37	52	59	17	8				
	76	76	719	110	2	65	6	1	226	19	9	10	14	29	6	15	3	6	10	4	8	16	4	8	0	12	6.9	14		
	76	76	721	18	169	67	21	54	6	14	67	17	2	14	17	2	15	3	30	11	1	7	8	13	2	2.99	1			
550 - 731	111	134	718		22	68	8	68	47	53	34	50	54	161	48	130		68	162	80	110	63	50	11	95.1	23				
	105	105	720		73	0	13	68	2	17	4	83	26	31	10	39	1	1	12	1	4	10	0	20		63.2	17			
	93	93	722	9	81	21	14	39	12	12	26	8	15	5	7	14	29	8	9	11	15	11	4	8	13	11.3	1			
732 - 914	.	105	764							75		12	21	36			4		11		41									
	.	99	768							18		7	18	38			4		1		5									
	.	135	772							173		62	49	29			50		22		26									
915 - 1097	.	124	765							24		3	20	55			10		11		25									
	.	138	769							17		5	28	59			20		16		26									
	.	128	773							4		13	32	89	12			8		10		5								
1098 - 1280	.	144	766							24		2	37			57		24		29										
	.	128	770							4		23	67			13		16		2										
	.	135	774							4		31	15	27			43		4		0									
1281 - 1463	.	158	767							15		0	0			0		3		2										
	.	175	771							0		17	0			10		0		0										
	.	155	775							0		0	0	28			21		3		13									
<b>Grand Total</b>				14671	4036	6884	7827	7013	10397	12117	3698	4356	12446	6396	5586	9619	8798	16510	14911	13512	6240	16036	24721	23733	14876	13601	9077		6526	6878
<b>Biomass &gt;366 m</b>				140	29	0	410	193	95	386	116	436	433	224	384	562	381	87	460	35	241	306	328	150	131	124	116		196	64
<b>Percent &gt;366 m</b>				1.0	0.7	0.0	5.2	2.8	0.9	3.2	3.1	10.0	3.5	3.5	6.9	5.8	4.3	0.5	3.1	0.3	3.9	1.9	1.3	0.6	0.9	1.3		3.0	0.9	



Table 10. Abundance (000s) of witch flounder (M+F) in each stratum from surveys in Div. 3N during spring of 1984-2016 (Engel 145 data converted to Campelen units for 1984-95).

Depth Range (m)	Old Stratum Area	New Stratum Area	Stratum	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16					
<=56	1593 1499	1593 1499	375 376	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	73 88	44 0	0 0	0 0	0 0	0 0	0 0	0 0									
57 - 92	2992 1853 2520 2520 931 674	2992 1853 361 2520 373 931 383	360 153 0 0 0 62	2234 129 25 0 0 31	728 32 27 0 0 0	741 36 173 0 0 0	2641 36 0 0 0	220 0 0 0 0 0	0 0 0 0 0	132 65 0 0 0	224 224 0 0 0	613 212 0 0 0	0 85 0 0 0	82 212 0 0 0	123 36 0 0 0	1555 255 0 0 0	480 173 0 0 0	741 0 0 0 0	103 51 0 0 0	0 85 0 0 0	823 203 0 0 0	288 170 64 0 0	165 0 0 0 0	329 170 64 0 0	206 0 0 0 0	0 0 0 0 0															
93 - 183	421 100 647	421 100 647	359 377 382	405 14 0	58 0 0	232 186 30	58 7 0	985 83 0	203 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	0 0 0	695 0 0	87 0 0	0 0 0	1448 1953 3475	115 115 115	1371 1158 174																	
184 - 274	225 139 182	225 139 381	358 378 25	77 48 13	557 48 42	93 86 163	31 75	46 0 0	93 0 0	0 96 0	93 0 0	294 0 0	232 0 0	31 0	77 0	83 0	261 0	15 0	41 0	325 11 0	28 0 0	296 0 0	0 0	110 0 0	681 0 0	151 0 0	542 0 0	303 25 33	566 33 0	186 22 51	330 38 38	230 38 438	50 50 50	1593 17 0							
275 - 366	164 106 116	164 106 380	357 379 8	23 66 88	180 36 0	553 68 247	11 102 32	237 44 8	56 0 0	0 22 0	90 0 0	124 109 0	102 44 0	23 0 0	40 22 0	30 18 0	373 6 0	259 102 0	293 28 0	63 13 0	55 0 0	150 16 0	45 0 0	0 0 0	23 0 0	0 7 0	29 29 49	98 284 0	361 192 515	317 15 7	45 172 0	64 192 0	180 17 0								
367 - 549	155 105 160	155 105 160	723 725 727	1 1 0	1 1 0	1 1 0	1 1 0	1 1 0	1 1 0	1 1 0	1 1 0	1 1 0	1 1 0	1 1 0	1 1 0	1 1 0	288 166 0	341 101 11 55	256 235 0	53 13 0	181 235 0	45 26 0	51 51 0	149 72 0	96 19 0	171 6 0	88 17 0	322 26 0	152 17 0	96 0 0	313 24 0	107 40 31	245 10 73	33 0 0	364 110 20	99 13 20	107 26 77	353 51 179	582 18 69	199 116 11	380 36 260
550 - 731	124 72 156	124 72 156	724 726 728	1134 213 182	580 59 21	597 30 139	188 114 29	119 5 172	128 33 134	432 183 134	144 322 64	550 213 158	500 198 145	516 346 145	267 65 145	283 134 258	145 63 136	171 63 143	171 18 143	645 59 70	407 73 319	262 112 1409	176 238 383	206 128 225	395 128 268	55 74 326	312 62 558	181 74 296	202 178 469												
732 - 914	.	134 106 154	752 756 760	37 87 95																																					
Grand Total				3053	1246	1837	2595	4180	954	320	1991	1394	1892	1110	567	470	1184	1491	1947	2701	1799	1027	1289	664	3440	981	2696	1057	4299	4484	5844	2446	2374	4489	2716	3573					
Abundance >366 m				0	0	0	0	0	0	0	1984	1013	1178	712	477	353	913	738	1100	1107	1236	797	728	496	898	0	952	719	2071	1279	737	577	1144	1937	857	1659					
Percent >366 m				0.0	0.0	0.0	0.0	0.0	0.0	0.0	99.6	72.7	62.3	64.1	84.1	75.0	77.1	49.5	56.5	41.0	68.7	77.6	56.5	74.6	26.1	0.0	35.3	68.1	48.2	28.5	12.6	23.6	48.2	43.2	31.6	46.4					



Table 11. Abundance (000s) of witch flounder (M+F) in each stratum from surveys in Div. 30 during spring of 1984-2016 (Engel 145 data converted to Campelen units for 1984-95).



Table 12. Abundance (000s) of witch flounder (M+F) in each stratum from surveys in Div. 3N during fall of 1990-2016 (Engel 145 data converted to Campelen units for 1990-94).

Depth Range (m)	Old Stratum Area	New Stratum Area	Stratum	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16
<=56	1593 1499	1593 1499	375 376	0 0	55 0	0 0	0 0	0 0	0 23	0 19	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	
57 - 92	2992 1853 2520 2520 931 674	2992 1853 2520 373 931 674	360 361 362 441	382 32 277 116 0 0	206 425 0 0 0 0	1646 701 0 0 0 0	320 0 0 0 0 0	103 42 0 0 0 0	1232 0 0 0 0 0	41 0 0 0 0 0	672 23 0 0 0 0	755 306 50 0 0 0	360 51 204 0 0 0	926 51 255 102 0 0	514 211 0 0 0 0	1080 1132 154 0 0 0	1022 4888 0 0 0 0	1132 154 0 0 0 0	4888 9290 0 0 0 0	154 9290 0 0 0 0	0 17639 0 0 0 0	3224 3224 2381 2381 22490 17384	1286 0 262 153 149 149	1029 51 50 0 0 0						
93 - 183	421 100 647	421 377 647	359 382	0 0	0 0	608 0	0 0	0 0	87 7	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0		
184 - 274	225 139 182	225 139 182	358 378 381	0 0 0	46 105 0	108 19 0	31 0 0	0 0 0	234 0 0	0 8 0	31 10 0	93 8 0	46 0 0	69 8 0	136 10 0	0 0 0	307 31 0	31 251 0	252 252 0	31 230 0	230 190 0	174 174 0	155 650 0	120 765 0	58 51 0					
275 - 366	164 106 116	164 106 116	357 379 380	0 7 0	384 15 0	23 0 0	338 0 0	135 19 0	180 22 0	0 0 0	60 8 0	0 8 0	124 8 0	33 24 0	20 0 0	102 0 0	34 0 0	98 16 0	242 250 0	116 15 0	259 350 0	29 24 0	72 81 0	11 1500 0	143 51 0	68 10 0	346 87 24	11 10 0		
367 - 549	155 105 160	155 105 160	723 725 727	53 36 0	330 701 44	394 173 11 11	117 49 0 0	21 29 55	88 101 11 11	313 71 0 0	85 22 0 0	104 14 0 0	190 71 0 0	57 22 0 0	347 14 0	43 29 0	329 72 0	72 38 0	38 227 0	227 239 0	239 94 0	94 153 0	153 87 0	87 96 0	2644 166 127	117 39 0				
550 - 731	124 72 156	124 72 156	724 726 728	443 669 268	1126 114 195	512 119 129	223 99 212	178 40 215	571 92 311	326 125 417	640 40 223	337 37 223	264 176 633	270 129 351	177 84 161	247 42 73	629 42 204	384 106 343	1651 106 428	771 102 303	381 91 860	432 20 118	245 44 245	213 78 354	213 11 204	26 113 230				
732 - 914	.	134 106 154	752 756 760								165 255 244	28 149 229	0 182 409	74 22 530																
915 - 1097	.	138 102 171	753 757 761								0 0 106	9 0 578	0 96 202	33 92 24																
1098 - 1280	.	180 99 212	754 758 762								0 0 0	0 0 483	0 8 0	12 0 58																
1281 - 1463	.	385 127 261	755 759 763								0 0 72	0 0 18	0 9 88	0 0 0																
Grand Total				863	1995	3272	3515	1793	2470	488	2046	5355	2073	3233	3756	3717	2912	3806	7017	2126	3289	14269	23473	6742	6251	25163	19725	6732	1943	
Abundance >366 m				0	497	36	2825	1506	714	427	1203	1755	1274	2033	2148	2032	866	475	1446	1026	1575	2773	2775	1421	841	762	863	3282	618	
Percent >366 m				0.0	24.9	1.1	80.4	84.0	28.9	87.6	58.8	32.8	61.5	62.9	57.2	54.7	29.7	12.5	48.3	47.9	19.4	11.8	21.1	13.5	3.0	4.4	48.8	31.8		



Table 13. Abundance (000s) of witch flounder (M+F) in each stratum from surveys in Div. 30 during fall of 1990-2016 (Engel 145 data converted to Campelen units for 1990-94).

Depth Range (m)	Old Stratum Area	New Stratum Area	Stratum	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16
57 - 92	2089	2089	330	131	144	72	0	0	517	0	96	335	383	192	575	588	766	123	479	718	671	1149	2062	899	1197	144	2086	2402	1006	
	456	456	331	42	502	125	0	0	408	0	0	596	4799	533	1066	1850	1004	31	1098	345	439	345	1296	3907	2729	215	2164	220	125	
	1898	1898	338	3264	627	1436	6893	4700	8459	522	2872	1723	7572	609	1984	2245	6893	11652	4774	1567	1044	3220	5817	13606	7989	1816	3290	2141	574	
	1716	1716	340	262	330	118	0	0	295	0	47	0	1652	189	378	189	94	47	243	1416	47	1014	320	140	236	1054	2041	202	330	
	2520	2520	351	1837	347	58	0	0	0	0	50	347	0	198	0	50	50	99	495	297	231	99	154	99	347	0	50	149		
	2580	2580	352	1597	1242	2011	1115	355	371	355	1141	754	1825	1668	1065	1448	2296	6584	2484	1787	811	2419	11915	3712	4817	2789	2563	862	152	
	1282	1282	353	2822	485	941	0	1176	999	882	573	5467	5996	6172	2954	9523	3395	5291	6525	3357	1950	2469	16690	17768	7186	11243	4144	2381	6922	
	93 - 183	1721	1721	329	132	101	0	47	0	663	0	616	852	0	0	805	1989	379	703	710	8181	0	10750	6155	300	4972	4856	2736	0	1184
1047	332	3625	396	5281	2064	960	5233	11954	1248	2544	7393	3249	1392	4342	3738	6145	8381	13093	2939	8910	2603	5770	1509	14968	1632	2016	3649			
	948	948	337	2347	424	2347	1043	5216	1434	717	1130	1613	3738	1623	348	714	1434	397	5067	696	1956	3775	1546	4482	782	1198	729	609	391	
	585	585	339	1556	241	724	121	966	2776	0	1086	356	3943	563	3822	684	7559	4507	2374	4064	2070	4529	5754	4547	1927	885	2052	885		
	474	474	354	1891	33	685	359	424	489	8955	489	782	391	2478	630	1415	1989	1150	978	1206	2195	663	4492	1992	978	261	978	1304	359	
	151	147	333	582	52	83	62	312	187	192	147	152	27	118	90	243	30	51	153	81	108	27	54	57	30	18	10	73		
121	336	222	466	216	633	42	549	208	100	215	300	141	150	58	75	50	300	150	422	518	72	83	50	72	50	50	50	50		
	103	103	355	1459	298	425	85	63	768	28	170	411	85	21	28	21	92	35	27	50	246	94	64	50	101	16	8	28		
	92	96	334	76	70	0	21	57	56	33	20	58	18	36	35	53	65	122	0	7	0	24	18	65	75	47	40	32		
58	335	371	100	112	68	52	64	64	4	40	48	37	8	39	12	18	7	24	18	18	0	11	0	27	0	7	4			
	61	61	356	25	8	1254	252	40	113	13	34	75	55	19	17	34	31	45	0	7	0	37	4	56	8	4	0	0		
	93	166	717	122	0	96	703	46	833	2166	0	91	203	351	117	10	93	41	1214	360	100	340	670	434	91	157				
76	719	209	42	277	10	52	612	183	178	99	75	183	37	96	96	78	95	14	41	167	50	43	12	132	47	58				
	76	76	721	47	444	183	102	131	17	125	311	98	10	84	81	11	135	9	273	68	19	62	38	161	24	30	10			
	111	134	718	107	428	164	535	618	581	396	488	1432	1483	575	1040	479	2013	959	1039	507	489	126	1155	374						
105	105	720	339	0	105	316	29	202	39	762	298	302	206	336	6	6	141	7	14	31	0	165	581	116						
	93	93	722	26	243	58	64	134	51	103	122	70	94	34	50	90	199	51	61	117	89	65	77	44	128	41	19			
	. .	105	764																											
732 - 914	. .	99	768																											
	. .	135	772																											
915 - 1097	. .	124	765																											
	. .	138	769																											
	. .	128	773																											
1098 - 1280	. .	144	766																											
	. .	128	770																											
	. .	135	774																											
1281 - 1463	. .	158	767																											
	. .	175	771																											
	. .	155	775																											
<b>Grand Total</b>				21086	7158	14515	15517	15369	23795	25731	10499	20054	38620	22908	15520	33557	26262	41114	39294	35843	18702	41498	60585	60036	38388	42483	24415.3		16298	16597
<b>Abundance &gt;366 m</b>				331	114	0	1411	774	1191	1193	831	4354	3480	1890	3210	5163	3103	1095	3390	254	1704	3593	2879	1331	1036	1376	1009.3		1945	734
<b>Percent &gt;366 m</b>				1.6	1.6	0.0	9.1	5.0	5.0	4.6	7.9	21.7	9.0	8.3	20.7	15.4	11.8	2.7	8.6	0.7	9.1	8.7	4.8	2.2	3.2	4.1		11.9	4.4	



Table 14. Mean numbers per tow for witch flounder (M=F) in each stratum from surveys in Div. 3N during spring of 1984-2016 (Engel 145 data converted to Campelen units for 1984-95).

Depth Range (m)	Old Stratum Area	New Stratum Area	Stratum	Strata 84-160																																		
				84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16		
<=56	1593	1593	375	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.33	0.20	0.00	0.20	0.00	0.00	0.00	0.00	0.00	0.00					
	1499	1499	376	0.00	0.00	0.00	0.13	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.17	0.17	0.00	0.00	0.00	0.00	0.00	0.00	0.43	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00			
57 - 92	2992	2992	360	5.43	0.31	1.77	1.80	6.42	0.53	0.00	0.00	0.14	0.55	0.00	0.00	0.32	0.16	0.55	1.49	0.00	0.00	0.20	0.30	3.78	1.17	1.80	0.25	0.00	2.00	0.70	0.40	0.80	0.50	0.00	0.00			
	1853	1853	361	0.60	0.00	0.00	0.13	0.14	0.00	0.11	0.00	0.00	0.00	0.14	0.00	0.00	0.00	0.00	0.83	0.33	0.00	0.00	0.00	0.00	0.14	1.00	0.00	0.20	0.33	0.00	0.80	0.00	0.67	0.25	0.00	0.00		
	2520	2520	362	0.00	0.27	0.07	0.08	0.50	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.50	0.00	0.00	0.00	0.11	0.00	0.00	0.00	0.00	0.00	0.00			
	2520	2520	373	0.00	0.00	0.14	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.11	0.11	0.00	0.00	0.00	0.00				
	931	931	374	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.33	0.33	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.67	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00					
	674	674	383	0.00	0.67	0.00	0.33	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00				
93 - 183	421	421	359	7.00	1.00	4.00	1.00	17.00	3.50	0.00	0.00	0.50	0.00	0.00	0.00	0.00	0.00	0.00	3.50	7.00	1.00	0.50	0.00	0.00	12.00	1.50	0.00	25.00	33.71	60.00	10.50	2.00	23.67	20.00	3.00			
	100	100	377	1.00	0.00	0.00	13.50	0.50	6.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	4.00	0.00	0.00	0.00	0.00	0.00			
	647	647	382	0.00	0.00	0.00	0.33	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00				
184 - 274	225	225	358	2.50	18.00	3.00	9.00	1.00	1.50	3.00	0.00	3.00	9.50	7.50	1.00	2.50	2.67	8.44	0.50	1.33	10.50	0.89	9.57	0.00	3.56	22.00	4.89	17.50	9.78	18.29	6.00	10.67	7.43	1.60	51.50			
	139	139	378	2.50	1.50	2.50	18.50	4.50	6.00	0.00	0.00	5.00	0.00	0.00	0.00	0.44	0.00	0.00	0.00	0.44	1.71	0.44	0.00	0.89	0.00	0.00	0.00	0.00	0.00	0.00	1.02	5.83	0.00	0.00	0.89	0.00		
	182	182	381	1.00	0.50	1.67	6.50	3.00	0.00	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.50	0.00	0.00	0.44	0.00	0.00	0.00	3.22	1.00	1.33	0.00	0.89	2.03	1.50	1.50	17.50	2.00				
275 - 366	164	164	357	1.00	8.00	24.50	0.50	10.50	2.50	0.00	4.00	5.50	4.50	1.00	1.78	1.33	16.56	11.50	13.00	2.78	2.44	6.67	2.00	0.00	1.00	0.00	1.00	4.33	16.00	14.07	2.00	2.83	8.00	4.29				
	106	106	379	4.50	2.50	4.67	29.00	7.00	3.00	7.50	0.50	3.00	0.00	1.50	0.00	0.00	1.24	0.44	7.00	1.94	0.89	0.00	1.07	0.00	2.73	0.00	0.00	0.50	29.16	3.37	19.50	13.07	35.33	0.50	0.00			
	116	116	380	0.50	5.50	0.00	15.50	2.00	0.50	0.50	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.50	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.44	0.00	0.00	0.00	3.36	0.67	1.33	0.50	0.00				
367 - 549	155	155	723							13.50	16.00	12.00	2.50	8.50	2.13	2.40	7.00	4.50	8.00	4.11	15.11	7.11	4.50	14.67	5.00	11.49	1.56	17.06	4.64	5.00	16.53	27.30	9.33	17.83				
	105	105	725							11.50		7.00	6.00	0.00	0.89	16.30	1.78	3.50	5.00	1.33	0.44	1.14	0.00	18.28	2.78	0.67	7.60	0.90	1.83	3.50	1.24	10.67	8.00	2.50				
	160	160	727							0.00	0.50	2.50	1.00	0.00	0.00	0.50	1.50	1.50	0.94	0.44	0.00	0.00	1.40	3.11	1.39	3.33	0.00	0.89	3.75	3.50	8.13	3.14	0.50	11.83				
550 - 731	124	124	724							66.50	34.00	35.00	11.00	7.00	7.50	25.35	8.44	32.22	29.33	30.22	15.67	16.61	8.50	10.00	37.83	23.87	15.36	10.30	12.10	23.17	3.21	18.29						
	72	72	726							21.50	6.00	3.00	11.50	0.50	3.33	18.49	32.50	21.50	20.00	34.94	6.58	13.50	6.35	1.78	6.00	7.33	11.36	24.00	12.93	7.50	6.29	18.00	18.29	20.36				
	156	156	728							8.50	1.00	6.50	1.33	8.00	6.22	3.00	7.37	6.76	12.00	6.33	6.67	7.50	3.00	3.28	14.86	65.65	17.83	10.50	12.50	15.21	26.00	13.78	21.83					
732 - 914	.	134	752															2.00																				
	.	106	756															6.00																				
	.	154	760															4.50																				
Grand Total				1.33	0.54	0.80	1.14	1.82	0.42	0.14	0.83	0.58	0.79	0.45	0.24	0.20	0.50	0.62	0.81	1.12	0.75	0.43	0.54	0.28	1.43	0.63	1.12	0.44	1.79	1.87	2.43	1.03	0.99	1.87	1.13	1.48		



Table 15. Mean numbers per tow for witch flounder (M+F) in each stratum from surveys in Div. 30 during spring of 1984-2016 (Engel 145 data converted to Campelen units for 1984-95).

Depth Range (m)	Old Stratum Area	New Stratum Area	Stratum	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16				
57 - 92	2089	2089	330	0.00	0.00	0.00	0.00	0.11	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.25	0.13	0.73	0.84	0.00	0.00	0.51	0.71	5.19	0.00	1.42	0.00	3.13	6.25	0.43	0.29	2.00	0.00	0.00						
	456	456	331	56.67	6.00	1.50	0.50	16.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	1.00	0.00	1.50	17.60	1.00	11.50	1.50	0.00	0.00	12.50	46.00	18.00	39.50	62.28	8.39	3.00	0.44	12.50	0.50	0.00					
	1898	1898	338	0.80	45.56	5.78	7.44	20.75	9.50	2.25	0.00	0.50	1.83	0.00	1.17	5.43	0.00	2.57	7.56	1.33	8.67	1.17	2.33	11.45	8.00	19.56	6.50	3.33	7.33	52.11	8.30	21.71	24.50	4.00	8.00	0.83				
	1716	1716	340	0.25	0.89	0.00	0.11	0.00	0.00	0.22	0.00	0.60	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.60	0.00	0.00	0.00	0.00	0.20	0.50	1.00	0.00	1.40	0.59	0.00	0.00	0.40	0.33	0.00					
	2520	2520	351	2.67	0.67	1.43	0.77	3.80	0.69	0.33	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.11	0.13	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00					
	2580	2580	352	0.29	5.09	1.21	5.77	5.18	2.62	5.00	0.14	0.25	0.14	0.13	0.20	0.22	0.56	0.10	5.11	0.56	0.13	5.50	0.13	3.33	3.00	7.00	3.38	2.38	0.43	10.46	3.53	12.39	1.50	0.40	0.14	0.00				
	1282	1282	353	53.00	7.00	9.71	12.17	74.00	22.00	16.50	0.00	1.50	2.00	0.00	0.20	0.20	1.50	2.60	28.66	14.40	5.11	4.71	6.25	5.43	4.94	43.19	4.50	6.00	1.75	100.75	13.64	36.25	12.56	13.50	4.67	0.00				
93 - 183	1721	1721	329	0.00	0.00	0.00	0.00	6.14	0.22	0.14	3.22	0.00	0.00	51.80	2.20	0.00	0.15	0.29	2.63	0.20	0.00	0.00	22.40	0.00	3.13	5.46	3.00	9.80	1.27	7.47	12.29	77.00	0.67	5.20	0.00					
	1047	1047	332	76.50	115.20	45.33	50.20	111.25	19.80	73.40	31.33	40.00	3.50	3.00	27.25	20.32	39.33	7.54	35.03	15.50	58.00	47.00	228.33	170.24	35.00	17.33	89.33	60.07	40.06	40.30	29.33	217.33	178.50	15.67	41.00					
	948	948	337	1.00	70.40	20.20	27.17	20.25	19.60	20.00	24.40	6.25	16.00	0.67	9.50	6.33	3.59	6.50	28.44	25.00	51.67	14.00	12.00	30.00	18.67	34.37	8.00	54.29	23.67	6.50	14.00	30.50								
	585	585	339	5.50	0.00	1.00	3.33	1.67	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	2.00	0.44	1.00	0.44	1.00	3.50	3.00	0.00	5.50	21.78	10.57	4.00	20.00	71.50	0.89	0.00	3.50	3.00	0.00					
	474	474	354	18.00	3.67	50.33	7.00	9.50	3.00	5.50	4.00	4.00	25.50	0.00	0.00	1.50	0.50	8.64	49.20	42.00	32.21	22.50	5.50	14.00	30.06	19.00	35.00	16.00	30.56	36.83	10.00	16.50	20.64	21.50	1.00					
184 - 274	151	147	333	1.00	7.50	1.67	0.00	7.00	2.50	16.00	65.50	9.00	14.50	665.00	21.00	1.50	13.72	6.93	13.20	12.93	28.50	46.50	10.61	11.14	13.50	8.61	3.56	12.50	2.67	2.67	1.81	9.50	1.50	0.50	26.50					
	121	121	336	1.50	1.00	10.50	4.00	12.50	0.00	9.50	82.00	197.50	16.00	182.00	7.50	25.93	41.00	9.00	10.40	13.14	35.00	76.50	31.50	15.50	22.11	14.00	16.50	12.83	4.33	8.67	2.00	13.56	5.50	3.00	17.50					
	103	103	355	6.50	29.50	9.00	9.50	0.00	27.00	36.00	24.00	2.00	7.00	7.00	11.83	13.78	11.11	2.67	2.89	15.50	40.13	66.67	17.36	4.00	7.50	6.00	12.22	4.50	3.73	5.21	11.00	1.50	3.50	13.00						
275 - 366	92	96	334	0.00	7.50	13.00	5.00	7.50	3.50	4.00	3.00	21.50	5.00	169.50	3.00	35.00	66.67	0.50	12.18	12.67	2.28	28.50	40.39	18.00	1.50	5.22	2.50	10.00	1.33	2.89	2.40	4.00	3.50	1.33	19.33					
	58	58	335	0.00	25.50	5.00	1.00	18.50	8.50	41.50	13.67	293.33	28.00	27.00	13.50	24.06	30.40	1.50	21.22	46.14	7.50	5.94	16.44	4.36	9.78	2.78	0.89	2.28	1.33	7.10	8.50	4.44	7.50	0.00	1.50					
	61	61	356	2.00	25.50	4.50	6.50	13.00	9.50	15.00	11.00	41.50	38.00	22.50	15.00	10.50	4.80	10.67	6.44	6.00	7.94	9.33	15.56	3.00	9.78	1.94	1.78	2.89	0.44	1.24	2.00	23.17	2.00	3.00	17.50					
367 - 549	93	166	717													2.50	29.00	13.00	261.00	10.00	59.67	506.49	31.11	10.40	7.11	11.94	28.50	20.50	2.00	7.93	4.00	5.14	29.87	4.39	2.60	2.00	12.17	3.71	12.43	
	76	76	719													27.50	242.50	25.50	3.50	4.00	34.84	111.07	14.33	10.67	21.80	9.27	25.67	8.50	1.80	12.50	7.78	7.67	2.67	4.80	27.14	9.71	4.80	1.50	7.11	0.57
	76	76	721													22.50	20.00	9.00	18.50	4.00	6.04	20.50	14.55	10.67	19.56	13.33	8.00	3.00	1.78	5.71	5.39	24.00	5.94	23.33	4.00	5.00	2.00	0.00	1.00	
550 - 731	111	134	718													18.50	8.00	33.50	63.00	29.00	28.09	27.52	28.02	17.56	7.50	28.50	64.50	31.33	3.56	9.59	13.00	19.36	111.24	56.34	35.39	9.24	70.00	21.00	16.43	
	105	105	720													25.00	26.00	71.00	34.50	3.00	6.98	35.83	12.84	7.20	24.28	21.39	3.43	7.22	2.83	52.94	4.31	5.21	5.00	1.00	1.50	1.71	35.17	3.67	0.67	
	93	93	722													3.50	13.00	40.00	40.50	47.00	21.44	64.06	13.83	28.44	16.21	28.22	15.49	16.44	4.14	12.00	13.78	10.39	7.50	5.11	19.14	8.00	5.71	5.07	2.00	4.79
732 - 914	.	105	764													15.00																								
	.	135	772													27.00																								
Grand Total				11.01	17.41	7.04	7.43	17.96	5.61	8.55	5.25	6.97	3.54	16.00	3.33	3.78	9.30	2.53	9.78	5.58	9.68	7.55	17.98	12.83	7.05	10.65	6.11	10.11	9.89	23.52	8.32	12.69	25.84	13.58	4.15	5.28				



Table 16. Mean numbers per tow for witch flounder (M+F) in each stratum from surveys in Div. 3N during fall of 1990-2016 (Engel 145 data converted to Campelen units for 1990-1994).

Depth Range (m)	Old Stratum Area	New Stratum Area	Stratum	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16		
<=56	1593	1593	375	0.00	0.25	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.25	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00			
	1499	1499	376	0.00	0.00	0.00	0.00	0.00	0.11	0.00	0.09	0.00	0.00	0.00	0.00	0.29	0.29	0.00	0.00	0.00	0.00	0.00	0.33	0.00	0.00	0.00	0.50	1.25	0.25	0.00		
57 - 92	2992	2992	360	0.93	0.50	4.00	0.78	0.25	2.99	0.10	1.63	1.83	0.88	2.25	1.25	2.63	2.48	2.75	11.88	0.38	0.00	22.57	42.86	7.83	5.79	54.64	42.24	3.13	2.50			
	1853	1853	361	0.13	1.67	2.75	0.00	0.17	0.00	0.00	0.09	0.00	1.20	0.20	0.80	1.00	0.40	0.00	0.83	0.20	4.00	0.33	0.00	2.20	0.98	1.03	0.60	0.00	0.20			
	2520	2520	362	1.27	0.80	0.33	0.00	0.00	0.00	0.00	0.00	0.00	0.14	0.00	0.00	0.00	0.57	0.00	0.00	0.14	0.00	0.00	0.17	0.86	0.29	0.43	0.43	0.00	0.14			
	2520	2520	373	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00			
	931	931	374	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00			
	674	674	383	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.50			
93 - 183	421	421	359	0.00	0.00	10.50	0.00	0.00	1.50	0.00	0.00	47.00	0.50	0.00	7.00	2.00	4.00	3.50	1.50	2.50	9.06	21.00	45.50	15.00	16.50	5.71	4.67	14.57	1.00			
	100	100	377	0.00	0.00	0.00	0.00	0.50	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	34.39	3.22	1.50	8.00	0.00	0.00	0.00	0.00		
	647	647	382	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00			
184 - 274	225	225	358	0.00	1.50	3.50	1.00	0.00	7.56	0.00	1.00	3.00	1.50	2.22	4.39	0.00	9.91	1.00	8.11	8.14	1.00	7.44	6.14	5.61	5.00	21.00	3.89	0.00	1.89			
	139	139	378	0.00	5.50	1.00	0.00	0.00	0.00	0.44	0.50	0.00	0.00	0.44	0.50	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.44	1.00	0.44	2.00	5.86	18.78	40.00	2.67		
	182	182	381	0.00	0.00	0.00	0.00	0.00	0.00	0.30	0.50	0.00	0.00	0.44	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.44	0.00	0.00	0.00	0.00	0.00	0.00			
275 - 366	164	164	357	0.00	17.00	1.00	15.00	6.00	8.00	0.00	2.67	0.00	5.50	1.44	0.89	4.50	1.50	4.33	10.72	5.14	11.50	1.29	3.20	0.50	6.33	3.00	15.33	0.50				
	106	106	379	0.50	1.00	0.00	0.00	0.00	1.33	1.50	0.00	0.00	0.44	20.33	6.21	1.78	131.36	0.89	0.44	1.00	24.00	1.64	5.52	102.86	3.50	0.67	6.00	0.67	0.00	1.50		
	116	116	380	0.00	0.00	0.00	0.00	0.00	0.00	0.50	0.50	1.50	0.00	0.00	0.00	1.00	1.50	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.44			
367 - 549	155	155	723	2.50	15.50	18.50	5.50	1.00	4.12	14.67	4.00	4.89	8.93	2.67	16.28	2.00	14.00	3.38	1.78	10.67	11.20	4.40	7.17	4.10	4.50	124.00	5.50					
	105	105	725	2.50	48.50	12.00	3.40	0.00	16.40	2.00	7.00	4.89	1.50	1.00	2.00	1.44	1.02	2.22	4.00	6.29	2.57	2.00	10.74	11.50	2.67	0.00	0.00	0.00	0.00	0.00		
	160	160	727	0.00	2.00	0.50	0.00	2.50	0.50	0.50	0.00	0.57	0.00	0.50	0.50	0.26	0.00	0.00	13.94	7.42	3.00	2.60	3.50	1.50	5.78	0.00	0.00	0.00	0.00			
550 - 731	124	124	724	26.00	66.00	30.00	13.07	10.44	33.50	19.11	37.50	19.73	15.50	15.82	10.35	14.49	36.89	22.50	96.76	45.20	22.36	25.33	14.39	12.50	1.50	7.00						
	72	72	726	67.50	11.50	12.00	10.00	4.00	9.33	12.67	4.00	3.71	17.78	13.00	8.50	4.28	10.73	12.57	10.33	30.60	2.00	4.40	7.91	1.14	11.67	11.39	0.00	0.00	0.00	0.00		
	156	156	728	12.50	9.07	6.00	9.90	10.00	14.50	19.43	10.40	29.50	16.33	7.50	3.42	9.50	16.00	19.94	41.60	40.06	5.50	11.43	16.49	9.50	10.72	0.00	0.00	0.00	0.00	0.00		
732 - 914	.	134	752									8.94	1.50	0.00	4.00																	
	.	106	756									17.50	10.21	12.50	1.50					12.00												
	.	154	760									11.50	10.79	19.29	25.00					2.50	16.00											
915 -1097	.	138	753									0.00	0.50	0.00	1.71																	
	.	102	757									0.00	0.00	6.86	6.57					0.50												
	.	171	761									4.50	24.57	8.57	1.00					17.50	1.00											
1098 -1280	.	180	754									0.00	0.00	0.00	0.50	0.50																
	.	99	758									0.00	0.00	0.00	0.57					0.00												
	.	212	762									0.00	0.00	16.57	0.00					2.00	3.33											
1281 -1463	.	385	755									0.00	0.00	0.00	0.00																	
	.	127	759									0.00	0.00	0.50	0.00					0.00												
	.	261	763									2.00	0.50	2.44	0.00					0.00												
Grand Total				0.38	0.87	1.79	1.48	0.75	1.03	0.20	0.85	2.04	0.87	1.20	1.40	1.38	1.22	1.59	2.73	0.89	1.22	5.94	9.67	2.82	2.60	10.48	8.22	2.80	0.81			



Table 17. Mean numbers per tow for witch flounder (M+F) in each stratum from surveys in Div. 30 during fall of 1990-2016 (Engel 145 data converted to Campelen units for 1990-1994).

Depth Range (m)	Old Stratum Area	New Stratum Area	Stratum	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16
57 - 92	2089	2089	330	0.46	0.50	0.25	0.00	0.00	1.80	0.00	0.33	1.17	1.33	0.67	2.00	2.05	2.67	0.43	1.67	2.50	2.33	4.00	7.17	3.13	4.17	0.50	7.26	8.36	3.50	
	456	331	331	0.67	8.00	2.00	0.00	0.00	6.50	0.00	0.00	9.50	76.50	8.50	17.00	29.50	16.00	0.50	17.50	5.50	7.00	5.50	20.67	62.28	43.50	3.43	34.50	3.50	2.00	
	1898	1898	338	12.50	2.40	5.50	26.40	18.00	32.40	2.00	11.00	6.60	29.00	2.33	7.60	8.60	26.40	44.63	18.29	6.00	4.00	12.33	22.28	52.11	30.60	6.96	12.60	8.20	2.20	
	1716	1716	340	1.11	1.40	0.50	0.00	0.00	1.25	0.00	0.20	0.00	7.00	0.80	1.60	0.80	0.40	0.20	1.03	6.00	0.20	4.30	1.36	0.59	1.00	4.47	8.64	0.86	1.40	
	2520	2520	351	5.30	1.00	0.17	0.00	0.00	0.00	0.00	0.00	0.14	1.00	0.00	0.57	0.00	0.14	0.14	0.29	1.43	0.86	0.67	0.29	0.44	1.00	0.00	0.14	0.43		
	2580	2580	352	4.50	3.50	5.67	3.14	1.00	1.05	1.00	3.21	2.13	5.14	4.70	3.00	4.08	6.47	18.55	7.00	5.04	2.29	6.81	33.57	10.46	13.57	7.86	7.22	2.43	0.43	
	1282	1282	353	16.00	2.75	5.33	0.00	6.67	5.67	5.00	3.25	31.00	34.00	35.00	16.75	54.00	19.25	30.00	37.00	19.04	11.06	14.00	94.68	100.75	40.75	63.75	23.50	13.50	39.25	
93 - 183	1721	1721	329	0.56	0.43	0.00	0.20	0.00	2.80	0.00	2.60	3.60	0.00	0.00	3.40	8.40	1.60	2.97	3.00	34.56	0.00	45.41	26.00	1.27	21.00	20.51	11.56	0.00	5.00	
	1047	1047	332	25.17	2.75	36.67	14.33	6.67	36.33	83.00	8.67	17.67	51.33	22.56	9.67	30.15	25.95	42.67	58.19	90.91	20.41	61.87	18.07	40.06	10.48	103.93	11.33	14.00	25.33	
	948	948	337	18.00	3.25	18.00	8.00	40.00	11.00	5.50	8.67	12.37	28.67	12.44	2.67	5.48	11.00	3.05	38.86	5.33	15.00	28.94	11.85	34.37	6.00	9.19	5.59	4.67	3.00	
	585	585	339	19.33	3.00	9.00	1.50	12.00	34.50	0.00	13.50	4.43	49.00	7.00	47.50	8.50	93.93	56.00	29.50	50.50	25.72	56.29	71.50	56.50	23.94	11.00	25.50	11.00		
	474	474	354	29.00	0.50	10.50	5.50	6.50	7.50	137.33	7.50	12.00	6.00	38.00	9.67	21.70	30.50	17.64	15.00	18.50	33.67	10.17	68.89	30.56	15.00	4.00	15.00	20.00	5.50	
184 - 274	151	147	333	28.00	2.50	4.00	3.00	15.00	9.00	9.50	7.28	7.50	1.33	5.83	4.44	12.00	1.50	2.50	7.56	4.00	5.33	1.33	2.67	2.83	1.50	0.89	0.50	3.61		
	121	121	336	13.33	28.00	13.00	38.00	2.50	33.00	12.50	6.00	12.94	18.00	8.50	9.00	3.50	4.50	3.00	18.00	9.00	25.33	31.11	4.33	5.00	3.00	4.33	3.00			
	103	103	355	103.00	21.00	30.00	6.00	4.44	54.20	2.00	12.00	29.00	6.00	1.50	2.00	1.50	6.50	2.44	1.89	3.56	17.33	6.67	4.50	3.50	7.11	1.14	0.57	0.50		
275 - 366	92	96	334	6.00	5.50	0.00	1.67	4.50	4.43	2.50	1.50	4.43	1.33	2.72	2.67	4.00	4.89	9.22	0.00	0.50	0.00	1.83	1.33	4.89	5.67	3.56	3.00	2.44		
	58	58	335	46.50	12.50	14.00	8.50	6.50	8.00	8.00	0.50	5.00	6.00	4.61	1.00	4.89	1.50	2.22	0.89	3.00	2.22	2.22	0.00	1.33	0.00	3.33	0.00	0.89	0.50	
	61	61	356	3.00	1.00	149.50	30.00	4.78	13.50	1.50	4.00	8.89	6.50	2.28	2.00	4.00	3.71	5.33	0.00	0.89	0.00	4.40	0.44	6.67	1.00	0.50	0.00	0.00		
367 - 549	93	166	717	9.50	0.00	7.50	54.95	2.00	36.50	94.83	0.00	4.00	8.89	15.39	5.14	0.44	4.06	1.78	53.14	15.78	4.39	14.90	29.33	19.00	4.00	6.89				
	76	76	719	20.00	4.00	26.50	1.00	5.00	58.50	17.50	17.00	9.50	7.15	17.50	3.56	9.14	9.14	7.50	9.07	1.33	3.94	16.00	4.80	4.11	1.17	12.64	4.50	5.50		
	76	76	721	4.50	42.50	17.50	9.80	12.50	1.60	12.00	29.71	9.33	1.00	8.00	7.72	1.02	12.89	0.89	26.10	6.50	1.77	5.94	3.67	15.36	2.29	2.86	1.00			
550 - 731	111	134	718	7.00	28.00	10.72	29.00	33.50	31.50	21.50	26.50	77.67	80.44	31.20	56.40	26.00	109.20	52.00	56.34	27.50	26.53	6.86	62.67	20.28						
	105	105	720	23.50	0.00	7.28	21.89	2.00	14.00	2.67	52.76	20.62	20.89	14.29	23.24	0.44	0.40	4.98	0.50	1.00	2.16	0.00	11.43	40.19	8.00					
	93	93	722	2.00	19.00	4.50	5.00	10.50	4.00	8.06	9.50	5.50	7.33	2.67	3.89	7.06	15.56	4.00	4.79	9.11	6.98	5.11	6.00	3.43	10.00	3.24	1.50			
732 - 914	.	105	764									24.71		5.00	10.00	15.00			2.00		5.00		24.57							
	.	99	768									15.94		1.78	12.00	27.43			2.50		0.44		2.50							
	.	135	772									81.50		36.00	20.63	10.22			21.00		6.00		8.72							
915 - 1097	.	124	765									9.67		1.83	7.00	16.93			4.50		3.78		9.22							
	.	138	769									9.50		2.00	12.50	20.00			7.50		7.00		11.50							
	.	128	773									2.00		7.71	19.67	40.20	5.35		3.50		4.50		2.10							
1098 - 1280	.	144	766									5.71		0.57	7.39			15.50		8.00		9.50								
	.	128	770									2.07		10.50	26.14			5.00		7.50		1.00								
	.	135	774									1.50		13.00	6.43	13.14			16.00		1.89		0.00							
1281 - 1463	.	158	767									3.00		0.00	0.00	0.00			0.00		0.44		0.57							
	.	175	771									0.00		5.50	0.00	0.00			2.50		0.00		0.00							
	.	155	775									0.00		0.00	0.00	10.00			5.00		1.33		4.50							
Grand Total				8.56	2.87	5.89	6.11	6.05	9.37	10.39	4.14	7.56	15.63	8.25	5.63	12.09	9.99	16.11	14.16	14.15	6.74	16.26	21.96	23.52	15.04	16.65	9.57	6.39	6.55	

Table 18. Mean weights (kg) per tow for witch flounder (M+F) in each stratum from surveys in Div. 3N during spring of 1984-2016 (Engel 145 data converted to Campelen units for 1984-95).

Depth Range (m)	Old Stratum Area	New Stratum Area	Stratum	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16												
<=56	1593	1593	375	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.19	0.16	0.10	0.00	0.00	0.00	0.00	0.00	0.00																
	1499	1499	376	0.00	0.00	0.00	0.09	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.04	0.09	0.00	0.00	0.00	0.00	0.00	0.00	0.43	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00															
57 - 92	2992	2992	360	4.17	0.22	1.53	1.12	3.69	0.43	0.00	0.00	0.07	0.40	0.00	0.00	0.00	0.28	0.08	0.29	0.65	0.00	0.00	0.05	0.24	2.39	0.64	1.32	0.21	0.00	0.96	0.38	0.17	0.46	0.33	0.00	0.00												
	1853	1853	361	0.47	0.00	0.00	0.15	0.20	0.00	0.08	0.00	0.00	0.00	0.15	0.00	0.00	0.00	0.95	0.18	0.00	0.00	0.00	0.14	0.55	0.00	0.07	0.28	0.00	0.51	0.00	0.36	0.03	0.00	0.00	0.00													
	2520	2520	362	0.00	0.24	0.07	0.05	0.42	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.38	0.00	0.00	0.00	0.00	0.05	0.00	0.00	0.00	0.00	0.00	0.00												
	2520	2520	373	0.00	0.00	0.12	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.04	0.03	0.00	0.00	0.00	0.00													
	931	931	374	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.14	0.27	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.23	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00														
	674	674	383	0.00	0.62	0.00	0.40	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00													
93 - 183	421	421	359	3.99	0.81	1.71	0.75	5.28	2.09	0.00	0.00	0.00	0.33	0.00	0.00	0.00	0.00	1.15	2.58	1.00	0.23	0.00	0.00	5.78	0.90	0.00	10.23	12.42	23.56	5.16	1.43	14.43	10.57	2.02														
	100	100	377	0.58	0.00	0.00	5.25	0.21	2.31	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00													
	647	647	382	0.00	0.00	0.00	0.14	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.45	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00												
184 - 274	225	225	358	1.29	9.94	1.37	4.42	0.64	0.95	1.86	0.00	1.44	4.26	3.42	0.24	1.65	1.58	4.33	0.21	0.29	4.98	0.47	5.43	0.00	1.36	10.23	2.20	7.67	5.03	7.78	2.78	6.11	4.35	0.78	28.56													
	139	139	378	1.17	1.00	1.69	8.10	1.64	2.19	0.00	0.00	1.51	0.00	0.00	0.00	0.00	0.18	0.00	0.00	0.00	0.24	0.41	0.07	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.33	0.00														
	182	182	381	0.82	0.28	1.27	4.04	2.77	0.00	1.14	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.27	0.00	0.00	0.00	0.00	2.10	0.50	0.71	0.00	0.01	1.19	0.01	0.92	10.66	0.01													
275 - 366	164	164	357	0.35	3.85	6.83	0.18	2.65	0.91	0.00	1.36	2.16	3.61	0.89	1.58	0.53	7.04	0.95	3.33	0.77	1.14	2.87	1.85	0.00	0.85	0.00	0.19	1.36	3.68	5.95	1.10	1.87	4.15	2.47														
	106	106	379	2.48	0.83	1.60	11.84	3.00	1.38	2.38	0.21	1.27	0.00	0.29	0.00	0.00	0.60	0.15	1.75	0.25	0.26	0.00	0.28	0.00	0.40	0.00	0.00	0.45	0.85	1.55	6.92	6.07	16.28	0.37	0.00													
	116	116	380	0.40	3.34	0.00	8.38	1.52	0.43	0.24	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.35	0.00	0.00	0.20	0.00	0.00	0.00	0.00	0.33	0.00	0.00	0.00	1.39	0.33	0.75	0.24	0.00														
367 - 549	155	155	723														4.21	4.80	3.71	1.68	2.41	0.77	1.16	2.48	1.53	1.70	1.08	6.08	2.80	1.58	5.04	2.34	3.87	0.62	6.43	2.55	1.96	5.86	11.49	4.10	8.01							
	105	105	725															4.26	2.78	3.08	0.00	0.37	1.91	0.31	1.40	2.20	0.56	0.20	0.51	0.00	7.15																	
	160	160	727															0.00	0.24	1.73	0.76	0.00	0.00	0.15	0.42	0.60	0.56	0.14	0.00	0.00	1.84	0.49	1.23	0.00	0.64	1.47	1.57	4.49	1.98	0.44	8.13							
550 - 731	124	124	724															19.18	10.63	12.81	3.02	2.11	1.73	9.22	3.11	6.17	6.22	7.47	5.62	5.91	3.18	3.80	12.15	8.55	4.78	3.60	4.47	8.79	0.59	7.11								
	72	72	726																8.21	2.52	2.24	2.81	0.35	1.20	4.25	9.65	5.93	6.58	8.46	1.78	5.07	2.15	0.82	1.90	2.49	4.14	10.59	4.60	3.23	2.31	7.73	9.41	10.45					
	156	156	728																4.31	0.88	3.84	1.02	7.07	0.97	0.70	1.51	2.08	4.56	2.01	2.49	3.51	1.95	1.57	8.17	34.83	7.64	5.43	6.62	8.70	17.28	9.42	12.41						
732 - 914	.	134	752																	1.47																												
	.	106	756																	2.29																												
	.	154	760																	1.22																												
Grand Total				0.96	0.33	0.47	0.62	0.97	0.21	0.07	0.27	0.20	0.36	0.21	0.13	0.07	0.19	0.24	0.22	0.43	0.26	0.16	0.23	0.14	0.75	0.37	0.60	0.21	0.81	0.76	1.00	0.48	0.49	1.04	0.61	0.80												



Table 19. Mean weights (kg) per tow for witch flounder (M+F) in each stratum from surveys in Div. 30 during spring of 1984-2016 (Engel 145 data converted to Campelen units for 1984-95).

Depth Range (m)	Old Stratum Area	New Stratum Area	Stratum	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16											
57 - 92	2089	2089	330	0.00	0.00	0.00	0.00	0.08	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.07	0.42	0.39	0.00	0.00	0.41	0.45	1.98	0.00	0.97	0.00	0.00	3.04	0.19	0.13	1.02	0.00	0.00													
	456	456	331	30.49	4.82	0.58	0.29	7.09	0.00	0.00	0.00	0.00	0.00	0.00	1.18	0.00	0.58	8.56	0.45	5.98	1.63	0.00	0.00	4.65	20.74	6.77	17.93	0.26	3.37	1.29	0.16	5.62	0.32	0.00													
	1898	1898	338	0.51	29.90	4.24	4.53	11.78	7.00	1.66	0.00	0.42	1.13	0.00	0.88	3.33	0.00	1.37	2.99	0.70	5.19	0.46	1.23	4.49	2.48	6.41	3.89	1.73	3.79	2.94	3.63	9.84	10.12	1.74	3.08	0.46											
	1716	1716	340	0.17	0.62	0.00	0.09	0.00	0.00	0.07	0.00	0.62	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.35	0.00	0.00	0.00	0.00	0.11	0.38	0.00	0.00	0.77	0.00	0.00	0.00	0.02	0.19	0.00												
	2520	2520	351	1.99	0.61	1.11	0.64	2.82	0.63	0.31	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.06	0.06	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00												
	2580	2580	352	0.23	2.68	0.63	3.59	3.75	1.87	4.02	0.11	0.30	0.17	0.11	0.18	0.17	0.28	0.15	3.37	0.37	0.15	1.95	0.08	1.77	1.55	3.38	2.07	1.56	0.29	1.58	2.23	4.89	0.84	0.24	0.09	0.00											
	1282	1282	353	25.63	6.36	6.05	9.12	40.87	14.10	9.28	0.00	1.38	1.19	0.00	0.24	0.13	0.01	1.54	12.53	7.37	2.66	3.90	2.66	3.24	2.44	19.22	3.27	3.00	0.98	1.70	6.11	16.91	7.17	7.17	2.34	0.00											
93 - 183	1721	1721	329	0.00	0.00	0.00	0.00	3.33	0.20	0.11	2.09	0.00	0.00	21.42	0.82	0.00	0.05	0.21	1.01	0.11	0.00	0.00	9.33	0.00	0.62	2.36	0.91	4.15	2.36	3.18	4.72	31.85	0.28	2.09	0.00	0.00											
	1047	1047	332	26.24	59.64	17.26	23.38	47.42	10.31	31.93	16.84	15.15	2.49	0.41	12.44	8.20	1.63	3.19	6.81	2.83	21.00	17.07	71.07	55.16	7.46	4.45	22.13	13.92	11.59	8.82	6.32	67.81	33.94	4.37	14.72	0.00											
	948	948	337	0.39	31.66	10.85	11.55	8.13	11.83	12.48	12.12	4.45	5.18	0.38	5.01	2.53	1.25	2.46	6.74	7.18	13.98	5.77	5.48	1.79	5.02	2.55	9.28	4.32	4.83	1.52	15.01	7.72	1.07	3.47	13.07	0.00											
	585	585	339	4.17	0.00	0.20	2.78	1.69	0.00	0.00	0.00	0.00	0.00	0.00	0.02	0.00	0.01	0.01	0.01	0.06	0.02	0.00	0.00	0.23	0.05	0.47	3.53	0.03	0.72	0.00	0.18	0.70	0.00	0.00	0.00												
	474	474	354	7.60	1.61	18.88	3.58	5.30	0.73	3.68	2.21	2.28	12.90	0.00	0.00	0.55	0.01	3.46	16.28	12.68	14.02	8.48	2.50	7.60	9.81	6.03	17.60	6.60	2.25	14.84	2.52	5.79	6.58	7.33	0.86												
184 - 274	151	147	333	0.50	2.30	0.48	0.00	3.23	0.75	6.20	23.96	3.83	3.83	256.95	8.00	0.33	5.38	1.22	1.32	1.47	6.03	18.55	3.12	1.80	1.95	1.36	0.47	1.58	0.97	0.31	0.47	2.07	0.01	0.09	7.65	0.00											
	121	121	336	0.75	0.43	2.60	1.48	3.79	0.00	3.16	29.55	82.55	6.00	63.53	3.73	10.83	17.58	1.38	2.82	1.65	9.78	35.90	12.65	3.65	3.07	2.62	3.65	0.96	0.94	1.56	0.60	2.28	1.10	0.90	4.46	0.00											
	103	103	355	3.19	12.77	2.69	5.05	0.00	6.82	8.89	9.58	1.14	2.38	9.12	3.03	6.09	3.40	3.56	1.27	1.01	6.18	13.63	23.99	8.25	0.87	1.93	2.42	4.72	3.12	0.82	1.82	0.97	0.23	1.72	56.27	0.00											
275 - 366	92	96	334	0.00	3.32	3.31	1.44	1.74	1.80	2.07	1.59	8.51	1.58	65.16	1.17	11.36	27.44	0.33	0.54	0.83	0.16	10.87	10.07	2.20	0.20	0.83	0.35	1.06	0.49	0.44	0.09	0.73	0.32	0.17	7.00	0.00											
	58	58	335	0.00	12.26	2.27	0.30	6.40	2.72	11.50	5.27	138.78	8.20	12.91	5.44	9.78	13.71	0.22	7.81	16.03	0.95	1.01	6.64	1.25	1.33	0.31	0.12	0.52	0.34	0.42	2.11	1.48	0.98	0.00	0.40	0.00											
	61	61	356	0.59	9.84	2.09	2.78	2.13	3.51	6.56	4.61	15.34	9.23	9.00	7.34	4.75	1.28	3.44	2.75	1.68	4.01	4.58	5.84	1.58	2.14	0.40	0.69	0.69	0.57	0.04	0.53	3.41	0.27	1.11	8.72	0.00											
367 - 549	93	166	717									0.83	9.41	2.77	104.01	2.32	20.37	190.65	1.91	0.82	0.76	1.80	8.80	6.23	0.24	0.73	0.43	0.53	2.40	0.54	0.26	0.69	0.69	0.29	1.24	0.00											
	76	76	719									14.16	97.97	4.65	1.31	1.74	13.13	57.44	1.44	1.49	2.39	1.16	9.12	3.70	0.26	1.33	1.44	1.07	0.53	0.65	3.60	0.79	0.70	0.28	1.64	0.11	0.00										
	76	76	721									7.24	4.64	2.93	6.91	1.76	1.55	1.80	3.65	3.50	2.64	8.17	3.60	2.47	0.85	0.40	0.98	1.01	2.35	1.03	1.45	0.58	0.38	0.28	0.00	0.45	0.00	0.45	0.00	0.00							
550 - 731	111	134	718									2.27	1.89	6.80	11.97	4.34	3.88	1.99	1.79	2.08	0.82	3.10	3.00	2.34	0.69	0.70	1.10	2.34	8.50	1.17	1.94	0.98	3.37	2.08	1.29	0.00	0.00	0.00	0.00	0.00	0.00						
	105	105	720									15.00	9.28	12.60	6.56	1.03	1.48	10.41	2.19	1.42	2.79	2.60	0.51	1.57	0.65	4.81	0.63	0.62	0.60	0.63	0.27	0.45	2.96	0.42	0.10	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00			
	93	93	722									1.42	3.86	11.72	16.93	16.11	6.94	6.78	2.42	5.54	3.69	9.42	4.84	4.97	0.94	2.13	0.86	1.62	1.36	1.16	2.31	1.41	0.63	0.67	0.56	0.87	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
732 - 914	.	105	764																																												
	.	135	772																																												
				4.92	9.67	3.30	3.98	9.11	3.45	4.41	2.53	3.05	1.33	6.10	1.47	1.53	2.62	0.83	3.30	1.74	3.44	2.81	6.00	4.50	1.98	4.44	2.25	3.26	2.82	1.89	2.90	4.59	9.09	3.22	1.35	2.08											



Table 20. Mean weights (kg) per tow for witch flounder (M+F) in each stratum from surveys in Div. 3N during fall of 1990-2016 (Engel 145 data converted to Campelen units for 1990-94).

Depth Range (m)	Old Stratum Area	New Stratum Area	Stratum	Mean Weight (kg) per Tow																															
				90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16					
<=56	1593	1593	375	0.00	0.33	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.16	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00				
	1499	1499	376	0.00	0.00	0.00	0.00	0.07	0.00	0.11	0.00	0.00	0.00	0.00	0.00	0.19	0.13	0.00	0.00	0.00	0.00	0.33	0.00	0.00	0.29	0.98	0.11	0.00	0.00						
57 - 92	2992	2992	360	0.64	0.42	3.15	0.42	0.18	2.16	0.06	1.04	1.05	0.43	1.30	0.79	1.26	1.42	2.03	5.74	0.24	0.00	11.63	25.11	3.95	3.19	29.14	17.72	1.79	1.38						
	1853	1853	361	0.11	1.83	1.82	0.00	0.13	0.00	0.00	0.06	0.00	1.05	0.11	0.67	0.58	0.39	0.00	0.66	0.15	2.29	0.10	0.00	1.61	0.75	0.74	0.31	0.00	0.11						
	2520	2520	362	1.15	0.64	0.25	0.00	0.00	0.00	0.00	0.00	0.00	0.09	0.00	0.00	0.00	0.39	0.00	0.00	0.11	0.00	0.00	0.13	0.55	0.16	0.20	0.26	0.00	0.09						
	2520	2520	373	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00					
	931	931	374	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00					
	674	674	383	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.28	0.00	0.00	0.00	0.00	0.29	0.25	0.00						
93 - 183	421	421	359	0.00	0.00	4.81	0.00	0.00	0.39	0.00	0.00	20.95	0.01	0.00	2.10	0.73	1.90	2.40	0.75	2.60	3.31	7.63	18.65	4.97	6.88	3.28	2.69	9.03	0.72						
	100	100	377	0.00	0.00	0.00	0.56	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	2.86	2.29	0.71	6.86	0.00	0.00	0.00	0.00	0.00					
	647	647	382	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00						
184 - 274	225	225	358	0.00	0.65	2.14	0.76	0.00	2.40	0.00	0.36	0.98	0.63	1.29	1.45	0.00	4.69	0.73	3.44	4.64	0.90	4.54	2.78	2.68	3.36	12.08	3.16	0.00	0.90						
	139	139	378	0.00	2.17	0.81	0.00	0.00	0.00	0.05	0.01	0.00	0.00	0.14	0.27	0.00	0.00	0.00	0.00	0.00	0.00	0.20	0.38	22.00	1.14	2.92	9.97	23.30	1.26						
	182	182	381	0.00	0.00	0.00	0.00	0.00	0.00	0.04	0.01	0.00	0.00	0.27	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.14	0.00	0.00	0.00	0.00	0.00	0.00	0.00						
275 - 366	164	164	357	0.00	10.39	0.42	8.27	1.91	3.76	0.00	1.18	0.00	2.33	0.82	0.91	1.80	1.20	1.64	4.55	2.60	3.98	0.77	1.75	0.22	4.13	1.36	7.37	0.30							
	106	106	379	0.27	0.25	0.00	0.00	0.00	0.04	0.45	0.00	0.00	0.13	7.59	2.24	0.58	59.46	0.00	0.20	0.01	10.70	0.88	2.01	45.39	1.23	0.27	2.73	0.41							
	116	116	380	0.00	0.00	0.00	0.00	0.00	0.00	0.05	0.14	0.30	0.00	0.00	0.55	0.68	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.78	0.00					
367 - 549	155	155	723	1.93	7.65	8.44	2.69	0.73	1.31	3.48	1.25	1.33	3.09	0.77	5.77	0.95	4.58	1.80	0.78	4.60	4.34	1.29	2.92	1.71	1.78	59.95	0.18								
	105	105	725	1.01	26.05	3.20	1.31	0.00	9.32	0.68	2.28	1.29	0.48	0.38	0.68	0.51	0.52	0.78	1.48	2.75	0.85	0.85	4.89	5.73	1.18										
	160	160	727	0.00	1.71	0.01	0.00	1.30	0.30	0.20	0.00	0.46	0.00	0.02	0.33	0.97	0.00	0.00	6.51	3.71	0.96	0.99	1.44	0.75	3.19	0.00									
550 - 731	124	124	724	10.11	24.29	10.57	6.09	3.54	11.58	4.21	10.60	5.08	4.13	5.28	4.09	5.54	12.09	7.45	26.69	11.97	6.85	8.40	4.20	4.65	0.58	2.37									
	72	72	726		31.26	5.47	4.80	4.03	2.08	3.80	3.39	1.59	2.18	5.96	5.20	3.20	1.92	4.93	4.50	4.24	10.63	0.63	1.69	2.35	0.35	5.72	5.40								
	156	156	728		7.11	1.62	1.00	3.53	3.65	4.95	7.11	4.80	13.33	8.31	4.34	0.90	5.70	8.91	12.56	18.84	20.20	2.37	5.82	9.91	5.05	6.75									
732 - 914	.	134	752									6.54	1.23	0.00	0.03										0.30										
	.	106	756									8.53	3.47	5.67	0.60										4.60										
	.	154	760									4.18	1.93	3.67	8.18										0.83	5.20	10.44								
915 - 1097	.	138	753									0.00	0.01	0.00	0.17										0.00										
	.	102	757									0.00	0.00	2.66	0.48										0.01	0.00									
	.	171	761									1.94	6.23	1.80	0.41									5.00	0.30	4.33									
1098 - 1280	.	180	754									0.00	0.00	0.00	0.01										0.00		0.00								
	.	99	758									0.00	0.00	0.00	0.01										0.00		0.00								
	.	212	762									0.00	0.00	3.74	0.00									0.50	0.97	1.38									
1281 - 1463	.	385	755									0.00	0.00	0.00	0.00										0.00										
	.	127	759									0.00	0.00	0.10	0.00									0.00		0.00									
	.	261	763									0.53	0.15	0.28	0.00									0.00		0.09									
Grand Total				0.31	0.63	1.22	0.69	0.34	0.56	0.07	0.41	0.89	0.37	0.46	0.53	0.56	0.64	0.89	1.25	0.46	0.55	2.79	5.17	1.38	1.28	5.60	3.58	1.49	0.41						



Table 21. Mean weights (kg) per tow for witch flounder (M=F) in each stratum from surveys in Div. 30 during fall of 1990-2016 (Engel 145 data converted to Campelen units for 1990-94).

Depth Range (m)	Old Stratum Area	New Stratum Area	Stratum	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16
57 - 92	2089	2089	330	0.43	0.23	0.28	0.00	0.00	0.86	0.00	0.25	0.58	0.73	0.17	0.99	1.19	1.53	0.26	1.09	1.33	1.26	1.77	3.78	1.20	2.46	0.17	2.91	3.42	1.50	
	456	456	331	0.35	5.02	2.14	0.00	0.00	1.73	0.00	0.00	4.08	15.08	3.88	7.46	12.35	4.88	0.23	6.28	1.73	2.30	1.82	8.99	19.43	12.64	1.20	10.98	1.33	0.77	
	1898	1898	338	8.53	1.68	3.21	15.19	8.40	17.94	1.93	5.09	1.85	10.48	1.44	3.61	3.74	10.21	14.94	7.40	2.32	2.08	5.39	7.86	21.00	9.78	2.46	4.68	3.39	0.88	
	1716	1716	340	0.73	1.19	0.27	0.00	0.00	0.86	0.00	0.09	0.00	1.76	0.44	0.73	0.52	0.24	0.12	0.49	2.77	0.01	2.09	0.49	0.34	0.60	2.43	4.06	0.56	0.65	
	2520	2520	351	4.87	0.82	0.21	0.00	0.00	0.00	0.00	0.00	0.11	0.59	0.00	0.50	0.00	0.07	0.10	0.16	1.06	0.46	0.48	0.08	0.22	0.19	0.68	0.00	0.10	0.26	
	2580	2580	352	3.99	2.52	3.81	2.66	0.64	1.07	0.77	1.62	1.06	4.20	2.59	1.21	2.22	2.72	9.51	4.69	3.13	1.57	3.97	16.67	6.50	7.32	3.76	4.61	1.34	0.18	
	1282	1282	353	13.64	1.94	2.70	0.00	4.15	3.05	4.48	0.95	6.04	16.99	13.49	7.71	8.45	6.83	15.07	21.04	9.00	6.35	8.12	45.57	46.69	17.57	24.51	8.20	6.83	20.92	
93 - 183	1721	1721	329	0.42	0.36	0.00	0.07	0.00	1.76	0.00	0.73	1.29	0.00	0.00	1.19	3.09	0.41	2.05	1.06	12.56	0.00	18.94	8.35	0.72	6.82	6.41	4.63	0.00	1.96	
	1047	1047	332	14.59	1.08	11.97	5.65	2.23	7.74	31.73	1.32	1.70	11.55	3.77	2.38	8.02	5.60	10.50	14.31	26.99	4.92	17.03	3.47	9.67	1.97	23.41	1.96	3.37	6.69	
	948	948	337	10.22	1.45	7.32	4.32	16.35	3.23	3.78	2.47	3.67	7.50	2.64	0.51	1.62	2.70	0.88	13.20	1.46	4.42	12.21	2.70	7.59	1.21	2.52	1.15	1.70	0.77	
	585	585	339	14.07	2.78	8.10	1.48	9.22	23.75	0.00	5.98	3.25	4.27	4.20	23.95	5.68	46.66	23.04	13.30	13.18	14.26	29.88	33.47	29.32	10.96	3.98	15.83	6.08		
	474	474	354	19.81	0.36	4.85	1.16	3.22	2.94	71.28	3.30	3.08	1.58	11.75	3.95	7.21	14.83	6.72	4.85	7.75	10.64	4.69	20.24	8.34	4.79	1.20	4.52	8.15	1.00	
184 - 274	151	147	333	10.65	0.52	1.07	1.46	4.43	1.24	0.19	0.29	1.65	0.18	0.99	0.84	2.38	0.01	0.17	1.17	0.15	0.10	0.25	0.32	0.71	0.01	0.16	0.07	0.31		
	121	121	336	4.92	9.10	4.57	17.93	0.78	2.08	1.93	1.13	1.14	4.03	1.88	2.20	1.36	0.58	0.33	2.10	0.15	3.16	8.54	1.34	1.06	0.51	0.76	1.05			
	103	103	355	35.07	6.59	8.44	1.76	1.16	24.22	0.45	0.99	7.75	2.48	0.35	0.45	0.46	1.50	0.18	0.39	1.20	5.10	1.61	1.42	1.09	2.88	0.18	0.13	0.56		
275 - 366	92	96	334	1.93	1.26	0.00	0.75	1.34	0.28	0.41	0.11	0.52	0.40	1.03	0.67	0.60	0.01	1.24	0.00	0.03	0.00	0.76	0.18	0.32	0.29	0.63	0.02	0.94		
	58	58	335	24.31	3.09	3.20	3.76	2.23	0.10	2.89	0.01	0.17	2.92	1.00	0.37	1.07	0.09	0.67	0.36	0.35	0.10	0.72	0.00	0.02	0.00	0.85	0.00	0.10	0.09	
	61	61	356	1.35	0.81	51.23	11.66	0.84	7.14	0.38	0.50	3.80	2.67	0.88	0.36	0.70	0.21	0.89	0.00	0.01	0.00	1.14	0.07	1.01	0.52	0.35	0.00	0.00		
367 - 549	93	166	717	2.31	0.00	2.50	2.87	0.53	1.83	11.37	0.00	0.58	0.50	2.38	0.40	0.11	0.61	0.40	4.45	1.76	0.59	1.62	2.30	2.59	0.74	0.33				
	76	76	719	10.53	0.23	6.24	0.58	0.13	21.58	1.78	0.85	0.93	1.29	2.80	0.56	1.46	0.32	0.55	0.92	0.35	0.79	1.52	0.40	0.75	0.02	1.12	0.66	1.35		
	76	76	721	1.69	16.19	6.39	2.02	5.15	0.54	1.32	6.43	1.59	0.20	1.35	1.62	0.23	1.42	0.28	2.83	1.05	0.09	0.67	0.80	1.25	0.15	0.29	0.11			
550 - 731	111	134	718					1.45	4.43	0.52	3.70	2.55	2.90	1.83	2.73	2.94	8.71	2.63	7.06	3.69	8.78	4.36	5.98	3.44	2.73	0.59	5.16	1.25		
	105	105	720					5.02	0.00	0.91	4.68	0.12	1.15	0.24	5.72	1.78	2.16	0.70	2.67	0.04	0.09	0.82	0.08	0.26	0.72	0.00	1.41	4.37	1.15	
	93	93	722	0.69	6.30	1.62	1.13	3.03	0.91	0.91	2.05	0.66	1.16	0.38	0.58	1.07	2.24	0.62	0.69	1.32	1.16	0.87	0.34	0.61	1.00	0.88	0.09			
732 - 914	.	105	764								5.21	0.80	1.43	2.50				0.26		0.79		2.81								
	.	99	768								1.34	0.49	1.35	2.80				0.27		0.06		0.38								
	.	135	772								9.29	3.33	2.65	1.54				2.68		1.19		1.41								
915 - 1097	.	124	765								1.40	0.21	1.18	3.25				0.59		0.65		1.48								
	.	138	769								0.92	0.26	1.45	3.13				1.08		0.83		1.39								
	.	128	773								0.23	0.73	1.80	5.08	0.71			0.45		0.55		0.27								
1098 - 1280	.	144	766								1.21	0.13	1.85				2.90		1.23		1.48									
	.	128	770								0.23	1.29	3.79				0.73		0.90		0.13									
	.	135	774								0.22	1.65	0.83	1.46				2.33		0.21		0.00								
1281 - 1463	.	158	767								0.68	0.00	0.00				0.00		0.13		0.11									
	.	175	771								0.00	0.73	0.00				0.41		0.00		0.00									
	.	155	775								0.00	0.00	0.00	1.29			0.98		0.13		0.60									
Grand Total				5.96	1.62	2.80	3.08	2.76	4.10	4.89	1.46	1.64	5.04	2.30	2.03	3.47	3.35	6.47	5.37	5.333	2.248	6.29	8.96	9.30	5.83	5.33	3.56	2.56	2.71	



Table 22. Estimated biomass (tonnes) of Witch flounder (M+F) in each stratum from Spanish rv surveys in Div. 30 during spring of 1984-2016. (Engel 145 data converted to Campelen Units for 1984-95).

Strata	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
357	5	439	12	33	114	227	615	0	249	139	24	31	12	46	82	19	29	15	6	8	13	72
358	797	595	114	51	108	437	2156	52	110	123	181	63	139	220	82	185	60	150	52	69	89	1000
359	155	118	190	511	259	3137	3249	26	64	400	44	209	54	434	42	78	157	386	388	62	630	142
360	328	391	190	234	2216	5061	724	38	75	598	456	1014	242	256	0	745	280	933	323	36	82	0
374	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
375	0	0	0	0	7	0	0	0	0	0	0	0	0	0	0	0	0	6	0	0	0	0
376	0	21	0	22	78	140	16	4	0	30	31	40	6	0	0	0	0	1	12	0	0	0
377	0	0	0	0	0	1	0	1	0	5	0	4	0	2	0	0	0	0	4	0	7	0
378	6	0	5	6	0	4	5	0	0	8	0	3	5	9	0	0	0	2	6	11	35	3
379	18	0	14	15	2	33	12	12	0	0	3	1	1	4	6	7	2	6	0	8	3	5
380	n.s.	0	0	2	15	2	1	2	0	3	1	1	2	0	0	7	2	6	3	10	6	10
381	n.s.	4	0	0	0	0	0	0	0	0	5	0	0	0	0	0	22	44	13	0	15	0
382	n.s.	0	0	0	0	2	0	0	0	0	5	0	0	0	0	0	0	0	0	16	6	0
723	n.s.	81	236	200	476	403	184	25	106	55	24	83	37	46	79	158	84	21	33	64	63	38
724	1200	181	49	245	534	757	278	56	133	223	67	40	258	224	167	109	88	165	95	41	86	77
725	108	22	11	27	73	646	35	6	2	173	67	33	60	16	45	29	18	14	46	21	65	93
726	418	14	0	87	205	1103	327	19	0	64	28	22	46	36	408	36	48	23	120	28	19	40
727	n.s.	5	0	9	23	5	0	0	0	41	29	6	12	53	29	94	30	53	86	22	7	101
728	n.s.	51	5	48	329	167	34	8	37	18	8	8	67	11	77	142	55	120	56	75	81	156
752	n.s.	276	86	106	196	459	390	5	59	4	0	0	0	0	8	2	14	21	55	86	115	160
753	n.s.	391	322	311	583	996	339	9	4	0	0	0	0	0	0	n.s.	13	0	8	14	10	0
754	n.s.	n.s.	27	19	0	96	353	3	3	0	0	0	0	0	0	0	0	0	0	0	0	0
755	n.s.	n.s.	0	0	0	4	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
756	n.s.	16	66	316	730	749	1565	10	40	33	25	31	41	77	154	228	76	43	46	82	46	152
757	n.s.	359	132	987	174	487	358	50	16	0	0	0	0	0	22	36	15	15	45	50	29	4
758	n.s.	n.s.	108	54	0	15	18	2	0	0	0	0	0	0	0	6	0	0	0	0	0	0
759	n.s.	n.s.	0	7	5	20	6	8	0	0	0	0	n.s.	0	0	0	0	0	0	0	0	0
760	n.s.	602	207	1262	1894	1853	1604	134	267	127	223	104	170	227	188	48	125	107	89	43	221	41
761	n.s.	1789	1219	1826	193	1243	816	284	91	23	81	99	1	0	17	146	81	93	35	13	37	14
762	n.s.	n.s.	55	0	1001	209	159	0	88	141	82	14	n.s.	0	0	20	0	0	0	0	8	17
763	n.s.	n.s.	0	8	51	136	12	0	0	13	0	0	n.s.	2	n.s.	n.s.	0	0	0	0	0	0
TOTAL	3033	5354	3049	6386	9265	18390	13255	752	1342	2223	1384	1805	1153	1663	1406	2095	1198	2224	1539	749	1667	2126



Table 23. Yield (t) and risk of  $F > F_{lim}$ ,  $B < Blim$  and  $B < B_{MSY}$  for projected  $F$  values of  $F2016$ ,  $75\% F2016$ ,  $125\% F2016$  2/3 FMSY,  $75\% FMSY$ , and  $85\% FMSY$  for witch flounder in NAFO Divs. 3NO.

	Yield 2018	Yield 2019	p> $F_{lim}$		p< $B_{lim}$			p< $B_{MSY}$			p2020>2016
			2018	2019	2018	2019	2020	2018	2019	2020	
$F2016=0.04$	1116	1175	24%	25%	19%	18%	17%	79%	76%	73%	65%
$75\%F2016 =0.03$	844	891	15%	16%	19%	18%	20%	80%	75%	72%	66%
$2/3 Fmsy=0.05 =125\%F2016$	1316	1384	31%	32%	19%	18%	19%	79%	76%	73%	63%
$75\%Fmsy=0.052$	1468	1555	36%	37%	18%	19%	19%	79%	76%	73%	62%
$85\% Fmsy=0.06$	1662	1745	42%	43%	19%	19%	20%	80%	77%	74%	60%
$F=0$					18%	16%	14%	79%	77%	70%	72%

Table 24. Medium-term projections for witch flounder. Estimates and 80% confidence interval for yield and relative biomass  $B_y/B_{msy}$ , are shown, for projected F values of F2016, 75% F2016, 125% F2016, 2/3 Fmsy, 75% Fmsy and 85%Fmsy.

Projections with catch in 2017 = 2 225 t		
	Projected Yield (t)	Projected Relative Biomass ( $B_y / B_{msy}$ )
$F_{2016}=0.04$	Median	Median (80% CI)
2018	1126	0.57 (0.22, 1.35)
2019	1175	0.61 (0.22, 1.47)
2020		0.64 (0.21, 1.57)
75% $F_{2016}=0.03$	Projected Yield (t)	Projected Relative Biomass ( $B_y / B_{msy}$ )
2018	844	0.57 (0.22, 1.35)
2019	891	0.61 (0.22, 1.47)
2020		0.65 (0.22, 1.59)
2/3 $F_{msy}=0.05$ (=125%F2016)	Projected Yield (t)	Projected Relative Biomass ( $B_y / B_{msy}$ )
2018	1316	0.57 (0.22, 1.33)
2019	1384	0.60 (0.21, 1.44)
2020		0.63 (0.20, 1.56)
75% $F_{msy}=0.052$	Projected Yield (t)	Projected Relative Biomass ( $B_y / B_{msy}$ )
2018	1468	0.57 (0.22, 1.35)
2019	1555	0.60 (0.21, 1.46)
2020		0.62 (0.20, 1.56)
85% $F_{msy}=0.06$	Projected Yield (t)	Projected Relative Biomass ( $B_y / B_{msy}$ )
2018	1662	0.57 (0.22, 1.34)
2019	1745	0.59 (0.20, 1.45)
2020		0.62 (0.19, 1.54)
$F=0$	Projected Yield (t)	Projected Relative Biomass ( $B_y / B_{msy}$ )
2018	0	0.57(0.22, 1.34)
2019	0	0.59(0.20, 1.45)
2020		0.69(0.25, 1.65)




Fig. 1. Commercial catch of witch flounder in NAFO Divs. 3NO from 1960-2016 and total allowable catch (TACs).




Fig. 2. Witch flounder length frequency (cm) distributions for Spain, Russia, and Portugal, and Canada commercial fisheries in NAFO Divs. 3NO in 2016.




Fig. 3. Biomass (tons), abundance (millions) for witch flounder from Canadian spring rv surveys in NAFO Divs. 3N and 30 during 1984-2016. Vertical lines represent 95% confidence intervals. White markers represent years where >50% of deep-water strata were surveyed. No data was available for spring 2006.




Fig. 4. Mean numbers and weights (kg) per tow for witch flounder from Canadian spring rv surveys in NAFO Divs. 3N and 3O during 1984-2016. Vertical lines represent 95% confidence intervals. White markers represent years where >50% of deep-water strata were surveyed. No data was available for spring 2006.




Fig. 5. Biomass (tons), abundance (millions) for witch flounder from Canadian fall surveys in NAFO Divs. 3N and 3O during 1990-2016 (no data was available for fall 2014). Vertical lines represent 95% confidence intervals. White markers represent years where >50% of deep-water strata were surveyed.




Fig. 6. Mean number per tow and mean weight (kg) per tow for witch flounder from Canadian fall surveys in NAFO Divs. 3N and 3O during 1990–2016 (no data was available for fall 2014). Vertical lines represent 95% confidence intervals. White markers represent years where >50% of deep-water strata were surveyed.




Fig. 7. Biomass (tons), abundance (millions) for witch flounder from Canadian spring and fall surveys in NAFO Divs. 3NO combined during 1984-2016 (no data was available for fall 2014 or spring 2016). Vertical lines represent 95% confidence intervals. White markers represent years where >50% of deep-water strata were surveyed.




Fig. 8. Mean number per tow and mean weight (kg) per tow for witch flounder from Canadian spring and fall surveys in NAFO Divs. 3NO combined during 1984-2016 (no data was available for fall 2014 or spring 2016). Vertical lines represent 95% confidence intervals. White markers represent years where >50% of deep-water strata were surveyed.




Fig. 9. Distribution of witch flounder (weight (kg) per tow) from Canadian spring surveys in NAFO Divs. 3NO during 2013 to 2016.




Fig. 10. Distribution of witch flounder (weight (kg) per tow) from Canadian fall surveys in NAFO Divs. 3NO during 2013 to 2016 (note there was no fall survey in 2014).




Fig. 11. Length frequency distributions of witch flounder from Canadian spring and fall surveys (1997-2016) using the Campelen 1800 shrimp trawl. Estimates represent abundance at length (cm) of the surveyed area. All distributions are for NAFO Divs. 3NO.




Fig. 12. Length frequencies (abundance at length) of witch flounder from spring Canadian (2000-2016), fall Canadian (2001-2016), and Spanish (2001-2016) rv surveys in NAFO Divs.3NO. Canadian rv survey data was not available in spring 2006 or fall 2014.




Fig.13. Recruitment index means proportioned to 1 of witch flounder <21cm in length from spring and fall Canadian rv surveys in NAFO Divs. 3NO 1995-2016. Note there was no fall survey in 2014 or spring survey in 2006.




Fig. 14. Witch flounder stratified mean catches (kg) and mean number by year. Spanish spring surveys in NAFO Divs. 3NO: 2002-2016.




Fig. 15. Witch flounder biomass calculated by the swept area method in tons  $\pm$ SD by year. Spanish spring surveys in NAFO Divs. 3NO: 2002-2016.




Fig. 16. Witch flounder length-frequencies (cm) for Canadian and Spanish spring rv surveys in NAFO Divs. 3NO: 2013-2016.




Fig. 17. Relative biomass (biomass divided by BMSY) for Div. 3NO witch flounder. The median with its 80<sup>th</sup> percent credible intervals are shown. The horizontal red dashed line is Blim (30% BMSY).




Fig. 18. Relative fishing mortality (fishing mortality divided by FMSY) for Div. 3NO witch flounder. The median with its 80th percent credible intervals are shown. The horizontal red dashed line is Flim (FMSY)