

NOT TO BE CITED WITHOUT PRIOR  
REFERENCE TO THE AUTHOR(S)



Northwest Atlantic

Fisheries Organization

**Serial No. N6923**

**NAFO SCR Doc. 19/008**

### **SCIENTIFIC COUNCIL MEETING – JANUARY 2019**

Biomass and Abundance of Demersal Fish Stocks off West and East Greenland estimated from the Greenland Institute of Natural resources (GINR) Shrimp and Fish Survey (SFW), 1990-2018.

By

Rasmus Nygaard and Adriana Nogueira  
Greenland Institute of Natural Resources, Box 570, 3900 Nuuk, Greenland

#### **Abstract**

Since 1988, the Greenland Institute of Natural Resources has annually conducted a bottom trawl survey off West Greenland (NAFO SA1). The survey was initially designed with the focus to evaluate the biomass and abundance of the Northern shrimp (*Pandalus borealis*), but fish catches have systematically been recorded since 1992. The gear was changed prior to the 2005 survey from a shrimp trawl with steel bobbins to a slightly larger shrimp trawl with rock-hoppers. In 2008, the survey was expanded to include The East Greenland shelf area using the same gear and depth range. This paper contain biomass and abundance indices, length distributions and maps of survey densities for Greenland halibut (*Reinhardtius hippoglossoides*), Atlantic cod (*Gadus morhua*), redfish (*Sebastes norvegicus* and *Sebastes Mentella*), Atlantic wolffish (*Anarhichas lupus*), Spotted wolffish (*Anarhichas minor*), American plaice (*Hippoglossoides platessoides*) and Thorny skate (*Amblyraja radiata*). The appendix contains total biomass and abundance estimates for Teleosts, Elasmobranchs and Cephalopods. In 2018, the survey was completed in West Greenland. In East Greenland, the survey was limited to shallow water in 2017, and no survey has been conducted in 2018. In 2018, the annual trawl survey was conducted in West Greenland with a chartered vessel, the Faroese trawler Sjurdarberg. All the standard gear from the research vessel Paamiut (such as cosmos trawl, doors, all equipment such as bridles ect, Marport sensors on doors and headlines) were used, in attempt to make the 2018 survey identical as possible with the previous years survey. No survey was conducted in East Greenland.

In West Greenland, the Greenland halibut biomass increased until 2004. After the Gear change in 2005, the biomass index has been in a decreasing trend, but slightly increasing since 2013. The shelf in division 1C-A is an important nursery area for Greenland halibut and in all years new Year Classes are present. The 2016 survey indicated a major decline in the offshore cod stock in West Greenland (NAFO 1A-1E) with 80% in abundance and 86% in biomass compared to 2015 and this was further



confirmed in 2017 and 2018. The biomass of golden redfish and deep-sea redfish in West Greenland, have increased in the past few years. The abundance of redfish, mainly juvenile redfish, is a mixture one or both specie, has decreased about a factor 20 since the beginning of the time series, indicating continued failing redfish recruitment in the area. Both the East Greenland and West Greenland shelf (1AB) are known nursery grounds for redfish, but new YC have virtually been absent in the since 2011 in both West Greenland and in East Greenland. In West Greenland abundance and biomass indices of spotted wolffish, Atlantic wolffish, American plaice and thorny skate have been in an increasing trend both before and after the gear change.

## Materials and Methods

### The Greenland Shrimp and Fish survey in West Greenland (SFW survey)

The survey has been conducted with the 722 GRT stern trawler M/Tr 'Pâmiut' since 1991, similar vessels were used in 1988-1990. However, in the beginning of 2018 it was decided that the old research vessel Paamiut has to be scraped owing to increasing expenses to maintenance. Instead, the survey was conducted with a chartered fishing vessel Sjúðarberg. All fishing gear were removed from Paamiut and installed on Sjúðarberg and fishing practice and handling of catch were exactly as used on the research ship Paamiut (Burmeister and Rigét, 2018) (Appendix I). The survey design, the area coverage and the trawl and its rigging has been unchanged since 2005. The survey period is June – July in West Greenland. For a more thorough description of swept area calculations and survey design see Burmeister and Rigét, 2018.

**The Survey Gear and trawl:** The survey initially used a Skjervoy 3000/20 trawl with steel bobbin gear and double bag. In 2005, the skjervoy trawl was replaced by a Cosmos trawl (Wieland and Bergström, 2005). Until 2003, *Greenland Perfect* trawldoors were used ( $9.25 \text{ m}^2$ , 2.4 tons), but they were replaced in 2004 by Injector International trawl doors ( $7.5 \text{ m}^2$ , 2.8 tons) to facilitate the trawl change in 2005. Calibration experiments were conducted in the main shrimp areas in 2004 and 2005 and a formal analysis of conversion factors were established for shrimp (Rosing and Wieland, 2005). Preliminary conversion factors for a few commercial fish species were derived as described by Rosing and Wieland (2005) and are given in table 3. However, the calibration factors were never finally evaluated and in some cases (particularly cod) the calibration factors relied on few data. Without calibration the two separate time series seems well connected for most species indicating that the trawl experiments overestimate the difference between the gears.

**Survey area and stratification:** The trawl survey initially covered the traditional offshore shrimp area, between  $60^\circ$  -  $72^\circ$  north, depth 150-600m. In 1991, the area was extended to include Disko Bay. The area is delimited by a line 3nm off the base line and the 600 m depth curve. Areas shallower than 150 m was initially rather unsystematically covered, but from 2004 two extra depth zones have been formally included (50-100m, and 100-150m).

The stratification is based on designated 'Shrimp Areas' that is divided into depth zones of: 151-200, 201-300, 301-400 and 401-600 m, as based on depth contour lines (figure. 1). The depth zones 0-100 m and 100-150m are delimited by the NAFO Subdivision boundaries. The "shrimp Areas" and their sizes are provided in table 1. The number of valid hauls by year and strata are listed in table 2. Fish species was prior to 2007 analysed using a re-stratification that followed the NAFO divisions. Re-stratification implies a potential bias and the survey information from 2005 and onwards has therefore been reanalysed in accordance with the shrimp strata actually used in the survey. If strata had no stations in a given year, the neighbouring strata with stations in that year is geographically

enhanced to include of the non-visited strata. This way the total area surveyed is maintained (since 2005). Tow duration was over the years gradually reduced from 60 min. (prior to 1997) to 30 and has been fixed to 15 min since 2005 (Wieland and Storr-Paulsen 2006). Towing speed has been about 2.5 knots throughout the years. Survey abundance and biomass is expressed per swept area: Wingspread\*towed distance, where wingspread is inferred from Scanmar recordings and the towed distance is measured by GPS.

**Allocation of stations per strata.** From 1988 to 1998, the survey followed a stratified random design where stations were allocated to strata in proportion to the stratum area. In 1999, the placing of stations independently and randomly was replaced by a buffered random sampling design. The method combines the use of a minimum between stations-distance rule (buffer zero) with a random allocation scheme (Kingsley et al., 2004, Nogueira et al., 2018). Hauls are allocated to strata based on the stratum area and on predictions, from past surveys, of the variability of the catch. About 50 % of the stations included in the preceding year's design have been repeated as fixed stations in the following year, the others being placed, using the buffered sampling rules (Burmeister and Riget, 2018).

### The Greenland Shrimp and Fish survey in East Greenland (SFE survey)

The survey is carried out with the same gear and survey protocols as used in West Greenland after 2005. Stratification is based on the "Q-areas" used for the East Greenland survey for Greenland halibut. The areas are further depth stratified into 0-200 m, 200-400m and 400-600 m zones, the areas are shown in figure 1 and the area sizes are given in table 1. In East Greenland, bottom conditions severely restrict the areas that can be trawled and in some strata, stations are randomly selected from historical known trawl-able sites. No survey has been carried out in 2018.

## Results

### Greenland halibut (*Reinhardtius hippoglossoides*).

Greenland halibut is widely distributed along NAFO 1A-F, but highest concentrations are found in the important nursery areas in division 1A, 1B-north and Disko Bay (table 4 and 5). The abundance indices is mainly driven by year to year variability in the number of one- and two-year old recruits, which by number typically constitute 80-90% of the catch.

The biomass and abundance indices increased gradually until the gear change in 2005. Since 2005, the general trend in the abundance indices has been decreasing until 2014, with the exception of record high numbers of one-year old recruits observed in 2011, 2013. Since then, the abundance has fluctuated reaching picking in 2011 and 2013 with the highest values of the time series. From 2017 to 2018 the biomass decreased (Figure 2). In West Greenland clear modes can be found in the length distribution at 12-15 and 23 cm every year corresponding to year-classes 1 and 2 (figure 14).

### Greenland halibut recruitment.

A recruitment index was estimated for the entire survey area. By means of the Petersen-method ages 1, 2 and 3+ were separated in the survey catches. To allow comparison of abundance throughout the time series, data from 1988 to 2004 were converted by a conversion factors to adjust the old Skjervoy to the new Cosmos trawl catches trawl catches. For Greenland halibut the calibration was length dependent (Table 3).

The number of one-year-old fish in the total survey area including Disko Bay (NAFO Divisions 1A-F ) increased gradually from 1996 to a peak in 2003 (470 millions indiv.)(Fig. 9). Since then, the

abundance decreased until 2014, with the exception of two exceptionally recruitment events of more than 1000 millions of indiv. in 2011 and 2013. The recruitment decreased from 2017 (449 millions indiv.) to 2018 (170 millions indiv.). Almost all the one year old fish were found in Div. 1AS and Div. BN (Fig. 10).

The recruitment offshore fluctuated since the good 2011 and 2013 year-classes. In 2011, it peaked in North Division 1A and Inshore, and in 2013, in South 1A and 1B. In 2014, the recruitment decreased in all divisions, and since then it increased (Fig. 11). More than 90 % of the one year old fish was found in the off shore areas (Fig.12).

Generally there is a steep decline between abundance at age 1 and age 2 and 3+ which also was observed in the 2017 survey (Fig. 13).

### **Cod (*Gadus morhua*)**

The Atlantic cod (*Gadus morhua*) stock complex in Greenland is considered to be composed of primarily three different stock components; inshore cod, West Greenland Cod and East Greenland/Iceland offshore cod. **Inshore cod** are believed to be relatively stationary, as tagging experiments showed that most (82-86 %) of the cod recaptured were found in the same area as tagged (Hovgård and Christensen 1990). **West Greenland offshore** cod that historically has been spawning along the banks of the West Greenland's coast (Wieland and Hovgaard 2002, Therkildsen et al. 2013). Larvae from the East Greenland/Iceland cod in some years drift with the current to West Greenland and gradually moves to Eastgreenland before reaching maturity. The 2016 survey indicated a major decline in the offshore cod stock in West Greenland (NAFO 1A-1E) with 80% in abundance and 86% in biomass compared to 2015 and the index and this was further confirmed in 2017 and 2018. For more information on cod see the ICES Report of the North-Western Working Group (Anon., 2019)

### **Demersal Redfish (*Sebastes sp.*) combined.**

Two species of redfish are common in the area, golden redfish, *Sebastes norvegicus* and deep-sea redfish *Sebastes mentella*. Both the East Greenland and West Greenland shelf (1AB) are known nursery grounds for redfish. However, the distinction between the species and the normal dominance of recruits in the survey, has had the effect that in some years redfish were classified as *Sebastes sp.* (prior to 2007). After 2007 redfish smaller than approximately ~18 cm has been classified as juvenile redfish *Sebastes sp.* and larger redfish are classified on a species level. A separation of redfish by species has been attempted since 2007 (table 10-15, figure 4b-d and 16).

In East Greenland, the survey has revealed a substantial decrease in the abundance index from 2008-2016 in redfish <20 cm indicating poor recruitment since 2011. The index was not evaluated in 2017 due to the low number of stations and lag of stations in the deeper strata and no survey was conducted in 2018.

The biomass of golden redfish and deep-sea redfish in West Greenland have increased in the past few years but the indices is about one order of magnitude lower than in East Greenland (fig 4a). The indices for redfish is often somewhat uncertain with a single or few hauls provide the majority of the annual biomass estimate. The increasing biomass observed in West Greenland during a period of decreasing or failing recruitment could be related to either connection with the stocks in East Greenland or perhaps increased survival of redfish since the implementation of sorting grids in the shrimp-fishery in 2002. Indeed, tow densities by haul reveal an almost continuous distribution of both species from East to West around Cape farewell (fig 23-26).

For further information on redfish in East Greenland see the ICES Report of the North-Western Working Group (Anon., 2019)

## **Redfish recruitment.**

The high numbers of redfish in the survey in the 1990's were mainly recruits concentrated in division 1AS and 1B. The combined abundance index for both redfish species can be viewed as a continuous decrease in the number of recruits since the 1990's (fig 4a). The decreasing numbers of recruits has continued in both East Greenland and West Greenland since 2008 with a almost complete lack of recruits since 2011 (fig 4b and 16). Annual growth increments of 4 cm are indicated by repeatedly pronounced peaks in length compositions at 7-8 cm and around 12 cm probably corresponding to age 1 and 2 (Nedreaas, 1990).

## **Deep-sea redfish *S. mentella***

In East Greenland, the deep-sea redfish abundance decreased substantially from 2008 to 2016, but a gradual shift in the length distribution from around 18-30 to 30-45 cm maintained the biomass at a higher level (fig 4c). In West Greenland, the abundance and biomass of deep-sea redfish increased from 2011 to 2016 (fig 4c). However, the high 2016 biomass index was caused by a single haul in division 1D of large redfish between 25 and 40 cm. In 2017 and 2018, there were no such large hauls in the survey, but the indices remain at a higher level.

## **Golden Redfish *S. Norvegicus***

In East Greenland, the biomass index for Golden redfish increased from the 2008-2010 period and remained higher until the most recent survey year in 2016 (fig 4d).

In West Greenland, golden redfish biomass was stable from 2006-2010 but increased gradually until 2016 (table 15 and fig 4d). The 2017 and 2018 biomass indices are however close to the 2006-2011 level. The increasing biomass observed from 2011-2016 occurred division 1E and 1F and was often caused by one or 2 hauls containing larger individuals (figure 26) contributing more than half the total West Greenland biomass. In 2016, a single haul in division 1E consisted of large golden redfish between 45-70 cm and provided 80% of the total biomass estimate.

## **American plaice (*Hippoglossoides platessoides*).**

In West Greenland, American plaice is common in all divisions (table 16, 17 and figure 27). The biomass and abundance indices have fluctuated substantially in recent years, but the general trend has been increasing both prior to and after the gear change (table 16, 17 and figure 5). Clear modes can be found at 5 and 15 cm indicating new incoming year-classes and individuals larger than 45 cm are rarely seen in Greenland (figure. 17). The highest concentrations are mainly found in West Greenland (figure 27). In East-Greenland, both abundance and biomass indices are much lower on the shallow parts of the East Greenlandic shelf areas.

## **Atlantic wolffish (*Anarhichas lupus*)**

Atlantic wolffish is common in all divisions, in both East and West Greenland (figure 28). Previously Atlantic wolffish had its main distribution south of 68°N, but it has shifted further north since the beginning of the time series (table 18, 19 and figure 6). Although the abundance and biomass indices tends to have extreme values in some years, the underlying trends has been increasing, since the beginning of the time series and the non-calibrated indices seems well connected. The length distribution reveals the dominance of smaller fish (figure. 18).

### **Spotted wolffish (*Anarhichas minor*)**

Spotted wolffish are common in all divisions in both East and West Greenland, inshore and offshore. Abundance and biomass indices have fluctuated with an increasing trend throughout the time series and the non-calibrated indices seems well connected (Table 20, 21 and figure 7). The length distribution ranges from 10-120 cm and modes at 13 and 100 cm likely represents one-year old recruits and individuals at  $L_{max}$  (figure 19). Recruits and juvenile individuals are mainly found in North-west Greenland (1A) (figure 19 and 29).

### **Thorny skate (*Amblyraja radiata*)**

In West Greenland, thorny skate is common in all divisions, but the majority of both the biomass and abundance is located in West Greenland (fig 8 and 30). Abundance and biomass indices have increased throughout the time series and the non-calibrated indices seems well connected (table 22, 23 and figure 8). Although  $L_{max}$  for thorny skates is reported to be more than 100 cm, it has been suggested that North American thorny skate grows to larger body size than East Atlantic individuals. Since individuals larger than 55 cm are rarely seen in Greenland waters and all individuals about 50 cm are fully mature, it seems likely that thorny skates in Greenland resemble East Atlantic stocks. During the most recent years, thorny skate length distributions have revealed clear modes at 10-15 cm and 35-50 cm probably corresponding to recruits and overlapping year-classes of adult individual (figure. 20).

### **Other species**

The Appendix II contains biomass and abundance estimates for elasmobranchs, teleosts, cephalopods and crustaceans excl. Shrimp for the West-Greenland part of the GINR shrimp fish survey including the West-Greenlandic Shelf part of NAFO div 0A.

### **Discussion**

Catch-ability is set at 1 for all species. However, since swept area is calculated for the trawl excluding doors and bridles, catchability may be higher than 1 for some species and below 1 for other species, implying that both biomass and abundance should be regarded as index values only, not absolute values.

Index changes from the end of the 1990's to 2001 for species related to shallow water and banks (50-150m) could reflect better coverage of these depths during the past decade. Changes from 2003 to 2004 could be influenced with the trawl door update from Greenland perfect to injector and may be species dependent, as no calibration experiments were made prior to the change. Index changes from 2004 to 2005 could be related to species specific data quality in the calibration experiments. If ignoring the calibration factors, the indices for most species seems better connected, indicating that there is little reason to calibrate in order to compare the time series. Indeed, even for Greenland halibut the length-dependent calibration factor has a 1:1 value around 12 cm which constitutes 80-90% of the individuals caught every year. Therefore, for the Greenland halibut abundance indices it matters little whether calibrating or not.

Since the main purpose of the survey is to evaluate the biomass of northern shrimp and the effort is concentrated in areas and depths where the commercial shrimp trawling is taking place, especially on the northern slopes of the bank Store Hellefiskebanke (67°50N 55°00W) and in the inshore area Disko Bay. As Store Hellefiskebanke and Disko Bay are important nursery areas for Greenland halibut and redfish, as well as other important species (Smidt, 1969; Tåning, 1949) it is likely, that the abundance estimates of the survey reflects the juvenile stock situation of these species.

## References

- Anon., 2016. Report of the North-Western Working Group (NWWG). *ICES CM 2016*.
- Burmeister, A. and Rigét F. The West Greenland trawl for *Pandalus borealis*, 2018, with reference to earlier results. NAFO Scientific Research Document 18/055.
- Cochran, W. G. 1977: Sampling Techniques, Third edition, Wiley & Sons.
- Kingsley, M.C.S., P. Kanneworff and D.M. Carlsson. 2004. Buffered random sampling: a sequential inhibited spatial point process applied to sampling in trawl survey for northern shrimp *Pandalus borealis* in West Greenland waters. *ICES J. Mar. Sci.* 61:12-24.
- Nedreaas, K. 1980: Age determination of Northeast Atlantic *Sebastes* species. *J. Cons. int. Mer.* 47: 47: 208-230.
- Nogueira, A and Jorgensen O. 2018. Results for Greenland halibut survey in NAFO Divisions 1C-1D for the period 1997-2017. NAFO Scientific Research Document 18/021.
- Rosing, M.& K. Wieland (2005): Preliminary results from shrimp trawl calibration experiments off West Greenland (2004, 2005) with notes on encountered design/analyses problems. NAFO SCR Doc. 05/92.
- Smidt, E.L.B., 1969: The Greenland Halibut *Reinhardtius hippoglossoides* (Walb.), Biology and Exploitation in the Greenland Waters. Meddelelser fra Danmarks Fiskeri- og Havundersøgelser, N.S.,6: 79-148.
- Sünksen, K. 2007: Discarded by-catch in shrimp fisheries in Greenlandic offshore waters 2006-2007. NAFO SCR Doc. 07/88
- Tåning, Å.V., 1949. On the breeding places and abundance of the redfish (*Sebastes*) in the North Atlantic. *Ibid. Journ. Cons.* Vol.16 No.1: 85-96.
- Wieland, K.& M. Storr-Paulsen, 2006: Effect of tow duration on catch rates and mean length of Northern shrimp (*Pandalus borealis*) and Greenland halibut (*Reinhardtius hippoglossoides*) in the West Greenland Bottom Trawl Survey. *Fish. Res.* 78: 276-285.

**Table 1.** The survey area (km<sup>2</sup>) in the Greenland Shrimp and Fish Survey.

West Greenland							
Area	Depthstrata						Total
	<100	100-150	150-200	200-300	300-400	400-600	
1A	3039	5220					8259
1B	11346	4966					16312
1C	4183	8169					12351
1D	4136	1538					5673
1E	494	2721					3215
1F	1497	5248					6745
C0				903	2179	1154	4236
I1			407	1963	2441	1499	6310
I2			419	815	1085	1338	3658
U1			2486	4633	4785	5129	17033
U2				6710	8481	7994	23185
U3			2012	3017	1675	2710	9413
W1			2873	6099	7520	816	17307
W2			1674	2612	1741	915	6941
W3			2122	4725	2085	2994	11926
W4			4119	1818	821	1961	8719
W5			3001	3648	1950	3021	11620
W6			1206	2006	1585	1234	6031
W7			2442	891	265	317	3915
W8			424	567	405	718	2114
W9			1711	938	516	430	3595
All strata	24695	27861	24895	41344	37534	32230	188559

East Greenland				
Area	Depthstrata			Total
	0001-0200	0201-0400	0401-0600	
Q1	217	35445	6975	42637
Q2	93	7657	1246	8996
Q3	3363	22547	9830	35740
Q4	1337	7770	2054	11161
Q5	469	2785	1819	5073
Q6	6307	6130	2063	14500
All strata				118107

**Table 2.** Numbers of valid hauls in West Greenland, since 1988. 1AX=Disko Bay. 0A= The West-Greenlandic shelf part in Canadian waters. (- area included in neighboring strata).

Year	West Greenland											Vessel
	0A	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	Total	
1990	29	63	*	68	17	35	16	*	*	228	Pa	
1991	18	39	41	44	18	11	16	*	*	187	Pa	
1992	20	33	39	36	8	18	18	11	15	198	Pa	
1993	16	22	31	39	10	21	15	12	13	179	Pa	
1994	16	33	27	49	9	23	8	9	9	183	Pa	
1995	17	33	33	48	13	29	13	14	11	211	Pa	
1996	18	20	33	46	11	29	12	9	11	189	Pa	
1997	17	33	34	47	9	32	12	12	19	215	Pa	
1998	10	34	33	66	14	27	19	14	14	231	Pa	
1999	10	40	34	63	18	33	16	14	17	245	Pa	
2000	8	25	23	45	17	37	23	14	29	221	Pa	
2001	9	28	23	59	16	36	24	15	26	236	Pa	
2002	2	26	22	68	12	32	18	20	27	227	Pa	
2003	11	21	19	51	12	30	18	15	22	199	Pa	
2004	15	25	14	41	14	24	22	20	34	209	Pa	
New survey gear introduced												
2005	6	20	30	16	45	10	26	19	23	23	212	Pa
2006	5	26	40	21	49	9	27	20	21	31	244	Pa
2007	8	18	38	18	47	9	27	27	31	39	254	Pa
2008	6	16	38	16	53	7	28	23	25	46	252	Pa
2009	8	21	31	24	60	13	28	22	24	48	271	Pa
2010	10	26	44	25	65	11	30	23	24	40	289	Pa
2011	-	17	31	26	54	9	24	18	12	25	216	Pa
2012	-	18	34	21	52	12	21	18	18	26	220	Pa
2013	4	19	37	17	44	8	20	13	21	28	211	Pa
2014	-	21	36	21	47	10	19	17	23	32	226	Pa
2015	-	24	29	17	42	7	24	22	20	36	221	Pa
2016	-	18	29	12	32	6	27	14	19	36	193	Pa
2017	3	27	42	30	43	9	25	18	25	35	257	Pa
2018	20	38	20	33	9	26	23	20	35	20	Sj	

Note.

2005. Strata 1A 50-100 no stations – area included in 1A 100-150.

2006. Strata 1C 50-100 no stations – area included in 1C 100-150.

2007. Strata 1C 50-100 no stations – area included in 1C 100-150.

2008. All strata surveyed.

2009. Strata 1A:50-100 no stations – area included in 1A 100-150.

2010. Strata 1A:50-100, 1A:100-150 no stations – area included in U1 150-200.

2011. Strata W1:400-600, C0:200-300, C0:300-400, C0:400-600 no stations - included in W1 300-400.

2012. Strata C0:200-300, C0:300-400, C0:400-600 no stations – area included in W1 300-400.

2013. Strata C0:400-600 no stations - included in C0 300-400. Strata W9:400-600 no stations – area included in W9 300-400.

2014. Strata C0:200-300, C0:300-400, C0:400-600 no stations – area included in W3 equivalent depth strata.

2015. Strata C0:200-300, C0:300-400, C0:400-600 no stations – area included in W3 equivalent depth strata. W1:400-600 included in W1 300-400. Northern area restricted to Eastern shelf due to sea ice.

2016. Strata C0:200-300, C0:300-400, C0:400-600 no stations – area included in W3 equivalent depth strata.

2017. Strata C0:400-600 no stations - included in C0 300-400. Strata W1:400-600 no stations – area included in W1 300-400.

2018. Strata C0:200-300, C0:300-400, C0:400-600 no stations – area included in W3 equivalent depth strata. W1:400-600 included in W1:200-300.



**Table 2 - continued.** Numbers of valid hauls in East Greenland since 2007.

East Greenland							
Year	Q1	Q2	Q3	Q4	Q5	Q6	Total
2007		5	12	9		9	35
2008	8	6	12	7	7	12	52
2009	21	12	26	19	6	13	97
2010	19	14	24	9	6	10	82
2011	20	11	21	12	7	14	85
2012	19	16	28	13	7	15	98
2013	25	12	22	14	5	14	92
2014	22	14	12	9	8	16	81
2015	26	11	24	12	8	14	95
2016	29	10	26	13	7	16	101
2017	2	4	7	6	6	11	36
2018	-	-	-	-	-	-	-

## Note:

2007 Survey startup year. Only 35 hauls. No results presented.

2008. Strata Q1,Q2,Q4:0-200 no stations - included Strata Q1,Q2,Q3:200-400.

2009. Strata Q1:0-200 m no stations – area included in Q1:200-400.

2010. Strata Q1:0-200 m no stations – area included in Q1:200-400.

2011. Strata Q1:0-200 m no stations – area included in Q1:200-400.

2012. Strata Q1:0-200 m no stations – area included in Q1:200-400.

2013. Strata Q1:0-200 m no stations – area included in Q1:200-400. Strata Q5:0-200 m no stations – included in Q4:0-200.

2014. Strata Q1:0-200 m no stations – area included in Q1:200-400. Strata Q3:0-200 included in Q4:0-200 (COD Q3:200-400.)

2015. Strata Q1:0-200 m no stations – area included in Q1:200-400. Stations in Icelandic EEZ included under strata in Grl.

2016. Strata Q1:0-200 m no stations – area included in Q1:200-400.

2017. Insufficient survey coverage for most species.

2018. No survey.

**Table 3.** Preliminary calibration factors tTTto adjust the Cosmos trawl catches to the former Skjervoy trawl standard. For Greenland halibut and American plaice the conversion were length dependent and for those species x in the equations represents the individual fish length.

Fishspecies	Greenland halibut	Redfish	American plaice	Atlantic wolffish	Spotted wolffish	Cod	Thorny skate
Conversion factor	0.0404x+0.6527	2.4	-0.0825x + 5.3307	2.3	2.3	1.78	5.1
Trawl size factor	1/1.1516	1/1.1516	1/1.1516	1/1.1516	1/1.1516	1/1.1516	1/1.1516
Final factor	0.035x+0.567	2.0	-0.072x + 4.629	2.0	2.0	1.5	4.4

**Table 4.** Greenland halibut *Reinhardtius hippoglossoides* abundance indices (millions).

Year	0A	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	W-Grl	CI	E-Grl
1992		50.7	8.2	96.7	231.5	1.5	1.2	0.8	0.6	0.0	391.2	27	
1993		29.1	24.3	34.0	152.6	9.8	4.3	4.6	0.4	0.5	263.5	28	
1994		22.9	35.2	63.0	130.7	10.0	18.1	2.4	0.1	0.4	282.8	22	
1995		39.5	28.9	89.4	98.9	18.3	12.2	5.7	0.1	0.3	293.4	32	
1996		92.6	13.7	102.5	265.9	11.8	14.0	10.7	0.2	0.4	511.7	25	
1997		41.3	17.5	112.1	97.6	2.2	13.0	0.9	0.1	0.3	285.0	22	
1998		78.5	30.4	209.6	47.2	23.3	7.2	3.6	1.4	0.6	401.6	29	
1999		100.8	58.2	95.4	91.0	5.8	5.3	7.7	1.3	0.9	366.3	35	
2000		81.7	19.3	172.8	126.7	4.1	4.8	6.7	0.1	0.6	416.8	30	
2001		145.9	97.9	223.7	111.5	1.7	4.8	2.6	0.7	1.6	590.2	28	
2002		78.0	75.2	148.1	42.5	2.7	6.2	6.5	0.6	1.3	360.9	38	
2003		154.6	37.7	227.0	116.7	2.3	2.0	2.4	0.1	1.0	543.8	36	
2004		154.8	20.6	199.1	84.8	2.4	2.9	2.8	0.1	0.4	468.0	34	
New survey gear introduced											ecv	cv	
2005	3.3	177.5	51.6	186.5	202.0	5.3	9.1	1.9	1.1	0.5	638.8	12	
2006	14.1	110.5	69.9	96.9	198.6	4.2	24.6	0.8	0.0	0.3	520.0	11	
2007	10.5	103.1	75.3	128.5	154.4	2.2	37.9	0.3	1.2	0.4	513.8	13	-
2008	13.2	184.2	106.5	64.6	88.6	0.7	0.6	0.9	0.3	0.3	459.9	12	3.0
2009	8.0	132.8	75.3	72.9	119.6	6.9	0.7	0.9	0.3	0.1	417.5	8	4.1
2010	9.1	154.5	117.7	123.3	115.9	4.5	0.7	1.2	0.1	0.1	527.1	9	3.0
2011	-	318.8	111.6	230.7	93.4	8.9	1.1	0.9	0	0.2	765.6	16	2.9
2012	-	80.2	48.8	105.9	52.3	5.0	0.4	0.1	0	0.1	292.8	8	1.7
2013	1.0	148.0	212.0	188.0	145.0	11.0	5.0	0.0	0.0	0.0	711.0	16	1.7
2014	-	79.1	117.0	103.7	32.6	1.7	1.6	0.3	0.6	0	336.1	15	0.8
2015	-	77.7	101.7	84.1	131.1	3.6	2.4	1.5	0.2	0.2	402.4	11	0.5
2016	-	99.9	134.3	120.1	108.1	5.8	4.5	1.3	0.2	0.3	474.5	16	1.1
2017	40.9	82	187.7	113.2	272.9	5.5	0.7	2.1	0.1	0.1	705.3	14	-
2018	0	66.6	62.9	95.3	95.1	46.7	3.1	0.6	0.9	0.2	371.4	15	-

**Table 5.** Greenland halibut *Reinhardtius hippoglossoides* biomass indices (tons).

Year	0A	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	W-Grl	CI	E-Grl
1992		3516	785	4992	4203	402	206	97	48	0	14250	22	
1993		2483	1286	2507	4255	747	595	539	333	60	12804	27	
1994		2007	1697	3598	4748	1665	1458	91	10	25	15199	26	
1995		4367	1291	5786	2567	825	971	502	12	45	16365	51	
1996		3682	1294	8593	5496	439	1248	899	9	118	21778	22	
1997		4972	1746	6456	4929	421	1754	180	25	84	20567	29	
1998		7025	4976	11874	2821	1724	863	275	117	278	29955	35	
1999		10205	6025	8060	5224	555	778	261	48	318	31473	44	
2000		3411	1713	9537	3985	454	692	567	38	280	20676	30	
2001		8433	2478	10161	3802	278	1208	289	33	443	27126	32	
2002		6158	2067	9070	3108	779	737	670	39	402	23055	40	
2003		8297	3399	16556	5693	478	589	297	4	355	35668	28	
2004		15182	2079	28229	11755	1147	420	319	2	201	59332	36	
New survey gear introduced											ecv	Cv	
2005	421	22894	7010	22580	17150	574	1129	347	263	412	72780	12	
2006	519	15179	4516	20246	13797	519	6693	93	0	206	61769	12	
2007	769	11603	5666	13137	6950	163	5920	3	82	246	44539	18	-
2008	402	19559	4417	16422	7822	147	302	120	59	179	49429	13	4850
2009	229	21764	2634	19902	7047	478	324	164	16	25	52584	20	4454
2010	489	25880	7071	17559	8658	599	54	320	1	58	60688	11	5729
2011	-	18213	6778	23977	13945	640	364	49	0	14	64547	10	3825
2012	-	10331	4327	16168	9194	1475	225	7	4	81	41813	7	2439
2013	210	13661	12510	15103	11508	689	812	3	95	105	54695	9	2664
2014	-	8577	7388	11463	6014	942	829	45	111	9	35377	12	28
2015	-	13651	7272	13180	12716	461	287	200	13	183	47962	15	1351
2016	-	14638	7567	11772	4799	685	270	78	42	246	40098	12	1181
2017	1378	14532	13377	12040	9015	804	139	483	15	44	51827	13	-
2018	-	10565	4520	17355	6894	3758	301	137	98	107	43735	11	-

**Table 6.** Cod *Gadus morhua* abundance indices (millions).

Year	0A	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	W-Grl.	CI	E-Grl
1991	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.5			0.8	51	
1992	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.3	0.0	0.0	0.6	49	
1993	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.3	0.0	0.5	66	
1994	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.1	47	
1995	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.1	0.0	0.6	55	
1996	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.3	54	
1997	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	68	
1998	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	54	
1999	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.2	29	
2000	0.0	0.0	0.2	0.4	0.3	0.2	0.1	0.1	0.0	0.2	1.3	23	
2001	0.0	0.0	0.0	0.0	0.2	0.1	0.1	0.4	0.3	0.3	1.5	26	
2002	0.0	0.0	0.0	0.0	0.1	0.1	0.1	3.3	0.1	0.5	4.2	50	
2003	0.0	0.0	0.1	0.4	1.4	0.0	0.4	0.7	0.2	0.2	3.4	22	
2004	0.0	0.0	0.0	0.2	0.0	0.1	0.4	2.6	1.5	1.6	6.5	29	
New survey gear introduced											cv	cv	
2005	0.1	0.0	0.0	0.1	0.7	0.1	1.8	4.8	6.8	94.1	108.7	52	
2006	0.5	0.0	0.2	0.2	2.8	1.4	2.5	16.5	3.3	46.3	73.8	27	
2007	0.7	0.0	1.0	0.3	2.5	0.7	7.1	3.0	2.6	38.1	56.1	46	-
2008	1.2	0.1	0.7	0.1	2.4	1.7	4.1	9.5	11.9	21.7	53.4	23	31.5
2009	0.9	0.0	0.8	0.1	3.2	1.0	3.2	2.8	1.4	1.7	15.1	11	54.7
2010	0.3	0.1	0.6	0.0	1.7	1.1	2.7	8.2	2.5	6.1	23.4	24	45.3
2011	-	0.0	3.4	3.7	35.8	7.8	2.1	19.5	1.0	7.4	80.9	17	42.7
2012	-	1.1	3.8	3.5	10.0	1.0	3.3	1.2	27.1	20.3	71.1	39	22.7
2013	4.7	1.0	4.0	3.8	11.1	1.6	6.1	7.5	30.0	55.5	125.2	36	67.2
2014	-	2.4	4.2	0.4	4.4	1.3	78.9	2.5	16.5	20.6	131.0	57	35.4
2015	-	2.0	1.4	3.1	9.1	2.2	27.2	31.7	19.0	40.4	136.1	28	29.1
2016	-	0.5	3.9	0.5	2.6	0.7	10.9	1.6	3.3	4.5	28.4	22	23.7
2017	0.5	0.4	1.5	0.7	2.3	2.1	15.7	4.9	5.5	13	46.4	28	-
2018	0	1,7	0.4	0,6	9,5	5,3	8	6,4	6	5,3	14		

**Table 7.** Cod *Gadus morhua* biomass indices (tons). () incomplete coverage of survey area.

Year	0A	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	W-Grl.	CI	E-Grl
1990	2	13	*	75	3	83	9005	*	*	*	(9180)	65	
1991	0	0	7	2	15	151	310	*	*	*	(485)	44	
1992	0	0	3	20	34	75	112	0	2	246	46		
1993	0	0	2	0	5	22	39	113	5	186	68		
1994	0	3	0	0	9	38	0	1	0	51	46		
1995	0	0	0	5	1	120	23	3	4	155	63		
1996	0	0	0	0	0	15	23	24	44	107	50		
1997	0	0	0	0	2	53	0	0	0	0	55	76	
1998	0	0	0	1	0	0	47	50	3	101	56		
1999	0	1	5	23	5	1	17	1	0	53	47		
2000	0	51	99	76	54	21	9	2	46	357	23		
2001	0	0	15	125	30	56	178	98	100	603	23		
2002	0	0	13	54	74	41	1489	42	150	1863	46		
2003	0	18	111	315	8	264	453	118	46	1332	26		
2004	0	0	496	46	7	176	680	685	305	2394	28		
New survey gear introduced											cv	cv	
2005	38	0	22	45	320	19	449	1077	1170	60546	63684	71	
2006	114	0	40	22	578	74	471	5513	536	19874	27221	32	
2007	247	13	317	56	711	121	1502	514	541	26843	30865	54	-
2008	421	99	227	46	1218	794	923	1730	3321	19702	28481	37	47864
2009	212	0	184	42	1046	199	688	453	282	499	3604	13	58141
2010	183	38	215	7	821	144	573	2417	835	2899	8133	31	110656
2011	-	1	537	726	7468	1493	398	196	3948	18730	16	81138	20
2012	-	194	1061	841	3000	313	1226	447	14104	15911	37098	39	64421
2013	2446	294	1026	1287	3443	447	1871	4361	19015	51622	85812	37	168771
2014	-	709	1644	286	1813	492	56061	2511	21714	27755	112984	50	153807
2015	-	1076	429	1958	4103	353	19705	33169	27532	60282	148606	30	104806
2016	-	319	1460	477	1068	106	9039	1345	2523	4755	21092	25	89413
2017	697	205	428	641	772	483	14110	3032	3975	8401	32744	39	-
2018	513	126	445	1770	338	2369	2796	2317	8498	19173	16		



**Table 8.** Redfish species combined, all sizes (*Sebastes sp.*) abundance indices (millions).

Year	0A	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	W-Grl	CI	E-Grl
1992		1	146	9	1006	187	69	34	6	4	1462	32	
1993		4	210	17	361	22	157	182	97	520	1568	69	
1994		12	187	12	1573	225	273	85	10	84	2461	26	
1995		1	67	11	559	33	183	93	5	5	957	23	
1996		1	8	8	1688	59	124	63	11	54	2017	29	
1997		5	43	7	348	58	156	57	23	22	719	24	
1998		1	24	20	236	15	115	71	10	32	522	24	
1999		2	69	9	287	11	11	61	4	23	574	21	
2000		0	27	6	27	32	28	10	18	159	165	23	
2001		2	77	3	25	15	20	12	1	180	183	60	
2002		1	23	1	50	84	43	44	2	10	257	27	
2003		2	45	2	210	32	79	25	2	26	423	23	
2004		2	11	1	52	39	47	30	2	69	253	37	
New survey gear introduced												cv	cv
2005	184	1	19	1	73	79	77	25	12	80	551	-	
2006	27	7	25	1	110	52	83	16	3	11	334	-	
2007	97	2	37	0	128	34	49	5	3	17	371	-	-
2008	99	2	40	0	121	33	24	3	2	11	334	-	1847
2009	37	2	29	0	115	26	12	1	2	9	234	-	1988
2010	27	6	27	0	69	42	23	8	3	8	213	-	1566
2011	-	10	23	1	121	18	31	13	1	9	225	-	2517
2012	-	7	10	0	42	13	20	10	2	16	120	-	1018
2013	12	2	10	0	29	4	30	6	44	7	145	-	1075
2014	-	1	3	0	23	5	3	39	20	3	96	-	559
2015	-	1	2	0	13	1	11	39	8	7	82	-	762
2016	-	1	1	0	4	4	15	199	34	6	265	-	429
2017	1	0	1	0	5	2	11	10	5	5	39	-	-
2018	0	1	0	0	9	0	15	5	21	20	71	-	-

**Table 9.** Redfish species combined, all sizes *Sebastes sp.* biomass indices (tons).

Year	0A	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	W-Grl.	CI	E-Grl.
1992		69	18117	437	13423	2832	1576	1124	169	147	37894	43	
1993		195	4994	710	6420	300	1549	3835	1923	2138	22065	38	
1994		590	5076	538	16064	1986	3886	995	179	1272	30586	24	
1995		52	1585	775	5029	869	2963	1952	358	123	13705	22	
1996		18	117	782	12178	1694	2552	1980	304	1788	21413	28	
1997		599	1481	337	4913	1597	6766	1901	1099	1229	19922	31	
1998		39	1467	1423	6193	2130	3274	1953	606	1198	18283	22	
1999		164	4021	742	5596	999	2742	2976	207	1124	18671	25	
2000		0	1790	793	1045	2185	2337	463	2411	1214	12237	36	
2001		192	5380	536	1746	1460	2637	1069	60	2256	15337	50	
2002		55	1917	397	2536	2386	1676	2654	272	998	12891	28	
2003		279	2886	702	6357	2319	6185	1918	187	2476	23308	32	
2004		369	462	368	2210	2274	2996	1679	101	1026	11486	41	
New survey gear introduced												cv	cv
2005	3491	134	1378	665	3370	6974	6212	2751	1388	2771	29132		
2006	591	1129	2196	759	4427	2717	6213	959	557	2350	21897		
2007	3367	248	2172	153	6886	1499	5166	358	282	1778	21908	-	-
2008	2845	433	2221	210	7411	4007	2542	224	286	2585	22761	-	290666
2009	1696	356	2472	193	5496	3688	1951	293	335	1339	17819	-	318517
2010	1348	761	3363	910	4765	4193	3073	1043	369	1069	20895	-	336840
2011	-	1917	1536	1486	8362	3538	5377	4026	226	2397	28865	-	511700
2012	-	1382	1224	998	4380	2438	3560	1942	239	10341	26505	-	234650
2013	965	429	1684	553	6063	1077	7327	1269	23081	3178	45626	-	454592
2014	-	370	394	150	5241	1074	1092	11621	11017	2009	32969	-	206406
2015	-	276	548	1000	3458	427	2833	12477	4590	4771	30381	-	288118
2016	-	502	317	420	964	1514	4894	61617	41426	3373	115028	-	486768
2017	61	282	291	134	2489	291	4255	3308	3399	1822	16332	-	-
2018	0	498	158	136	3856	30	5305	2029	14000	10565	36577	-	-



**Table 10.** Juvenile redfish <20cm *Sebastes sp.* abundance indices (millions).

Year	0A	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	W-Grl	cv	E-Grl	cv
2006	26.7	6.4	19.5	0	107.1	47.1	75.8	14.4	1.6	7.3	305.8	16	-	-
2007	94.7	0	28.2	0	121.8	32.5	45.2	4.7	2.4	13.6	343.3	15	-	-
2008	95.6	0.4	35.7	0	120.9	0.2	8.6	1.9	0.9	5.2	269.4	26	452	53
2009	34.8	0	14	0	110	1	4.4	0.5	1.2	3.7	169.6	24	954	28
2010	23.7	0	9.5	0	34.5	0	3.5	3.7	1.9	4.5	81.2	21	507	31
2011	0	0.2	14.7	0.1	74.9	0.8	0.4	2.4	0.4	0.4	94.1	23	102	61
2012	0	0	1.8	0	20.2	0.2	1	0.4	0.6	0.3	24.5	23	253	56
2013	4.5	0.1	3.1	0	18.4	0.3	2.1	1	0.1	0.2	29.7	17	48	36
2014	0	0	0.9	0	8.1	0.2	0.1	0.1	0.1	0.1	9.5	31	21	33
2015	0	0	0.2	0	4.2	0	0	0.7	0.1	0.1	5.3	36	21	52
2016	0	0	0.1	0	0.6	0	0.1	0.1	0.1	0.4	1.5	18	5	22
2017	0.3	0	0	0	2	0.3	0.2	0.4	0.3	1.9	5.6	19	-	-
2018	0	0.4	0	0	4.8	0.1	0.2	0.3	0.2	0.5	6.4	59	-	-

**Table 11.** Juvenile redfish <20cm *Sebastes sp.* biomass indices (tons).

t	0A	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	W-Grl	cv	E-Grl	cv
2006	480	651	1400	0	3340	2128	4483	741	113	1210	14546	18	-	-
2007	2866	0	1449	0	4088	1132	4457	173	98	733	14996	16	-	-
2008	2322	2	1665	0	5360	2	143	43	12	49	9598	29	9011	53
2009	1315	0	471	3	3569	34	152	9	25	89	5668	29	41341	63
2010	895	0	401	0	1598	0	105	155	71	148	3372	29	24135	31
2011	-	9	291	1	2639	45	20	86	26	16	3132	32	3781	56
2012	-	0	52	0	735	8	31	13	22	14	875	24	36567	79
2013	223	2	159	1	811	10	113	12	3	6	1340	21	1244	33
2014	-	0	22	0	227	7	4	3	4	2	268	28	550	26
2015	-	0	6	0	213	0	0	11	5	2	236	44	796	45
2016	-	0	3	0	15	0	0	3	2	7	31	33	329	30
2017	10	0	0	0	33	6	5	11	9	41	114	19	-	-
2018	-	378	0	0	122	3	6	10	8	17	544	72	-	-

**Table 12.** Deep-sea redfish *Sebastes mentella* abundance indices (millions).

Year	0A	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	W-Grl	cv	E-Grl	cv
2006	0	0.6	1.4	0.2	0.9	0.8	1.5	0.5	0.3	1.6	7.8	24	-	-
2007	0.4	0	0.1	0.1	6.5	0	0.1	0.1	0.1	0.8	8.1	52	-	-
2008	0.9	1.4	3.6	0	2.2	31.6	13.6	0.4	0.2	2.2	56.2	56	1286	49
2009	0.7	1.5	13.7	0.1	3.9	23.4	7.7	0.5	0.1	4.1	55.7	45	895	29
2010	2.2	5.5	16.1	0	35.9	40.1	17.8	4	0.4	2.8	124.7	36	995	36
2011	0	8.7	8.3	0	55.6	15.4	28.6	6.9	0.4	5.3	129.1	22	400	22
2012	0	6.4	6.8	0.2	19.8	12	16.1	8	1.4	4.1	74.7	10	305	27
2013	7	0.9	6.1	0.1	9.7	3.4	24.1	2.3	41.9	2.6	98.2	85	338	33
2014	0	0.4	2.1	0	13.1	4.8	1.5	37.5	18.6	1.8	79.7	25	172	25
2015	0	0.4	2	0.1	6.4	0.8	9.6	36	5.7	4.1	65.1	38	130	23
2016	0	0	0.6	0	3.1	4.1	14.7	197.3	24.5	4.4	248.8	77	188	67
2017	0.3	0.5	0.5	0	2	1.2	10.4	8.5	4.2	1.6	29.2	20	-	-
	0	0.3	0.1	0	2.4	0.1	14.1	5	19.4	18	59.4	52	-	-

**Table 13.** Deep-sea *Sebastes mentella* biomass indices (tons).

Year	0A	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	W-Grl	cv	E-Grl	cv
2006	10	169	66	82	279	268	542	66	40	489	2010	22	-	-
2007	82	0	27	29	1592	0	8	29	16	315	2099	84	-	-
2008	141	244	390	10	473	3859	2247	72	52	430	7916	49	247557	47
2009	77	238	1679	47	650	2891	1499	215	12	894	8202	38	236680	35
2010	138	606	1819	12	2346	3906	2358	622	79	589	12475	36	283924	36
2011	0	1290	1011	4	4127	3051	4355	1533	93	1330	16794	24	122949	26
2012	0	1041	840	356	1906	2024	2316	1441	155	999	11078	9	100342	29
2013	640	124	931	232	2768	807	5833	479	21213	910	33938	59	162268	50
2014	0	71	264	8	1949	991	379	10466	9559	838	24525	22	67611	24
2015	0	103	543	32	1031	181	2287	10302	2800	2104	19382	35	60589	28
2016	0	7	86	0	515	1138	4282	59250	15241	2311	82831	72	163750	79
2017	51	282	156	9	771	210	3174	2386	2772	735	10546	24	-	-
2018	-	95	91	6	940	18	4958	1638	12732	9073	29550	64	-	-



**Table 14.** Golden redfish *Sebastes norvegicus* abundance indices (millions).

Year	0A	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	W-Grl	cv	E-Grl	cv
2006	0.1	0.5	3.5	0.4	2.6	0.2	2.4	0.4	1.1	1.4	12.6	23	-	-
2007	0.7	2.5	7	0.1	2.1	1	4.9	0.4	0.4	1.2	20.2	43	-	-
2008	1.4	0.3	0.4	0.1	1.3	0.1	0.7	0.3	0.4	3.5	8.5	25	108	29
2009	0.5	0.3	0.9	0.1	1.1	0.7	0.5	0.2	0.5	1.7	6.5	15	139	27
2010	0.4	0.3	1.1	0.4	0.7	0.9	0.7	0.3	0.7	0.5	5.9	17	64	36
2011	-	0.9	0.4	0.7	2.3	0.7	3.2	3.3	0.7	2.1	14.4	12	2015	36
2012	-	0.5	0.5	0.3	1.9	0.4	2	1.9	0.1	11.5	19	54	460	36
2013	0.3	0.6	0.8	0.1	2.1	0.4	2.4	3	3.5	3.6	17	15	689	34
2014	-	0.4	0.4	0	1.5	0.1	0.9	0.9	0.9	1.5	6.7	17	366	40
2015	-	0.2	0	0.4	1.8	0.2	0.8	2	2.5	3.1	11.1	19	611	37
2016	0.6	0.2	0.1	0.3	0.2	0.5	1.7	9.6	1.6	14.8	80	236	38	
2017	0	0	0.1	0.1	0.8	0.1	0.7	0.6	0.5	1.5	4.4	11	-	-
			0,1	0	1,5	0,1	0,3	0,2	0,9	1,7	4,7	18		

**Table 15.** Golden redfish *Sebastes norvegicus* biomass indices (tons).

Year	0A	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	W-Grl	cv	E-Grl.	cv
2006	91	311	696	724	986	197	1049	145	491	554	5242	16	-	-
2007	358	249	596	136	1598	373	825	180	227	674	5216	27	-	-
2008	339	186	157	211	1970	14	189	108	235	2030	5438	41	34098	32
2009	244	117	317	156	1434	642	383	70	315	426	4103	18	40496	27
2010	278	154	1207	954	1024	149	674	275	272	276	5264	28	28781	32
2011	-	608	257	1578	2828	338	1270	2483	165	777	10304	21	384970	36
2012	-	342	274	706	2049	327	1140	566	72	8925	14400	61	97741	28
2013	81	304	520	357	3064	228	1239	842	2439	2025	11096	31	291080	57
2014	-	299	108	142	3066	77	709	1152	1454	1169	8176	19	138245	41
2015	-	173	0	968	2214	246	546	2165	1785	2665	10762	22	226724	37
2016	-	495	228	420	434	376	612	2364	26182	1055	32166	106	322690	73
2017	0	0	134	125	1685	75	1077	912	618	1046	5672	13	-	-
2018	-	0	24	67	129	2795	10	341	381	1260	1475	23		

**Table 16.** American plaice *Hippoglossoides platessoides* abundance indices (millions).

t	0A	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	W-Grl.	CI	E-Grl
1992		1.4	1.0	2.8	1.4	0.6	1.7	1.4	0.5	0.1	10.8	22	
1993		1.3	1.7	1.2	3.0	0.8	2.0	1.7	0.6	0.9	13.2	24	
1994		2.2	3.7	3.3	14.9	7.0	9.5	0.7	0.3	0.5	42.0	32	
1995		1.0	0.6	1.8	6.3	0.9	2.7	3.0	0.3	0.5	17.2	29	
1996		1.6	3.4	7.3	4.6	1.7	4.2	3.1	0.1	0.7	26.6	18	
1997		6.6	2.0	2.7	15.1	1.0	10.4	2.0	0.3	0.7	40.8	47	
1998		1.6	1.9	2.4	3.6	1.2	1.5	6.4	0.9	5.6	25.2	27	
1999		0.5	1.7	2.0	6.8	1.2	5.3	1.9	1.0	0.6	20.9	18	
2000		1.8	4.8	6.7	14.8	1.9	3.5	3.8	0.5	0.5	38.5	23	
2001		1.3	1.3	2.2	13.6	1.5	3.5	2.3	0.6	0.6	26.8	31	
2002		0.0	3.8	4.7	8.8	1.8	5.1	31.8	1.5	1.8	59.5	49	
2003		2.2	5.2	5.5	25.7	1.6	13.7	15.0	0.8	2.0	71.7	22	
2004		0.7	1.4	5.3	11.9	2.8	10.2	8.9	0.6	1.6	43.4	27	
New survey gear introduced													
2005	3.5	2.1	13.7	5.8	59.9	17.3	74.5	19.5	6.1	5.6	208.0	9	
2006	3.9	1.3	15.0	10.9	40.3	10.0	38.1	21.6	4.4	4.5	150.0	10	
2007	6.5	0.6	12.3	6.3	46.8	9.4	31.2	11.9	2.6	2.7	130.8	12	
2008	7.1	3.7	10.8	4.0	29.5	2.2	13.5	11.4	1.3	1.3	84.8	8	17.2
2009	2.9	5.6	18.0	14.3	41.4	8.1	11.6	7.6	1.1	2.3	112.8	9	20.2
2010	9.5	2.8	40.7	14.3	50.7	9.2	34.4	20.3	1.2	2.1	185.2	8	17.4
2011	-	8.6	40.7	17.1	103.6	6.2	30.3	21.6	0.8	1.9	230.7	11	17.5
2012	-	2.3	18.3	15.3	51.3	8.3	19.4	11.4	0.9	1.1	129.4	9	10.4
2013	4.4	19.0	39.0	25.4	84.1	9.1	34.1	27.9	4.7	0.9	248.6	13	13.1
2014	-	7.0	22.0	11.0	36.0	6.0	15.0	10.0	2.0	1.0	110.0	7	10.0
2015	-	41.0	55.0	28.0	89.0	39.0	25.0	17.0	2.0	1.0	296.0	10	10.8
2016	-	10.1	21.6	11.3	43.1	2.4	16.2	7.6	2.2	1.2	115.7	8	10.1
2017	5	6.3	23.3	26.9	64.1	18.8	23.4	11.2	1.4	1.1	181.5	9	-
2018	0	11,9	20,4	21,8	126	11,4	20,7	8,7	2,4	2,1	225,4	10	

**Table 17.** American plaice *Hippoglossoides platessoides* biomass indices (tons).

Year	0A	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	W-Grl.	CI	E-Grl
1992	57	54	213	78	51	137	128	55	6	779	23		
1993	56	72	87	90	28	107	141	69	43	694	26		
1994	112	293	277	487	308	284	60	22	64	1906	22		
1995	65	54	279	191	51	87	130	19	18	895	18		
1996	119	264	670	231	74	142	119	7	27	1654	18		
1997	323	150	287	398	87	367	135	31	25	1803	21		
1998	154	178	328	185	48	82	398	97	102	1573	20		
1999	81	136	170	287	43	202	145	65	44	1173	17		
2000	175	278	408	551	74	178	227	89	40	2021	18		
2001	169	79	140	403	65	162	153	38	67	1276	17		
2002	0	184	327	414	151	275	1061	92	67	2570	23		
2003	196	352	338	1013	125	680	1048	59	171	3980	20		
2004	138	143	192	537	128	715	747	38	150	2789	27		
New survey gear introduced													
2005	246	346	944	722	3049	1136	4511	2196	470	569	14190	8	
2006	268	210	608	1148	2288	702	3534	2163	399	380	11699	13	
2007	356	112	544	731	2877	731	2418	810	230	308	9118	12	
2008	371	437	648	382	1889	212	1067	898	128	149	6181	9	2495
2009	183	556	649	643	1977	585	826	825	102	127	6473	7	2577
2010	355	235	1539	1214	2436	748	2128	1460	105	224	10442	9	3181
2011	-	463	1364	1459	5464	792	1391	2020	99	208	13256	7	4385
2012	-	236	756	730	3508	662	1350	777	121	178	8317	9	2702
2013	325	920	1472	1457	4066	521	2818	1425	568	121	13694	10	2269
2014	-	311	746	990	2245	337	1089	1027	203	130	7078	7	1203
2015	-	2046	1320	2956	4547	1364	1635	1642	246	188	15944	9	2351
2016	-	563	1428	1117	2735	201	1152	841	324	146	8508	7	2385
2017	303	734	965	1726	2759	1122	1632	1218	195	148	10802	8	-
2018	1699	997	1988	5928	719	1164	1055	282	178	14009	10		

**Table 18.** Atlantic wolffish *Anarhichas lupus* abundance indices (millions).

Year	0A	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	W-Grl	CI	E-Grl
1992		0.0	0.0	0.0	0.0	0.1	0.3	0.1	0.1	0.1	0.8	33	
1993		0.0	0.0	0.0	0.0	0.0	0.2	0.2	0.3	0.3	1.2	41	
1994		0.0	0.1	0.0	0.3	0.2	1.9	0.4	0.4	1.6	4.9	36	
1995		0.0	0.0	0.0	0.1	0.1	0.4	0.1	0.4	0.2	1.3	34	
1996		0.0	0.1	0.0	0.1	0.1	0.4	0.2	0.6	0.6	2.0	50	
1997		0.0	0.0	0.0	0.2	0.0	0.8	0.2	0.3	0.1	1.7	28	
1998		0.0	0.3	0.0	0.3	0.1	0.7	0.4	0.5	0.7	3.0	30	
1999		0.0	0.2	0.1	0.1	0.1	0.4	0.2	0.4	0.4	2.0	44	
2000		0.0	0.5	0.1	0.3	0.4	0.4	0.6	1.7	0.0	4.0	69	
2001		0.0	0.1	0.2	0.4	0.1	0.2	0.4	0.2	0.1	1.5	40	
2002		0.0	0.4	0.1	0.5	0.0	0.9	0.6	1.1	0.5	4.1	33	
2003		0.0	0.1	0.6	2.3	0.2	1.5	0.5	2.3	1.1	8.8	27	
2004		0.0	0.1	0.3	1.6	0.9	0.2	0.3	0.7	0.3	4.4	39	
New survey gear introduced											cv	cv	
2005	0.1	0.0	0.5	0.0	1.5	0.6	6.9	2.4	5.5	1.5	19.1	16	
2006	0.0	0.1	0.8	0.1	0.9	0.6	2.4	1.2	1.0	2.8	9.8	14	
2007	0.0	0.0	0.7	0.0	1.3	0.6	0.7	0.3	0.4	0.7	4.8	15	-
2008	0.2	0.0	1.0	0.1	0.7	0.5	1.0	0.5	0.4	0.3	4.6	15	4.7
2009	0.0	0.1	1.0	0.1	0.7	0.4	1.5	0.2	1.1	0.4	5.6	19	12.2
2010	0.1	0.4	2.6	0.1	2.3	0.6	3.1	0.8	0.9	0.6	11.4	8	5.3
2011	-	0.1	1.1	0.1	1.3	0.3	0.5	0.9	0.3	0.4	5.0	14	6.4
2012	-	0.0	2.2	0.2	1.6	0.3	0.7	1.2	0.3	0.6	7.0	18	3.8
2013	0.1	0.1	3.1	0.0	1.0	0.3	1.7	0.4	0.5	0.8	7.9	18	6.5
2014	-	0.4	2.3	0.3	0.7	0.8	0.3	0.1	0.2	0.3	5.5	16	2.6
2015	-	0.6	4.3	0.3	3.1	4.6	6.3	1.1	0.8	0.3	21.5	23	2.9
2016	-	0.5	1.8	0.2	1.2	1	1.4	0.2	0.4	1	7.7	10	2.3
2017	0.1	0.5	5.8	0.4	2.3	2.9	3.6	0.4	0.2	0.4	16.7	20	-
2018	0	1,2	4,4	0	4,9	2,1	3	0,7	0,3	0,6	17,2	15	-

**Table 19.** Atlantic wolffish *Anarhichas lupus* biomass indices (tons).

Year	0A	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	W-Grl	CI	E-Grl
1992	0	7	0	8	21	47	22	28	31	163	33		
1993	0	5	6	1	2	26	35	29	188	292	64		
1994	0	12	9	40	39	198	30	65	249	644	38		
1995	0	0	0	22	9	38	24	90	36	219	40		
1996	0	1	3	17	23	41	35	103	101	324	53		
1997	0	3	0	21	1	115	16	58	15	228	30		
1998	0	3	2	13	9	60	34	104	133	358	38		
1999	0	4	13	21	12	8	6	202	62	329	79		
2000	0	20	3	52	31	55	54	396	15	626	90		
2001	0	1	3	11	1	16	21	42	23	117	40		
2002	0	9	2	77	5	73	78	216	118	579	37		
2003	0	2	41	267	64	361	60	205	148	1148	24		
2004	0	6	19	160	56	96	21	162	100	620	27		
New survey gear introduced											cv	cv	
2005	11	0	76	5	201	123	1564	348	890	640	3858	16	
2006	3	4	58	60	127	134	359	168	172	1241	2326	20	
2007	0	0	73	6	357	326	147	107	122	533	1670	21	-
2008	15	0	76	31	124	55	348	88	136	226	1098	16	1091
2009	0	3	109	30	155	72	275	92	211	648	1595	31	2870
2010	10	12	509	51	350	144	668	134	104	345	2327	17	1295
2011	-	46	96	64	197	69	121	233	63	205	1094	17	1633
2012	-	0	239	82	506	44	79	252	64	444	1708	23	1172
2013	9	27	239	20	136	81	555	183	115	636	2000	25	1652
2014	-	9	79	69	158	62	107	78	100	458	1121	30	520
2015	-	56	450	168	478	762	1543	752	547	310	5066	22	1585
2016	-	60	270	174	271	228	440	156	119	708	2425	14	952
2017	2	58	437	215	381	454	960	321	168	586	3583	15	-
2018	109	291	13	730	294	491	451	88	691	3158	15		



**Table 20.** Spotted wolffish *Anarhichas minor* abundance indices (millions).

Year	0A	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	W-Grl	CI	E-Grl
1992		0.0	0.1	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.3	28	
1993		0.0	0.1	0.0	0.1	0.0	0.1	0.0	0.0	0.1	0.4	36	
1994		0.0	0.3	0.0	0.2	0.1	0.4	0.0	0.0	0.0	1.1	33	
1995		0.0	0.2	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.4	22	
1996		0.1	0.1	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.5	24	
1997		0.1	0.3	0.0	0.2	0.0	0.0	0.1	0.0	0.0	0.8	23	
1998		0.1	0.2	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.6	25	
1999		0.9	0.3	0.1	0.4	0.1	0.0	0.0	0.0	0.0	1.7	42	
2000		0.0	0.8	0.0	0.5	0.1	0.0	0.0	0.1	0.0	1.6	28	
2001		0.3	0.4	0.1	0.3	0.0	0.0	0.0	0.0	0.0	1.1	27	
2002		0.3	2.1	0.1	0.4	0.1	0.1	0.1	0.0	0.0	3.1	42	
2003		0.5	1.5	0.2	1.4	0.0	0.6	0.0	0.0	0.1	4.3	24	
2004		0.2	0.7	0.2	0.6	0.0	0.1	0.1	0.0	0.1	2.1	30	
New survey gear introduced											cv	cv	
2005	0.0	1.2	1.9	0.0	0.7	0.2	0.4	0.3	0.2	0.1	5.1	23	
2006	0.0	0.4	1.0	0.1	1.4	0.3	0.3	0.2	0.1	0.2	3.9	23	
2007	0.1	0.5	1.0	0.1	1.5	0.4	0.2	0.1	0.0	0.1	3.9	31	-
2008	0.1	0.5	0.6	0.1	0.4	0.0	0.1	0.1	0.0	0.1	2.1	15	1.1
2009	0.0	0.4	1.7	0.1	0.6	0.1	0.0	0.0	0.0	0.1	3.1	26	2.3
2010	0.2	1.0	1.8	0.2	0.7	0.2	0.2	0.1	0.1	0.2	4.5	12	2.9
2011	-	0.9	1.8	0.2	0.9	0.1	0.2	0.1	0.0	0.1	4.4	14	2.3
2012	-	0.8	1.0	0.1	0.7	0.1	0.1	0.2	0.1	0.2	3.2	15	2.9
2013	0.1	1.3	2.2	0.2	1.7	0.1	0.3	0.2	0.4	0.4	6.9	20	2.5
2014	-	1.4	2.3	0.1	0.2	0	0.1	0.2	0.2	0.2	4.7	21	1.5
2015	-	2.2	2.2	0.2	0.7	0.4	0.5	0.4	0.2	0.4	7.4	14	2.0
2016	-	1.7	1.2	0.1	0.5	0.1	0.6	0.2	0.3	0.2	4.9	17	1.3
2017	0	2.7	2.1	0.3	0.7	0.5	0.3	0.1	0.3	0.4	7.5	16	-
2018	-	1.7	2.5	0.3	1.6	0.3	0.3	0.2	0.3	0.1	7.5	11	-

**Table 21.** Spotted wolffish *Anarhichas minor* biomass indices (tons).

Year	0A	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	W-Grl	CI	E-Grl
1992		4	76	65	110	3	34	33	6	19	351	28	
1993		55	0	100	47	16	66	4	0	282	571	53	
1994		223	180	0	81	40	119	28	11	1	683	25	
1995		0	60	15	68	16	22	19	11	164	377	49	
1996		169	77	12	193	15	6	31	0	50	554	26	
1997		193	72	37	81	0	16	124	0	5	530	34	
1998		2	64	0	143	18	6	125	100	7	465	32	
1999		131	121	23	28	36	13	2	0	0	354	31	
2000		0	188	31	133	36	19	1	593	0	1000	114	
2001		523	30	25	310	80	4	0	0	10	982	52	
2002		135	194	20	169	81	74	233	71	126	1104	28	
2003		299	1416	195	978	22	741	107	0	226	3985	22	
2004		124	1270	623	567	2	78	603	352	545	4164	35	
New survey gear introduced											cv	cv	
2005	150	764	1182	6	1058	155	741	2514	568	137	7275	26	
2006	0	472	1257	243	1345	1066	1336	716	350	1145	7930	19	
2007	14	543	705	196	1249	678	874	372	178	634	5442	17	-
2008	63	1487	2050	74	730	24	347	995	425	372	6567	44	5262
2009	0	280	627	653	1453	154	35	129	189	160	3662	18	2890
2010	66	2363	1508	1195	1006	167	222	238	212	1715	8691	18	3877
2011	-	2537	2244	742	2460	1210	2294	479	218	769	12955	23	6133
2012	-	1227	683	464	3166	325	468	390	555	1104	8383	21	6871
2013	7	2026	2402	544	3135	632	1494	658	3163	4845	18906	22	4622
2014	-	779	3038	381	753	9	427	2740	1496	1916	11538	23	10468
2015	-	3994	3814	843	3655	518	1451	3208	1127	3646	22256	15	3227
2016	-	700	1755	279	1333	382	2726	1164	2999	1099	12436	22	4623
2017	0	4184	5387	804	858	937	212	604	3269	3701	19957	21	-
2018	0	4009	1811	508	2795	801	1040	1358	2406	1083	15812	19	




**Table 22.** Thorny skate *Amblyraja radiata* abundance indices (millions).


	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	W-Grl	CI	E-Grl
1992	2.9	1.3	1.3	1.4	0.2	0.6	1.0	0.1	0.0	9.0	25	
1993	0.7	0.4	0.8	1.6	0.5	0.9	0.5	0.6	0.2	6.1	21	
1994	1.9	1.8	1.2	3.9	1.8	2.8	0.4	0.2	0.0	14.1	21	
1995	2.7	1.3	0.8	2.8	1.8	0.7	2.4	0.5	0.1	13.1	26	
1996	4.6	2.2	1.5	4.0	0.4	0.7	0.6	0.0	0.6	14.6	23	
1997	4.6	1.4	0.6	4.4	0.2	2.3	0.7	0.1	0.0	14.4	26	
1998	2.8	4.1	3.2	3.0	0.3	0.6	0.9	0.4	0.3	15.5	25	
1999	1.7	3.2	0.9	2.8	0.3	1.0	0.8	0.2	0.2	11.0	23	
2000	2.1	3.4	2.0	4.0	0.5	0.6	0.5	0.1	0.6	13.9	23	
2001	3.2	1.2	0.6	2.3	0.3	0.4	0.5	0.1	0.2	8.8	32	
2002	0.5	1.1	0.9	2.2	0.4	0.9	2.6	0.2	0.5	9.3	25	
2003	3.9	2.4	1.4	7.2	0.1	0.7	0.8	0.2	0.7	17.4	26	
2004	2.5	1.3	1.3	1.2	0.2	0.2	0.7	0.2	0.1	7.9	24	
New survey gear introduced										cv	cv	
2005	0.1	4.1	2.8	2.2	3.7	0.5	0.8	0.6	0.4	15.4	10	
2006	0.2	3.8	2.2	3.0	2.8	0.2	2.3	2.8	0.5	1.1	18.8	10
2007	0.1	3.3	1.4	1.6	3.8	0.1	1.3	0.9	1.4	0.4	14.3	20
2008	0.2	5.4	1.4	1.0	1.9	0.1	0.5	1.1	0.1	0.2	11.9	16
2009	0.0	10.9	1.1	3.9	1.9	0.3	0.3	0.6	0.2	0.4	19.7	20
2010	0.1	5.6	4.0	4.1	3.3	0.8	2.7	1.2	0.1	0.0	21.8	12
2011	-	5.5	2.3	3.6	6.7	0.4	4.0	4.9	1.7	0.1	29.2	16
2012	-	2.7	2.2	2.4	4.7	0.5	1.3	2.1	0.1	0.0	16.1	15
2013	0.4	4.9	2.1	3.2	9.1	0.6	1.7	1.1	0.1	0.1	23.4	27
2014	-	2.4	2.4	1.6	1.9	0.5	0.7	0.2	0.1	0	9.8	13
2015	-	7.8	8.7	6.5	7.6	5.7	2.5	3.2	0.2	0.2	42.3	15
2016	-	3.3	2.1	3	3.6	0.6	0.9	0.8	0.1	0.1	14.3	11
2017	0	2.3	3.5	4.7	2.3	4.1	2	0.5	0	0	19.4	10
2018	0	2.3	3.5	4.7	2.3	4.1	2	0.5	0	0	19.4	9

**Table 23.** Thorny skate *Amblyraja radiata* biomass indices (tons).


Year	1AN	1AS	1AX	1BN	1BS	1C	1D	1E	1F	W-Grl	CI	E-Grl
1992	370	268	162	226	37	57	113	32	5	1271	20	
1993	60	65	199	171	87	116	128	40	22	887	24	
1994	494	283	182	465	275	311	55	61	3	2129	23	
1995	253	227	301	451	327	121	300	78	24	2083	21	
1996	631	554	623	509	61	105	65	0	207	2755	23	
1997	830	411	322	566	56	156	187	25	7	2559	26	
1998	392	839	535	427	78	38	114	81	76	2580	26	
1999	278	931	253	247	45	94	96	25	49	2019	34	
2000	323	1178	345	428	122	84	120	3	197	2799	23	
2001	325	215	222	248	52	52	89	10	60	1272	28	
2002	13	246	320	280	101	86	687	63	177	1973	29	
2003	1005	902	567	1481	11	107	174	24	206	4478	25	
2004	598	520	791	197	47	33	333	98	78	2694	23	
New survey gear introduced										cv	cv	
2005	26	776	953	676	558	219	145	249	125	96	3822	15
2006	66	836	364	662	361	91	477	807	224	303	4193	14
2007	55	897	319	566	709	50	258	152	164	87	3258	15
2008	98	1411	315	400	353	20	45	222	25	47	2937	21
2009	7	2267	411	904	374	90	81	97	78	55	4365	11
2010	20	1092	1036	1062	623	293	434	368	16	19	4962	11
2011	-	970	556	1129	1152	84	477	1172	80	42	5661	11
2012	-	738	635	722	910	107	192	145	31	16	3496	12
2013	117	1222	756	1671	1453	219	408	255	57	40	6198	18
2014	-	742	896	713	410	249	200	73	24	3	3311	17
2015	-	1467	1473	1830	1814	1582	273	791	90	145	9465	13
2016	-	987	638	1011	937	290	406	395	53	67	4784	13
2017	0	528	1072	1614	417	876	319	235	0	14	5076	10
2018	-	528	1072	1614	417	876	319	235	0	14	5076	10




**Figure 1.** The Stratification areas used in the Greenland Shrimp and Fish survey. In West Greenland each strata is divided in depth strata of 150-200m, 200-300m, 300-400m and 400-600m. "Shallow" water strata of 0-100m and 100-150m delimited by the 3 nm line and the NAFO Div. Borders of the shallow water stratas are not shown. In East Greenland each strata is divided in depth strata of 200-400m and 400-600m. "Shallow" water strata of 0-200m is delimited by the 3 nm line.




**Figure 2.** Greenland halibut abundance (million) and biomass (Kt) for West Greenland.




**Figure 3.** Atlantic cod abundance (million) and biomass (Kt) for West Greenland.




**Figure 4a.** Redfish species (*S. mentella* and *S. norvegicus* combined) abundance (million) and biomass (Kt) for West Greenland.




**Figure 4b.** Juvenile redfish >20 cm (*S. mentella* and *S. norvegicus* combined) abundance (million) and biomass (Kt).




**Figure 4c.** Deep-sea redfish *S. mentella* abundance (million) and biomass (Kt).




**Figure 4d.** Golden redfish *S. norvegicus* abundance (million) and biomass (Kt).




**Figure 5.** American plaice abundance (million) and biomass (Kt) for West Greenland.




**Figure 6.** Atlantic wolffish abundance (million) and biomass (Kt) for West Greenland.




**Figure 7.** Spotted wolffish abundance (million) and biomass (Kt) for West Greenland.




**Figure 8.** Thorny skate abundance (million) and biomass (Kt) for West Greenland.




**Figure 9.** Total recruitment of age one offshore SA1 and inshore 1A from 1988 to 2018.




**Figure 10.** Distribution of one year old Greenland halibut in 2018




**Figure 11.** Number of one-year of Greenland halibut by division and year from 1991 to 2018.




**Figure 12.** Relative distribution of one-year old Greenland halibut between offshore areas and inshore Disko Bay from 1992 to 2019.




**Figure 13:** Length frequencies for Greenland halibut offshore for the period 1988-2018.




**Figure 14.** Greenland halibut (*Reinhardtius hippoglossoides*). Length frequencies for West Greenland.




**Figure 15.** Atlantic cod (*Gadus morhua*). Length frequencies for West Greenland (left) and length frequencies per division (right).




**Figure 16.** Stacked Length frequencies for golden redfish (*Sebastes Norvegicus*), deep-sea redfish (*Sebastes mentella*) and juvenile redfish (*Sebastes sp. <18cm*) for West Greenland.




**Figure 17.** American plaice (*Hippoglossoides platessoides*). Length frequencies for West Greenland (left) and length frequencies per division in (right).




**Figure 18.** Atlantic wolffish (*Anarhichas lupus*). Length frequencies for West Greenland by year (left) and length frequencies per division (right).




**Figure 19.** Spotted wolffish (*Anarhichas minor*). Length frequencies for West Greenland (left) and length frequencies per division (right).






**Figure 20.** Thorny skate (*Amblyraja radiata*) length frequencies for West Greenland (left) and length frequencies per division (right).




**Figure 21.** Greenland halibut survey biomass in kg/km<sup>2</sup> and abundance in numbers/km<sup>2</sup>.




**Figure 22.** Atlantic cod survey biomass in kg/km<sup>2</sup> and abundance in numbers/km<sup>2</sup>.







**Figure 25.** Golden redfish survey biomass in kg/km<sup>2</sup> and abundance in numbers/km<sup>2</sup>.




**Figure 26.** Golden redfish and deep-sea redfish survey kg/km<sup>2</sup> and numbers/km<sup>2</sup>.




**Figure 27.** American plaice survey biomass in kg/km<sup>2</sup> and abundance in numbers/km<sup>2</sup>.




**Figure 28.** Atlantic wolffish survey biomass in kg/km<sup>2</sup> and abundance in numbers/km<sup>2</sup>.



**Figure 29.** Spotted wolffish survey biomass in kg/km<sup>2</sup> and abundance in numbers/km<sup>2</sup>.



**Figure 30.** Thorny skate survey biomass in kg/km<sup>2</sup> and abundance in numbers/km<sup>2</sup>.



## Appendix I.

### Grønlands Naturinstitut

Greenland Institute of Natural Resources

R/V Paamiut

OYZC – GR6-251

MMSI 331 102 00

engine.paamiut@gmail.com



#### Statement regarding using M/Tr Sjúrðarberg to carry out the same surveys as R/V Paamiut

In November 2017 R/V Paamiut failed to comply with DNVGL standards for working in Artic waters. The management of Greenland Institute of Natural Resources (GN) decided not to spend more money on the vessel, but to go for building a new ship.

For the 2018-season, GN decided to charter Sjúrðarberg, a Faroese trawler of almost same dimensions as Paamiut, doing the normal surveys on the Greenland west coast.

To make the surveys as identical as possible this equipment was used from R/V Paamiut:

- Cosmos trawls
- All other equipment, such as bridles etc.
- Doors
- Marport sensors on doors and headline

Other steps taken ensuring the validity of received data:

- The wires/warps on Sjúrðarberg were same dimension (26mm) as used on Paamiut
- The distance between the hanging blocks was the same
- The Marport equipment on the bridge was set up and calibrated as on Paamiut
- All data from the tows were logged as normal procedure on Paamiut
- Skipper on Paamiut (Birgir Sivertsen) was on the bridge as supervisor, taking care of that all of the trawling was carried out as on Paamiut
- Chief Engineer from Paamiut was on board ensuring that all technical equipment performed as normal
- Crew from Paamiut worked together with the rest of the crew, ensuring that all maintenance of trawls etc. were carried out exactly as normal

To my best conviction regarding comparism, the surveys were executed in the best possible way, and I have absolutely no thoughts that this could be done otherwise or better.

Best regards

Kári Hansen, Chief Engineer



## Appendix II.

2018 biomass (in Kilotonnes) and abundance (in million individuals) indices for Elasmobranchs. Teleosts. Cephalopods and crustaceans excl. Shrimp species for the West and East-Greenland part of the GINR shrimp fish survey 0-600m including the West-Greenlandic Shelf part of NAFO div 0A (Canada).

Species	Year	NameLatin	'000 tons	millions	Remarks
GHL	2018	Reinhardtius hippoglossoides	43,735	371,369	
REB	2018	Sebastes mentella	29,55	59,397	
GSK	2018	Somniosus microcephalus			Errors...
POC	2018	Boreogadus saida	19,924	2251,824	
COD	2018	Gadus morhua	19,173	43,259	
CAS	2018	Anarhichas minor	15,812	7,452	
PLA	2018	Hippoglossoides platessoides	14,009	225,393	
MSC	2018	Myoxocephalus scorpius	9,299	50,641	
RRD	2018	Raja radiata	7,916	25,147	
TRN	2018	Triglops nybelini	7,467	1126,134	
REG	2018	Sebastes marinus	6,482	4,688	
SAN	2018	Ammodytes sp.	5,014	506,541	
CAD	2018	Anarhichas denticulatus	4,95	1,226	
CAA	2018	Anarhichas lupus	3,158	17,219	
CAP	2018	Mallotus villosus	3,035	298,978	
LEM	2018	Leptoclinus maculatus	2,842	704,074	
HYS	2018	Hyas sp.	2,75	57,748	
TRI	2018	Triglops sp.	1,186	147,228	
BEG	2018	Benthosema glaciale	1,047	580,962	
EUM	2018	Eumicrotremus spinosus	0,968	61,061	
LIG	2018	Liparis gibbus	0,821	26,53	
MYC	2018	Myctophidae	0,546	0,265	
RED	2018	Sebastes sp.	0,544	6,445	
LSE	2018	Lycodes seminudus	0,46	31,455	
ARA	2018	Artediellus atlanticus	0,447	43,823	
ELZ	2018	Lycodes sp.	0,447	13,555	
EPR	2018	Eumesogrammus praecisus	0,408	13,083	
LYV	2018	Lycodes vahlii	0,369	6,686	
LYN	2018	Lycodes eudipleurostictus	0,345	5,716	
GON	2018	Gonatus	0,323	27,934	
LLA	2018	Lumpenus lampretaeformis	0,316	16,478	
ARZ	2018	Arctozenus risso	0,289	10,179	
RHG	2018	Macrourus berglax	0,271	1,136	
LYP	2018	Lycodes pallidus	0,22	5,558	
GRC	2018	Gadus ogac	0,179	0,635	
MXS	2018	Myoxocephalus scorpioides	0,171	14,553	
LIF	2018	Liparis fabricii	0,168	18,325	
LUM	2018	Cyclopterus lumpus	0,153	0,212	
HAL	2018	Hippoglossus hippoglossus	0,143	0,075	
CAR	2018	Careproctus reinhardti	0,134	3,024	
TRM	2018	Triglops murrayi	0,129	13,369	
USK	2018	Brosme brosme	0,129	0,106	
LIT	2018	Liparis tunicatus	0,124	4,112	



RNG	2018	<i>Coryphaenoides rupestris</i>	0,122	1,817
ARU	2018	<i>Artediellus uncinatus</i>	0,119	10,93
LMC	2018	<i>Lampanyctus macdonaldi</i>	0,119	10,585
SQU	2018	<i>Cephalopoda</i>	0,1	1,469
TRP	2018	<i>Triglops pingelii</i>	0,078	19,058
ARS	2018	<i>Argentina silus</i>	0,076	1,727
RJJ	2018	<i>Raja jensenii</i>	0,072	0,093
RHB	2018	<i>Raja hyperborea</i>	0,071	0,054
CFB	2018	<i>Centroscyllium fabricii</i>	0,065	0,37
GYT	2018	<i>Gymnophanths tricuspidis</i>	0,063	2,428
COM	2018	<i>Cottunculus microps</i>	0,056	0,403
LPA	2018	<i>Lycodes paamiuti</i>	0,055	1,622
EUD	2018	<i>Leptagonus decagonus</i>	0,053	7,629
LYE	2018	<i>Lycodes esmarkii</i>	0,044	0,153
RBT	2018	<i>Raja bathyphila</i>	0,039	0,108
RFL	2018	<i>Raja fyllae</i>	0,038	0,107
HAD	2018	<i>Melanogrammus aeglefinus</i>	0,029	0,196
NOK	2018	<i>Notoscopelus kroyeri</i>	0,029	1,51
EDR	2018	<i>Eumicrotremus derjugini</i>	0,028	0,629
ICE	2018	<i>Icelus sp.</i>	0,027	5,021
WHB	2018	<i>Micromesistius poutassou</i>	0,026	0,096
ASP	2018	<i>Aspidophoroides monopterygius</i>	0,025	10,094
ROS	2018	<i>Rossia</i>	0,021	0,846
OCT	2018	<i>Octopoda</i>	0,02	1,683
MXQ	2018	<i>Myoxocephalus quadricornis</i>	0,018	0,104
MYX	2018	<i>Myxine glutinosa</i>	0,018	0,457
ONN	2018	<i>Gaidropsarus ensis</i>	0,016	0,361
ACT	2018	<i>Arctogadus glacialis</i>	0,015	0,307
ONA	2018	<i>Gaidropsarus argentatus</i>	0,015	0,208
SPU	2018	<i>Stichaeus punctatus</i>	0,014	0,866
ATP	2018	<i>Anotopterus pharao</i>	0,011	0,017
ANI	2018	<i>Anisarchus medius</i>	0,01	1,188
BAT	2018	<i>Bathylagus euryops</i>	0,01	1,322
KCT	2018	<i>Lithodes maja</i>	0,007	0,051
ULO	2018	<i>Ulcina olrikii</i>	0,006	1,712
SYN	2018	<i>Synaphobranchus kaupii</i>	0,005	0,191
GOF	2018	<i>Gonatus fabricii</i>	0,004	1,773
LEP	2018	<i>Lepidion eques</i>	0,004	0,017
STO	2018	<i>Stomias boa</i>	0,004	0,27
GYV	2018	<i>Gymnelus viridis</i>	0,003	0,627
ICB	2018	<i>Icelus bicornis</i>	0,003	0,37
LIP	2018	<i>Liparidae</i>	0,003	0,172
LYS	2018	<i>Lycenchelys sarsi</i>	0,003	0,474
SER	2018	<i>Serrivomer beanii</i>	0,003	0,11
MYJ	2018	<i>Myxine jespersenae</i>	0,002	0,074
RMA	2018	<i>Rossia macrosoma</i>	0,002	0,16
CAT	2018	<i>Anarhichas sp.</i>	0,001	0,793
LFA	2018	<i>Lumpenus fabricii</i>	0,001	0,08
LYR	2018	<i>Lycodes reticulatus</i>	0,001	0,047
MUG	2018	<i>Munida tenuimana</i>	0,001	0,073
CLM	2018	<i>Cyclothona microdon</i>	0	0,032
GAD	2018	<i>Gadidae</i>	0	0,024



GTA	2018	Gasterosteus aculeatus	0	0,087
GYM	2018	Gymnelus sp.	0	0,004
GYR	2018	Gymnelus retrodorsalis	0	0,006
ICS	2018	Icelus spatula	0	0,021
LYK	2018	Lycenchelys kolthoffi	0	0,033
MAV	2018	Maurolicus muelleri	0	0,033
MYP	2018	Myctophum punctatum	0	0,012
NY1	2018	New Species No 1	0	0,016
PAB	2018	Paraliparis bathybius	0	0,017
POP	2018	Polyipnus polli	0	0,018
SKA	2018	Raja sp.	0	0,065
SQT	2018	Decapodiformes	0	0,034