

NOT TO BE CITED WITHOUT PRIOR  
REFERENCE TO THE AUTHOR(S)

Northwest Atlantic



Fisheries Organization

Serial No. N6959

NAFO SCR Doc. 19/038

### SCIENTIFIC COUNCIL MEETING – JUNE 2019

An Assessment of the Status of Redfish in NAFO Division 30

by

D. W. Ings and R.M. Rideout

Science Branch, Department of Fisheries and Oceans  
P. O. Box 5667, St. John's, NL, Canada A1C 5X1

#### **Abstract**

There are two species of redfish, the deep sea redfish (*Sebastes mentella*) and the Acadian redfish (*Sebastes fasciatus*) that have been commercially fished and reported collectively in fishery statistics in Div. 30. Nominal catches have ranged between 3 000 t and 35 000 t since 1960. Up to 1986 catches averaged 13 000 tons and subsequently increased to a maximum value of 35 000 tons in 1988, exceeding the TAC by 21,000 tons. Following 1988 catches generally declined to a low of 3,000 tons in 1995, partly due to reductions in foreign allocations within the Canadian fishery zone since 1993. There was an overall increasing trend in catch over the next five years to a high value of 20 000 tons in 2001. Then, catch declined to 4000 t in 2008 and has remained below 9500 t since that time with a mean of approximately 7500 t taken annually during 2016-2018. Assessment of this stock has been based primarily on research survey data due to variable commercial indices and fleets prosecuting different areas of the stock. It is difficult to reconcile year to year changes in the indices, but generally, the Canadian spring survey biomass index suggests the stock may have increased between the early and mid-1990s, fluctuated over 100,000 tons from 1994 to 1999 and declined to 20,000 t in 2002. Then, the index increased to 230,000 t in 2012, which approximates the time series high, but it has declined to lower levels since then. The index mean was 62,000 t for 2016-2018. The Canadian autumn survey, while more stable in the early 1990s, generally supports the pattern of the spring survey index indicating a gradual increase from 18,000 t in 2003 to 150 000 t in 2010 with similar, high values until 2012. Index values were considerably lower from 2015 to 2018 ranging between 30,000 t and 53,000 t. Canadian RV surveys do not adequately sample fish greater than 25 cm, which in some years comprise a large portion of the fishery, making interpretation of survey estimates difficult. A strong year class born in the early 2000s has grown sufficiently to provide some individuals exceeding the small fish protocol of 22 cm. There is concern that there has been little sign in recent surveys of size groups smaller than 17 cm despite using a shrimp trawl, which is very effective at catching small fish.

#### **Introduction**

Two species of redfish have been commercially fished in Div. 30, the deep sea redfish (*Sebastes mentella*) and the Acadian redfish (*Sebastes fasciatus*). The two species are difficult to distinguish based on external characteristics, and as a consequence are reported collectively as “redfish” in the commercial fishery statistics. Most of the habitable redfish area in Div. 30 lies within Canada’s 200 mile exclusive fishery zone and has been subject to management regulation since 1974. Approximately 8% of the habitable redfish area within Div. 30 lies within the NAFO Regulatory Area (NRA) and was brought under TAC regulation starting in 2005.



### Nominal Catches and TACs

Nominal catches have ranged between 3 000 tons and 35 000 tons since 1960 (Table 1, Fig. 1). Catches averaged 13 000 t up to 1986 and then increased to 27 000 t in 1987 and 35 000 t in 1988 (exceeding TACs by 7 000 t and 21 000 t, respectively). Catches declined to 16 000 t by 1993 then to about 3 000 t in 1995, partly due to reductions in foreign allocations within the Canadian fishery zone since 1993. Catches increased to 20 000 t by 2001 and subsequently declined to 4000 t in 2008 and have been in the 6000 to 9000 t range since 2008.

The large redfish catches in 1987 and 1988 were due mainly to increased activity in the NRA by South Korea and non-Contracting parties (NCPs), primarily Panama. There has been no activity by NCPs in the NRA since 1994. Estimates of under-reported catch which have occurred primarily before 1995, have ranged from 200 tons to 23 500 tons. There have also been estimates of over-reported catch in the recent period since 2000, with a maximum value of 4 300 t in 2003. There were no alternate sources of catch estimates available until 2016 to compare with the reported catch

A TAC of 16 000 tons was first implemented by Canada within its 200-mile limit in 1974. The TAC was increased in 1978 to 20 000 tons and generally remained at that level through to 1987. The TAC for 1988 was reduced to 14 000 tons and remained unchanged until 1994 when it was reduced to 10 000 tons as a precautionary measure and maintained at that level to 2003. In September 2004, the NAFO Fisheries Commission adopted TAC regulation for redfish in 30, implementing a level of 20 000 tons for the entire division in 2005 and remaining in effect up to 2019.

### Description of the Fishery

Russia predominated in the 30 redfish fishery up until 1993 (Table 2) and generally caught about 50% of the total non-Canadian allocation, which accounted for about 2/3 of the Canadian TAC. Russia and Cuba, impacted by the reduction and eventual elimination of foreign allocations by Canada, ceased directed fishing in 1994. Russia resumed directed fishing in 2000, rapidly increasing their catch from 2 200 tons to about 11 000 tons from 2001-2003 before a large reduction in catch to only 240 t in 2004. Generally, catches increased from 50 t in 2007 to about 2000 tons in 2016, but decreased to 350 t in 2018. Portugal began fishing redfish in Div. 30 in 1992 and averaged about 1 800 tons between 1992 and 1998. Their catches escalated to 5 500 tons in 1999 and have ranged between 3 200 – 6 400 tons thereafter with 3000 tons taken in 2018. Spain, which had taken less than 50 tons before 1996, increased catches from 1 200 tons in 1997 to a peak of 4 500 tons in 1999 with a subsequent decline to 300 tons in 2004. Since then, Spanish catch ranged between 600 – 1 800 with about 1800 tons taken in 2018.

Canada has had limited interest in a fishery in Div. 30 because of small sizes of redfish encountered in areas suitable for trawling. Canadian landings were less than 200 tons annually from 1983-1991. In 1994, Canada took 1 600 tons due to improved markets related to lobster bait, but reduced catch to less than 200 tons in 1995. Between 1996 and 1999 Canadian catches alternated between levels of about 8 000 tons and 2 500 tons based on market acceptance for redfish near the 22 cm size limit regulated within Canada. From 2000-2006 Canada averaged about 3 600 tons, followed by a decrease to 1000 tons in 2007. Canadian landings decreased further in 2008 and have been below 500 tons since then with only 411 tons reported for 2018.

The redfish fishery in Div. 30 has generally occurred throughout the year (Table 3a). The vast majority (>90%) of catch has been taken via bottom trawling by Canadian, Portuguese, Russian and Spanish fleets (Table 3b). Catches via midwater trawl prior to 2005 were taken predominantly by Russia.

### **Commercial Fishery Data**

#### Commercial fishery sampling

Sampling of the redfish fisheries was conducted by Spain (González-Costas *et al.*, MS 2019), Portugal (Vargas *et al.*, MS 2019) and Estonia (Hubel, MS 2019) from the 2018 trawl fishery (Fig. 2). The Portuguese fleet fished between 214 and 606 m. Sampling details for the Canadian fleets over time are given in Table 4. Lengths between 19-23 cm (range 15-35 cm) dominated the Portuguese catch. The Spanish catch was



dominated by 21-28 cm fish (range 10-35 cm) while the Estonian catch had a modal length of 24-26 cm (range 16-40 cm).

A compilation of catch at length from various fleets from 1995 to 2004 suggested that the size composition has changed over the time period with fleets catching a larger portion of fish >25 cm prior to 1998 (Power, MS 2005). These size compositions were converted to catch at length for 2011 to 2018 and compared to Canadian RV survey numbers at length in Figure 3.

### **Research Survey Data**

#### Abundance and Biomass Indices

Stratified random groundfish surveys have been conducted by Canada in the spring and autumn in Div. 30 since 1991, with regular coverage of depths to 730 m (and sporadic coverage of deeper strata in the autumn). In addition, a summer survey was conducted in 1993. Surveys utilized an Engel 145 otter trawl (1.75 n. mi. standard tow) from 1991 to spring 1995 and a Campelen 1800 shrimp trawl (0.75 n. mi. standard tow) from autumn 1995 to the present. The Engel 145 data were converted into Campelen 1800 trawl equivalent data based on comparative fishing trials (see Power and Atkinson MS 1998a). Vessel problems during the 2006 spring survey resulted in the completion of only a single tow in redfish depths. There was no fall survey in 2014. See Rideout and Ings (2019) for details of recent Canadian surveys.

Abundance (Tables 5 and 7) and biomass (Tables 6 and 8) estimates based on spring and autumn data from the Canadian surveys demonstrate large fluctuations between seasons and years for some strata. This is usually accounted for by the influence of one or two large sets on the survey. It is difficult to reconcile year to year changes in the indices, but generally, the spring survey biomass index (Fig. 3) suggests the stock may have increased between the early and mid-1990s, and subsequently declined to 2002 (20 000 t). The low 1997 value is considered a sampling anomaly. The biomass index increased in 2003 and again in 2004 (85,000 t) but in 2004 was influenced by one large set in a stratum that represented 40% of the biomass index. The biomass index has alternated between 60 000 t and 90 000 t between 2005 and 2009 (no data for 2006) and increased steadily to an estimate of 232 000 t in 2012. During 2013 to 2018, the biomass index was variable at lower values with 34,000 t observed in 2018. Biomass estimates from the autumn surveys, while more stable in the early 1990s, generally support the pattern of the spring survey index with a gradual increase in the biomass index from 2003 (18 600 t) to 2012 (154 000 t). Lower values (< 90,000 t) were observed from 2013 to 2018 with a value of 53,000 t in 2018. It should be noted that the 1996 autumn estimate does not include important strata that were not sampled due to problems on the survey. Also, there was no autumn survey in 2014.

Density estimates per stratum were generally lower in the NAFO Regulatory Area (denoted in Tables 5 - 8 as strata 354, 355, 356, 721, 722) compared to strata inside the Canadian 200 mile exclusive fishery zone, although it should be pointed out that part of these NRA strata overlap the Canadian zone. Estimates of the proportion of survey biomass within the NRA have ranged from 3% to 53% (average 15.6%) for the spring survey and 4% to 36% (average 18.1%) for the autumn survey.

Data were available from EU-Spain spring surveys conducted in the NAFO regulatory area (NRA) of Div. 30 from 1997 to 2018. These surveys use the same stratification scheme as the Canadian surveys and the area of redfish habitat covered in Div. 30 is less than 8% compared to the Canadian surveys for strata <732m. The surveys covered depths to 1500m (800 fathoms). Until 2001, these surveys were conducted with a Pedreira type bottom trawl and thereafter with a Campelen trawl similar to that used in Canadian surveys. The data prior to 2001 were converted into Campelen equivalent units.

The biomass indices for Div. 30 from the EU-Spain survey generally decreased from a time-series peak of 773 kg per tow in 2010, to 129 kg per tow in 2014. The 2015 value was somewhat higher at 341 kg per tow, but the index decreased further to 32 kg per tow in 2018 (Fig. 4). Generally, indices are consistent from Canadian spring, Canadian autumn and EU-Spain surveys (Fig. 5).

### Recruitment

Size distributions from the Canadian spring and autumn surveys plus the EU-Spain survey in terms of mean number per tow at length indicated that there was a new relatively large pulse at 17cm in the 2007 surveys corresponding to a year class born in the early 2000s that has remained the dominant mode to 2018 (Fig. 6). Although their presence was detected at smaller sizes in previous surveys, the sudden increase in density at 17cm in 2007 is unusual. At a modal length of 23-27 cm in 2018, this pulse is now recruiting fully to the fishery. No strong recruitment pulses are apparent at smaller sizes in the Canadian or EU-Spain surveys.

The size distributions of the survey catches indicate only a narrow range of sizes caught each year in Div. 30. Generally fish smaller than about 10 cm and larger than about 25 cm are absent in survey catches from 1991-2000 which cover strata down to 732 m (400 fathoms). It is well documented that the Engel survey gear (e.g. Power MS 1995) and the Campelen survey gear (e.g. Power and Atkinson, MS 1998b) can catch both smaller (than 10 cm) and larger (than 25 cm) redfish. Length sampling from the commercial fisheries in the mid-1990s reveals a higher proportion of fish greater than 25 cm compared to the survey catches (see Power, MS 2005). Therefore, it appears that fish sizes outside this range, especially fish greater than 25 cm, are generally unavailable to the gear in this area. The reasons for this are unknown but may be related to distribution relative to trawlable bottom.

### **Estimation of Stock Parameters**

#### Catch/Biomass ratio

A fishing mortality proxy was derived by simple catch to survey biomass ratios. In deriving a fishing mortality proxy, and because most of the catch is taken in the last three quarters of the year, the catch in year "n" was divided by the average of the Canadian Spring (year = n) and Autumn (year = n-1) survey biomass estimates to better represent the relative biomass at the time of the year before the catch was taken. Survey catchability ( $q$ ) for redfish is not known but assumed to be less than one. All fish sizes were included in the survey biomass estimate. The results (Fig. 7) suggest that relative fishing mortality increased from 1998 to the highest estimate in the series in 2002. This relatively high value was maintained in 2003 but declined substantially in 2004. In 2005, relative fishing mortality increased once more and was around the series average. The 2006 estimate of fishing mortality was calculated using only the autumn survey biomass. In 2007-2008 the estimate of fishing mortality dropped to some of the lowest levels since the mid 1990s and remained at similar levels up to 2014. The 2018 estimate returned to low values following a slight increase from 2014 to 2016.

#### Size at maturity

No new maturity at length data were analysed for this assessment. However, based on previous analyses of size at maturity for this stock estimated L<sub>50</sub> is about 28 cm for females and 21 cm for males (Power and Atkinson MS 1998). Based on current catches dominated by lengths between 18cm-27 cm, it is clear that the fishery is based predominantly on immature fish.

### **State of the Stock**

It is still not possible to determine absolute size of the stock. RV survey estimates suggest that stock size has decreased from near time series highs in the late 2000s. Using the ratio of catch:biomass as a proxy for fishing mortality suggests a value less than 0.17 for 2007-2018, among the lowest levels observed since the mid 1990s. The appearance of a relatively strong year class (born in the early 2000s) in the 2007 and subsequent surveys constitutes the best sign of recruitment to the population since the relatively strong 1988 year-class. The bulk of the catches in recent years are comprised of fish less than 27 cm, suggesting that these fisheries continue to be comprised of predominantly immature fish (Power and Atkinson 1998 MS).

## References

- González-Costas, F., G. Ramilo, E. Román, J. Lorenzo, A. Gago, D. González-Troncoso, J. L. del Rio and M. Sacau. Spanish Research Report for 2018. SCS Doc. 19/10. Serial No. N6922, 44 p.
- González-Troncoso, D., I. Garrido and A. Gago MS 2019. Yellowtail flounder, redfish (*Sebastes spp.*), and witch flounder indices from the Spanish survey conducted in Divisions 3NO of the NAFO Regulatory Area. NAFO SCR Doc. 19/18. Serial No. N6934, 35 p.
- Power, D. MS 1995. Status of redfish in Subarea 2 + Division 3K. DFO Atl. Fish. Res. Doc. 95/25, 25 p.
- Power, D., and D. B. Atkinson. MS 1998a. Update on the status of Redfish in 3O. CSAS Res. Doc. 98/110, 20 p.
- Power, D., and D. B. Atkinson. MS 1998b. The status of Redfish in Unit 2 (Laurentian Channel Management Unit). CSAS Res. Doc. 98/21, 41 p.
- Power, D., MS 2005. An Assessment of the status of Redfish in NAFO Division 3O. NAFO SCR Doc. 05/59. Serial No. N5145, 19 p.
- Rideout, R. M., and D. W. Ings. 2019. Temporal and spatial coverage of Canadian (Newfoundland and Labrador region) spring and autumn multi-species RV bottom trawl surveys, with an emphasis on surveys conducted in 2018. NAFO SCR Doc. 19/15. Serial No. N6931, 59 p.
- Hubel, K. Estonian research report for 2018. SCS Doc. 19/6. Serial No. N6917, 17 p.
- Vargas, J., R. Alpoim, E. Santos and A. M. Ávila de Melo MS 2019. Portuguese Research Report for 2018. NAFO SCS Doc. 19/09. Serial No. N6921, 32 p.
- Vascov, A. A. MS 2003. Distribution of Redfish in Division 3O Based on Data from Russian Trawl Surveys in 1983-1993. NAFO SCR Doc. 03/12. Serial No. N4818, 13pp.

Table 1. Estimated catches (t) and TACs of redfish in Div. 3O.

Year	Canada	Others	Catch <sup>a</sup>	TAC	Year	Canada	Others	Catch <sup>a</sup>	TAC
1960	100	4900	5000		1990	155	8887	14242	14000
1961	1000	10000	11000		1991	28	7533	8461	14000
1962	1046	6511	7557		1992	1219	12149	15268	14000
1963	2155	7025	9180		1993	698	12522	15720	14000
1964	1320	14724	16044		1994	1624	3004	5428	10000
1965	203	19588	19791		1995	177	2637	3214	10000
1966	107	15198	15305		1996	7255	2390	9845	10000
1967	645	18392	19037		1997	2554	2558	5112	10000
1968	52	6393	6445		1998	8972	4380	14052	10000
1969	186	15692	15878		1999	2344	10249	12593	10200
1970	288	12904	13192		2000	2206	10584	10003	10000
1971	165	19627	19792		2001	4893	17681	20274	10000
1972	508	15609	16117		2002	3000	16453	17234 <sup>b</sup>	10000
1973	133	8664	8797		2003	3125	18466	17246	10000
1974	91	13033	13124	16000	2004	2616	3848	3753	10000 <sup>c</sup>
1975	103	15007	15110	16000	2005	5501	6409	11305	20000
1976	3664	11684	15348	16000	2006	3580	7455	12610	20000
1977	2972	7878	10850	16000	2007	1053	6472	5179	20000
1978	1841	5019	6860	16000	2008	203	4816	4020	20000
1979	6404	11333	17737	20000	2009	255	6233	6431	20000
1980	1541	15765	17306	21900	2010	260	6285	5234	20000
1981	2577	10027	12604	20000	2011	97	5875	5972	20000
1982	491	10869	11360	20000	2012	0	6967	6967	20000
1983	7	7133	7340	20000	2013	75	7720	7795	20000
1984	167	9861	16978	20000	2014	374	7150	7524	20000
1985	104	8106	12860	20000	2015	283	8073	8356 <sup>d</sup>	20000
1986	141	10314	11055	20000	2016	199	8818	9017 <sup>d</sup>	20000
1987	183	12837	27170	20000	2017	245	7267	7512 <sup>d</sup>	20000
1988	181	11111	34792	14000	2018	411	5710	6120 <sup>d</sup>	20000
1989	27	11029	13256	14000					

<sup>a</sup> Totals from 1983 to 2010 may include adjustments for estimated catches from various sources<sup>b</sup> Midpoint of estimates ranging between 16100-18400<sup>c</sup> Prior to 2005 TACs were set by Canada within its fisheries jurisdiction<sup>d</sup> Based on daily catch reports from the NRA and Statlant 21A data for Canada

Table 2. Reported and estimated catches (t) of redfish in Div. 3O by country and year since 1992.

Year	Canada (M)	Canada (N)	France (SPM)	Japan	Portugal	Spain	Russia	Cuba	Ukraine	Estonia	Lithuania	Korea (S)	Faroe Islands	Latvia	Total	TAC	a
1992	27	1192	-	125	1468	-	5845	2776	-	-	-	1935	-	-	15268	14000	
1993	21	677	-	159	4794	-	6887	665	-	-	-	17	-	-	15720	14000	
1994	779	845	-	-	2918	26	60	-	-	-	-	-	-	-	5428	10000	
1995	4	173	-	264	1935	22	416	-	-	-	-	-	-	-	3214	10000	
1996	2124	5131	-	417	1635	338	-	-	-	-	-	-	-	-	9845	10000	
1997	693	1861	134	285	894	1245	-	-	-	-	-	-	-	-	5112	10000	
1998	2851	6121	266	355	1875	1884	-	-	-	-	-	-	-	-	14052	10000	
1999	317	2027	-	-	5469	4549	231	-	-	-	-	-	-	-	12593	10200	
2000	1326	880	-	-	4555	3747	2233	-	-	49	-	-	-	-	10003	10000	
2001	336	4557	-	-	3537	2792	11343	-	-	9	-	-	-	-	20274	10000	
2002	12	2988	-	-	4610	660	11182	-	-	-	1	-	-	-	17234	10000	
2003	32	3093	-	-	6382	1289	10794	-	1	-	-	-	-	-	17246	10000	
2004	276	2340	-	2	3279	320	242	-	3	2	-	-	-	-	3753	10000	
2005	137	5364	-	1	4555	1683	170	-	-	-	-	-	-	-	11305	20000	
2006	-	3580	-	0	5184	1294	977	-	-	-	-	-	-	-	12610	20000	
2007	-	1053	-	61	4755	1502	54	-	-	100	-	-	-	-	5179	20000	
2008	0	203	-	-	3850	603	82	-	-	42	139	-	100	-	4020	20000	
2009	-	255	-	-	4273	1691	169	-	-	100	-	-	-	-	6431	20000	
2010	218	42	-	-	3853	1692	474	-	-	103	-	-	163	-	6545	20000	
2011	-	97	-	-	4006	1661	570	-	-	121	5	-	-	82	6542	20000	
2012	-	0	-	-	4142	1572	971	-	-	181	-	-	101	-	6967	20000	
2013	75	0	-	-	4820	1135	1438	-	-	269	-	-	58	-	7795	20000	
2014	340	34	-	-	4720	932	1271	-	-	227	-	-	-	-	7524	20000	
2015	252	31	-	-	4659	1510	1086	-	-	817	-	-	-	-	8355	20000	
2016	178	21	-	29	4348	1678	2031	31	-	701	-	-	-	-	9017	20000	
2017	213	32	-	6	4079	1826	753	-	-	603	-	-	-	-	7512	20000	
2018	357	54	-	5	3019	1813	350	-	-	522	-	-	-	-	6120	20000	

<sup>a</sup> Prior to 2005 TACs were set by Canada within its fisheries jurisdiction

Table 3a. Nominal reported catches (t) of redfish in Div. 3O by month and year since 1992.

Year	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Unk	Total
1992	0	57	14	10	635	3262	2520	1808	896	1261	797	2108		13368
1993	226	14	754	817	2089	1601	1887	2068	1809	829	630	496		13220
1994	60	93	742	1609	236	83	-	68	1000	540	19	178		4628
1995	7	125	145	2	45	28	56	765	645	879	107	10		2814
1996	0	0	89	119	166	46	773	882	1685	2864	1539	1482		9645
1997	4	0	10	34	86	417	1298	909	622	1274	409	49		5112
1998	40	193	216	279	1329	2723	1924	953	1280	1964	2275	176		13352
1999	100	139	262	463	527	942	1644	2513	2298	2056	1434	215		12593
2000	80	92	943	739	1077	1844	1088	1254	1545	2068	1814	246		12790
2001	31	193	1228	1909	1958	2750	1257	1421	2020	4048	3472	2287		22574
2002	1850	1269	2356	1904	1490	1423	300	2085	2000	2309	1402	1064		19452
2003	453	1212	910	1392	2361	3232	2826	961	2294	2212	2484	1149		21486
2004	323	343	597	794	318	180	336	400	651	1393	859	270		6464
2005	100	12	241	169	436	371	2114	2115	1100	1288	1933	2029		11908
2006	743	485	49	1044	617	654	885	1436	1303	1786	1566	467		11035
2007	225	132	214	475	858	657	950	1298	966	859	358	479		7471
2008	124	328	56	289	187	72	97	433	350	689	1247	966		4838
2009	223	660	136	307	525	901	310	1118	778	368	754	234		6314
2010	242	298	303	211	435	1148	1246	201	1205	879	189	186		6543
2011	331	574	466	734	764	275	405	768	528	380	373	374		5972
2012	242	60	253	130	50	135	170	163	91	202	117	240		1853
2013	185	483	538	316	593	748	76	831	1112	923	990	923		7718
2014	130	181	665	896	1069	946	655	359	937	770	236	550		7394
2015	25	269	395	422	1259	1444	403	987	659	896	226	934		7919
2016	59	68	533	856	211	92	0	225	1	204	0	16		2265

Table 3b. Nominal reported catches (t) of redfish in Div. 3O by gear since 1992.

Year	Otter Trawls				Total
	Bottom	Midwater	Gillnets	Misc.	
1992	10046	3292	1	29	13368
1993	11997	1214	-	9	13220
1994	3085	1498	26	19	4628
1995	2221	525	26	42	2814
1996	9303	335	7	-	9645
1997	5091	10	2	9	5112
1998	13352				13352
1999	11623	970			12593
2000	12750	39		1	12790
2001	21945	629			22574
2002	16586	2866			19452
2003	19226	2260			21486
2004	6308	156	0	0	6464
2005	11908	0	0	0	11908
2006	10058	0	0	977	11035
2007	7525	0	0	0	7525
2008	4880	0	0	0	4880
2009	6314				6314
2010	6545		0		6545
2011	5890	82			5972
2012	1853		0		1853
2013	7718				7718
2014	7394				7394
2015	7919				7919
2016	1229	1036			2265



Table 4. Commercial sampling of redfish catches from CAN (N) in 2007-2018.

Year	Observer samples					Port samples							
	Month	Samples	n	depth	min	depth	max	Samples	n	depth	min	depth	max
2007	APR	2	489	411		457		-	-	-	-	-	-
	MAY	5	1325	278		508		2	602	365		457	
2008	APR	-	-	-		-		2	622	487		487	
	MAY	2	488	361		450		-	-	-		-	
2009	NOV	6	572	357		505		-	-	-		-	
2010	OCT	5	1190	362		477		-	-	-		-	
2011		-	-	-		-		-	-	-		-	
2012		-	-	-		-		-	-	-		-	
2013		-	-	-		-		-	-	-		-	
2014		-	-	-		-		-	-	-		-	
2015		-	-	-		-		-	-	-		-	
2016		-	-	-		-		-	-	-		-	
2017	MAR	1	266	538		538		-	-	-		-	
2018	MAR	1	255	277		277		-	-	-		-	
	APR	2	501	348		466		-	-	-		-	

**Table 5.** Mean number per standard tow from Canadian SPRING surveys in Div. 30 covering strata from 93 to 731 m (400ftm.). Dashes (---) represent unsampled strata. Number of successful sets in brackets. Data from 1991-1995 are Campelen trawl equivalent units (see text). Data from 1996 to present are actual Campelen data. G=Gadus Atlantica, W=Wilfred Templeman, A=Alfred Needler.

Stratum	Depth Range (M)	Area sq mi	Area within NRA sq mi	Area within NRA		May3-11	May2-13	May5-18	May14-22	May13-27	May22-30	May-Jun 1997-Q2	May-Jun 1998-Q2	May-Jun 1999-Q2	May-Jun 2000-Q2	May-Jun 2001-Q2
				W105	W119-20	W136-7	W153	W168-69	W188	W204	W221-2	W238	W315-16	W365,367		
329	093-183	1721	0	0.00	13.3 (9)	0.0 (8)	0.0 (6)	169.6 (5)	19.6 (5)	0.0 (6)	33.5 (6)	0.0 (7)	0.3 (6)	0.0 (5)	0.0 (5)	
332	093-183	1047	0	0.00	35.5 (6)	1.4 (5)	0.0 (4)	0.0 (4)	1177.8 (4)	181.8 (4)	7.3 (3)	348.0 (4)	899.0 (4)	43.5 (4)	44.0 (3)	
337	093-183	948	0	0.00	607.2 (5)	6.5 (4)	3.0 (2)	0.0 (3)	3462.8 (4)	5.0 (3)	2.0 (3)	703.5 (4)	339.0 (3)	207.5 (4)	48.7 (3)	
339	093-183	585	0	0.00	0.0 (3)	0.0 (2)	0.0 (2)	0.0 (2)	0.0 (2)	0.0 (2)	0.0 (2)	0.0 (2)	0.0 (2)	0.0 (2)	0.0 (2)	
354	093-183	474	246	0.52	0.0 (3)	0.0 (2)	2537.0 (2)	0.0 (2)	0.0 (3)	2.5 (2)	0.0 (2)	422.9 (2)	1006.5 (2)	4.5 (2)	81.1 (2)	
333	185-274	151(147)	0	0.00	1089.0 (2)	3240.0 (2)	8184.5 (2)	50275.0 (2)	979.5 (2)	870.1 (2)	231.9 (2)	4321.3 (2)	5502.4 (2)	1355.9 (2)	1525.5 (2)	
336	185-274	121	0	0.00	187.5 (2)	688.5 (2)	4496.5 (2)	9955.5 (2)	83150.0 (2)	1360.6 (2)	139.1 (2)	34839.0 (2)	1682.7 (2)	1714.3 (2)	1742.0 (2)	
355	185-274	103	74	0.72	119.5 (2)	111.0 (2)	7307.0 (2)	5829.0 (2)	1928.0 (2)	36488.9 (2)	306.2 (2)	5152.0 (2)	2191.6 (2)	4161.1 (2)	407.5 (2)	
334	275-366	92(96)	0	0.00	733.0 (2)	223.0 (2)	837.0 (2)	1179.0 (2)	159.0 (2)	1206.8 (2)	286.2 (2)	733.5 (2)	2515.2 (2)	3960.3 (2)	730.9 (2)	
335	275-366	58	0	0.00	39.7 (3)	265.3 (3)	582.5 (2)	6992.0 (2)	2267.0 (2)	15196.4 (2)	531.6 (2)	5796.0 (2)	8671.3 (2)	957.6 (2)	4730.6 (2)	
356	275-366	61	47	0.77	444.0 (2)	805.5 (2)	2552.5 (2)	883.0 (2)	3980.0 (2)	4347.0 (2)	133.6 (2)	3990.2 (2)	9384.4 (2)	24603.5 (2)	503.2 (2)	
717	367-549	93(166)	0	0.00	1461.5 (2)	324.0 (2)	279.0 (2)	1269.0 (2)	312.5 (2)	597.0 (2)	3398.6 (2)	483.6 (2)	3239.6 (2)	740.9 (2)	139.5 (2)	
719	367-549	76	0	0.00	277.0 (2)	88.5 (2)	497.5 (2)	1985.0 (2)	331.0 (2)	440.5 (2)	374.3 (2)	1098.0 (2)	1487.6 (2)	1685.1 (2)	1755.4 (2)	
721	367-549	76	58	0.76	176.0 (2)	4369.0 (2)	449.0 (2)	108.0 (2)	7596.5 (2)	575.5 (2)	262.6 (2)	543.0 (2)	3263.2 (2)	687.8 (2)	541.1 (2)	
718	550-731	111(134)	0	0.00	56.5 (2)	17.5 (2)	174.0 (2)	349.0 (2)	15.5 (2)	47.8 (2)	60.8 (2)	79.3 (3)	35.4 (3)	369.0 (3)	22.5 (2)	
720	550-731	105	0	0.00	35.5 (2)	113.0 (2)	24.0 (2)	34.5 (2)	40.0 (2)	284.6 (2)	63.2 (2)	35.6 (2)	221.3 (2)	53.6 (2)	52.1 (2)	
722	550-731	93	71	0.76	186.5 (2)	79.0 (2)	76.0 (2)	327.5 (2)	17.0 (2)	80.0 (2)	91.8 (2)	334.0 (2)	47.5 (2)	640.2 (2)	447.9 (2)	
Total:		6011	496	8.25												
Upper ( 95% CI )					465.3	495.8	1955.9	3238.5	4318.0	8884.4	1255.6	10277.2	1348.6	895.5	288.5	
Weighted mean ( by area )					190.99	180.3	698.4	1748.5	2662.6	953.2	141.7	1250.0	869.5	571.3	204.7	
Lower ( 95% CI )					-83.3	-135.1	-559.1	258.6	1007.2	-6978.1	-972.1	-7777.3	390.4	247.1	121.0	
<b>SURVEY ABUNDANCE(x10<sup>6</sup>)</b>					155.4	146.7	568.3	1445.8	2201.7	788.2	117.2	1033.6	719.0	472.4	169.3	
<b>ABUNDANCE within NRA</b>					7.3	42.0	181.1	69.1	106.1	405.0	7.0	100.2	143.6	213.3	18.8	
%within NRA					4.7	28.6	31.9	4.8	4.8	51.4	6.0	9.7	20.0	45.2	11.1	



**Table 5** continued.

Stratum	Depth Range (M)	Area sq mi	Area within NRA sq mi	Area within NRA		% May											
				May 2002-Q2 W419,421	May 2003-Q2 W479-480	May 2004-Q2 W546-547	May 2005-Q2 W618-621	May 2006-Q2 W693,A729	May 2007-Q2 W759-761	May 2008-Q2 W827	May 2009-Q2 A904-05	May 2010-Q2 A932-33	May 2011-Q2 A403-04	May 2012-Q2 A417-19			
329	093-183	1721	0	0.00	0.0 (5)	80.0 (5)	0.0 (5)	0.2 (5)	—	25.0 (5)	399.8 (5)	0.0 (5)	3.2 (5)	1271.1 (5)	3.4 (5)		
332	093-183	1047	0	0.00	23.7 (3)	79.7 (3)	94.8 (3)	69.3 (3)	—	83.7 (3)	0.7 (3)	8.5 (3)	78.8 (3)	798.2 (3)	3020.2 (3)		
337	093-183	948	0	0.00	2.7 (3)	429.7 (3)	1048.8 (3)	18.5 (3)	—	2886.0 (3)	29.0 (3)	13.0 (3)	3314.6 (3)	78.8 (3)	1037.0 (3)		
339	093-183	585	0	0.00	0.5 (2)	0.0 (2)	0.0 (2)	0.5 (2)	0.4 (2)	0.0 (2)	0.0 (2)	0.0 (2)	0.0 (2)	0.0 (2)	0.5 (2)		
354	093-183	474	246	0.52	0.0 (2)	3.0 (2)	1.0 (2)	433.3 (2)	—	27.5 (2)	0.0 (2)	0.0 (2)	3465.0 (2)	3050.8 (2)	3745.5 (2)		
333	185-274	151(147)	0	0.00	941.5 (2)	534.3 (2)	2759.2 (2)	5329.0 (2)	—	1683.1 (2)	4605.1 (2)	614.0 (2)	8921.3 (2)	8974.2 (2)	4727.6 (2)		
336	185-274	121	0	0.00	1048.0 (2)	1456.5 (2)	12646.5 (2)	4701.9 (2)	—	694.2 (2)	8781.5 (2)	726.9 (2)	10791.5 (2)	7300.4 (2)	20363.5 (2)		
355	185-274	103	74	0.72	515.2 (2)	1191.0 (2)	1321.6 (2)	643.6 (2)	—	540.0 (2)	3929.5 (2)	9261.3 (2)	2168.0 (2)	7279.1 (2)	4179.7 (2)		
334	275-366	92(96)	0	0.00	916.5 (2)	3154.1 (2)	1387.1 (2)	2364.5 (2)	—	617.4 (2)	9238.5 (2)	16761.3 (2)	1790.8 (2)	4545.8 (2)	5887.9 (2)		
335	275-366	58	0	0.00	4291.9 (2)	1155.1 (2)	1037.1 (2)	2563.6 (2)	—	3760.2 (2)	2560.9 (2)	13134.5 (2)	10498.7 (2)	28306.9 (2)	7583.5 (2)		
356	275-366	61	47	0.77	2020.9 (2)	521.3 (2)	658.0 (2)	3515.6 (2)	—	924.8 (2)	1489.8 (2)	3027.2 (2)	1580.3 (2)	5820.2 (2)	10444.5 (2)		
717	367-549	93(166)	0	0.00	242.0 (2)	584.0 (2)	1349.7 (2)	1211.4 (2)	—	2633.9 (2)	2404.6 (2)	2612.7 (2)	1325.7 (2)	4881.9 (2)	3101.3 (2)		
719	367-549	76	0	0.00	208.8 (2)	602.5 (2)	326.5 (2)	1346.0 (2)	—	2653.8 (2)	1508.8 (2)	3134.0 (2)	4963.6 (2)	375.2 (2)	8938.9 (2)		
721	367-549	76	58	0.76	94.7 (2)	304.4 (2)	116.5 (2)	566.2 (2)	—	470.0 (2)	253.6 (2)	1688.0 (2)	1122.4 (2)	501.0 (2)	2148.0 (2)		
718	550-731	111(134)	0	0.00	79.0 (2)	0.0 (2)	30.2 (2)	55.1 (2)	—	20.5 (2)	18.0 (2)	58.7 (2)	0.0 (2)	39.9 (2)	70.7 (2)		
720	550-731	105	0	0.00	93.1 (2)	31.5 (2)	42.0 (2)	23.1 (2)	—	112.4 (2)	32.6 (2)	270.5 (2)	191.3 (2)	—	239.0 (2)		
722	550-731	93	71	0.76	86.7 (2)	71.9 (2)	69.5 (2)	60.9 (2)	—	21.8 (2)	2.0 (2)	52.0 (2)	3.7 -3	15.2 (2)	202.4 (2)		
Total:		6011	496	8.25													
Upper ( 95% CI )					234.6	490.9	1861.6	1909.1		2655.9	1302.7	1033.3	3985.2	3157.5	4094.6		
Weighted mean ( by area )					149.3	263.6	610.9	443.6		714.0	752.9	757.0	1546.1	1830.9	2091.0		
Lower ( 95% CI )					64.1	36.3	-639.8	-1021.9		-1228.0	203.2	480.7	-893.0	504.4	87.5		
<b>SURVEY ABUNDANCE(x10<sup>6</sup>)</b>					123.5	218.0	505.1	366.8		590.4	622.6	625.9	12784.4	1487.5	1729.0		
<b>ABUNDANCE within NRA</b>					19.9	18.7	19.4	49.1		16.4	51.7	127.8	158.5	219.1	256.0		
<b>% within NRA</b>					16.1	8.6	3.8	13.4		2.8	8.3	20.4	1.2	14.7	14.8		



**Table 5** continued.

Stratum	Depth Range (M)	Area sq mi	Area within NRA sq mi	% Area within NRA		Apr-May 2013-Q2 A432-33	May-Jun 2014-Q2 T139	May 2015-Q2 A452-53	Apr-May 2016-Q2 A169	May 2017-Q2 A479-80	May-Jun 2018-Q2 A496 T194
				2013-Q2	A432-33						
329	093-183	1721	0	0.00	147.83 (5)		0 (3)	188 (5)	0.8 (5)	70.8 (5)	0 (5)
332	093-183	1047	0	0.00	245 (3)		1969 (2)	243 (3)	2 (3)	503.67 (3)	134.33 (3)
337	093-183	948	0	0.00	3.3333 (3)		202 (2)	716.33 (3)	1730.5 (2)	2161 (2)	170.33 (3)
339	093-183	585	0	0.00	0.5 (2)		0 (2)	0 (2)	0 (2)	0 (2)	0 (2)
354	093-183	474	246	0.52	5.5 (2)		6759.07 (2)	1638 (2)	0 (2)	0 (2)	5.1429 (2)
333	185-274	151(147)	0	0.00	4168.5 (2)		985.5 (2)	2305.5 (2)	1065.5 (2)	662.56 (2)	856 (2)
336	185-274	121	0	0.00	1792.9 (2)		2148.5 (2)	1333.5 (2)	4425.6 (2)	2074.2 (2)	407 (2)
355	185-274	103	74	0.72	4640.3 (2)		1862 (2)	1551.5 (2)	3633 (2)	2345.7 (2)	612 (2)
334	275-366	92(96)	0	0.00	2254.2 (2)		8832.5 (2)	3098.1 (2)	1217.8 (2)	1587 (2)	2729.5 (2)
335	275-366	58	0	0.00	3716 (2)		4680.5 (2)	3701.5 (2)	909 (2)	622 (2)	1073.3 (2)
356	275-366	61	47	0.77	658.17 (2)		1704.5 (2)	810.5 (2)	159 (2)	627 (2)	502.5 (2)
717	367-549	93(166)	0	0.00	687.09 (2)		2742.79 (2)	1320.1 (2)	--	1810.5 (2)	928.5 (2)
719	367-549	76	0	0.00	12735 (2)		1510 (2)	629.44 (2)	538.86 (2)	487.43 (2)	1559.3 (2)
721	367-549	76	58	0.76	1493.7 (2)		987 (2)	1787.5 (2)	473.67 (2)	68.5 (2)	3992.5 (2)
718	550-731	111(134)	0	0.00	2 (2)		5.71429 (2)	30 (2)	--	62.667 (2)	191.17 (2)
720	550-731	105	0	0.00	23.143 (2)		113 (2)	18.667 (2)	298.67 (2)	5.7143 (2)	114.5 (2)
722	550-731	93	71	0.76	3.4286 (2)		5.14286 (2)	12.5 (2)	346.64 (2)	53.5 (2)	72 (2)
Total:		6011	496	8.25							
Upper ( 95% CI )					922.9		4403.8	899.6	3664.9	3915.5	416.0
Weighted mean ( by area )					581.5		1320.1	609.7	530.6	644.0	252.5
Lower ( 95% CI )					240.0		-1763.6	319.9	-2603.7	-2627.4	89.1
<b>SURVEY ABUNDANCE(x10<sup>6</sup>)</b>					481		1092	504	417	533	209
<b>ABUNDANCE within NRA</b>					64		267	91	45	29	42
<b>% within NRA</b>					13		24	18	11	5	20



**Table 6.** Mean weight (kg) per standard tow from Canadian SPRING surveys in Div. 30 covering strata from 93 to 731 m (400ftm.). Dashes (---) represent unsampled strata. Number of successful sets in brackets. Data from 1991-1995 are Campelen trawl equivalent units (see text). Data from 1996 to present are actual Campelen data. G=Gadus Atlantica, W=Wilfred Templeman, A=Alfred Needler.

Stratum	Depth Range (M)	Area sq mi	Area within NRA sq mi	% within NRA		May3-11	May2-13	May5-18	May14-22	May13-27	May22-30	May-Jun 1997-Q2	May-Jun 1998-Q2	May-Jun 1999-Q2	May-Jun 2000-Q2	May-Jun 2001-Q2
				W105	W119-20	W136-7	W153	W168-69	W188	W204	W221-2	W238	W315-16	W365,367		
329	093-183	1721	0	0.00	0.3 (9)	0.0 (8)	0.0 (6)	11.2 (5)	0.5 (5)	0.0 (6)	1.0 (6)	0.0 (7)	0.0 (6)	0.0 (5)	0.0 (5)	
332	093-183	1047	0	0.00	0.7 (6)	0.2 (5)	0.0 (4)	0.0 (4)	148.5 (4)	11.9 (4)	0.3 (3)	49.1 (4)	238.5 (4)	1.7 (4)	2.3 (3)	
337	093-183	948	0	0.00	16.0 (5)	1.5 (4)	0.9 (2)	0.0 (3)	335.0 (4)	0.1 (3)	0.1 (3)	75.9 (4)	29.5 (3)	14.5 (4)	4.7 (3)	
339	093-183	585	0	0.00	0.0 (3)	0.0 (2)	0.0 (2)	0.0 (2)	0.0 (2)	0.0 (2)	0.0 (2)	0.0 (2)	0.0 (2)	0.0 (2)	0.0 (2)	
354	093-183	474	246	0.52	0.0 (3)	0.0 (2)	284.6 (2)	0.0 (2)	0.0 (3)	0.0 (2)	0.0 (2)	109.4 (2)	28.7 (2)	0.1 (2)	8.4 (2)	
333	185-274	151(147)	0	0.00	120.8 (2)	404.0 (2)	1339.7 (2)	5428.5 (2)	113.5 (2)	120.4 (2)	20.2 (2)	696.3 (2)	797.6 (2)	236.2 (2)	225.7 (2)	
336	185-274	121	0	0.00	11.6 (2)	81.2 (2)	630.9 (2)	1032.9 (2)	8543.1 (2)	161.8 (2)	7.7 (2)	5068.7 (2)	198.9 (2)	226.1 (2)	222.9 (2)	
355	185-274	103	74	0.72	2.7 (2)	2.8 (2)	972.9 (2)	608.3 (2)	178.4 (2)	4916.3 (2)	7.5 (2)	741.6 (2)	314.7 (2)	502.8 (2)	44.2 (2)	
334	275-366	92(96)	0	0.00	103.3 (2)	36.5 (2)	202.9 (2)	171.1 (2)	29.4 (2)	220.0 (2)	33.9 (2)	140.3 (2)	478.9 (2)	733.0 (2)	146.4 (2)	
335	275-366	58	0	0.00	4.3 (3)	54.3 (3)	118.3 (2)	1210.4 (2)	263.7 (2)	2445.8 (2)	58.7 (2)	1053.9 (2)	1460.3 (2)	138.7 (2)	741.6 (2)	
356	275-366	61	47	0.77	26.6 (2)	113.0 (2)	462.4 (2)	135.8 (2)	468.0 (2)	515.8 (2)	7.5 (2)	651.6 (2)	1600.5 (2)	4317.8 (2)	73.3 (2)	
717	367-549	93(166)	0	0.00	452.4 (2)	74.3 (2)	83.2 (2)	395.3 (2)	91.4 (2)	191.2 (2)	534.7 (2)	143.1 (2)	670.0 (2)	310.6 (2)	30.2 (2)	
719	367-549	76	0	0.00	33.7 (2)	12.3 (2)	150.0 (2)	669.7 (2)	71.8 (2)	79.5 (2)	59.6 (2)	291.6 (2)	289.0 (2)	326.3 (2)	366.5 (2)	
721	367-549	76	58	0.76	24.7 (2)	183.6 (2)	110.5 (2)	22.0 (2)	1220.5 (2)	68.2 (2)	20.9 (2)	153.0 (2)	651.6 (2)	129.6 (2)	90.7 (2)	
718	550-731	111(134)	0	0.00	42.2 (2)	7.5 (2)	87.7 (2)	156.0 (2)	7.3 (2)	27.2 (2)	15.0 (2)	35.5 (3)	16.7 (3)	174.5 (3)	7.4 (2)	
720	550-731	105	0	0.00	11.7 (2)	57.7 (2)	9.7 (2)	15.9 (2)	14.6 (2)	129.1 (2)	21.0 (2)	14.5 (2)	103.6 (2)	17.7 (2)	18.2 (2)	
722	550-731	93	71	0.76	118.4 (2)	12.6 (2)	33.2 (2)	126.1 (2)	6.3 (2)	25.4 (2)	12.2 (2)	137.0 (2)	19.7 (2)	261.0 (2)	114.2 (2)	
Total:		6011	496	8.25												
Upper (95% CI)					100.7	104.2	277.6	848.6	451.0	1081.0	189.5	1504.1	268.3	145.8	45.7	
Weighted mean (by area)					18.8	19.6	103.1	208.3	283.8	124.2	19.0	192.7	148.2	101.0	31.7	
Lower (95% CI)					-63.2	-65.0	-71.5	-431.9	116.6	-832.6	-151.5	-1118.8	28.1	56.2	17.6	
SURVEY BIOMASS(tons)		15278	15961		83874	172264	234648	102695	15699	159313	122550	83508	26183			
BIOMASS within NRA		1553	2347		23733	8478	14641	54177	410	18024	19914	36624	3048			
%within NRA		10.2	14.7		28.3	4.9	6.2	52.8	2.6	11.3	16.2	43.9	11.6			



**Table 6** continued.

Stratum	Depth Range (M)	Area sq mi	Area within NRA	% within NRA											
				May 2002-Q2	May 2003-Q2	May 2004-Q2	May 2005-Q2	May 2006-Q2	May 2007-Q2	May 2008-Q2	May 2009-Q2	May 2010-Q2	May 2011-Q2	May 2012-Q2	
329	093-183	1721	0	0.00	0.0 (5)	3.0 (5)	0.0 (5)	0.0 (5)	—	0.1 (5)	15.3 (5)	0.0 (5)	0.1 (5)	109.0 (5)	0.1 (5)
332	093-183	1047	0	0.00	3.1 (3)	10.3 (3)	5.5 (3)	3.6 (3)	—	0.7 (3)	0.0 (3)	0.1 (3)	7.9 (3)	193.5 (3)	456.6 (3)
337	093-183	948	0	0.00	0.0 (3)	58.3 (3)	152.1 (3)	2.4 (3)	—	405.0 (3)	0.3 (3)	0.2 (3)	332.3 (3)	9.1 (3)	121.6 (3)
339	093-183	585	0	0.00	0.0 (2)	0.0 (2)	0.0 (2)	0.2 (2)	0.0 (2)	0.0 (2)	0.0 (2)	0.0 (2)	0.0 (2)	0.0 (2)	0.0 (2)
354	093-183	474	246	0.52	0.0 (2)	0.7 (2)	0.2 (2)	43.2 (2)	—	7.5 (2)	0.0 (2)	0.0 (2)	341.6 (2)	446.1 (2)	463.3 (2)
333	185-274	151(147)	0	0.00	154.9 (2)	71.3 (2)	426.0 (2)	705.6 (2)	—	276.7 (2)	301.8 (2)	47.3 (2)	1025.7 (2)	1055.4 (2)	568.3 (2)
336	185-274	121	0	0.00	133.7 (2)	202.3 (2)	2033.3 (2)	698.5 (2)	—	82.1 (2)	739.2 (2)	63.0 (2)	1012.6 (2)	901.0 (2)	2527.7 (2)
355	185-274	103	74	0.72	78.3 (2)	154.9 (2)	232.5 (2)	80.3 (2)	—	41.0 (2)	496.2 (2)	849.2 (2)	217.4 (2)	883.4 (2)	429.3 (2)
334	275-366	92(96)	0	0.00	142.3 (2)	447.8 (2)	284.9 (2)	418.2 (2)	—	102.5 (2)	957.5 (2)	2325.4 (2)	200.6 (2)	659.9 (2)	795.1 (2)
335	275-366	58	0	0.00	740.4 (2)	164.1 (2)	192.7 (2)	496.1 (2)	—	605.2 (2)	381.6 (2)	1351.7 (2)	1144.9 (2)	3452.4 (2)	1014.5 (2)
356	275-366	61	47	0.77	302.7 (2)	66.3 (2)	133.5 (2)	713.4 (2)	—	127.9 (2)	255.6 (2)	360.0 (2)	241.6 (2)	837.3 (2)	1330.5 (2)
717	367-549	93(166)	0	0.00	45.3 (2)	135.8 (2)	452.0 (2)	352.7 (2)	—	658.3 (2)	720.5 (2)	746.4 (2)	235.5 (2)	1007.3 (2)	528.4 (2)
719	367-549	76	0	0.00	52.4 (2)	113.0 (2)	99.1 (2)	312.1 (2)	—	618.7 (2)	396.5 (2)	634.3 (2)	875.5 (2)	115.7 (2)	1342.0 (2)
721	367-549	76	58	0.76	17.2 (2)	43.0 (2)	30.0 (2)	154.0 (2)	—	147.0 (2)	64.0 (2)	289.9 (2)	330.9 (2)	90.4 (2)	298.7 (2)
718	550-731	111(134)	0	0.00	18.1 (2)	0.0 (2)	9.3 (2)	33.3 (2)	—	7.1 (2)	7.9 (2)	34.9 (2)	0.0 (2)	9.3 (2)	18.8 (2)
720	550-731	105	0	0.00	30.9 (2)	5.8 (2)	15.0 (2)	6.9 (2)	—	47.5 (2)	11.0 (2)	84.5 (2)	48.3 (2)	—	64.5 (2)
722	550-731	93	71	0.76	26.6 (2)	16.3 (2)	23.2 (2)	25.4 (2)	—	8.7 (2)	1.2 (2)	15.5 (2)	2.2 (2)	4.1 (2)	43.2 (2)
Total:		6011	496	8.25											
Upper ( 95% CI )					37.4	75.9	298.4	282.2		387.4	126.4	179.9	419.7	418.8	568.7
Weighted mean ( by area )					24.3	35.8	103.0	72.7		111.5	82.9	105.7	169.3	248.0	280.9
Lower ( 95% CI )					11.3	-4.2	-92.4	-136.7		-164.4	39.3	31.4	-81.2	77.1	-6.8
SURVEY BIOMASS(tons)		20126	29642	85170	60138				92202	68519	87362	139960	201458	232298	
BIOMASS within NRA		3151	2529	3702	8369				2754	7226	13437	17997	30263	31456	
% within NRA		15.7	8.5	4.3	13.9				3.0	10.5	15.4	12.9	15.0	13.5	



**Table 6** continued.

Stratum	Depth	Area	%	Area within NRA	Area within NRA	Apr-May 2013-Q2 A432-33	May-Jun 2014-Q2 T139	May 2015-Q2 A452-53	Apr-May 2016-Q2 A169	May 2017-Q2 A479-80	May-Jun 2018-Q2 A496 T194
	Range (M)	Area sq mi	Area sq mi								
329	093-183	1721	0	0.00	19.238 (5)		0 (3)	22.032 (5)	0.008 (5)	10.604 (5)	0 (5)
332	093-183	1047	0	0.00	30.977 (3)		294.06 (2)	30.28 (3)	0.1433 (3)	67.627 (3)	3.72 (3)
337	093-183	948	0	0.00	0.03 (3)		24.035 (2)	84.217 (3)	253 (2)	334.23 (2)	1.8733 (3)
339	093-183	585	0	0.00	0.01 (2)		0 (2)	0 (2)	0 (2)	0 (2)	0 (2)
354	093-183	474	246	0.52	0.975 (2)		1138.66 (2)	301.09 (2)	0 (2)	0 (2)	0.0457 (2)
333	185-274	151(147)	0	0.00	597.97 (2)		140.46 (2)	311.71 (2)	160.43 (2)	114.11 (2)	142.52 (2)
336	185-274	121	0	0.00	240.27 (2)		271.02 (2)	177.81 (2)	657.13 (2)	351.17 (2)	52.79 (2)
355	185-274	103	74	0.72	727.31 (2)		258.82 (2)	221.48 (2)	580.83 (2)	400.48 (2)	80.97 (2)
334	275-366	92(96)	0	0.00	307.54 (2)		1330.16 (2)	492.03 (2)	193.47 (2)	323.88 (2)	486.43 (2)
335	275-366	58	0	0.00	456.01 (2)		616.63 (2)	526.6 (2)	140.94 (2)	112.85 (2)	182.99 (2)
356	275-366	61	47	0.77	88.038 (2)		253.19 (2)	122.27 (2)	25.86 (2)	126.42 (2)	92.035 (2)
717	367-549	93(166)	0	0.00	159.98 (2)		516.188 (2)	256.44 (2)	---	461.1 (2)	222.86 (2)
719	367-549	76	0	0.00	2153.3 (2)		296.97 (2)	138.48 (2)	104 (2)	134.71 (2)	335.78 (2)
721	367-549	76	58	0.76	231.57 (2)		178.14 (2)	454.18 (2)	116.66 (2)	15.989 (2)	878.26 (2)
718	550-731	111(134)	0	0.00	0.4533 (2)		1.72 (2)	8.74 (2)	---	15.9 (2)	54.653 (2)
720	550-731	105	0	0.00	7.1676 (2)		28.76 (2)	3.9778 (2)	72.96 (2)	1.5429 (2)	35.83 (2)
722	550-731	93	71	0.76	1.2914 (2)		1.64 (2)	2.11 (2)	107.2 (2)	17.075 (2)	28 (2)
Total:		6011	496	8.25							
Upper ( 95% CI )					146.2		762.1	134.6	530.7	597.9	76.4
Weighted mean ( by area )					87.8		208.7	92.7	81.5	107.1	41.1
Lower ( 95% CI )					29.4		-344.7	50.7	-367.6	-383.7	5.8
<b>SURVEY BIOMASS(tons)</b>					72634		172583	76615	64056	88540	34011
<b>BIOMASS within NRA</b>					9867		44244	16879	8058	5189	8702
<b>% within NRA</b>					14		26	22	13	6	26

**Table 7.** Mean number per standard tow from Canadian AUTUMN surveys in Div. 30 covering strata from 93 to 731 m (400ftm.). Dashes (---) represent unsampled strata. Number of successful sets in brackets. Data from 1991-1995 are Campelen trawl equivalent units (see text). Data from 1996 to present are actual Campelen data. G=Gadus Atlantica, W=Wilfred Templeman, A=Alfred Needler.

Stratum	Depth Range (M)	Area sq mi	% within NRA	Area within NRA											
				Oct27-Nov10 1991-Q4 W113-4	Oct26-Nov5 1992-Q4 W128-9	Nov1-12 1993-Q4 W144-5	Oct29-Dec13 1994-Q4 W160-61	Sep28-Oct26 1995-Q4 W176-77	Nov25-Dec13 1996-Q4 W200	Oct-Dec 1997-Q4 W212-13	Sep-Oct 1998-Q4 W229-230	Sep-Oct 1999-Q4 W246-247	Sep-Oct 2000-Q4 W319-320	Sep-Oct 2001-Q4 W372	Sep-Oct T338
329	093-183	1721	0	0.00	1.1 (7)	0.0 (3)	0.0 (5)	0.0 (6)	47.8 (5)	0.2 (5)	421.4 (5)	0.8 (5)	0.0 (5)	0.0 (5)	746.8 (5)
332	093-183	1047	0	0.00	0.0 (4)	88.3 (3)	49.7 (3)	118.0 (3)	403.0 (3)	11.5 (2)	89.0 (3)	45.3 (3)	32.0 (3)	65.5 (3)	8.7 (3)
337	093-183	948	0	0.00	175.5 (4)	667.5 (2)	35.3 (3)	41.5 (2)	515.0 (2)	0.0 (2)	149.3 (3)	273.8 (3)	28.7 (3)	50.6 (3)	37.3 (3)
339	093-183	585	0	0.00	0.0 (2)	0.0 (2)	0.0 (2)	0.0 (2)	0.0 (2)	0.0 (3)	0.0 (2)	0.0 (2)	1.0 (2)	1.0 (2)	
354	093-183	474	246	0.52	0.0 (2)	628.0 (2)	0.0 (2)	0.0 (2)	8100.0 (3)	427.3 (2)	6357.5 (2)	226.5 (2)	695.5 (2)	0.0 (2)	272.5 (2)
333	185-274	151(147)	0	0.00	314.5 (2)	1365.0 (2)	479.0 (2)	2073.0 (2)	923.5 (2)	—	217.0 (2)	155.2 (2)	230.5 (2)	488.8 (2)	320.7 (2)
336	185-274	121	0	0.00	364.5 (2)	2760.0 (2)	3298.5 (2)	3807.0 (2)	450.0 (2)	161.5 (2)	918.0 (2)	691.7 (2)	3481.0 (2)	802.0 (2)	131.0 (2)
355	185-274	103	74	0.72	9957.0 (2)	6381.0 (2)	1317.5 (2)	2310.5 (2)	2317.3 (2)	391.4 (2)	215.0 (2)	124.5 (2)	2333.5 (2)	1020.5 (2)	879.1 (2)
334	275-366	92(96)	0	0.00	8774.0 (2)	3290.0 (2)	2603.7 (3)	975.0 (2)	3474.0 (2)	—	1670.0 (2)	1110.5 (2)	178.1 (2)	378.7 (2)	1441.2 (2)
335	275-366	58	0	0.00	3853.0 (2)	5346.0 (2)	2541.5 (2)	5648.0 (2)	1667.0 (2)	2895.5 (2)	8352.5 (2)	2459.5 (2)	2748.0 (2)	2403.4 (2)	740.5 (2)
356	275-366	61	47	0.77	678.5 (2)	3828.0 (2)	568.5 (2)	2671.0 (2)	3637.1 (2)	868.4 (2)	735.5 (2)	5602.0 (2)	3452.9 (2)	5888.0 (2)	2481.2 (2)
717	367-549	93(166)	0	0.00	—	—	6079.5 (2)	1172.5 (2)	2247.5 (2)	—	13031.5 (2)	8428.5 (2)	603.2 (2)	5420.1 (2)	1401.5 (2)
719	367-549	76	0	0.00	813.5 (2)	—	4854.0 (2)	2715.5 (2)	2892.6 (2)	5015.5 (2)	5311.5 (2)	1953.0 (2)	3604.0 (2)	8204.0 (3)	2407.5 (2)
721	367-549	76	58	0.76	315.5 (2)	—	543.5 (2)	82.5 (2)	9946.8 (2)	575.5 (2)	3882.0 (2)	1872.5 (2)	905.6 (2)	1502.7 (2)	1970.5 (2)
718	550-731	111(134)	0	0.00	—	—	520.0 (2)	1051.5 (2)	863.8 (2)	—	95.0 (2)	12.5 (2)	169.5 (2)	102.0 (2)	289.5 (2)
720	550-731	105	0	0.00	—	—	147.0 (2)	306.0 (2)	43.2 (2)	1560.6 (2)	—	471.0 (2)	103.5 (2)	160.0 (2)	88.4 (2)
722	550-731	93	71	0.76	11.5 (2)	—	371.5 (2)	56.5 (2)	365.5 (2)	324.0 (2)	13.8 (2)	278.0 (2)	15.0 (2)	156.4 (2)	282.3 (2)
764	732-914	105	105	1.00	—	—	—	—	—	—	—	5.0 (2)	—	4.5 (2)	0.0 (2)
768	732-914	99	0	0.00	—	—	—	—	—	—	—	0.5 (2)	—	0.0 (2)	0.0 (2)
772	732-914	135	0	0.00	—	—	—	—	—	—	—	0.0 (2)	—	6.3 (2)	—
Total:		6350	601	9.46											
Upper ( 95% CI )					3059.2	1217.7	587.0	672.0	9437.2	445.6	7592.4	3138.3	686.7	515.7	1000.7
Weighted mean ( by area )					436.0	572.0	371.5	388.6	1233.7	203.8	1304.5	455.7	359.5	411.0	416.0
Lower ( 95% CI )					-2187.1	-73.7	156.0	105.2	-6969.8	-25.1	-4983.5	-2226.9	32.3	306.3	-168.7
<b>SURVEY ABUNDANCE(x10<sup>6</sup>)</b>					336.3	421.8	302.3	321.3	1020.1	153.3	1059.8	398.0	268.3	359.0	355.6
<b>ABUNDANCE within NRA</b>					108.4	111.0	25.1	42.0	404.2	31.8	253.2	62.9	77.0	62.0	52.7
<b>% within NRA</b>					32.2	26.3	8.3	13.1	39.6	20.8	23.9	15.8	28.7	17.3	14.8



**Table 7** continued.

Stratum	Depth (M)	Range	Area sq mi	Area within NRA	% within NRA	Sep-Oct	Sep-Oct	Nov	Oct	Oct	Oct	Oct	Sep-Oct	Sep-Oct	Sep-Oct
						2002-Q4 W427	2003-Q4 W485-6	2004-Q4 W557	2005-Q4 W627-628	2006-Q4 W704	2007-Q4 W770-771	2008-Q4 W835-836	2009-Q4 A913-15	2010-Q4 A942-43	2011-Q4 A409-10
329	093-183	1721	0	0.00	405.8 (5)	0.4 (5)	0.0 (5)	14.2 (5)	74.2 (5)	0.0 (5)	3.0 (3)	91.8 (5)	0.0 (5)	86.2 (5)	634.8 (5)
332	093-183	1047	0	0.00	12.8 (3)	37.4 (3)	29.7 (3)	41.2 (3)	0.3 (3)	27.3 (3)	0.7 (3)	261.9 (3)	107.9 (3)	84.9 (3)	36.7 (3)
337	093-183	948	0	0.00	61.9 (3)	55.3 (3)	54.9 (3)	90.3 (3)	38.3 (3)	402.0 (3)	0.0 (2)	383.5 (3)	1564.2 (3)	56.0 (3)	125.6 (3)
339	093-183	585	0	0.00	0.5 (2)	0.0 (2)	0.6 (2)	3.0 (2)	0.0 (2)	1.5 (2)	0.0 (2)	0.0 (2)	0.5 (2)	0.0 (2)	0.0 (2)
354	093-183	474	246	0.52	150.9 (2)	0.0 (2)	171.6 (2)	69.5 (2)	6.0 (2)	1124.9 (2)	363.6 (2)	1172.9 (2)	263.6 (2)	776.0 (2)	42.5 (2)
333	185-274	151(147)	0	0.00	31.6 (2)	96.5 (2)	77.5 (2)	674.0 (2)	103.8 (2)	159.6 (2)	963.1 (2)	389.3 (2)	1362.7 (2)	148.9 (2)	796.2 (2)
336	185-274	121	0	0.00	87.5 (2)	85.5 (2)	273.5 (2)	255.0 (2)	744.0 (2)	722.2 (2)	7145.3 (2)	—	149.3 (2)	342.5 (2)	630.9 (2)
355	185-274	103	74	0.72	614.5 (2)	61.5 (2)	527.0 (2)	643.4 (2)	963.8 (2)	2225.3 (2)	11598.4 (2)	1738.7 (2)	1148.0 (2)	970.0 (2)	2039.1 (2)
334	275-366	92(96)	0	0.00	106.2 (2)	872.5 (2)	256.3 (2)	816.5 (2)	569.8 (2)	1474.5 (2)	754.2 (2)	9905.8 (2)	1576.9 (2)	1971.6 (2)	3259.8 (2)
335	275-366	58	0	0.00	781.7 (2)	1051.0 (2)	2291.6 (2)	626.7 (2)	898.0 (2)	7626.0 (2)	8416.0 (2)	4935.4 (2)	2917.0 (2)	6197.7 (2)	6804.8 (2)
356	275-366	61	47	0.77	692.0 (2)	828.0 (2)	603.4 (2)	2484.4 (2)	5727.5 (2)	1123.1 (2)	8649.7 (2)	3298.4 (2)	65720.0 (2)	497.4 (2)	732.0 (2)
717	367-549	93(166)	0	0.00	488.9 (2)	675.7 (2)	2530.3 (2)	1382.2 (2)	1756.3 (2)	2576.0 (2)	2205.3 (2)	8353.8 (2)	8394.6 (2)	20038.4 (2)	8641.1 (2)
719	367-549	76	0	0.00	6420.9 (2)	1265.0 (2)	1844.2 (2)	3854.5 (2)	14161.4 (2)	7449.1 (2)	5984.2 (2)	15486.3 (2)	11698.5 (2)	12566.3 (2)	55983.3 (2)
721	367-549	76	58	0.76	4210.8 (2)	3567.9 (2)	927.6 (2)	648.0 (2)	410.2 (2)	2007.9 (2)	1472.6 (2)	8545.8 (2)	4935.2 (2)	4239.3 (2)	2881.4 (2)
718	550-731	111(134)	0	0.00	545.3 (2)	16.0 (2)	120.5 (2)	45.2 (2)	—	928.7 (2)	64.7 (2)	725.3 (2)	207.1 (2)	25.0 (2)	751.9 (2)
720	550-731	105	0	0.00	12.7 (2)	236.4 (2)	478.9 (2)	4489.7 (2)	1761.3 (2)	786.6 (2)	1005.3 (2)	424.5 (2)	669.2 (2)	2606.9 (2)	1048.0 (2)
722	550-731	93	71	0.76	336.4 (2)	112.4 (2)	106.9 (2)	35.1 (2)	9.0 (2)	35.6 (2)	7.5 (2)	9.6 (2)	9.5 (2)	391.5 (2)	443.5 (2)
764	732-914	105	105	1.00	0.5 (2)	---	0.0 (2)	---	0.0 (2)	---	73.7 (2)	---	---	---	---
768	732-914	99	0	0.00	0.0 (2)	---	0.0 (2)	---	0.0 (2)	---	2.5 (2)	---	---	---	---
772	732-914	135	0	0.00	0.6 (2)	1.3 (2)	---	0.0 (2)	---	0.0 (2)	---	0.0 (2)	---	---	---
Total:		6350	601	9.46											
Upper ( 95% CI )					618.7	457.2	915.3	394.0	2367.8	894.6	2855.5	1524.0	9812.3	7379.0	9783.7
Weighted mean ( by area )					317.1	135.9	193.8	268.1	410.5	518.6	751.1	1083.6	1521.2	1051.0	1428.4
Lower ( 95% CI )					15.5	-185.4	-527.7	142.3	-1546.7	142.6	-1353.3	643.2	-6770.0	-5277.0	-6926.9
<b>SURVEY ABUNDANCE(x10<sup>6</sup>)</b>					277.0	114.9	160.2	234.2	331.9	453.0	621.1	878.0	1257.0	869.1	1181.1
<b>ABUNDANCE within NRA</b>					52.7	35.5	23.5	30.5	50.4	84.4	198.1	147.0	485.0	77.0	54.3
% within NRA					19.0	30.9	14.7	13.0	15.2	18.6	31.9	16.7	38.6	8.9	4.6

**Table 7** continued.

Stratum	Depth Range (M)	Area sq mi	Area	%	Sep	Sep-Oct	Sep	Sep	Sep	
			within NRA sq mi	Area within NRA	2013-Q4 A438	2014-Q4	2015-Q4 A458-60	2016-Q3 A466-67	2017-Q3 A484	2018-Q3 A500
329	093-183	1721	0	0.00	0.4 (5)		0.0 (5)	1.2 (5)	38.2 (5)	0.2 (5)
332	093-183	1047	0	0.00	102.0 (3) no		121.0 (3)	0.0 (3)	21.7 (3)	26.7 (3)
337	093-183	948	0	0.00	314.9 (3) survey		208.0 (3)	2.0 (3)	101.7 (3)	73.0 (3)
339	093-183	585	0	0.00	0.0 (2) in		0.0 (2)	0.5 (2)	0.0 (2)	0.0 (2)
354	093-183	474	246	0.52	4.0 (2) redfish		513.0 (2)	4.0 (2)	42.9 (2)	1.5 (2)
333	185-274	151(147)	0	0.00	562.2 (2) strata		296.5 (2)	145.5 (2)	39.5 (2)	5.5 (2)
336	185-274	121	0	0.00	394.7 (2)		456.0 (2)	—	180.0 (2)	89.0 (2)
355	185-274	103	74	0.72	364.0 (2)		531.7 (2)	278.3 (2)	352.0 (2)	62.0 (2)
334	275-366	92(96)	0	0.00	6271.1 (2)		1000.5 (2)	638.3 (2)	243.0 (2)	253.1 (2)
335	275-366	58	0	0.00	7026.2 (2)		1728.9 (2)	443.0 (2)	3957.8 (2)	1920.4 (2)
356	275-366	61	47	0.77	3405.5 (2)		179.9 (2)	139.0 (2)	257.5 (2)	101.0 (2)
717	367-549	93(166)	0	0.00	4169.0 (2)		1751.8 (2)	1650.3 (2)	2044.0 (2)	5412.6 (2)
719	367-549	76	0	0.00	10118.3 (2)		1616.5 (2)	6113.1 (2)	1319.3 (2)	4504.9 (2)
721	367-549	76	58	0.76	4973.4 (2)		4672.6 (2)	197.5 (2)	1399.1 (2)	1922.2 (2)
718	550-731	111(134)	0	0.00	961.7 (2)		146.0 (2)	73.5 (2)	2804.0 (2)	574.8 (2)
720	550-731	105	0	0.00	1336.6 (2)		588.9 (2)	1815.0 (2)	343.4 (2)	749.0 (2)
722	550-731	93	71	0.76	1021.0 (2)		1089.7 (2)	504.5 (2)	(2)	6.7 (2)
764	732-914	105	105	1.00	—		—	—	—	—
768	732-914	99	0	0.00	—		—	—	—	—
772	732-914	135	0	0.00	—		—	—	—	—
Total:		6350	601	9.46						
Upper ( 95% CI )					926.9		525.2	309.7	602.4	429.8
Weighted mean ( by area )					664.8		312.7	195.7	252.7	299.7
Lower ( 95% CI )					402.8		100.1	81.7	-97.0	169.5
<b>SURVEY ABUNDANCE(x10<sup>6</sup>)</b>					549.7		258.5	158.5	205.7	247.8
<b>ABUNDANCE within NRA</b>					75.5		71.9	10.4	17.9	16.7
<b>%within NRA</b>					13.7		27.8	6.5	8.7	6.8



**Table 8.** Mean weight (kg) per standard tow from Canadian AUTUMN surveys in Div. 30 covering strata from 93 to 731 m (400ftm.). Dashes (---) represent unsampled strata. Number of successful sets in brackets. Data from 1991-1995 are Campelen trawl equivalent units (see text). Data from 1996 to present are actual Campelen data. G=Gadus Atlantica, W=Wilfred Templeman, A=Alfred Needler.

Stratum	Depth Range (M)	Area sq mi	% within NRA	Area within NRA											
				Oct27-Nov10 1991-Q4 W113-4	Oct26-Nov5 1992-Q4 W128-9	Nov1-12 1993-Q4 W144-5	Oct29-Dec13 1994-Q4 W160-61	Sep28-Oct26 1995-Q4 W176-77	Nov25-Dec13 1996-Q4 W200	Oct-Dec 1997-Q4 W212-13	Sep-Oct 1998-Q4 W229-230	Sep-Oct 1999-Q4 W246-247	Sep-Oct 2000-Q4 W319-320	Sep-Oct 2001-Q4 W372	Sep-Oct T338 T357
329	093-183	1721	0	0.00	0.0 (7)	0.0 (3)	0.0 (5)	0.00 (6)	1.0 (5)	0.0 (5)	22.6 (5)	0.0 (5)	0.0 (5)	0.0 (5)	42.1 (5)
332	093-183	1047	0	0.00	0.0 (4)	13.3 (3)	2.7 (3)	15.59 (3)	31.5 (3)	0.2 (2)	7.7 (3)	2.7 (3)	0.8 (3)	0.1 (3)	
337	093-183	948	0	0.00	30.8 (4)	64.7 (2)	7.0 (3)	5.04 (2)	55.5 (2)	0.0 (2)	17.9 (3)	34.6 (3)	1.9 (3)	12.7 (3)	2.9 (3)
339	093-183	585	0	0.00	0.0 (2)	0.0 (2)	0.0 (2)	0.00 (2)	0.0 (2)	0.0 (3)	0.0 (2)	0.0 (2)	0.2 (2)	0.2 (2)	
354	093-183	474	246	0.52	0.0 (2)	171.5 (2)	0.0 (2)	0.00 (2)	785.3 (3)	15.6 (2)	915.0 (2)	31.5 (2)	69.0 (2)	0.0 (2)	35.2 (2)
333	185-274	151(147)	0	0.00	27.1 (2)	168.0 (2)	46.5 (2)	257.7 (2)	107.0 (2)	—	26.5 (2)	20.0 (2)	18.0 (2)	24.4 (2)	31.0 (2)
336	185-274	121	0	0.00	18.5 (2)	374.3 (2)	378.8 (2)	357.8 (2)	49.7 (2)	9.1 (2)	117.4 (2)	103.8 (2)	548.7 (2)	98.9 (2)	13.5 (2)
355	185-274	103	74	0.72	352.2 (2)	450.7 (2)	77.9 (2)	264.2 (2)	237.0 (2)	37.9 (2)	25.9 (2)	11.9 (2)	387.8 (2)	127.8 (2)	119.0 (2)
334	275-366	92(96)	0	0.00	1317.9 (2)	480.7 (2)	380.5 (3)	171.1 (2)	506.8 (2)	—	289.5 (2)	188.3 (2)	22.6 (2)	54.6 (2)	188.8 (2)
335	275-366	58	0	0.00	512.6 (2)	850.9 (2)	351.8 (2)	877.1 (2)	187.7 (2)	332.2 (2)	1114.4 (2)	362.1 (2)	443.2 (2)	355.4 (2)	89.0 (2)
356	275-366	61	47	0.77	59.4 (2)	684.6 (2)	60.1 (2)	303.8 (2)	387.6 (2)	145.5 (2)	106.1 (2)	914.5 (2)	592.9 (2)	801.6 (2)	370.6 (2)
717	367-549	93(166)	0	0.00	—	—	1391.3 (2)	340.4 (2)	588.8 (2)	—	2281.8 (2)	1834.0 (2)	135.7 (2)	1143.7 (2)	229.2 (2)
719	367-549	76	0	0.00	268.9 (2)	—	930.5 (2)	536.2 (2)	414.0 (2)	656.4 (2)	880.2 (2)	321.3 (2)	691.0 (2)	1313.7 (3)	373.6 (2)
721	367-549	76	58	0.76	53.7 (2)	—	100.4 (2)	16.57 (2)	1666.7 (2)	87.3 (2)	732.5 (2)	410.5 (2)	177.5 (2)	230.2 (2)	319.2 (2)
718	550-731	111(134)	0	0.00	—	—	169.3 (2)	442.1 (2)	409.4 (2)	—	37.1 (2)	4.4 (2)	48.0 (2)	24.8 (2)	79.5 (2)
720	550-731	105	0	0.00	—	—	50.0 (2)	118.7 (2)	16.5 (2)	572.6 (2)	—	162.6 (2)	21.3 (2)	52.3 (2)	16.1 (2)
722	550-731	93	71	0.76	7.7 (2)	—	164.0 (2)	22.71 (2)	125.8 (2)	103.9 (2)	4.0 (2)	108.6 (2)	5.3 (2)	34.9 (2)	125.2 (2)
764	732-914	105	105	1.00	—	—	—	—	—	—	—	1.6	—	2.6 (2)	0.0 (2)
768	732-914	99	0	0.00	—	—	—	—	—	—	—	0.3	—	0.0 (2)	0.0 (2)
772	732-914	135	0	0.00	—	—	—	—	—	—	—	0.0	—	2.2 (2)	—
Total:		6350	601	9.46											
Upper ( 95% CI )					306.5	147.4	105.2	109.0	972.0	86.2	1182.1	664.3	106.8	83.3	75.6
Weighted mean ( by area )					44.9	76.3	63.6	64.5	151.9	30.5	190.3	86.6	56.4	68.7	43.6
Lower ( 95% CI )					-216.7	5.2	22.1	20.0	-668.2	-25.1	-801.5	-491.0	6.0	54.0	11.6
<b>SURVEY BIOMASS(tons)</b>					34618	56247	51782	53324	125578	22974	154622	75676	42100	60004	37286
<b>BIOMASS within NRA</b>					4473	14818	3584	5008	46022	3565	37798	11459	11585	8700	8567
% within NRA					12.9	26.3	6.9	9.4	36.6	15.5	24.4	15.1	27.5	14.5	23.0



**Table 8** continued.


Stratum	Depth Range (M)	Area sq mi	% within NRA	Area	%	Sep-Oct	Sep-Oct	Nov	Oct	Oct	Oct	Oct	Sep-Oct	Sep-Oct	Sep-Oct
				within NRA	Area within NRA	2002-Q4 W427	2003-Q4 W485-6	2004-Q4 W557	2005-Q4 W627-628	2006-Q4 W704	2007-Q4 W770-771	2008-Q4 W835-836	2009-Q4 A913-15	2010-Q4 A942-43	2011-Q4 A409-10
329	093-183	1721	0	0.00	32.2 (5)	0.0 (5)	0.0 (5)	0.2 (5)	4.5 (5)	0.0 (5)	0.0 (3)	3.3 (5)	0.0 (5)	4.0 (5)	61.1 (5)
332	093-183	1047	0	0.00	1.7 (3)	2.9 (3)	0.1 (3)	0.7 (3)	0.0 (3)	0.8 (3)	0.0 (3)	32.2 (3)	7.1 (3)	3.7 (3)	1.5 (3)
337	093-183	948	0	0.00	3.9 (3)	3.8 (3)	0.3 (3)	1.7 (3)	1.1 (3)	49.9 (3)	0.0 (2)	16.2 (3)	140.7 (3)	6.3 (3)	10.1 (3)
339	093-183	585	0	0.00	0.1 (2)	0.0 (2)	0.0 (2)	0.1 (2)	0.0 (2)	0.1 (2)	0.0 (2)	0.0 (2)	0.1 (2)	0.0 (2)	0.0 (2)
354	093-183	474	246	0.52	10.9 (2)	0.0 (2)	7.2 (2)	3.6 (2)	0.6 (2)	131.2 (2)	26.9 (2)	102.3 (2)	18.2 (2)	108.2 (2)	2.8 (2)
333	185-274	151(147)	0	0.00	3.9 (2)	11.3 (2)	3.0 (2)	53.4 (2)	8.0 (2)	12.5 (2)	81.7 (2)	37.0 (2)	150.2 (2)	15.9 (2)	128.2 (2)
336	185-274	121	0	0.00	9.0 (2)	10.0 (2)	31.9 (2)	25.0 (2)	51.0 (2)	47.3 (2)	613.8 (2)	—	15.7 (2)	37.9 (2)	76.9 (2)
355	185-274	103	74	0.72	64.2 (2)	6.3 (2)	67.2 (2)	59.3 (2)	117.1 (2)	161.8 (2)	853.7 (2)	146.1 (2)	119.5 (2)	116.0 (2)	231.6 (2)
334	275-366	92(96)	0	0.00	13.7 (2)	146.6 (2)	54.9 (2)	162.7 (2)	105.4 (2)	256.1 (2)	122.2 (2)	1044.9 (2)	156.2 (2)	244.6 (2)	424.3 (2)
335	275-366	58	0	0.00	82.5 (2)	136.1 (2)	334.0 (2)	92.2 (2)	126.4 (2)	669.5 (2)	1250.5 (2)	549.1 (2)	311.6 (2)	700.7 (2)	816.7 (2)
356	275-366	61	47	0.77	96.4 (2)	91.3 (2)	82.3 (2)	437.8 (2)	1176.1 (2)	157.0 (2)	1903.0 (2)	328.0 (2)	7178.6 (2)	58.6 (2)	98.3 (2)
717	367-549	93(166)	0	0.00	75.9 (2)	115.4 (2)	540.9 (2)	253.7 (2)	355.2 (2)	497.1 (2)	235.5 (2)	1424.6 (2)	1230.5 (2)	3007.8 (2)	1205.6 (2)
719	367-549	76	0	0.00	889.9 (2)	194.9 (2)	385.7 (2)	627.9 (2)	2137.5 (2)	1297.0 (2)	1034.3 (2)	1679.4 (2)	1689.1 (2)	1592.6 (2)	7348.1 (2)
721	367-549	76	58	0.76	762.0 (2)	718.8 (2)	184.1 (2)	119.8 (2)	73.8 (2)	381.7 (2)	339.8 (2)	892.8 (2)	924.5 (2)	593.1 (2)	384.3 (2)
718	550-731	111(134)	0	0.00	118.0 (2)	3.9 (2)	34.8 (2)	12.0 (2)	—	320.0 (2)	24.3 (2)	277.0 (2)	75.8 (2)	9.2 (2)	212.3 (2)
720	550-731	105	0	0.00	2.9 (2)	49.3 (2)	134.1 (2)	1013.5 (2)	403.9 (2)	217.1 (2)	279.2 (2)	103.4 (2)	180.5 (2)	710.2 (2)	290.3 (2)
722	550-731	93	71	0.76	68.1 (2)	33.8 (2)	46.7 (2)	11.2 (2)	3.5 (2)	13.8 (2)	3.5 (2)	2.9 (2)	3.4 (2)	103.3 (2)	116.7 (2)
764	732-914	105	105	1.00	0.4 (2)	—	—	0.0 (2)	(2)	0.0 (2)	—	—	—	—	—
768	732-914	99	0	0.00	0.0 (2)	—	—	0.0 (2)	(2)	0.0 (2)	—	—	—	—	—
772	732-914	135	0	0.00	0.1 (2)	0.2 (2)	—	0.0 (2)	(2)	0.0 (2)	—	—	—	—	—
Total:		6350	601	9.46											
Upper ( 95% CI )					68.1	86.6	179.4	65.5	356.1	133.9	144.0	220.5	1069.1	1093.0	1275.9
Weighted mean ( by area )					38.9	22.0	33.4	43.6	65.9	76.3	93.8	128.4	181.2	150.6	186.8
Lower ( 95% CI )					9.7	-42.6	-112.6	21.8	-224.3	18.8	43.6	36.3	-706.7	-791.8	-902.9
<b>SURVEY BIOMASS(tons)</b>		33976	18604	27631	38125	53291	66682	77562	104013	149819	124539	154234			
<b>BIOMASS within NRA</b>		8396	6720	3385	4619	9439	10285	24651	14222	55658	10964	7294			
<b>%within NRA</b>		24.7	36.1	12.3	12.1	17.7	15.4	31.8	13.7	37.2	8.8	4.7			




**Table 8** continued.

Stratum	Depth Range (M)	Area sq mi	Area within NRA sq mi	% Area within NRA	Sep 2013-Q4 A438	2014-Q4	Sep-Oct 2015-Q4 A458-60	Sep 2016-Q3 A466-67	Sep 2017-Q3 A484	Sep 2018-Q3 A500
329	093-183	1721	0	0.00	0.0 (5)		0.0 (5)	0.0 (5)	1.1 (5)	0.0 (5)
332	093-183	1047	0	0.00	7.9 (3) no		17.2 (3)	0.0 (3)	0.2 (3)	0.1 (3)
337	093-183	948	0	0.00	56.9 (3) survey		31.0 (3)	0.2 (3)	17.2 (3)	4.1 (3)
339	093-183	585	0	0.00	0.0 (2) in		0.0 (2)	0.0 (2)	0.0 (2)	0.0 (2)
354	093-183	474	246	0.52	0.6 (2) redfish		85.1 (2)	0.0 (2)	3.7 (2)	0.0 (2)
333	185-274	151(147)	0	0.00	74.2 (2) strata		13.6 (2)	17.6 (2)	5.1 (2)	0.5 (2)
336	185-274	121	0	0.00	44.3 (2)		29.7 (2)	--	13.3 (2)	7.5 (2)
355	185-274	103	74	0.72	48.1 (2)		65.5 (2)	34.8 (2)	44.9 (2)	10.0 (2)
334	275-366	92(96)	0	0.00	785.7 (2)		172.0 (2)	123.9 (2)	40.8 (2)	55.3 (2)
335	275-366	58	0	0.00	858.8 (2)		243.6 (2)	64.8 (2)	685.0 (2)	357.3 (2)
356	275-366	61	47	0.77	450.7 (2)		26.8 (2)	26.1 (2)	40.8 (2)	16.0 (2)
717	367-549	93(166)	0	0.00	625.2 (2)		321.0 (2)	301.4 (2)	411.7 (2)	1225.9 (2)
719	367-549	76	0	0.00	1442.2 (2)		249.0 (2)	1061.1 (2)	237.3 (2)	993.0 (2)
721	367-549	76	58	0.76	666.5 (2)		838.5 (2)	33.7 (2)	253.6 (2)	352.4 (2)
718	550-731	111(134)	0	0.00	284.0 (2)		38.5 (2)	23.7 (2)	741.7 (2)	207.9 (2)
720	550-731	105	0	0.00	287.2 (2)		153.7 (2)	470.4 (2)	103.0 (2)	192.9 (2)
722	550-731	93	71	0.76	225.9 (2)		257.1 (2)	123.6 (2)	--	1.6 (2)
764	732-914	105	105	1.00	--		--	--	--	--
768	732-914	99	0	0.00	--		--	--	--	--
772	732-914	135	0	0.00	--		--	--	--	--
Total:		6350	601	9.46						
Upper ( 95% CI )					130.4		91.2	62.6	174.9	94.6
Weighted mean ( by area )					98.1		52.1	37.5	48.9	64.4
Lower ( 95% CI )					65.8		13.0	12.4	-77.2	34.2
<b>SURVEY BIOMASS(tons)</b>					81122.8		43117.6	30394.2	39768.1	53238.8
<b>BIOMASS within NRA</b>					10948.7		12921.2	2000.0	2868.9	3033.4
<b>%within NRA</b>					13.5		30.0	6.6	7.2	5.7







**Figure 1.** Nominal catches and TACs of Redfish in Div. 3O. TAC to 2004 was only for Canadian fishery zone.




**Figure 2.** Catch numbers at length ('000s) and Canadian RV survey numbers at length.




**Figure 3.** Survey biomass index for redfish in Div. 30 for spring and autumn surveys from 1991 - 2018 (upper panels) with 95% CI (lower panels). Surveys prior to 1995 utilized an Engel trawl. Estimates were converted to Campelen equivalents based on comparative fishing trials.




**Figure 4.** EU-Spain survey biomass index for redfish in Div. 30 for spring 1997 to 2018. Surveys prior to 2001 used a Pedreira trawl. The data prior to 2001 were converted to Campelen equivalent units. Dashed line is the time-series mean.



**Figure 5.** Biomass indices from the Canadian spring and autumn plus the EU-Spain research vessel surveys for redfish in Div. 30. Survey indices were normalized based on average values during the period 1997 to 2018.



**Figure 6.** Length distributions from Canadian RV surveys in Div. 30 during spring and autumn plus the EU-Spain survey from 2004-2018. Plotted are mean per standard tow.



**Figure 7.** Catch/Biomass ratios for Div. 30 based on Canadian RV surveys. Plotted are average survey biomass between spring (n) and autumn (n-1) for year (n) in which catch was taken. The 2006 value of biomass is only from the autumn survey and the 2014 value is from the spring survey.