



SCIENTIFIC COUNCIL MEETING -2019

A Compilation of Research Vessel Surveys on a Stock-by-stock Basis

by

NAFO Secretariat

Discussions at the Scientific Council Meeting in June 1990 resulted in the compilation of information on research vessel surveys on a stock-by-stock basis (SCS Doc. 90/22, Serial No. N1820). This inventory was judged to be very helpful for the establishment of future inventories. It was agreed that the Designated Experts for the stocks be responsible for future compilations and that Designated Experts should update the information annually for review by STACFIS (*NAFO Sci. Coun. Rep.*, 1991, p. 114). The following tabulations were received at the Secretariat from the Designated Experts during the year 2019.



| Stock name: Greenland halibut in Subarea 0 +Division 1A (offshore) and Divisions 1B-F | | | | | | | | | | | | | | | | |
|---|-----------------|------------------|---|---|-------------------|-------------|---------------|--|-------------|---|---|---|---|--------------------|-----------|------------------|
| Name of survey | Location (Div.) | Season months(s) | Years of survey from, to or since | Type of survey and gears | Target species | No. of sets | 24 hr fishing | Area of coverage in relation to stock distribution (%) | Depth range | Research data related to stock size estimates | | | | Individual weights | Country | SCS/ SCR Doc. No |
| Greenland Deep Sea | 1CD | Sep-Nov | 1997 - 2017 | Stratified Random bottom Trawl, Alfredo trawl | Greenland halibut | 53 | Y | ? | 400-1500 m | Y | Y | N | Y | Y | Greenland | SCR 18/21 |
| Greenland Shrimp Trawl | 1A-F | Jul-Sep | Since 1988 | Stratified Random bottom Trawl, Cosmos trawl | Shrimp Groundfish | 257 | N | ? | 50-600m | Y | Y | N | Y | Y | Greenland | SCR 19/08 |
| Canada Deep Sea | 0A | Oct-Nov | 1999 2001 2004 2006 2008 2010 2012 2014-2017 | Stratified Random bottom Trawl, Alfredo trawl | Greenland halibut | 74 | Y | ? | 400-1500 m | Y | Y | N | Y | Y | Canada | SCR 18/15 |
| Canada Deep Sea | 0B | Oct-Nov | 2000 2001 2011 2013 2014-2016 | Stratified Random bottom Trawl, Alfredo trawl | Greenland halibut | 81 | Y | ? | 400-1500 m | Y | Y | N | Y | Y | Canada | SCR 17/28 |

| Stock name: Greenland halibut in Division 1A inshore | | | | | | | | | | | | | | | | |
|--|---------------------|------------------|-----------------------------------|--------------------------|-------------------|------------|---------------|--|-------------|---|----------|-----|----|--------------------|-----------|------------------|
| Name of survey | Location (Div.) | Season months(s) | Years of survey from, to or since | Type of survey and gears | Target species | No.of sets | 24 hr fishing | Area of coverage in relation to stock distribution (%) | Depth range | Research data related to stock size estimates | | | | Individual weights | Country | SCS/ SCR Doc. No |
| | | | | | | | | | | Abundance | Bio-mass | ALK | LF | | | |
| Disko bay Inshore | Disko Bay | Jun-Jul | 1993-01 2001-18 | Longline Gillnet | Greenland halibut | 0 41 | Y | - | 200-700m | Y | Y | Y | Y | Y | Greenland | 19/033 |
| Uummannaq Inshore | Uummannaq | Aug-Sep | 1993-15 2011-18 | Longline Gillnet | Greenland halibut | 0 54 | Y | - | 200-900m | Y | Y | Y | Y | Y | Greenland | 19/033 |
| Upernivik Inshore | Upernivik | Aug-Sep | 1998-14 2012-18 | Longline Gillnet | Greenland halibut | - 52 | Y | - | 200-900m | Y | Y | Y | Y | Y | Greenland | 19/033 |
| Greenland Shrimp and Fish survey | Disko bay + (1A-1F) | Jul-Sep | 1991-18 | Stratified-random trawl | Shrimp & fish | 20 | N | - | 50-600m | Y | Y | Y | Y | Y | Greenland | 19/008 |

| Stock name: Redfish in Subarea 1 | | | | | | | | | | | | | | | | |
|----------------------------------|-----------------|------------------|-----------------------------------|--------------------------------|--|------------|---------------|--|-------------|---|----------|-----|----|--------------------|------------|------------------|
| Name of survey | Location (Div.) | Season months(s) | Years of survey from, to or since | Type of survey and gears | Target species | No.of sets | 24 hr fishing | Area of coverage in relation to stock distribution (%) | Depth range | Research data related to stock size estimates | | | | Individual weights | Country | SCS/ SCR Doc. No |
| | | | | | | | | | | Abundance | Bio-mass | ALK | LF | | | |
| West Greenland Groundfish | 1B - 1F | Oct-Nov | since 1982 | Stratified random bottom trawl | Cod, redfish, other finfish | 0 | N | ? | 0-400m | Y | Y | N | Y | Y | EU-Germany | SCR 14/28 |
| Greenland – Deepsea | 1C - 1D | Aug-Nov | since 1997 | Stratified random bottom trawl | Greenland halibut, Roundnose Grenadier | 53 | Y | ? | 400-1500m | Y | Y | N | Y | N | Greenland | SCR 18/021 |
| Greenland Shrimp and Fish survey | 1A - 1F | Jul-Aug | since 1992 | Stratified random bottom trawl | Shrimp & fish | 244 | N | ? | 50-600m | Y | Y | N | Y | N | Greenland | SCR 19/008 |



| Stock name: Wolffish in SA1 and American plaice in SA 1 | | | | | | | | | | | | | | | | |
|---|-----------------|------------------|-----------------------------------|--------------------------------|----------------------|------------|---------------|--|-------------|---|--------------|-----|----|-------------------------|------------|------------------------|
| Name of survey | Location (Div.) | Season months(s) | Years of survey from, to or since | Type of survey and gears | Target Species | No.of sets | 24 hr fishing | Area of coverage in relation to stock distribution (%) | Depth range | Research data related to stock size estimates | | | | Indiv- idual weights | Country | SCS/ SCR Doc. No |
| | | | | | | | | | | Abun- dance | Bio- mass | ALK | LF | | | |
| West Greenland groundfish | 1B - 1F | Oct-Nov | since 1982 | Stratified random bottom trawl | Cod, redfish finfish | 0 | N | ? | 0-400m | Y | Y | N | Y | Y | EU-Germany | SCS 14/28 |
| Greenland Shrimp and Fish survey | 1A - 1F | Jul-Aug | since 1992 | Stratified random bottom trawl | Shrimp, finfish | 244 | N | ? | 50-600m | Y | Y | N | Y | N | Greenland | SCR 17/015 |

| Stock name: Cod in Division 3M | | | | | | | | | | | | | | | | |
|--------------------------------|-----------------|------------------|-------------------------------------|---|----------------|------------|---------------|--|----------------------------------|---|----------|-----|----|--------------------|---------------------|------------------|
| Name of survey | Location (Div.) | Season months(s) | Years of survey from, to or since | Type of survey and gears | Target species | No.of sets | 24 hr fishing | Area of coverage in relation to stock distribution (%) | Depth range | Research data related to stock size estimates | | | | Individual weights | Country | SCS/ SCR Doc. No |
| | | | | | | | | | | Abundance | Bio-mass | ALK | LF | | | |
| EU-bottom trawl survey | 3M | Jun-Jul | since 1988 | Stratified random | All | 181 | N | 100% | <750 m (88-03) <1400m (04-18) | Y | Y | Y | Y | Y | EU | SCR 19/26 |
| Canadian Flemish Cap survey | 3M | Jan-Feb | 1978-1995 | Stratified random | All | 132 | Y | 100% | <730 m | Y | Y | Y | Y | Y | Canada Newfoundland | SCR 92/76 |
| Russian trawl survey | 3M | May-June | 1983-1993 1995-1996 2001-2002 | Stratified Random (and acoustic survey 1987-93) | All | 94 | Y | 100% | <1280 m | Y | Y | N | N | N | Russia | SCS 03/9 |
| Canadian groundfish survey | 3M | Sept-Oct | 1996 | Stratified random | All | 68 | Y | 100% | <730 m | Y | Y | N | N | N | Canada Newfoundland | SCR 97/42 |



| Stock name: American plaice in Division 3M | | | | | | | | | | | | | | | | |
|--|-----------------|------------------|-----------------------|--------------------------------|-----------------------|-------------|---------------|--|-------------|---|---|---|---|--------------------|----------------------|-------------------|
| Name of survey | Location (Div.) | Season months(s) | Years of survey since | Type of survey and gears | Target Species | No. of Sets | 24 hr fishing | Area of coverage in relation to stock distribution (%) | Depth range | Research data related to stock size estimates | | | | Individual weights | Country | SCS/ SCR Doc. No. |
| EU - bottom Trawl | 3M | Summer Jun/Jul | 1988 | Stratified-random bottom trawl | Groundfish and shrimp | 181 | N | 100 | 127-1460 m | Y | Y | Y | Y | Y | EU-Spain EU-Portugal | 19/21 |



| Stock name: Cod in Divisions 3NO | | | | | | | | | | | | | | | | |
|-------------------------------------|-----------------------|------------------|-----------------------------------|--------------------------------|----------------|------------|---------------|--|-------------|---|----------|-----|----|--------------------|---------|------------------|
| Name of survey | Location (Div.) | Season months(s) | Years of survey from, to or since | Type of survey and gears | Target species | No.of sets | 24 hr fishing | Area of coverage in relation to stock distribution (%) | Depth range | Research data related to stock size estimates | | | | Individual weights | Country | SCS/ SCR Doc. No |
| | | | | | | | | | | Abundance | Bio-mass | ALK | LF | | | |
| Canadian Spring Bottom Trawl Survey | 3NO | Apr-Jun | 1972-2017 (except 1983, 2006) | Stratified-random Bottom trawl | Multi-species | 134 | Y | 100 | 38-666 m | Y | Y | Y | Y | Y | Canada | SCR 19/15 |
| Juvenile flatfish | 3NO | Sep | 1989-94 | Stratified-random Bottom trawl | Groundfish | | Y | 100 | 30-150 m | Y | Y | Y | Y | Y | Canada | SCR 96/80 |
| Canadian Autumn Bottom Trawl Survey | 3NO | Oct-Dec | 1990-2017 (except 2014) | Stratified-random Bottom trawl | Multi-species | 121 | Y | 100 | 39-641 m | Y | Y | Y | Y | Y | Canada | SCR 19/15 |
| EU-Spain | 3NO - Regulatory Area | Jun | 1995-2017 | Stratified-random Bottom trawl | Multi-species | 122 | N | 20 | 44-1450 m | Y | Y | Y | Y | Y | Spain | SCR 19/19 |

| Stock name: Redfish in Divisions 3LN | | | | | | | | | | | | | | | | |
|--|---------------------|----------------------|-----------------------------------|--------------------------------|--------------------------|----------------------------|---------------|--|-----------------------------|---|---|---|---|--------------------|----------|-----------------------|
| Name of survey | Location (Div.) | Season months(s) | Years of survey from, to or since | Type of survey and gears | Target species | No. of sets | 24 hr fishing | Area of coverage in relation to stock distribution (%) | | Research data related to stock size estimates | | | | Individual weights | Country | SCS/ SCR Doc. No. |
| USSR groundfish | 3L, 3N | Apr-Jul (mostly) | 1972-82 | Fixed station Bottom trawl | Multi-species Groundfish | 50-60 per Div. | Y | Unknown | 1972-76: to 367m | Y | Y | Y | Y | Sometimes | USSR | 86/66 |
| | | | | | | | | | 1977-82: to 732 m | Y | Y | Y | Y | | | |
| USSR ground-fish | 3L, 3N | Apr-Jun (mostly) | 1983-91, 1993, 1994 | Stratified-random Bottom trawl | Multi-species | approx. 48-3L | Y | Unknown | to 732 m | Y | Y | Y | Y | Sometimes | USSR | 86/66 |
| | | | | | Groundfish | approx. 42-3N | Y | Unknown | to 732 m | Y | Y | Y | Y | | | |
| USSR acoustic | 3LN | Apr-Jun | 1987-91, 1993 | Acoustic | Multi-species | unknown midwater sampl. | Y | Unknown | to 732 m | Y | Y | | | Sometimes | USSR | 88/24 |
| Canadian groundfish (covering deep-water strata) | 3L | various | 1978-86 | Stratified-random Bottom Trawl | Multi-species Groundfish | min. of 40 in depths >183m | Y | Unknown | to 732 m | Y | Y | Y | Y | Sometimes | Canada | NAFO Studies No. 2 |
| Canada redfish | 3L | Jan | 1990 | Stratified-random Bottom trawl | Redfish | 55 | Y | Unknown | 183-732 m | Y | Y | Y | Y | N | Canada | NAFO Studies No. 2 |
| Canada ground-fish | 3LN | Apr-Jun | 1990-93 | Stratified-random Bottom trawl | Multi-species Groundfish | 90 in redfish depths | Y | Unknown | to 732 m | Y | Y | Y | Y | Y | Canada | NAFO Studies No. 2 |
| Canada redfish | 3L, 3N (since 1990) | Aug | 1990-91, 1993 | Stratified-random Bottom trawl | Redfish | 59-3L 47-3N | Y | Unknown | 183-732 m | Y | Y | Y | Y | Y | Canada | NAFO Studies No. 2 |
| Canadian Groundfish (Autumn) | 3LN | Autumn (Sep-Dec) | 1991-18 | Stratified-random Bottom trawl | Multi-species Groundfish | 49 in 3L 30 in 3N | Y | Unknown | 183-668 in 3L 93-634 in 3N | Y | Y | N | Y | Y | Canada | 19/015 and pers. com. |
| Canadian Groundfish (Spring) | 3LN | Spring (May-Jun) | 1991-18 | Stratified-random Bottom trawl | Multi-species Groundfish | 35 in 3L 30 in 3N | Y | Unknown | 194--665 in 3L 96-725 in 3N | Y | Y | N | Y | Y | Canada | 19/015 and pers. com. |
| Spanish Groundfish (Spring) | 3N | Spring (Jun) | 1995-18 | Stratified-random Bottom trawl | Multi-species Groundfish | 92 | N | Unknown | 44-1450 | Y | Y | N | Y | Y | EU-Spain | 19/018 and pers. com. |
| Spanish Groundfish (Summer) | 3L | Summer (July-August) | 2006-18 | Stratified-random Bottom trawl | Multi-species Groundfish | 100 | N | Unknown | 104-1478 | Y | Y | N | Y | Y | EU-Spain | 19/013 and pers. com. |



| Stock name: American plaice in Divisions 3LNO | | | | | | | | | | | | | | | | |
|---|------------------------------|--------------------------------|-----------------------------------|---|--|-------------------|----------------|--|-------------|---|-----------|-----|----|----------------------|---------|------------------|
| Name of survey | Location (Div.) | Season months(s) | Years of survey from, to or since | Type of survey and gears | Target species | No. o f sets | 24 hr fishin g | Area of coverage in relation to stock distribution (%) | Depth range | Research data related to stock size estimates | | | | Individ- ual weights | Country | SCS/ SCR Doc. No |
| | | | | | | | | | | Abundance | Bio- mass | ALK | LF | | | |
| Canadian spring | 3LNO | May-Jun | 1971-2018 (no 2006 or 2015, 2017) | Stratified random Campelen | Multispecies | 171 | Y | 95 | 44-732 m | Y | Y | Y | Y | Y | Canada | 19/05 18/39 |
| Canadian autumn | 3LNO | Sep-Dec | 1990-2018 (no 2004 or 2014) | Stratified random Campelen | Multispecies | 286 | Y | 100 | 42-1379 m | Y | Y | Y | Y | Y | Canada | 18/17 18/39 |
| EU-Spain | 3NO Reg. Area | Jun | 1995-2018 | Stratified random Campelen | Groundfish | 122 | N | 11 | 44-1450 m | Y | Y | N | Y | N | Spain | 19//19 |
| EU-Spain | 3L Reg. Area | Jul-Aug | 2003-2018 (no 2005) | Stratified random Campelen | Groundfish | 99 | N | unknown | 106-1433 | Y | Y | N | Y | N | Spain | 18/12 |
| Canadian DFO-Industry Co-operative | 3NO – expanded to 3L in 2000 | Jul-Aug (quarterly in 1997-98) | 1996-2004 | Grid survey with commercial Engel trawl (unlined) | Yellowtail (bycatches of cod, A. plaice) | 61 - Jul 80 - Aug | Y | 22 | 45-90 | N | N | N | N | N | Canada | 05/9 |

| Stock name: Yellowtail flounder in Divisions 3LNO | | | | | | | | | | | | | | | | |
|---|-----------------|------------------|-----------------------------------|----------------------------------|------------------------|------------|---------------|--|-------------|---|----------|-----|----|--------------------|---------|------------------|
| Name of survey | Location (Div.) | Season months(s) | Years of survey from, to or since | Type of survey and gears | Target species | No.of sets | 24 hr fishing | Area of coverage in relation to stock distribution (%) | Depth range | Research data related to stock size estimates | | | | Individual weights | Country | SCS/ SCR Doc. No |
| | | | | | | | | | | Abundance | Bio-mass | ALK | LF | | | |
| Canadian Groundfish Spring | 3LNO | May-June | Since 1971 | Stratified Random Campelen Trawl | Groundfish & shellfish | 269 | Y | 100 | 41-725 | Y | Y | N | Y | Y | Canada | 19/25 |
| Canadian Groundfish autumn | 3LNO | Sep- Nov. | Since 1990 | Stratified Random Campelen Trawl | Groundfish & shellfish | 286 | Y | 100 | 41-725 | Y | Y | N | Y | Y | Canada | 19/15 |
| Spanish Groundfish | 3NO | June | Since 1995 | Stratified Random Campelen Trawl | Groundfish & shellfish | 114 | N | | 44-1450 | Y | Y | N | Y | N | Spain | 19/18 |

| Stock name: Witch flounder in Divisions 3N and 3O | | | | | | | | | | | | | | | | |
|---|--------------------------|------------------|-----------------------------------|-----------------------------------|----------------|----------------|---------------|--|-------------|---|----------|------------|----|----------------------|---------|------------------|
| Name of survey | Location (Div.) | Season months | Years of survey from, to or since | Type of survey and gears | Target species | No.of sets | 24 hr fishing | Area of coverage in relation to stock distribution (%) | Depth range | Research data related to stock size estimates | | | | Indiv- idual weights | Country | SCS/ SCR Doc. No |
| | | | | | | | | | | Abundance | Bio-mass | ALK | LF | | | |
| Canadian Multispecies (Spring) | Div. 3NO | Spring (Apr-Jun) | 1971-2018 | Stratified Random - Bottom trawl | Multispecies | 79-3N 70-3O | Y | 80-90 | 41 - 725 m | Y | Y | until 1994 | Y | Y | Canada | 19/34 |
| Canadian Multispecies (Autumn) | Div. 3NO | Autumn (Oct-Dec) | 1990-2018 | Stratified- Random - Bottom trawl | Multispecies | 70-3N 75-3O | Y | 80-90 | 40 - 665 m | Y | Y | until 1994 | Y | Y | Canada | 19/34 |
| Spanish Groundfish (Spring) | Div. 3NO Regulatory Area | Spring (Apr-Jun) | 1995-2018 | Stratified- Random - Bottom trawl | Groundfish | 114 | N | 20 | 45 to 1480m | Y | Y | N | Y | N | Spain | 19/19 |



| Stock name: Capelin in Divisions 3NO | | | | | | | | | | | | | | | | |
|--------------------------------------|-----------------|------------------|-----------------------------------|----------------------------------|-----------------|-------------|---------------|--|-------------|---|-----------|-----|----|----------------------|---------|-------------------|
| Name of survey | Location (Div.) | Season months(s) | Years of survey from, to or since | Type of survey and gears | Target species | No. of sets | 24 hr fishing | Area of coverage in relation to stock distribution (%) | Depth range | Research data related to stock size estimates | | | | Indiv- idual weights | Country | SCS/ SCR Doc. No. |
| | | | | | | | | | | Abun- dance | Bio- mass | ALK | LF | | | |
| Multi-species bottom trawl survey | 3NO | 6 | 1977-2006 | Stratified-random, Campelen 1800 | Demersal fishes | 54 | + | 50 | 45-105 m | Y | Y | N | Y | Y | Canada | 07/12 |

| Stock name: Redfish in Divisions 3O | | | | | | | | | | | | | | | | |
|-------------------------------------|-----------------|---------------------------------|------------------------------------|--------------------------------|--------------------------|---------------------------------|---------------|--|-----------------------|---|--------------|-----|----|----------------------|---------|--------------------|
| Name of survey | Location (Div.) | Season months(s) | Years of survey from, to or since | Type of survey and gears | Target species | No. of sets | 24 hr fishing | Area of coverage in relation to stock distribution (%) | Depth range | Research data related to stock size estimates | | | | Indiv- idual weights | Country | SCS/ SCR Doc. No. |
| | | | | | | | | | | abun- dance | bio- mass | ALK | LF | | | |
| USSR ground-fish | 3O | Apr-Jun (mostly) | 1983-91, 1993, 1994 | Stratified-random Bottom trawl | Multi-species Groundfish | 75-100 | Y | Unknown | to 732 m | Y | Y | Y | Y | Sometimes | Russia | 86/66 |
| Canada ground-fish | 3O | Apr-Jun | 1971-90 | Stratified-random Bottom trawl | Multi-species Groundfish | 30-36 in redfish depths | Y | Unknown | to 367m | Y | Y | Y | Y | N | Canada | NAFO Studies No. 2 |
| EU-Spain | 3O | Jun | 1995-2018 | Stratified-random Bottom trawl | Multi-species Groundfish | 6-23 | N | Unknown | 93 - 1410 m | Y | Y | N | Y | Some Surveys | Spain | SCR 19/18 |
| Canada ground-fish | 3O | Sep-Dec series + Apr-May series | 1991-2013, 2015-18 1991-2015-18 | Stratified-random Bottom trawl | Multi-species Groundfish | 24-40 + 35-51 in redfish depths | Y | Unknown | To 1464 m To 732 m | Y | Y | N | Y | Y | Canada | SCR 19/15 |



| Stock name: Thorny skate in Divisions 3L, 3N, 3O, and 3Ps | | | | | | | | | | | | | | | | |
|---|--------------------------|----------------------|-----------------------------------|----------------------------------|------------------------|--------------------|---------------|--|---------------|---|---|---|---|--------------------|---------|-------------------|
| Name of survey | Location (Div.) | Season months(s) | Years of survey from, to or since | Type of survey and gears | Target Species | No. of Sets (2018) | 24 hr fishing | Area of coverage in relation to stock distribution (%) | Depth range | Research data related to stock size estimates | | | | Individual weights | Country | SCS/ SCR Doc. No. |
| Canadian Groundfish (Spring) | Div. 3LNOPs | Spring (March-June) | 1971-2018 | Stratified random - bottom trawl | Groundfish & shellfish | 439 | Y | 100 | 39 to 725 m | Y | Y | N | Y | Y | Canada | SCR 19/15 |
| Canadian Groundfish (Autumn) | Div. 3LNO | Autumn (Sept-Nov) | 1990-2018 | Stratified-random - bottom trawl | Groundfish & shellfish | 286 | Y | 42 | 40 to 668m | Y | Y | N | Y | Y | Canada | SCR 19/15 |
| EU-Spain Groundfish | Div. 3NO Regulatory Area | Spring (May-June) | 1997-2018 | Stratified-random - bottom trawl | Groundfish & shellfish | 114 | N | unknown | 44 to 1 450m | Y | Y | N | Y | Y | Spain | SCR 19/20 |
| EU-Spain Groundfish | Div. 3L Regulatory Area | Summer (July-August) | 2003-2018 | Stratified-random - bottom trawl | Groundfish & shellfish | 100 | N | unknown | 116 to 1 442m | Y | Y | N | Y | Y | Spain | SCR 19/13 |

| Stock name: White hake in Divisions 3N, 3O, and 3Ps | | | | | | | | | | | | | | | | |
|---|--------------------------|--------------------|-----------------------------------|----------------------------------|------------------------|--------------------|---------------|--|--------------|-----------|----------|-----|----|--------------------|---------|-------------------|
| Name of survey | Location (Div.) | Season months(s) | Years of survey from, to or since | Type of survey and gears | Target Species | No. of Sets (2018) | 24 hr fishing | Area of coverage in relation to stock distribution (%) | Depth range | abundance | bio-mass | ALK | LF | Individual weights | Country | SCS/ SCR Doc. No. |
| Canadian Groundfish (Spring) | Div.3NOPS | Spring (March-May) | 1971-2018 | Stratified random - bottom trawl | Groundfish & shellfish | 328 | Y | 100 | 39 to 725m | Y | Y | N | Y | Y | Canada | SCR 19/15 |
| Canadian Groundfish (Autumn) | Div. 3NO | Autumn (Sept-Oct) | 1990-2018 | Stratified-random - bottom trawl | Groundfish & shellfish | 145 | Y | unknown | 40 to 6665 | Y | Y | N | Y | Y | Canada | SCR 19/15 |
| EU-Spain Groundfish (Spring) | Div. 3NO Regulatory Area | Spring (May-June) | 1997-2018 | Stratified-random - bottom trawl | Groundfish & shellfish | 114 | N | unknown | 44 to 1 450m | Y | Y | N | Y | Y | Spain | SCR 19/020 |



| Stock name: Roughhead Grenadier in Subareas 2 and 3 | | | | | | | | | | | | | | | | |
|---|-----------------|------------------------|-----------------------|---|----------------------|-------------|---------------|--|-------------|---|---|---|---|---------------------|------------------|-------------------------------|
| Name of survey | Location (Div.) | Season Months(s) | Years of Survey since | Type of survey and gears | Target Species | No. of Sets | 24 hr fishing | Area of coverage in relation to stock distribution (%) | Depth Range | Research data related to stock size estimates | | | | Individual weight s | Country | SCR Doc. No. |
| Canadian Autumn Survey | 2GHJ 3MLNO | Oct-Dec | 1978 | Stratified random <= 1994 Engel >=1995 Campelen | Groundfish Shellfish | 513 in 2018 | Yes | | <1500 | + | + | | + | No | CAN | SCR 19/15 |
| Canadian Spring Survey | 3LNO | Spring May-July | 1978 | Stratified random <=1995 Engel >= 1996 Campelen | Groundfish Shellfish | 478 in 2018 | Yes | | <730 m | | + | | | No | CAN | SCR 19/15 |
| Canadian deepwater | 3MLN | August February Spring | 1991 1994 1995 | Bottom trawl | Groundfish | | | | <1500 | + | + | | + | No | CAN | SCR 95/51 |
| Spanish Surveys in Div. 3NO | 3NO | June | 1995 | Stratified Random <=2001 Pedreira >=2001 Campelen | Groundfish | 114 in 2018 | No | Low | <1500 | + | + | | + | + | EU-ESP | SCR 19/20 |
| Spanish Surveys in Div. 3L | 3L | August | 2003 | Stratified Random Campelen | Groundfish | 101 in 2018 | No | Low | <1500 | + | + | | + | + | EU-ESP | SCR 19/13 |
| EU Flemish Cap Surveys | 3M | July | 1988 | Stratified Random (Lofoten) | Groundfish Shrimp | 184 in 2018 | No | Flemish Cap | <1500 | + | + | + | + | + | EU-ESP EU-PRT | SCR 19/21 |
| Russian | 3M | May/June | 2001 and 2002 | Stratified random bottom trawl | Groundfish | 94 | Yes | Flemish Cap | 120-1280 m | + | + | | + | | RUS | SCR 96/34, 97/19, 97/20 |
| EU Deepwater | 3LMN | April-May | 1996 | Longline | Groundfish | 64 | Yes | | 700-3100m | | | | + | + | EU-ESP EU-PRT | SCR 03/09 SCR 07/10 |



| Stock name: Greenland halibut in Subareas 2+3KLMNO | | | | | | | | | | | | | | | | |
|--|-----------------|------------------|--|---|----------------------|-------------------------------|---------------|---|--|---|----------|-----|----|--------------------|---------|------------------|
| Name of survey | Location (Div.) | Season months(s) | Years of survey from, to or since | Type of survey and gears | Target species | No.of sets | 24 hr fishing | Area of coverage in relation to stock distribution (%) | Depth range | Research data related to stock size estimates | | | | Individual weights | Country | SCS/ SCR Doc. No |
| | | | | | | | | | | Abundance | Bio-mass | ALK | LF | | | |
| Canadian groundfish spring | 3LNO | Apr-June | 1995-2018 2006 & 2015 incomplete | Stratified-random Campelen trawl | Multi-species survey | 260 | Y | | 42-725 m | Y | Y | Y | Y | Y | Canada | 19/36 |
| Canadian autumn groundfish | 2J,3K, 3L | Autumn | 1978-2018 2J3K 1981-2017 3L | Stratified-random Engel trawl 1978-94 2J3K Yankee trawl 1981-82 3L Engel trawl 1984-94 3L Campelen Trawl all areas 1995-2018 | Multi-species survey | 2J -91 3K -111 3L - 141 | Y | This column not appropriate for this species given differences in depth coverage and areas covered in the surveys | 2J 129-1325 m 3K 139-658 m 3L 62-668 m | Y | Y | Y | Y | Y | Canada | 19/36 |
| EU bottom trawl | 3M | Jun-Jul | 1988-2018 | Stratified-random Lofoten trawl | Groundfish & shrimp | 181 | N | | 128-1460 m | Y | Y | Y | Y | Y | EU | 19/21 |
| EU-Spain bottom trawl | Reg. Area 3LNO | Apr-May | 3NO 1995-2018 3L 2003-18; Ex.2005 & 2008 | Stratified-random Bottom trawl | Groundfish | 3NO - 114 3L - 100 | N | | 3NO 44-1450m 3L 126-1447 m | Y | Y | N | Y | Y | Spain | 19/19 19/12 |



| | | | | | | | | | | | | | | | | |
|---------------------|------------------|-------------|---|---|----------------------|----------------|---|-------------|-----------------------------|---|---|---|---|---|--------|-------|
| Canadian groundfish | 2GH | Sep-Oct | 2G no longer covered; 2H annualy | Stratified-random Campelen in 1996-2018 | Multi-species survey | 2H-83 | Y | | 98-1399 m 2H | Y | Y | Y | Y | Y | Canada | 19/36 |
| Canadian groundfish | 3MNO | Sep-Dec | 3NO 1990-2018 3M 1996-2003, 2006-07 | Stratified-random Campelen | Multi-species survey | 70-3N 75-3O | | | 3N 40-634 m 3O 64-7665 m | | | | | | Canada | 19xx |
| Russian | 3M | May | 1987-96 | Stratified-random Bottom trawl | Groundfish | 90 | Y | Flemish Cap | 914 m in 1996 | Y | Y | Y | Y | N | Russia | 98/13 |
| Russian | 3LM Flemish Pass | Feb Apr-May | 1995-96 2000 | Stratified-random Bottom trawl | Groundfish | 84 | Y | | 732-1280 m | Y | Y | Y | Y | N | Russia | 01/10 |
| Russian | 3M | May-Jun | 2001-02 | Stratified-random Bottom trawl | Groundfish | | Y | | 127-1280m | Y | Y | Y | Y | N | Russia | 03/09 |



| Stock name: Short-finned Squid in Subareas 3 and 4 | | | | | | | | | | | | | | | | |
|--|-----------------|---|-----------------------------------|--------------------------------|----------------|----------------------------------|---------------|--|--|---|----------|-----|----------|----------------|------------|--------------------------|
| Name of survey | Location (Div.) | Season months(s) | Years of survey from, to or since | Type of survey and gears | Target species | No. of sets | 24 hr fishing | Area of coverage in relation to stock distribution (%) | Depth range | Research data related to stock size estimates | | | | Indiv. weights | Country | SCS/SCR Doc.No |
| | | | | | | | | | | Abundance | Bio-mass | ALK | LF | | | |
| July multispecies bottom trawl | SA 4 | July | Since 1970 | Stratified random bottom trawl | Multi-species | ~190 | Y | Unknown | 20-366 m | Y | Y | N | Y | N | Canada | SCR 19/42REV |
| CA Autumn multispecies bottom trawl | 4T | Sep | Since 1971 | Stratified random bottom trawl | Multi-species | 63-202; increased over the years | Y, after 1986 | Unknown, likely < 20% | | Y | Y | N | Un-known | N | Canada | SCR 19/42REV |
| EU Flemish Cap multispecies bottom trawl | 3M | July | Since 1988 | Stratified random bottom trawl | Multi-species | 101-121 | N, 0600-2200 | Unknown, likely < 5% | 127-730 m, up to 1,400 m in 2005 | Y | Y | N | Y | N | EU (Spain) | SCR 16/21 SCR19/42REV |
| CA Spring multispecies bottom trawl | 3LNOP | April-June | Since 1996 | Stratified random bottom trawl | Multi-species | 470-597 | Y | Unknown | 20-870 m, but inconsis-tent inshore | Y | Y | N | Y | N | Canada | SCR 19/42REV |
| CA Autumn multispecies bottom trawl | 3KLNO | Mainly Sept-Dec, but also Jan in some years | Since 1995 | Stratified random bottom trawl | Multi-species | 608-840 | Y | Unknown | 30-1 500 m, va ri able inshore and > 730 m | Y | Y | N | Y | N | Canada | SCR 19/42REV |

*The data in this table is only updated every three years (e.g., 2016 and 2019) when a full assessment is conducted

| Stock name: Northern Shrimp in Divisions 3LNO | | | | | | | | | | | | | | | | |
|---|-----------------|---------------------|-----------------------------------|--------------------------------|--|-------------------|---------------|--|-------------|---|----------|-----|----|--------------------|----------|------------------|
| Survey | Location (Div.) | Season months(s) | Years of survey from, to or since | Type of survey and gears | Target species | No. of sets | 24 hr fishing | Area of coverage in relation to stock distribution (%) | Depth range | Research data related to stock size estimates | | | | Individual weights | Country | SCS/ SCR Doc. No |
| | | | | | | | | | | Abundance | Bio-mass | ALK | LF | | | |
| Multi-species Spring Survey | 3LNO 3Ps | May to June | 1999 to present | Stratified-random bottom trawl | Groundfish Shellfish (crab, shrimp)/ capelin | ~300 sets in 3LNO | Yes | 100% of area covered | 30-732 m | Y | Y | N | Y | N | Canada | SCR 16/58 |
| Multi-species Autumn Survey | 2HJ 3KLNO | October to December | 1995 to present | Stratified-random bottom trawl | Groundfish Shellfish (crab, shrimp)/ capelin | ~300 sets in 3LNO | Yes | 100% of area covered | 30-1500 m | Y | Y | N | Y | N | Canada | CSAS 2017/12 |
| EU Bottom Trawl Survey | 3NO | May June | 1995 - present | Stratified-random bottom trawl | Groundfish Shellfish (crab, shrimp)/ capelin | 95 | No | | 60-1500 | Y | Y | | | | EU-Spain | SCR 16/51 |
| EU Bottom Trawl Survey | 3L | August | 2003 - present | Stratified-random bottom trawl | Groundfish Shellfish (crab, shrimp)/ capelin | 97 | No | | 60-1500 | Y | Y | | Y | Y | Spain | SCR 16/51 |

| Stock name: Northern Shrimp in Division 3M | | | | | | | | | | | | | | | | |
|--|-----------------|------------------|-----------------------------------|--------------------------------|----------------|-------------|---------------|--|---------------------------------|---|----------|-----|----|--------------------|---------|------------------|
| Name of survey | Location (Div.) | Season months(s) | Years of survey from, to or since | Type of survey and gears | Target species | No. of sets | 24 hr fishing | Area of coverage in relation to stock distribution (%) | Depth range | Research data related to stock size estimates | | | | Individual weights | Country | SCS/ SCR Doc. No |
| | | | | | | | | | | Abundance | Bio-mass | ALK | LF | | | |
| EU-Flemish Cap Survey | 3M | Jul | 1988 to present | Stratified-random bottom trawl | All | 181 | N | 100% | <750m (88-02) <1400m (03-19) | Y | Y | Y | Y | Y | EU | 19/51 19/53 |



| Stock name: : Northern Shrimp in Subareas 0+1 | | | | | | | | | | | | | | | | |
|--|-----------------|------------------|-----------------------------------|---|--------------------------------|-------------------------|---------------|--|-------------|---|--------------|------------------|----|---------------------------------|---------|---------------------------|
| Name of survey | Location (Div.) | Season months(s) | Years of survey from, to or since | Type of survey and gears | Target species | No.of Sets ¹ | 24 hr fishing | Area of coverage in relation to stock distribution (%) | Depth range | Research data related to stock size estimates | | | | Indiv- idu- al weights | Country | SCS/ SCR Doc. No |
| | | | | | | | | | | Abundance | Bio- mass | ALK ² | LF | | | |
| West Greenland Trawl Survey (West Greenland Shelf) | SA 0+1 | Jun-Jul | Since 1988 | Stratified semi-systematic; trawl; Cosmos 2000 w/rockhopper | Northern shrimp and groundfish | 166 | N | 95 | 150-600m | Y | Y | N/A | Y | Y | GRL | 19/043 19/046 |

¹ First number is number of stations trawled at depths between 150 and 600 m targeting shrimp; second number is number of stations in water shallower than 150 m targeting principally Atlantic cod; Third number is number of stations in water deeper than 600 m targeting principally Greenland halibut.

² Lack of satisfactory method of ageing shrimp precludes generating age-length keys.

| Stock name: : Northern Shrimp in Denmark Strait* | | | | | | | | | | | | | | | | |
|--|-----------------|------------------|--|---|-------------------------------|------------|---------------|--|-------------|---|--------------|-----|----|---------------------------------|-----------|---------------------------|
| Name of survey | Location (Div.) | Season months(s) | Years of survey from, to or since | Type of survey and gears | Target species | No.of sets | 24 hr fishing | Area of coverage in relation to stock distribution (%) | Depth range | Research data related to stock size estimates | | | | Indiv- idu- al weights | Country | SCS/ SCR Doc. No |
| | | | | | | | | | | Abundance | Bio- mass | ALK | LF | | | |
| East Greenland Trawl Survey | ICES XIV | July- Aug | Since 2008 No surveys in 2017, 2018, and 2019 | Stratified semi-systematic; trawl; Cosmos 2000 w/rockhopper | Northern shrimp Groundfish | 101 | N | unknown | 0-600m | Y | Y | N | Y | Y | Greenland | 16/45 |

*No update for 2017 and 2018 due to a cancelled survey.

