



SCIENTIFIC COUNCIL MEETING – JUNE 2020

**Assessment of Thorny Skate (*Amblyraja radiata* Donovan, 1808)
in NAFO Divisions 3LNO and Subdivision 3Ps**

by

M.R. Simpson, and C.M. Miri

Fisheries and Oceans Canada
Northwest Atlantic Fisheries Centre, P.O. Box 5667
St. John's, NL, Canada A1C 5X1

Abstract

Available information on the fishery, management, biology, and assessment of Thorny Skate in NAFO Divisions 3LNO and Subdivision 3Ps was reviewed to determine the status of this stock. Based on the continuous distribution of this species, and lack of physical barriers between Divs. 3LNO and Subdiv. 3Ps, Thorny Skate in Divs. 3LNOPs is considered to constitute a single stock. In 2012-2018, an average of 3 805 tons of Thorny Skate from Divs. 3LNO was commercially landed. STACFIS-agreed total landings from Divs. 3LNO were 3 697 t in 2019. Reported landings in Subdiv. 3Ps averaged 460 t in 2012-2018, and totaled 889 t in 2019. An Index of Fishing Mortality for Divs. 3LNO increased from the late 1980s to a peak of 32% in 1997, then decreased and stabilized at approximately 17% during 1998-2004. In 2005-2011, this Index averaged 6%, and remained around 4% since 2012. The Fishing Mortality Index in Subdiv. 3Ps was $\leq 2\%$ since 2010. After a drastic decline over 1985-1995, Canadian spring research surveys indicated that biomass and abundance of Thorny Skate in Divs. 3LNO were relatively stable at low levels. Thorny Skate distribution in Divs. 3LNOPs over 2010-2019 continued to be concentrated on the southwestern Grand Banks, in Subdiv. 3Ps, and northward along the edge of the Bank.

Introduction

Thorny Skates (*Amblyraja radiata* Donovan, 1808) are widely distributed in temperate and Arctic waters of the North Atlantic. In the western Atlantic, Thorny Skate are distributed from Greenland to South Carolina, with the centre of distribution on the Grand Banks (Fig. 1) in NAFO Divisions 3LNO. Commercial catches of skates consist mainly of Thorny Skate, with about 95% in Canadian skate catches (Kulka and Miri 2007); which is similar to the 97-99% of Thorny Skates observed in the Canadian spring survey (see **Survey Catch Composition – Other Skates** below). In the EU-Spain research survey over 2014-2016 in Divs. 3NO, Thorny Skate comprised approximately 90% of the skate taken in the survey (González-Costas *et al.* 2020). Thus, the skate fishery on the Grand Banks can be considered a directed fishery for Thorny Skate.



Fishery and Management

TAC Regulation

Thorny Skate came under quota regulation in 1995, after a directed skate fishery was established in 1994 by Canada in its Exclusive Economic Zone (EEZ). A Total Allowable Catch (TAC) of 5 000 tons for Divisions 3LNO and 1 000 t for Subdivision 3Ps were adopted by Canada in 1995. In 1996, the TAC was raised to 6 000 t for Divs. 3LNO and 2 000 t for Subdiv. 3Ps. In 1997, the TAC was reduced to 1 950 t for Divs. 3LNO, and 1 050 t for Subdiv. 3Ps.

Outside Canada's EEZ, catch was unregulated until September 2004, when the Northwest Atlantic Fisheries Organization (NAFO) Fisheries Commission set a TAC of 13 500 t for 2005-2009 in the NAFO Regulatory Area (NRA) of Divs. 3LNO (Fig. 2). This TAC was lowered to 12 000 t for 2010-2011, and to 8 500 t for 2012. The TAC was further reduced to 7 000 t for 2013-2020. The TAC for Subdiv. 3Ps has been maintained at 1 050 t.

Landings Trends

On the Grand Banks, Kulka and Mowbray (1998) reported that significant bycatch of skates have been taken since commencement of offshore fishing in the late 1940s, initially by non-Canadian fleets and later by Canadian vessels. Prior to the mid-1980s, non-Canadian fleets comprised the largest component of offshore fisheries on the Grand Banks, and took several thousand tons of skate as bycatch each year. The bycatch derived primarily from the Greenland Halibut fishery and from the Canadian mixed fishery for Thorny Skate, White Hake (*Urophycis tenuis*), and Monkfish (*Lophius americanus*). Kulka and Mowbray (1998) estimated that approximately 5 000 t, on average, were discarded annually by Canadian fleets during the 1980s and early 1990s, although only a few hundred tons were recorded in Canada's annual landings statistics during that period.

Catches for Divs. 3LNO (Table 1; Fig. 2) increased in the mid-to-late 1980s with the commencement of a directed fishery for Thorny Skate. In 1985, Spain began targeting skate in a non-regulated fishery in the NRA (Junquera and Paz 1998; del Río and Junquera 2001). During 1985-1991, landings averaged 17 058 t and peaked at approximately 28 400 t in 1991 (STATLANT-21A). This fishery was mainly prosecuted by Spain, Portugal, USSR, and the Republic of Korea. Non-Canadian landings declined significantly to 5 059 t in 1992 (Table 1). In 2000, Russia joined the directed fishery for Thorny Skate. Due to a new Canadian-directed fishery that began in 1994, Canadian landings increased during 1994-1999, to an average of 1 590 t (Simpson and Miri 2012). Since 2000, total reported landings of skate by all countries in Divs. 3LNO declined (Fig. 2). In 2012-2018, an average of 3 831 t of Thorny Skate was landed from Divs. 3LNO. STACFIS-agreed landings from Divs. 3LNO in 2019 totaled 3 697 t.

In Subdiv. 3Ps, STATLANT-21A data indicated that Canadian fleets reported the majority of Thorny Skate landings, while St. Pierre and Miquelon (EU-France) annually reported smaller amounts of landings (Table 2; Fig. 2). Prior to 1994, Canadian landings of Thorny Skate in Subdiv. 3Ps rarely exceeded a few hundred tons. Total reported landings in Subdiv. 3Ps averaged 1 328 t from 1994-2008, 500 t from 2009-2011, and 460 t over 2012-2018. In Subdiv. 3Ps, reported Canadian landings totaled 889 t in 2019.

Commercial Size

Sampling of Canadian commercial catches by Canadian At-Sea Fisheries Observers indicated that Thorny Skates trawled in the Div. 3L redfish (*Sebastes* sp.) fishery varied between 27-93 cm in 2016-2019, including modes of 35-40, 56, 62-66, and 72-82 cm (Fig. 3a; see Simpson and Miri 2012 for previous years). Skates caught by Canadian trawlers in the Divs. 3NO Yellowtail Flounder (*Limanda ferruginea*) fishery in 2016-2019 were 24-101 cm, with no dominant size classes observed. In 2017, Thorny Skates trawled in the Div. 3O Witch Flounder (*Glyptocephalus cynoglossus*) fishery ranged between 42-100 cm, with a mode of 80 cm. Skates trawled in the

Divs. 2J3KL Greenland Halibut (*Reinhardtius hippoglossoides*) fishery in 2018 varied between 31-88 cm in length, with modes of 48, 53, and 63 cm.

From skate-directed trawl fisheries (280 mm mesh) in the NRA of Divs. 3LNO over 2013-2019, EU-Spain reported a range of 15-100 cm TL skates (no distinct modes), with a small number of young-of-the-year (≤ 21 cm) caught in 2013-2014 and 2017-2018 (Fig. 3b; see Simpson and Miri 2012 for previous years). In 2013, EU-Portugal caught 26-85 cm TL skates (mode of 49-50 cm) in Div. 3N (280 mm mesh).

In other directed trawl fisheries (130-135 mm mesh) of Divs. 3LNO (NRA) in 2013-2019, EU-Portugal reported skate bycatch ranging from 25-88 cm TL, including modes of 46-49, 60-64, and 72-76 cm (Fig. 3c; see Simpson and Miri 2012 for previous years). EU-Portugal did not sample Divs. 3LNO skate bycatch in 2014-2016 and 2018, while EU-Spain have not done so since 2009). Russia sampled Thorny Skates during the Div. 3LN Greenland Halibut fishery in 2012, ranging from 24-78 cm TL (Fig. 3c; see Simpson and Miri 2012 for previous years). In the Divs. 3LO redfish fishery over 2013-2016, Russia reported skate bycatch of 35-89 cm. Russia sampled only 5 and 14 specimens in 2017 and 2018 (respectively). In 2019, Russia sampled 494 skates as bycatch in Divs. 3LNO ranging from 31-95 cm TL, with mean lengths of 56, 64, and 71 cm (respectively per Division).

Research Survey Data

Canadian Spring Surveys

Stratified-random surveys have been conducted by Canadian research vessels in the spring (April-June) of each year from 1971 to 2019. A summary of the stratified-random survey design adopted by the DFO-NL Region can be found in Doubleday (1981). While survey design has remained constant, additional strata have been included, and some of the original strata have been modified (Bishop 1994). A significant change in the surveys was the addition of shallower and deeper strata after 1993. Additional causes of variation in spring survey coverage are discussed in detail most recently by Rideout (2020), and Rideout and Ings (2020). The spring survey can be split into three time series, based on the trawl used in each period: 1971-1982 (Yankee 41.5), 1983-1995 (Engel 145), and 1996-2019 (Campelen 1800; McCallum and Walsh 1996). Conversion factors exist for the Engel to Campelen gear change (Simpson and Kulka 2005), but not for the Yankee to Engel gear change. In 2006, most of Subdiv. 3Ps was not surveyed, and only shallow strata in Divs. 3NO were surveyed, due to Canadian research vessels' mechanical difficulties. Therefore, survey estimates for that year are not comparable to others in the Campelen time series. In 2015 and 2017, Div. 3L was not completely surveyed, thus impacting biomass and abundance estimates of Thorny Skate.

Historical abundance and biomass indices from Canadian spring surveys in Divs. 3LNOPs are provided in Table 3(a,b) and Figures 4a and 5. Since the mid-1990s, spring biomass indices for Thorny Skate in Divs. 3LNOPs were generally increasing, following a declining trend over 1985-1995. In 2018, biomass estimates for Thorny Skate in Div. 3L, 3N, and 3O were 16 983 t, 40 868t, and 76 215 t (respectively). The biomass index in Div. 3N had increased since 2015 but declined in 2019 to 19 019 t. The biomass index in 2019 was 72 605 t in Div. 3O, slightly above the 2013-2017 average of 69 354 t. However, the 2019 biomass estimate of 11 860 t in Div. 3L was well below the 2018 value and the 2013-2017 average of 34 743 t. In Subdiv. 3Ps, the biomass index was 48 339 t and 37 211 t in 2018 and 2019 respectively. In 2019 the index declined and was below the 2014-2018 average of 39 903 t.

Canadian Fall Surveys

Annual stratified-random fall surveys have been conducted by Canada in Div. 3L from 1981 to 2019. In 1990-2019, fall surveys also extended onto the southern Grand Banks in Divs. 3NO. Canadian surveys were originally conducted with an Engel 145 trawl, then a Campelen 1800 trawl in 1995-2019. Although fall surveys sample deeper maximum depths (~1 450 m) than those in spring (~730 m), they do not cover the entire stock area, and thus are not considered spatially complete. For this reason, Canadian spring surveys are used as the primary source of biomass and abundance estimates for Thorny Skate. However, fall indices are still considered

in assessments of this stock, because this survey is conducted when a greater proportion of Thorny Skate is available to survey trawls. During fall, Thorny Skates are concentrated on the shelf; in spring, part of this population has moved to the shelf edge, and a proportion apparently moves outside of the survey area (Kulka *et al.* 2004). When using spring estimates of biomass and abundance to examine trends in this stock, it is assumed that the proportion of skate that moves outside of the surveyed area remains consistent between years. Additional causes of variation in fall survey coverage are discussed in detail most recently by Rideout (2020), and Rideout and Ings (2020).

Historical abundance and biomass indices from Canadian fall surveys in Divs. 3LNO are provided in Table 4(a,b) and Figure 4b. Fall abundance indices of Thorny Skate remained stable at very low levels since 1995. Divs. 3NO were not surveyed in fall 2014, as well as deep-water strata (>732 m) of Div. 3L in 2015, 2017-2018; thereby impacting abundance and biomass estimates of this species. In 2018, biomass estimates in Div. 3L, 3N, and 3O were 28 142 t, 71 267 t, and 72 500 t (respectively). In Div. 3N, the 2018 biomass estimate was above the 2013-2017 average of 59 268 t, while that for Div. 3O was above the five-year average of 65 354 t. The 2019 biomass estimate of 147 348 t for Divs. 3NO was above the 2014-2018 average of 128 451 t; increasing since 2015. In Div. 3L, the 2018 (28 142 t) and 2019 (24 403 t) biomass estimates were above the 2013-2017 average of 21 413 t.

EU-Spain 3NO Survey

Spain initiated a stratified-random survey in the NRA of Divs. 3NO in 1995. Initially, this survey was carried out in spring with the C/V *Playa de Mendiña* using a Pedreira bottom trawl. Since 2001, the R/V *Vizconde de Eza* replaced that research vessel, and a Campelen trawl replaced Pedreira gear (González-Troncoso *et al.* 2020). Abundance and biomass of Thorny Skate were estimated from EU-Spain surveys in the NRA of Divs. 3NO from 1997-2019. Although the EU-Spain biomass index indicated an increase in 2012, it appears to be following a declining trend since 2006 from 55 810 t to its lowest estimate of 2 758 t in 2019.

A comparison of the Canadian Campelen spring 3NO biomass indices to those of the Spanish Divs. 3NO surveys in 1997-2019 indicated that the trends have diverged since 2007: the EU-Spain index declined until 2011, while the Canadian index has followed a generally increasing trend (Fig. 6). It should be noted that the Canadian spring survey covers the entire area of Divs. 3NO, whereas the Spanish survey is limited to the NRA of Divs. 3NO.

EU-Spain 3L Survey

Spain initiated a stratified-random summer bottom trawl survey in the NRA of Div. 3L in 2003. This survey was conducted with the R/V *Vizconde de Eza* using a Campelen trawl. Due to the vessel's mechanical difficulties, the survey was not conducted in 2005, and some strata were missed in 2003-2004 (Román-Marcote *et al.* 2020). Abundance and biomass estimates of Thorny Skate were available from EU-Spain surveys in the NRA of Div. 3L from 2003-2019 (excluding 2005).

A comparison of the Spanish Div. 3L biomass and Canadian spring Div. 3L indices in 2003-2019 indicated generally similar patterns until 2014 when they diverged: the EU-Spain index followed an increasing trend to 2015 (Fig. 7). The EU-Spain index has since declined from 18 753 t to its lowest values in 2018 (7 479 t) and 2019 (7 820 t). The Canadian Campelen Div. 3L fall biomass indices reflected a pattern similar to that from the Spanish survey until 2016, when they diverged: the EU-Spain index followed the declining trend, while Canadian biomass estimates continued along a gradually increasing trend from 3 660 t to 5 302 t. It should be noted that the Canadian surveys cover the entire area of Div. 3L, whereas the Spanish survey is limited to the NRA of Div. 3L.

Index of Fishing Mortality

A relative Fishing Mortality Index (Relative F=STACFIS-agreed commercial landings/Canadian spring survey biomass) was calculated for Thorny Skate in Divs. 3LNO and Subdiv. 3Ps for 1996-2019. Relative F for Divs. 3LNO increased from the late 1980s to a peak of 32% in 1997, then decreased and remained stable at approximately 17% during 1998-2004 (Fig. 8). In 2005-2011, this Divs. 3LNO index averaged 6%, then remained around 4% since 2012. Since 2010, relative fishing mortality in Subdiv. 3Ps was $\leq 2\%$.

Survey Catch Composition – Other Skates

Canadian spring survey catches of skates in Divs. 3LNOPs over 1996-2019 were dominated by Thorny Skate: comprising 97-99% (Fig. 9). The remaining 2-3% of skates were mainly Smooth Skate (*Malacoraja senta*) and Spinytail Skate (*Bathyrhaja spinicauda*), with lesser amounts of Winter Skate (*Leucoraja ocellata*), Jensen's Skate (*Amblyraja jenseni*), Round Skate (*Rajella fyllae*), Barndoor Skate (*Dipturus laevis*), and Soft Skate (*Malacoraja spinacidermis*). The Smooth Skate spring biomass index was highest in 2009, then followed a declining trend to 2019 (Fig. 10). Although highly variable, Spinytail Skate spring biomass has been increasing since 2003; while Winter Skate appeared generally in decline since then. The Round Skate spring biomass index was also highly variable, but at very low levels (except for high estimates in 2010, 2018). The remaining skate species were occasionally caught by the Canadian Campelen trawl.

The Canadian fall survey of Divs. 3LNO in 1995-2019 consistently caught Smooth Skate, Spinytail Skate, and Round Skate; with biomass estimates fluctuating throughout this period (Fig. 10). In addition to the skate species caught in spring surveys, Bigelow's Skate (*Rajella bigelowi*) was captured occasionally at low levels in fall surveys.

Survey Size Structure

Lengths of Thorny Skates caught in Canadian Campelen spring surveys of Divs. 3LNO and Subdiv. 3Ps in 2008-2019 ranged from 7-105 cm TL (Fig. 11). For most areas and years, a peak of young-of-the-year skates (YOY: 7-21 cm TL) was observed, and averaged 15 cm. In 2008-2017, modes between 65-77 cm were usually found, with a few years also indicating modes within 23-28 cm. Similarly, the 2018 and 2019 spring surveys showed modes between 66-79 cm, and also 33-36 cm, while a 50-cm peak was seen in 2018. The Canadian Campelen fall surveys of Divs. 3LNO in 2008-2019 caught Thorny Skates from 9-105 cm TL (Fig. 12). Peaks of YOY skates were also found in most years, averaging 14-15 cm. Also notable were two dominant peaks at 44 cm and 70 cm in 2010. Although larger Thorny Skates (82-92 cm) were similar in all years, they were absent in 2014 while numbers of 73-81 cm skates were greatly reduced due to the lack of a fall survey in Divs. 3NO.

Lengths of Thorny Skate caught in the EU-Spain 3NO survey during 1997-2019 ranged from 11-104 cm. There has been no recruitment visible in many years (González-Costas *et al.* 2020). In the EU-Spain 3L survey over 2006-2019, Román-Marcote *et al.* (2020) reported a 10-95 cm range of Thorny Skates. In 2018-2019, they observed modes between 32-40 cm, and 62-76 cm; similar to the Canadian spring surveys.

Recruitment

A standardized index of Thorny Skate recruitment, based on the number of skates ≤ 21 cm TL, is illustrated in Figure 13. This recruitment index declined from 1.5 in 1996 to below the 1996-2019 average, with the lowest value in 1999 and 2004. Although this index remained above 1.0 over 2010-2013, it was followed by a steep decline to in 2014-2015. The recruitment index increased in 2016-2017, then dropped below average in 2018. In 2019, the recruitment index was at a level comparable to the long-term average. The 2006 index is not presented, because Canadian survey coverage was incomplete over that spring, and the Div. 3L survey was incomplete in 2015 and 2017.

Distribution

In Divs. 3LNO and Subdiv. 3Ps, the distribution of Thorny Skate has changed significantly since the 1980s. In the early 1980s, Thorny Skates were widely distributed over the entire Grand Banks in moderate to high concentrations (Kulka and Miri 2007). By the late 1990s, much of the biomass was concentrated in the southwest. In 2001-2005, the area of high concentration expanded northward and along the Bank edge. It is important to note that part of this population moves to the shelf edge in spring, thereby moving outside of the Canadian spring survey area; they are concentrated on the shelf during the fall (Kulka *et al.* 2004). In 2010-2019, Thorny Skate distribution in Divs. 3LNOPs continued to be concentrated on the southwestern Grand Banks, in Subdiv. 3Ps, and northward along the edge of the Bank in the spring survey (Fig. 14a,b). Strata-biomass plots for 2008-2019 show the expansion of skate biomass along the southwestern slopes and into Subdiv. 3Ps (Fig. 15a,b). In the Canadian fall survey over 2010-2019, Thorny Skate continued to be concentrated along the shelf edge; although higher densities were also apparent on the Grand Banks (Fig. 16a,b).

Summary

Thorny Skate underwent a decline in Divs. 3LNO over the late 1980s, suddenly increased in 1990-1991, declined again over 1992-1996, then stabilized at a low level (except for a slight increase in 2007-2008). Thorny Skate distribution in Divs. 3LNOPs for 2010-2019 continued to be concentrated on the southwestern Grand Banks, in Subdiv. 3Ps, and northward along the edge of the Bank. An Index of Fishing Mortality for Divs. 3LNO increased from the late 1980s to a peak of 32% in 1997, then stabilized at approximately 17% during 1998-2004. In 2005-2011, this Index averaged 6%, and remained around 4% since 2012. From a peak of 18 277 tons in 2000, total reported landings of skate by all countries in Divs. 3LNO declined to an average of 5 357 t in 2005-2011. In 2012-2018, average reported landings from Divs. 3LNO further declined to 3 805 t. In 2019, STACFIS-agreed landings totaled 3 697 t. The 2020 TAC of 7 000 t for skates in the NRA of Divs. 3LNO continues to exceed the average commercial catch during a period when minimal or no rebuilding of this stock has occurred.

Acknowledgments

We thank Fisheries and Oceans Canada staff, who participated in Canadian research surveys, Canadian At-Sea Fisheries Observers, who collected data and specimens aboard commercial vessels under adverse conditions over many years, and the Instituto Español de Oceanografía in Vigo (Spain) for data from Spanish research surveys of Divs. 3LNO (in the NRA); thereby allowing us to estimate trends and examine various biological aspects of Thorny Skate.

References

- Bishop, C.A. MS 1994. Revisions and additions to stratification schemes used during research vessel surveys in NAFO Subareas 2 and 3. NAFO SCR Doc. 94/043. 10p.
- del Río, J.L., and S. Junquera. MS 2001. Spanish skate (*Raja radiata* Donovan, 1808) fishery in the Grand Bank (NAFO Division 3N): 1997-2000. NAFO SCR Doc. 01/031, Ser. No. N4408. 10p.
- Doubleday, W.G. 1981. Manual on groundfish surveys in the Northwest Atlantic. NAFO Sci. Counc. Stud. No. 2.
- González-Costas, F., G. Ramilo, E. Román, J. Lorenzo, A. Gago, D. González-Troncoso, M. Sacau, P. Duran, M. Casas, and J.L. del Río. 2020. Spanish Research Report for 2019. NAFO SCS Doc. 20/07, Ser. No. N7045. 41p.
- González-Troncoso, D., I. Garrido, and A. Gago. 2020. Biomass and length distribution for roughhead grenadier, thorny skate, white hake and squid from the surveys conducted by Spain in NAFO 3NO. NAFO SCR Doc. 20/010, Ser. No. N7054. 34p.

- Junquera, S., and X. Paz. MS 1998. Non-traditional resources: Skate fishery and survey results in Division 3NO. NAFO SCR Doc. 98/026, Ser. No. N3011. 6p.
- Kulka, D.W., and C.M. Miri. 2007. Update on the status of Thorny Skate (*Amblyraja radiata* Donovan, 1808) in NAFO Divisions 3L, 3N, 3O, and Subdivision 3Ps. NAFO SCR Doc. 07/033, Ser. No. N5385. 30p.
- Kulka, D.W., C.M. Miri, M.R. Simpson, and K.A. Sosebee. 2004. Thorny Skate (*Amblyraja radiata* Donovan, 1808) on the Grand Banks of Newfoundland. NAFO SCR Doc. 04/035, Ser. No. N4985. 108p.
- Kulka, D.W., and F.K. Mowbray. MS 1998. The status of Thorny Skate (*Raja radiata*), a non-traditional species in NAFO Divisions 3L, 3N, 3O and 3Ps. CSAS Res. Doc. 98/131. 70p.
- McCallum, B.R., and S.J. Walsh. 1996. Groundfish survey trawls used at the Northwest Atlantic Fisheries Centre, 1971-present. NAFO SCR Doc. 96/050.
- Rideout, R.M. 2020. Do spatial coverage issues in the 2019 Canadian (NL) RV bottom trawl surveys influence the suitability of survey indices for use in NAFO stock assessments? NAFO SCR Doc. 20/xxx, Ser.No. Nxxxx. p.
- Rideout, R.M., and D.W. Ings. 2020. Temporal and spatial coverage of Canadian (Newfoundland and Labrador region) Spring and Autumn multi-species RV bottom trawl surveys, with an emphasis on surveys conducted in 2019. NAFO SCR 20/002, Ser. No. N7041. 59p.
- Román-Marcote, E., D. González-Troncoso, and M. Alvarez. 2020. Results for the Atlantic cod, roughhead grenadier, redfish, thorny skate and black dogfish of the Spanish Survey in the NAFO Division 3L for the period 2003-2019. NAFO SCR Doc. 20/014, Ser. No. N7059. 76p.
- Simpson, M.R., and D.W. Kulka. 2005. Development of Canadian research trawl gear conversion factors for Thorny Skate on the Grand Banks based on comparative tows. NAFO SCR Doc. 05/049. 14p.
- Simpson, M.R., and C.M. Miri. 2012. Assessment of Thorny Skate (*Amblyraja radiata* Donovan, 1808) in NAFO Divisions 3LNO and Subdivision 3Ps. NAFO SCR Doc. 12/028, Ser. No. N6054. 32p.

Table 1. NAFO-reported landings (tons) of skates in Divs. 3LNO, 1960-2019 (STATLANT-21A).

Year	Canada	Other	Total	Year	Canada	Other	Total
1960	0	73	73	1990	44	14,726	14,770
1961	0	119	119	1991	18	28,390	28,408
1962	0	99	99	1992	78	5,059	5,137
1963	0	65	65	1993	78	5,992	6,070
1964	0	145	145	1994	1,554	6,601	8,155
1965	17	199	216	1995	2,412	4,912	7,324
1966	75	347	422	1996	1,314	4,804	6,118
1967	212	188	400	1997	2,165	9,903	12,068
1968	128	31	159	1998	1,013	8,501	9,514
1969	68	1,123	1,191	1999	1,081	10,864	11,945
1970	99	539	638	2000	498	17,779	18,277
1971	125	77	202	2001	354	14,507	14,861
1972	64	487	551	2002	1,107	10,648	11,755
1973	10	413	423	2003	671	13,592	14,263
1974	638	1,690	2,328	2004	352	11,476	11,828
1975	180	2,535	2,715	2005	685	2,853	3,538
1976	260	1,006	1,266	2006	249	5,255	5,504
1977	551	1,266	1,817	2007	101	6,110	6,211
1978	816	1,015	1,831	2008	243	6,867	7,110
1979	382	657	1,039	2009	435	5,286	5,721
1980	351	1,027	1,378	2010	50	5,314	5,364
1981	244	1,467	1,711	2011	69	5,479	5,548
1982	52	756	808	2012	185	4,144	4,329
1983	4	1,277	1,281	2013	21	4,323	4,344
1984	0	2,013	2,013	2014	2	4,472	4,474
1985	9	10,390	10,399	2015	4	3,322	3,326
1986	52	14,277	14,329	2016	1	3,480	3,481
1987	195	18,301	18,496	2017	5	4,157	4,162
1988	91	18,675	18,766	2018	2	127	129
1989	15	14,222	14,237	2019	7	3267	3274

Table 2. NAFO-reported landings (tons) of Thorny Skate in Subdiv. 3Ps, 1960-2019 (STATLANT-21A).

Year	Canada	Other	Total		Year	Canada	Other	Total
1960	0	11	11		1990	5	549	554
1961	0	17	17		1991	1	639	640
1962	0	11	11		1992	13	46	59
1963	0	58	58		1993	22	11	33
1964	0	145	145		1994	1,566	3	1,569
1965	0	85	85		1995	1,866	4	1,870
1966	0	126	126		1996	603	2	605
1967	0	162	162		1997	829	3	832
1968	86	67	153		1998	1,251	6	1,257
1969	0	353	353		1999	1,102	4	1,106
1970	35	229	264		2000	935	21	956
1971	303	213	516		2001	1,769	39	1,808
1972	8	184	192		2002	1,413	238	1,651
1973	7	231	238		2003	1,705	82	1,787
1974	122	641	763		2004	1,190	87	1,277
1975	9	490	499		2005	967	15	982
1976	91	230	321		2006	910	78	988
1977	521	360	881		2007	1,347	491	1,838
1978	454	256	710		2008	763	632	1,395
1979	545	121	666		2009	645	354	999
1980	554	609	1,163		2010	342	529	871
1981	558	520	1,078		2011	513	228	741
1982	117	395	512		2012	354	224	578
1983	0	516	516		2013	286	271	557
1984	21	602	623		2014	201	145	346
1985	21	944	965		2015	168	75	243
1986	7	1,576	1,583		2016	404	250	654
1987	52	787	839		2017	413	192	605
1988	2	781	783		2018	944	115	1,059
1989	0	1,685	1,685		2019	889	0	889

Table 3a. Abundance of Thorny Skate from Canadian spring research surveys in Divs. 3LNOPs, 1984-2019. Surveys were conducted with an Engel trawl (1984-spring 1995; converted to Campelen-equivalents), and a Campelen trawl (spring 1996-2019). The deeper (>103 m) portion of Divs. 3NO, as well as Subdiv. 3Ps, were not surveyed in 2006; and Div. 3L was not completely surveyed in 2015 and 2017.

Year	3L	3N	3O	3Ps	3LNOPs
Engel series - Campelen Equivalents					
1984	7,574	25,226	24,615	9,417	66,832
1985	63,081	45,278	50,123	55,214	213,697
1986	51,231	53,394	21,134	36,153	161,911
1987	39,151	33,538	34,041	28,113	134,844
1988	35,030	26,474	42,991	19,043	123,538
1989	40,350	30,030	17,678	25,863	113,921
1990	43,938	71,656	40,118	21,344	177,055
1991	34,779	44,549	35,195	50,254	164,777
1992	37,475	20,645	35,567	21,510	115,198
1993	27,765	17,068	15,025	21,580	81,437
1994	15,999	17,565	19,105	19,221	71,891
1995	9,320	7,017	26,781	19,493	62,611
Campelen Series					
1996	10,418	10,636	22,731	25,591	69,376
1997	6,804	13,554	25,635	18,379	64,372
1998	7,764	10,140	34,130	22,781	74,815
1999	8,263	15,967	36,042	20,212	80,484
2000	12,512	16,027	28,525	18,574	75,638
2001	8,521	16,276	33,321	17,606	75,724
2002	5,920	8,469	32,902	17,560	64,851
2003	6,737	9,645	34,734	24,615	75,732
2004	4,762	8,925	21,153	24,256	59,095
2005	11,011	15,986	26,621	26,399	80,016
2006	8,450	23,618	17,778		49,846
2007	11,357	24,065	23,317	11,440	70,180
2008	10,572	14,477	22,738	31,239	79,027
2009	5,810	15,560	18,132	19,128	58,629
2010	10,964	20,714	32,747	26,447	90,872
2011	7,226	12,731	31,576	23,409	74,942
2012	13,342	15,866	24,268	21,848	75,324
2013	10,681	20,682	35,416	35,067	101,846
2014	7,668	27,140	23,675	23,081	81,564
2015		10,500	23,590	21,535	60,889
2016	9,154	10,154	27,085	45,191	91,584
2017		25,020	23,173	40,199	88,476
2018	14,380	19,628	28,626	25,431	88,066
2019	11,649	12,979	37,315	19,740	81,684

Table 3b. Biomass of Thorny Skate from Canadian spring research surveys in Divs. 3LNOPs, 1984-2019. Surveys were conducted with an Engel trawl (1984-1995; converted to Campelen-equivalents), and a Campelen trawl (1996-2019). The deeper (>103 m) portion of Divs. 3NO, as well as Subdiv. 3Ps, were not surveyed in 2006; and Div. 3L was not completely surveyed in 2015 and 2017.

Year	3L	3N	3O	3Ps	3LNOPs
Engel series - Campelen Equivalents					
1984	17,269	57,720	61,026	20,318	156,333
1985	102,351	86,438	110,322	36,954	336,065
1986	69,864	110,325	46,634	47,728	274,551
1987	82,037	60,535	51,007	40,697	234,276
1988	70,143	49,686	87,375	29,993	237,197
1989	73,291	49,142	40,172	44,271	206,875
1990	45,312	47,479	61,946	24,264	179,002
1991	22,197	28,925	99,003	61,534	211,659
1992	11,945	23,047	57,929	38,693	131,615
1993	8,546	18,550	35,113	16,256	78,465
1994	3,920	10,193	28,874	16,539	59,526
1995	2,798	2,824	32,323	24,924	62,869
Campelen Series					
1996	4,993	11,010	35,529	21,851	73,382
1997	3,969	9,703	28,293	20,705	62,669
1998	5,807	13,186	42,351	28,629	89,972
1999	7,266	26,254	54,045	32,062	119,626
2000	14,011	27,861	40,917	22,528	105,317
2001	10,383	29,197	59,078	24,566	123,223
2002	8,580	13,986	38,025	22,127	82,718
2003	8,411	18,216	49,707	37,072	113,406
2004	7,806	20,425	39,740	38,354	106,325
2005	19,266	33,757	46,515	32,702	132,240
2006	16,193	56,698	25,252		98,143
2007	25,044	54,188	48,369	21,080	148,682
2008	23,344	32,196	42,220	38,509	136,270
2009	7,765	29,478	52,619	27,788	117,651
2010	14,944	34,303	68,435	39,968	157,650
2011	10,046	21,239	57,020	44,310	132,615
2012	14,828	38,621	53,443	33,699	140,592
2013	11,713	43,547	76,358	34,455	166,072
2014	11,497	53,229	59,285	43,580	167,591
2015		23,031	69,703	34,788	134,701
2016	9,324	17,422	86,328	40,403	153,477
2017		36,484	55,096	46,290	138,146
2018	16,983	40,868	76,251	48,339	182,440
2019	11,860	19,019	72,605	37,211	140,695

Table 4a. Abundance of Thorny Skate from Canadian fall research surveys in Divs. 3LNO, 1981-1994. Surveys were conducted with an Engel trawl.

Year	Div. 3L	Div. 3N	Div. 3O	3LNO
Engel Series				
1981	33,523			
1982	36,223			
1983	103,303			103,303
1984	70,979			70,979
1985	86,070			86,070
1986	75,424			75,424
1987	80,879			80,879
1988	86,633			86,633
1989	76,793			76,793
1990	116,758	43,855	53,191	213,803
1991	73,576	61,128	29,680	164,384
1992	94,058	33,854	24,675	152,587
1993	61,501	31,073	41,382	133,957
1994	44,205	50,141	30,748	125,094

Table 4b. Abundance of Thorny Skate from Canadian fall research surveys in Divs. 3LNO, 1995-2019. Surveys were conducted with a Campelen trawl. Divs. 3NO were not surveyed in 2014; as well as deep-water strata (>732 m) of Div. 3L in 2015, 2017-2018.

Year	Div. 3L	Div. 3N	Div. 3O	3LNO
Campelen Series				
1995	23,299	37,322	30,582	91,203
1996	23,483	22,694	45,145	91,321
1997	13,448	30,540	50,047	94,035
1998	8,917	21,132	29,785	59,834
1999	10,448	25,116	31,847	67,411
2000	12,536	31,419	39,918	83,873
2001	12,655	21,352	42,095	76,103
2002	7,541	30,925	24,488	62,954
2003	9,363	19,203	34,556	63,121
2004	6,369	21,068	32,343	59,780
2005	11,346	20,027	30,553	61,927
2006	8,888	23,211	27,688	59,787
2007	13,372	36,453	29,768	79,594
2008	15,856	48,011	40,944	104,811
2009	17,145	28,813	42,965	88,922
2010	18,429	30,859	28,137	77,426
2011	16,841	26,907	36,711	80,459
2012	21,202	30,226	51,813	103,240
2013	15,693	25,196	35,785	76,674
2014	24,555			24,555
2015	12,555	30,530	20,287	63,372
2016	17,646	33,054	26,358	77,058
2017	20,499	34,883	31,239	86,621
2018	19,432	44,457	31,515	95,403
2019	18,322	29,031	44,760	92,113

Table 5a. Biomass of Thorny Skate from Canadian fall research surveys in Divs. 3LNO, 1981-1994. Surveys were conducted with an Engel trawl.

Year	Div. 3L	Div. 3N	Div. 3O	3LNO
Engel Series				
1981	36,467			
1982	65,293			
1983	165,500			165,500
1984	149,061			149,061
1985	141,054			141,054
1986	113,170			113,170
1987	87,843			87,843
1988	107,910			107,910
1989	67,877			67,877
1990	95,586	67,459	97,496	260,540
1991	52,655	103,959	75,526	232,141
1992	40,289	52,980	42,383	135,652
1993	24,096	35,528	64,294	123,918
1994	16,212	50,950	31,929	99,090

Table 5b. Biomass of Thorny Skate from Canadian fall research surveys in Divs. 3LNO, 1995-2019. Surveys were conducted with a Campelen trawl. Divs. 3NO were not surveyed in 2014; as well as deep-water strata (>732 m) of Div. 3L in 2015, 2017-2018.

Year	Div. 3L	Div. 3N	Div. 3O	3LNO
Campelen Series				
1995	11,306	40,775	44,653	96,734
1996	14,459	28,629	36,969	80,057
1997	7,534	43,075	58,160	108,770
1998	9,205	34,279	39,280	82,764
1999	13,614	32,609	42,608	88,831
2000	17,722	61,202	40,861	119,786
2001	16,420	34,311	62,156	112,886
2002	11,068	52,855	40,593	104,517
2003	14,463	36,829	46,123	97,416
2004	11,327	45,678	26,361	83,366
2005	20,107	37,442	61,595	119,143
2006	18,610	54,372	50,605	123,587
2007	30,089	70,198	56,976	157,263
2008	27,182	83,861	75,892	186,935
2009	22,848	40,801	63,200	126,849
2010	21,051	27,270	54,857	103,178
2011	16,150	51,955	69,053	137,158
2012	19,124	43,739	107,777	170,640
2013	20,752	44,160	92,042	156,954
2014	20,283			20,283
2015	19,540	66,638	48,720	134,898
2016	22,921	61,263	55,231	139,415
2017	23,568	65,010	65,422	154,000
2018	28,142	71,267	72,500	171,909
2019	24,403	56,070	91,278	171,750




Figure 1. Map of NAFO Divisions 3LNO and Subdivision 3Ps (thick black dashed lines) in relation to Canada's Exclusive Economic Zone (thin black dotted curve).




Figure 2. Total reported landings and Total Allowable Catch (TAC) of Thorny Skate in Divs. 3LNO (STACFIS) and Subdiv. 3Ps (STATLANT-21A), 1985-2019.




Figure 3a. Length distributions of Canadian commercial catches (sexes combined) in Subareas 2+3 from skate bycatch fisheries, 2016-2019. Note the Y-axis for Shrimp in 2018 is almost double the value of all others. Data are from Canadian At-Sea Fisheries Observers, and include discards.




Figure 3b. Length distributions of commercial catches (sexes combined) in Divs. 3LNO by country for the skate-directed (280 mm) trawl fishery, 2012-2019.




Figure 3c. Length distributions of commercial catches (sexes combined) in Divs. 3LNO by country for skate bycatch (130-135 mm) trawl fisheries in 2012-2013, 2015-2017, and 2019.




Figure 4a. Relative biomass and abundance indices for Thorny Skate from Canadian spring research surveys in NAFO Divs. 3LNO and 3LNOPs, 1971-2019. Divs. 3LNO were not surveyed in 1983; the deeper (>103 m) portion of Divs. 3NO, as well as Subdiv. 3Ps, were not surveyed in 2006; and Div. 3L was not completely surveyed in 2015 and 2017.




Figure 4b. Canadian fall research survey biomass and abundance indices for Thorny Skate in NAFO Divs. 3LNO, 1990-2019. Deep strata of Divs. 3NO were not surveyed in 2003, 2004, 2006, 2008; both Divisions were not surveyed in 2014; as well as deep strata of Div. 3L in 2015, 2017-2018.




Figure 5. Mean numbers and weights (kg) per tow (+/- 95% CI) of Thorny Skate from Canadian spring surveys in NAFO Divs. 3LNO and 3LNOPs, 1983-2019. Divs. 3LNO were not surveyed in 1983; the deeper (>103 m) portion of Divs. 3NO, as well as Subdiv. 3Ps, were not surveyed in 2006; and Div. 3L was not completely surveyed in 2015 and 2017




Figure 6. Comparison of Thorny Skate biomass indices from the Canadian Campelen spring survey and the Spanish spring survey in Divs. 3NO, 1997-2019. Note that Divs. 3NO were not surveyed by Canada in 2014, and Spanish surveys occur only in the NAFO Regulatory Area (NRA) of Divs. 3NO.




Figure 7. Comparison of Thorny Skate biomass indices from Canadian Campelen surveys and the Spanish summer survey in Div. 3L, 2003-2019. The Canadian spring survey was incomplete in 2015 and 2017; as well as Canada's fall survey of Div. 3L (deep-water) in 2015, 2017-2018; and Spanish surveys occur only in the NAFO Regulatory Area (NRA) of Div. 3L.




Figure 8. Fishing Mortality Index (reported landings/spring survey biomass) for Divs. 3LNO and Subdiv. 3Ps, 1985-2019. Commercial landings are STACFIS-agreed; biomass indices are from Canadian spring research surveys. The deeper (>103 m) portion of Divs. 3LNO, as well as Subdiv. 3Ps, were not surveyed in 2006. The Div. 3L survey was incomplete in 2015 and 2017.




Figure 9. Percent of each skate species caught in Canadian Campelen spring surveys of Divs. 3LNOPs, 1996-2019. Note the inset graph represents proportions of species with Thorny Skate removed. In 2006, the deeper portion of Divs. 3NO, as well as Subdiv. 3Ps, were not surveyed. Div. 3L was not completely surveyed in 2015 and 2017.




Figure 10. Biomass indices (in tons) for various skate species caught in Canadian Campelen spring surveys (black columns) of Divs. 3LNOPs in 1996-2019, and in fall surveys (gray columns) of Divs. 3LNO over 1995-2019. Spring: Subdiv. 3Ps was not surveyed in 2006; and Div. 3L was not completely surveyed in 2015, 2017. Fall: Divs. 3NO were not surveyed in 2014.




Figure 11. Length distributions of Thorny Skate from Canadian Campelen spring surveys in Divs. 3LNOPs, 2008-2019. Note the Y-axis for 2016-2017 is double the value of all others. The Div. 3L survey was incomplete in 2015 and 2017.




Figure 12. Length distributions of Thorny Skate from Canadian Campelen fall surveys in Divs. 3LNO, 2008-2019. Divs. 3NO were not surveyed in 2014.





Figure 13. Standardized recruitment index (Thorny Skate ≤ 21 cm) from Canadian spring surveys in Divs. 3LNOPs, 1996-2019. The deeper portion (>103 m) of Divs. 3NO, as well as Subdiv. 3Ps, were not surveyed in 2006. The Div. 3L survey was incomplete in 2015 and 2017. The red, horizontal line depicts the standardized average recruitment for 1996-2019.






Figure 14b. Distribution (number of skates per tow) of Thorny Skate on the Grand Banks (Divs. 3LNOPs) during Canadian spring surveys, 2015-2019. The Div. 3L survey was incomplete in 2015 and 2017.




Figure 15a. Distribution (biomass in tons) of Thorny Skate on the Grand Banks (Divs. 3LNOPs) during Canadian spring surveys, 2008-2013.




Figure 15b. Distribution (biomass in tons) of Thorny Skate on the Grand Banks (Divs. 3LNOPs) during Canadian spring surveys, 2014-2019. The Div. 3L survey was incomplete in 2015 and 2017.




Figure 16a. Distribution (number of skates per tow) of Thorny Skate on the Grand Banks (Divs. 3LNO) during Canadian fall surveys, 2010-2014. Divs. 3NO were not surveyed in 2014.




Figure 16b. Distribution (number of skates per tow) of Thorny Skate on the Grand Banks (Divs. 3LN0) during Canadian fall surveys, 2015-2019.