



**SCIENTIFIC COUNCIL MEETING - JUNE 2023**

Trawl and gillnet survey results from the Disko Bay, NAFO Division 1A Inshore

by

Rasmus Nygaard

Greenland Institute of Natural Resources, P.O. Box 570, 3900 Nuuk, Greenland

**Abstract**

This paper presents the updated indices for the surveys performed by the Greenland Institute of Natural Resources (GINR) in the Disko Bay located in NAFO division 1A inshore. The Disko Bay has been part of the Greenland Shrimp and Fish survey (trawl) since 1992. Greenland halibut in the Disko Bay was previously also surveyed with longlines, but in 2001 the longline survey was replaced by a gillnet survey directed at pre fishery recruits. A larger meshed section (90mm halfmesh) was added to the gillnet survey from 2016 to survey the commercial part of the stock and increase the number of large Greenland for ageing.

**Introduction**

Greenland halibut is a dominant fish species in the North-west Greenlandic fjords and of major importance to the people living in the area. In the Disko Bay, targeted species include Greenland halibut, shrimp, cod and snow crab, whereas other species like wolffish and redfish are mainly landed bycatch from the longline and gillnet fishery directed at Greenland halibut.

*Survey area*

The *Disko Bay* is characterized by areas of smooth bottom and depths are mostly less than 600 meters (Figure 1). Glaciers are located in the North-Eastern part of the bay at *Torssukattak* and in the central eastern part the *Ilulissat Icefjord*, where deeper waters are located (+900m). *Ilulissat Icefjord (Kangia)* was declared a UNESCO World Heritage Site in 2004 because of its natural beauty and the importance of the fast-moving *Jakobshavn Glacier*. The glacier often fill the icefjord with massive icebergs that strand at the *Ilulissat Icefjord Bank* just south of the city *Ilulissat*. Therefore the icefjord are rarely accessible during the summer months. The *Disko Bay* is connected to the *Baffin bay* through the western part and in the North Western part through the narrow fjord *Vejgat*. A map of the *Disko bay* is provided in figure 1.

**Materials and methods**

Longline surveys (not presented here)

Surveys have been conducted in the area since the mid 1970's, using different types of longlines. The longline surveys were highly variable from year to year and not easily interpreted. (for further details see *Simonsen et al. 2000*).

Trawl surveys

The central part of the *Disko Bay* and the slopes of *vejgat* are trawlable with a substantial shrimp fishery occurring. The *Disko Bay* has been part of the *Greenland Shrimp and Fish survey (SFW)* since 1991. From 1991



to 2017 the survey was conducted with the 722 GRT stern trawler M/Tr 'Pâmiut'. In 2005 the gear was changed in this survey, but since then the area coverage and the trawl and its rigging has been unchanged. RV Pâmiut was decommissioned from 2018 to 2020 the survey was updated with a chartered vessel using Pâmiut's gear. No survey was conducted in 2021. From 2022 the survey is annually updated with the new GINR research vessel R/V Tarajoq.

### Gillnet survey

In 2001, a gillnet survey replaced the poorly performing longline survey in the Disko bay. The main objective for starting up the gillnet survey was a well-estimated selectivity and the possibility for targeting pre-fishery sized Greenland halibut (35-55cm TTL). The gillnet survey uses fixed positions of stations arranged in transects towards the important fishing grounds West of Ilulissat city and Torssukattak icefjord in the northeastern part of the Disko Bay. The gillnets are composed of 60m long sections with mesh sizes 46, 55, 60 and 70 mm (knot to knot or half mesh). Sections are separated with a 2m open space to prevent catchability interactions. Soak time is approximately 6-18 hours and fishing takes place both day and night. Stations are paired two and two, close to each other (0,5-1 NM) to allow for analysis of within station variability. From 2016 and forward a 90mm section (halfmesh) was added to the gillnet survey to increase the number of large Greenland for halibut and also survey the commercial part of the stock. From 2012 the survey is annually updated with the GINR research vessel R/V Sanna.

### Biological samples

Length, weight, gutted weight, otoliths and occasionally DNA samples are regularly collected during the surveys. Otoliths are collected from individual Greenland halibut and frozen in a plastic bag with a printed plastic label with individual information and an automatically created number. At the GINR, otoliths are read after a method developed in Norway. In the Lab otoliths are photographed with translucent light with a leica S9i stereomicroscope in a 5 mp TIF image. After imaging the otoliths are archived. Digitally archived Images are then "read on screen" using imageJ. In ImageJ both contrast and brightness can easily be adjusted and a calibration beam allows for digital measurements of proportions of the otolith. Images are standardized and attempts for automated digital reading are being tested.

### ALK

An Age Length Key is produced from the aged otoliths for each cm group for the Disko bay for each year. If the ALK is incomplete for certain lengths, a backup ALK (derived from all northern inshore areas) is used for the missing length combinations. The backup-ALK produced from all inshore areas in a given year, is screened for the missing length-age combinations. To produce a complete backup-ALK, missing ages for certain lengths are estimated from the von bertalanffy growth equation.

## Results

An overview of the most recent surveys and successful stations by year, vessel and gear is given in table 1.

The number of otoliths collected in the survey has in general been between 500 and 1000 in most years (Table 2). Ageing of the otoliths was in the past done by looking at the dried otolith through a stereomicroscope. However from 2007 to 2009 the method changed to looking at fresh frozen otoliths through a stereomicroscope. Uncertainty about the method however led to a lack of reading until 2017. In 2017 the ageing was done by taking a High resolution (14mb) digital image under a Leica S9i digital stereomicroscope.

The Greenland Shrimp and fish survey in the Disko Bay, identified increasing abundance during the 1990s and high abundances were found from 1998 to 2005 when the old (skjaervoy) trawl was used (fig 2). In 2005 a new trawl (cosmos) was introduced and the timeseries was broken. After 2006 the abundance indices returned to the lower levels with the exception of the high abundances identified in 2011, 2013 (fig 2). Although such large recruitment events of age 1, has not been observed since then, the abundance indices have remained stable or slightly increasing. The length distribution in the survey shows higher than average numbers of ages 2 and 3 in most of the recent 5 years, indicating good recruitment in recent years. Especially the abundance of 2 year old Greenland halibut in 2017 and 2019 (2015 YC, 2017 YC respectively) and 3 year old in 2018, 2020 and 2022(2015 and 2017 YC respectively), seems far above average (fig 3). Greenland halibut are found all over the Disko Bay (figure 4 and 5)

The biomass indices in the trawl survey indicate a steady increase during the 1990's, with a substantial increase observed in 2003 and 2004 (fig 2 right). After the gear change in 2005, the biomass index gradually decreased until 2014. From the 2014 low, the biomass index has gradually increased and the 2022 biomass estimate is the highest since 2011 (fig 2).

#### Gillnet survey in the Disko Bay

The gillnet survey CPUE and NPUE indicated low levels of pre-fishery recruits in 2006 and 2007, but returned to above average levels in 2008 and 2010 (figure 6). The increase in 2011 NPUE's was observed in the northern area of the Bay, while in the main fishing grounds at the Ice fjord bank around Ilulissat the NPUEs remained low (Figure 6). The high numbers of larger fish in 2011, did not seem to have any origin in the previous year estimated populations. From 2013 to 2017 the gillnet CPUE and NPUE, indices gradually decreased to below average. However, from 2017 to 2021 the NPUE and CPUE gradually increased to the highest levels observed in the timeseries. (Figure 6). The substantial decrease observed in the 2022 gillnet survey seems far outside the trend in the most recent years and may be an outlier.

The 90mm section added in 2016, has in general not impacted the overall length distribution in the Disko bay, indicating few larger individuals (55-70 cm) in the surveyed area (fig 7).

Although the distribution of the stations varies from year to year, the increasing number of Greenland halibut seems evenly distributed over all stations (figure 8).

The strong year-classes observed in the trawl survey are also visible in the Gillnet survey from 2018 and onwards (figure 9). Over all the gillnet survey implies very good recruitment in recent years with the 2015 YC being particularly large, potentially followed by good 2017 and 2018 Year classes.

A CAA for the survey is calculated on basis of the new digital age readings (Table 3 and figure 9). The method holds a number of advantages that will not be mentioned here. The first initial years with untrained readers show unrealistic Mean-weight-At-Age (Figure 10). However the newest readings (2020 and 2021) and years prior to 2017 has less variability with a realistic and stable MWAA. LW relationship for the sampled Greenland halibut is given in table 6.

#### Cod

From 2013, significant numbers of cod have also been taken in the survey. Only length frequencies are shown but not the index (figure 11).

### Discussion

Trawl calibration experiments indicated that the difference in catchability between the gears was length dependant for Greenland halibut and was at equilibrium at lengths around 12 cm, but twice as high at 40 cm. Since the abundance is highly influenced by 1 year old recruits (15 cm) which normally constitute 80-90% of the abundance in the survey, there is little impact the abundance index whether calibrating or not. This is not true for the biomass where calibrating has a higher impact on the indices. However, since the calibration experiments revealed an almost 1:1 relationship between the most abundant individuals, but a stronger

difference individuals that must have been rare in the experiments, this could also imply that the catchability difference between the trawls are overestimated. The indices prior to and non-calibrated indices after the gear change are at the same level for Greenland halibut (but also other species) indicate that the effect of the gear change on the indices may be overestimated.

The correlation between the abundance of Greenland halibut larger than 35 cm in the trawl survey and the NPUE indices from the gillnet survey, provides an increased credibility in the survey indices of both surveys. The surveys generally occur separated by a month or less and in the same overall areas at the same depth intervals. The trawl survey covers most of the bay and relies on randomly distributed stations, whereas the gillnet survey relies on fixed stations.

If comparing the gillnet NPUE (all sizes) to the trawl survey indices of Greenland halibut larger than 35 cm, the surveys seems to be correlated to some extent. This correlation between the surveys seems to disappear in the 2019 and 2020 survey. The correlation between the surveys could be caused by an evenly distributed stock with a high overlap in size selectivity of the two very different gears in relation to the present length distribution of the stock.

In the beginning of the Disko Bay gillnet survey time series commercial landed Greenland halibut were larger in the area and fish smaller than 55 cm were regarded pre-fishery recruits. However, in the recent decade smaller and smaller Greenland halibut have been landed and nowadays the fishery targets Greenland halibut as small as 30 cm with smaller hooks and illegal use of finer meshed gillnets (80mm) selecting Greenland halibut around 1 kg most efficiently. Therefore, the gillnet survey is increasingly surveying both recruits and the fished stock.

#### Recruitment

Although recruitment seems to vary from year to year at age 1, In most years this does not seem to be the case at age 2 or 3. Often estimates strong or weak YC at age one seems close to average levels a year or two later. It has been suggested that this may be related to density dependant mortality at in years of extra high recruitment (Sünksen et al. 2009). However, the recent year classes (2015 YC and later) that can be followed in the length distribution in the trawl survey from 2016 to 2018 and in the gillnet survey from 2018 to 2021 tells a different story. The increase in the Gillnet survey indicates an unusually good recruitment in recent years. These year classes are showing up in the CAA plot as age 5 and 6. Age 6 in this case has reached a size of close to 50 cm and 1 kg corresponding to the mean size in the commercial catches.

#### References

- Boje, J. and Lyberth, B. (2005) Survey Calibration for Greenland Halibut in Division 1A Inshore. NAFO Scr. Doc.05/57 (N5143)
- Holland, D. M., Thomas, R. H., de Young, B., Ribergaard, M. H., & Lyberth, B. (2008). Acceleration of Jakobshavn Isbræ triggered by warm subsurface ocean waters. *Nature Geoscience*, 1, 659. Retrieved from <http://dx.doi.org/10.1038/ngeo316>
- Motyka, R. J., M. Truffer, M. Fahnestock, J. Mortensen, S. Rysgaard, and I. Howat (2011), Submarine melting of the 1985 Jakobshavn Isbræ floating tongue and the triggering of the current retreat, *J. Geophys. Res.*, 116, F01007, doi: 10.1029/2009JF001632.
- Nygaard, R. and Nogueira 2020. Biomass and Abundance of Demersal Fish Stocks off West Greenland Estimated from the Greenland Shrimp and Survey, 1988-2018. NAFO Scr. doc.20/008.
- Simonsen, C.S., Boje, J. and Kingsley, M.C.S., 2000. A Review Using Longlining to Survey Fish Populations with Special Emphasis on an Inshore Longline Survey for Greenland Halibut (*Reinhardtius hippoglossoides*) in West Greenland, NAFO Division 1A. NAFO Scr.Doc., 00/29

Sünksen, K., Stenberg, C., & Grønkjær, P. (2010). Temperature effects on growth of juvenile Greenland halibut (*Reinhardtius hippoglossoides* Walbaum) in West Greenland waters. *Journal of Sea Research*, 64(1), 125–132. <http://doi.org/https://doi.org/10.1016/j.seares.2009.10.006>



**Table 1.** Number of stations by gear and Area (Table is incomplete)

Year	Trawl	vessel	Longline	Gillnet	Vessel	Notes
1990	*	Pa			AJ	
1991	41	Pa			AJ	
1992	39	Pa			AJ	
1993	31	Pa	31	-	AJ	
1994	27	Pa	32	-	AJ	
1995	33	Pa	-	-	--	
1996	33	Pa	29	-	AJ	
1997	34	Pa	24	-	AJ	
1998	33	Pa	-	-	-	
1999	34	Pa	26	-	AJ	
2000	23	Pa	35	-	AJ	
2001	23	Pa	15	8	AJ	46,55,60,70
2002	22	Pa		55	AJ	46,55,60,70
2003	19	Pa		56	AJ	
2004	14	Pa	8	50	AJ	
2005	16	Pa	9	47	AJ	
2006	21	Pa	3	44	AJ	
2007	18	Pa	2	30	AJ	
2008	16	Pa	0	35	Ch	
2009	24	Pa	-	-	-	
2010	25	Pa	0	48	AJ	
2011	26	Pa	0	50	AJ	
2012	21	Pa	0	41	Sa	60 mm gillnet section defect
2013	17	Pa	0	27	Sa	
2014	21	Pa	0	37	Sa	
2015	17	Pa	0	26	Sa	Gillnets changed from rings to danline.
2016	12	Pa	0	54 (76)	Sa	46,55,60,70,90. 22 ekstra stations to survey cod.
2017	30	Pa	0	36	Sa	
2018	20	SJ	0	42	Sa	
2019	17	HM	0	32	Sa	
2020	22	HM	0	40	Sa	
2021	0	-	0	44	Sa	
2022	25	TA	0	46	Sa	

Note: Research vessels: RV Paamiut (Pa), RV Adolf Jensen (AJ), RV Sanna (Sa), CV Sjurdaberg (Sj), CV Helga Maria (HM), Chartered vessel (Ch), R/V Tarajoq (TA).

**Table 2.** Number of Greenland halibut otoliths collected in the surveys.

Year	Area	Survey number of Otoliths			Aged	Method
		Gillnet	Trawl	Total		
2003	Disko Bay	?	212	212		Dried whole
2004	Disko Bay	?	120	120		Dried whole
2005	Disko Bay	442	95	537		Dried whole
2006	Disko Bay	207	47	254		Dried whole
2007	Disko Bay	285	67	352	59	Frozen whole
2008	Disko Bay	0	307	307	59	Frozen whole
2009	Disko Bay	0	527	527	476	mixed
2010	Disko Bay	363	347	710	91	Frozen image
2011	Disko Bay	344	565	909	0	Frozen image
2012	Disko Bay	320	194	514	78	Frozen image
2013	Disko Bay	264	300	564	235	Frozen image
2014	Disko Bay	343	286	629	53	Frozen image
2015	Disko Bay	233	290	523	125	Frozen image
2016	Disko Bay	347	431	778	345	Frozen image
2017	Disko Bay	346	467	813	179	Frozen image
2018	Disko Bay	741	472	1213	212	Frozen image
2019	Disko Bay	364	371	735	343	Frozen image
2020	Disko Bay	372	401	773	750	Frozen image
2021	Disko Bay	465	0	465	440	Frozen image
2022	Disko Bay	747	518	1265	289	Frozen image

**Table 3.** Greenland halibut *Reinhardtius hippoglossoides* abundance and biomass indices from the GINR shrimp and fish trawl survey.

Year	Stations	Abundance			Biomass		
		skjarevoy	cosmos		skjarevoy	cosmos	
		Million	Million	CI	1.000 tonnes	1.000 tonnes	
1992	39	96.73		27	4.992		22
1993	31	33.96		28	2.507		27
1994	27	62.96		22	3.598		26
1995	33	89.41		32	5.786		51
1996	33	102.5		25	8.593		22
1997	34	112.1		22	6.456		29
1998	33	209.6		29	11.874		35
1999	34	95.36		35	8.06		44
2000	23	172.8		30	9.537		30
2001	23	223.7		28	10.161		32
2002	22	148.1		38	9.07		40
2003	19	227		36	16.556		28
2004	14	199.1		34	28.229		36
Gear change				CV			CV
2005	16		186.5255	12		22.58046	12
2006	21		96.92568	11		20.24643	12
2007	18		128.4625	13		13.13739	18
2008	16		64.61175	12		16.42221	13
2009	24		72.86297	8		19.90215	20
2010	25		123.3412	9		17.55925	11
2011	26		230.7465	16		23.97662	10
2012	21		105.8918	8		16.16816	7
2013	17		187.9086	16		15.10297	9
2014	21		103.7475	15		11.46336	12
2015	17		84.0865	11		13.17978	15
2016	12		115.073	16		11.77193	12
2017	30		113.2452	14		12.04002	13
2018	20		95.34112	15		17.35497	11
2019	17		81.68444	16		11.6016	12
2020	22		138.2684	8		15.13024	9
2021							
2022	25		125.999957	9		21.05213	9



**Table 4.** CPUE and NPUE from the Gillnet survey in the Disko Bay.

Year	Number of stations	CPUE	SE	NPUE	SE	remark
2001	8	18.08	4.35	19.96	5.38	
2002	55	9.44	1.68	8.97	1.35	
2003	58	12.94	4.98	14.55	5.24	
2004	51	14.86	2.76	18.94	4.16	
2005	47	16.32	3.12	17.88	2.45	
2006	44	8.89	2.46	10.3	2.38	
2007	30	7.12	1.97	8.36	2.04	
2008	35	14.8	3.88	18.6	4.76	
2009	0		0		0	No survey
2010	48	14.01	2.83	15.78	3.72	
2011	51	25.1	2.83	24.96	3.72	
2012	41		0		0	Defect 60 mm
2013	27	12.53	2.38	13.53	2.86	
2014	37	12.42	2.5	13.69	2.34	
2015	25	10.11	1.78	10.48	1.57	
2016	52	9.69	1.01	10.14	0.97	
2017	36	9.14	1.21	10.61	1.63	
2018	42	10.91	1.41	16.34	1.66	
2019	32	16.44	3.43	19.84	3.62	
2020	40	18.86	1.79	24.84	1.94	
2021	44	27.4	3.04	35.56	3.04	
2022	46	12.87	1.34	18.7	1.56	

**Table 5.** Catch At Age table for the gillnet survey. (index=Number/100 hrs)

Year	Index val	Age2	Age3	Age4	Age5	Age6	Age7	Age8	Age9	Age10	Age11	Age12	Age13	Age14	Age15	Age16
2012	157	0	1	24	41	34	26	23	4	1	2	0	0	0	0	0
2013	225	0	7	41	85	48	27	13	4	1	0	0	0	0	0	0
2014	228	0	0	7	15	68	48	68	19	3	0	0	0	0	0	0
2015	168	0	3	15	47	41	30	20	7	4	0	0	0	0	0	0
2016	163	0	2	25	47	42	23	14	5	2	0	0	0	0	0	0
2017	177	1	3	3	28	45	42	29	14	11	0	1	0	0	0	0
2018	272	7	28	66	55	46	36	20	7	3	3	0	0	0	0	0
2019	331	1	2	33	127	92	42	19	9	2	1	2	0	0	0	1
2020	414	2	26	144	169	51	17	3	1	0	0	0	0	0	0	0
2021	593	1	14	125	251	149	42	8	4	1	0	0	0	0	0	0
2022	312	0	0	1	12	79	155	44	13	5	1	0	0	0	0	0


**Table 6.** modelled Length-Weight relationship for Greenland halibut.

Year	Area	Number fish	Log a	b	R <sup>2</sup>
2008	Disko Bay	578	-12.616	3.244	0.989
2009	Disko Bay	827	-12.516	3.224	0.997
2010	Disko Bay	348	-12.687	3.270	0.994
2011	Disko Bay	563	-12.595	3.246	0.990
2012	Disko Bay	513	-12.541	3.235	0.987
2013	Disko Bay	564	-12.479	3.212	0.995
2014	Disko Bay	628	-12.275	3.144	0.992
2015	Disko Bay	523	-12.515	3.221	0.995
2016	Disko Bay	778	-12.626	3.256	0.996
2017	Disko Bay	813	-12.344	3.189	0.986
2018	Disko Bay	1210	-12.557	3.241	0.994
2019	Disko Bay	733	-12.589	3.249	0.995
2020	Disko Bay	771	-12.584	3.252	0.996
2021	Disko Bay	461	-12.949	3.345	0.988
2022	Disko Bay	1258	-12.361	3.176	0.992

Weight = exp(log (a)) \* Length<sup>b</sup>



**Figure 1.** Map of the Disko Bay.



**Figure 2.** Trawl survey (SFW) in the Disko Bay: Abundance (left) and biomass (right). (2021 no survey)






Figure 3. Greenland halibut trawl survey length frequencies (SFW).







**Figure 4.** Abundance distribution (individuals/Km<sup>2</sup>) from the Disko Bay part of the GINR Shrimp and fish survey




**Figure 5.** Biomass distribution (Kg/Km<sup>2</sup>) from the Disko Bay part of the GINR Shrimp and fish survey




**Figure 6.** Disko Bay gillnet survey NPUE (right) of Greenland halibut (all sizes) combined with SFW trawl survey abundance estimate of Greenland halibut sizes 35-100 cm and gillnet survey CPUE (left).




**Figure 7.** Disko bay gillnet survey: Observed LF (N/100hr) for Greenland halibut.



**Figure 8.** Gillnet survey in Disko Bay by year. Distribution of NPUE (Number/gillnet/100 hrs).



**Figure 9.** Disko bay survey CAA bubble plot.



**Figure 10.** Disko bay survey MWA (Mean-Weight-At-Age) Greenland halibut.






Figure 11. Disko bay gillnet survey: Observed LF (N/100hr) for cod (right).