



Serial No. N7412

NAFO SCS Doc. 23/14REV

SCIENTIFIC COUNCIL MEETING - JUNE 2023

Denmark/Greenland Research Report for 2022

Compiled by

Adriana Nogueira and Ramus Nygaard

Greenland Institute of Natural Resources
P.P. Box 570, DK-3900 Nuuk, Greenland

This report presents information on catch statistics from the commercial Greenland fishery in 2022 at West Greenland. Catches for main species by NAFO division based on STATLANT 21A are presented in Table 1. Information on length frequency samples and biological samples taken by technicians from the Greenland Institute of Natural Resources (GINR) or by fishermen working onboard the commercial vessels, from Greenland halibut is also presented. Length distribution of the Greenland halibut samples from the catches offshore are in Tables 3. A total of 158 length samples were taken, and 939 564 individuals, including Greenland halibut, cod, roundnose grenadier were measured, in NAFO Div. 1A to 1F. A total of 504 otoliths were collected from Greenland halibut and Atlantic cod Div. 1A to 1F (Tables 4). Length distribution from Greenland halibut in 1AB and 1CD are presented in Table 5 and 6.

Furthermore, the report gives a brief overview of the research carried out by the GINR (Figure 1). For further information on GINR survey activities planned in 2023, visit www.natur.gl. For future research activities, education, collaboration opportunities, infrastructure, logistics, and much more, visit Isaaffik – the Arctic gateway www.isaaffik.org.

WEST GREENLAND (NAFO SUBAREA 1)**A. Status of the fisheries**

In 2022, Greenlandic commercial vessels were not engaged in fishery in other NAFO subareas than Subarea 1.

Provisional STATLANT 21A statistics for the fisheries are presented in Table 1.

1. Shrimp

The shrimp stock off West Greenland is distributed mainly in NAFO Subarea 1 (Greenland EEZ), but a small part of the habitat, and of the stock, intrudes into the eastern edge of Div. 0A (Canadian



EEZ). Canada has defined 'Shrimp Fishing Area 1' (Canadian SFA1), to be the part of Div. 0A located east of 60°30'W, i.e. east of the deepest water in this part of Davis Strait.

At the West coast of Greenland Northern shrimp is found mainly at depths between 150 and 600 m. The stock is assessed as a single population. The Greenland fishery exploits the stock in Subarea 1 (Div. 1A– 1F). The Canadian fishery has been limited to Div. 0A (SFA1)

Four fleets, one from Canada and three from Greenland (Kongelige Grønlandske Handel (KGH) fleet fishing from 1976 to 1990, the offshore fleet and coastal fleet) have participated in the fishery since the late 1970s. The Canadian fleet and the Greenland offshore fleet (> 75 GRT) have been restricted by areas and quotas since 1977. The fishery by the Greenland coastal fleet (< 75 GRT) was unrestricted until 1997 when quota regulation was imposed. Mesh size is at least 44 mm in Greenland, 40 mm in Canada. Sorting grids to reduce bycatch of fish are required in both of the Greenland fleets and in the Canadian fleet. Discarding of shrimps is prohibited.

Greenland requires that logbooks should record catch live weight. A former allowance for crushed and broken shrimps in reckoning quota draw-downs was abolished in 2011 to bring the total catch live weight into closer agreement with the enacted TAC.

Catches of shrimp gradually increased throughout the 1980s and 1990s and reached a level around 157000 t by 2005-2008 but have since decreased to 72 256 t in 2015. Since 2016 the catches have been increasing in conjunction with increasing TACs.

2. Greenland halibut

The stocks of Greenland halibut in the North West Atlantic are assessed in several management units. Greenland halibut in East Greenland (ICES 14) is considered to be a part of a stock also distributed in Icelandic and Faroese waters. Greenland halibut in the Baffin Bay and the Davis Strait, (NAFO SA 0 + 1 offshore) is assessed as one stock while the inshore stock in NAFO Div. 1A-F are considered isolated from the offshore stock and assessed by fjord area. In 1994 analysis of tagging and other biological information resulted in the creation of separate management areas for inshore Div. 1A (Disko Bay, Uummannaq and Upernavik districts). In 2020 the inshore Div.1B-F were decided also to be separated in 3 other different management units 1BC, 1D and 1EF.

In 2022, total catches of Greenland halibut (*Reinhardtius hippoglossoides*) in NAFO Subarea 1 Increased to 42 288 t, of which 17 669t were taken offshore by large vessels, and 24 419 t were landed from small boats operating inshore in the fjords from South Greenland to Qaanaaq.

The offshore catches were mainly taken by trawlers at the traditional fishing grounds in Div.1CD and west of the Disko Island in division 1A, while the inshore catches are from sea-ice, small vessels, and open boats using gillnets and longlines.

Greenland vessels operating offshore caught 13 629t (8052 t in division 1AB + 5577 t in division 1CD) and other nations caught 4240 t (542 t in division 1AB + 3698 t in division 1CD). Inshore landings in

Div. 1A were distributed from the Qaanaaq fjord (127t), the fjords near Upernavik (7738 t), the Uummannaq fjord (9007 t) and the Disko Bay (10 324 t). In the fjords south of West Greenland (1B-F) catches increased 1 465t of which 689 t were taken inshore in 1D. Trawl fishery is banned inshore, with the exception of shrimp trawl fishery in the Disko Bay and a small area inshore in division 1B. Sorting grids in the shrimp fishery have been mandatory offshore since 2002 with a sorting grid dispensation given to the smaller shrimp vessels operating inshore until 2011.

3. Cod

Cod (*Gadus morhua*) found in Greenland is a mixture of four separate “stocks” that are defined by their spawning areas: I) offshore West Greenland waters; II) West Greenland fjords cod III) offshore East Greenland and offshore Icelandic waters and IV) inshore Icelandic waters (Therkildsen et al. 2013). Therkildsen et al. (2013) showed a relatively stable spatial and temporal distribution of these spawning stocks during actual spawning events, but the proportional contribution of the different components to commercial and survey catches in different areas, seasons and years and the associated variation is unclear. However, Icelandic inputs are believed to have been responsible for the previous large year classes in Greenland (i.e. 1984 and 2003). A proportion of these cod returns to Iceland when reaching maturity. Previously the stocks have been assessed together. From 2012 the inshore component (West Greenland, NAFO Subarea 1) was assessed separately from all offshore components (Therkildsen, 2013). From 2016 the offshore components have been assessed separately with the West Greenland offshore component being comprised in the offshore area corresponding to NAFO subdivisions 1A, 1B, 1C, 1D, and 1E. The East Greenland offshore component is comprised in the offshore area corresponding to NAFO subdivision 1F in South West Greenland and East Greenland (ICES subarea 14b). The stocks are assessed by the ICES North-Western Working Group (NWWG), see ICES (2022), and ACFM (2022) report. Total catches inshore and offshore 1A-F in 2022 were 14 604 t.

5. Capelin

The capelin (*Mallotus villosus*) fishery in West Greenland is carried out inshore and in the spawning season only (May-July). Only part of the catches are reported, as capelin are used directly by fishermen for bait and dog food during the capelin season. Reported catches of capelin amounted to 247 t in 2022 and comprise a mixture of factory-landed capelin (326 t) for bait, human and animal consumption landed from small open boats mostly, and logbook reported bycatch in other fisheries.

6. Redfish

Two species of redfish of commercial interest occur off West Greenland inshore and offshore, golden redfish (*Sebastes norvegicus*), and deep-sea redfish (*Sebastes mentella*). Relationships to other North Atlantic redfish stocks are unclear, but the nearest stocks are the demersal and pelagic stocks in East Greenland and the Irminger Sea. Redfish catches in West Greenland are reported as redfish (unspecified, mainly bycatch), golden redfish and beaked redfish (deep-sea redfish).

Demersal redfish

In 2021, logbook reported catch and bycatch of redfish in the Greenlandic fleet targeting shrimp increased to 258 t. which fits well with the increasing recruitment observed in surveys. Inshore reported factory landings of commercially sized redfish amounted to 177t. Golden redfish (REG) landed to factories amounted to 12 t and 3 tonnes were reported as bycatch offshore from Greenland vessels.

Pelagic redfish

The aggregations of pelagic redfish *S. mentella* found in the NAFO Convention Area likely belong to the same stock of pelagic redfish from the Irminger Sea. The stock is assessed by ICES (ICES-NWWG report 2019), and the assessment covers the pelagic redfish in ICES Divisions 5a, 5b, and 14 and in the NAFO Div. 1F, 2H and 2J. The pelagic fishery on *S. mentella* in NAFO Div. 1F started in 1999 and from 2000 - 2009, significant catches with up to 20% of total catches as in 2003 were taken in NAFO Divisions 1F outside Greenland EEZ and 2J. In 2013, 3113 t were taken in the NAFO 1F, but since then no catches have been reported from 1F.

7. Grenadiers

There are two species of grenadiers of commercial interest in Greenland, roundnose grenadier (*Coryphaenoides rupestris*) and roughead grenadier (*Macrourus berglax*). Grenadiers are mainly taken as a bycatch in the Greenland halibut fishery inshore and offshore. In 2022, 10 t of roundnose grenadier were reported from offshore vessels targeting Greenland halibut. Less than 1 t were reported as bycatch offshore or landed to factories inshore.

8. Snow Crab

Snow crab (*Chionoecetes opilio*) is distributed along the west coast of Greenland from division 1A to 1F. The fishery is conducted only by Greenland vessels. From 2005-2020, the crab resource has been managed in 6 areas (from North to South: Upernavik, Uummannaq-Disko Bay, Sisimiut, Maniitsoq-Kangaamiut, Nuuk-Paamiut, and Narsaq-Qaqortoq). From 2020, the management areas of Sisimiut and Nuuk – Paamiut were furthermore divided in inshore as well as an offshore part, separated by the 3-nautical mile line. The fishing fleet is dominated by small vessels (less than 75 GRT), which have exclusive rights for fishing inshore as well as offshore. No large vessels (greater than 75 GRT), which only was allowed to fish offshore, have been in the fishery since 2005. Total allowable catch (TAC) restrictions have been imposed since 1995, but have only limited the catch in some years and management areas since 2004.

The number of vessels with licenses to participate in the snow crab fishery increased from 1999 to 2002 from approximately 120 vessels to 392 vessels. Since then, the number of both of vessels have decreased substantially as the abundance of the resource has also declined. Since 2008, approximately less than 40 vessels have been active in the snow crab fishery.

The total catch in NAFO Subarea 1 peaked in 2001 with approximately 15.100 t. From 2001 to 2011

total landings decreased markedly to 2,000 t. In the subsequent years total annual catches fluctuated at around this level, but increased again from 2017 and amounted approximately 3000 tons in 2021 (table 1). Most of the landings are based on the fishery in the management areas of Nuuk-Paamiut, Disko Bay-Uummannaq, and Sisimiut. In 2022, they are not reported in the STATLANT.

9. Wolffish

There are three species of wolffish in subarea 1, Atlantic wolffish (*Anarhichas lupus*), spotted wolffish (*Anarhichas minor*), and Northern wolffish (*Anarhichas denticulatus*). Only the two first are of commercial interest. In the past, these stocks have mainly been taken as a bycatch in the offshore fisheries targeting Cod, Greenland halibut, and shrimp, but occasionally are directly targeted. In 2022, 157 t of wolffish were caught in NAFO subarea 1.

10. Scallops

Total catches of Icelandic scallops (*Chlamys islandica*) in NAFO Subarea 1 increased to 569 t in 2021. All catches are taken in inshore areas in Div. 1A, 1B, 1C, and 1D. New fishing grounds near Sisimiut (1B) were found in 2003, and quotas for two new areas were introduced in 2004. Only one vessel is currently involved in this fishery.

11. Lumpfish

Total landings of lumpfish (*Cycloperus lumpus*) in NAFO Subarea 1 increased from 1.200 t in 2000 to almost 9.000 t in 2003 and remained at a high level until 2011 where catches reached 11.443 t. Catches are taken in inshore areas in Div. 1A, 1B, 1C, 1D, 1E, and 1F, with the majority being caught in 1D. The fishery is conducted over a short time period of one to two months, and over a vast coastline from 59° N to 72° N. Total landings of lumpfish (for roe) in 2021 was converted from roe to 4547 t whole weight. Lumpfish is not reported in the STATLANT21A

12. Greenland cod

Greenland cod (*Gadus macrocephalus*/ *Gadus ogac*) is mostly bycatch in other fisheries or direct fishery in fjords where it is more abundant. Greenland cod is mostly used for human consumption as dried or frozen fish for the local Greenlandic market. Total reported landings in 2022 amounted to 168 t of which the majority was landed to factories inshore in division 1B.

13. Arctic char

Arctic char (*Salvinus alpinus*) is taken in gillnets when returning to natal rivers during their annual feeding migrations in coastal areas. Factory landings were 10 t in 2021, and the production is mainly for the Greenlandic market. There is no reporting required for private fisheries, which presumably is considerably larger. Arctic char is not reported in STATLANT21A.

14. Atlantic halibut

Catches of Atlantic halibut (*Hippoglossus hippoglossus*) peaked at the beginning of the 1960s and the mid 1980s at a level of 600 to 1000 t per year. With the implementation of sorting grids in the shrimp fishery, bycatch of Atlantic Halibut is rare offshore. In 2021, few hundred kg were reported in logbooks and 12 t were landed in factories. There is no reporting required for Atlantic halibut landed and consumed locally. Atlantic halibut is not reported in STATLANT 21A.

15. Polar cod / Arctic cod

Reported catches of polar cod (*Boreagadus saida*) is mainly taken as bycatch in the shrimp fishery and directed fishery from small boats near glaciers and used directly for bait in the longline fishery targeting Greenland halibut. In 2021, 87 t were reported of which 46 were landed to factories mainly from small boats. Arctic cod is not reported in STATLANT 21A.

16. Large sharks

Large sharks (FAO: SHX/GSK) are without doubt exclusively Greenland sharks *somniosus microcephalus*. In 2022, 100 t of large sharks were reported. Shrimp-trawls are equipped with sorting grids, and no sharks were reported via shrimp logbooks.

17. Seacucumber

A trial fishery for orange-footed sea cucumber (*Cucumaria frondosa*) (FAO: CUX) was initiated in 2019, and total catches amounted to 106 t. In 2020, only 2 t were reported and no fishery occurred since 2021 due to lack of permits.

B. Special Research Studies

a. Hydrography Studies

Hydrographic conditions are yearly monitored at 10 hydrographic standard sections and stations in June/July across the continental shelf off West Greenland. Data are uploaded to the ICES database (Mortensen, 2022).

b. Trawl Surveys in Greenland

The Greenland shrimp and fish survey in NAFO SA 1 and ICES 14b:

Since 1988, an annual buffered stratified random trawl survey SFW (Shrimp and Fish West Greenland) has been conducted by the GINR on the West Greenlandic shelf between 59°15'N and 72°30'N from the 3-mile limit down to the 600 m and the Disko Bay. The main purpose of the survey is to evaluate the biomass and abundance of the Northern shrimp (*Pandalus*

borealis), but since 1992 data on fish species have been included. In 2007 this survey was expanded to include the East Greenland shelf to Dohrn Bank at 67°N. The survey was conducted with RV Paamiut from 1988 to 2017 and with the chartered commercial vessel Sjurdarberg in 2018 and Helga Maria, in 2019 and 2020. No survey was performed in 2021. In 2022, the survey was carried out with a new vessel owned by the GINR, R/V Tarajoq (Nygaard and Nogueira, 2023). Comparative analysis of the Cosmos gear performance and other analysis for the 4 vessels (R/V Paamiut 2005-2017, C/V Sjurdarberg 2018, C/V Helga Maria 2019-2020 and R/V Tarajoq 2022) that have been used to carry out the 1AF offshore shrimp and fish survey were performed (Nogueira et al. 2023). SC concluded that the 2018, 2019-2020 and 2022 surveys could be comparable to estimates from previous surveys conducted by R/V Paamiut and results could be used for NAFO Subarea 1 shrimp and groundfish assessments.

Greenland halibut trawl survey in NAFO 1CD:

A Greenland offshore trawl survey for Greenland halibut was initiated in 1997. The survey is a continuing of the joint Japanese/Greenland survey carried out in the period 1987-95. From 1997 the survey was conducted with RV Paamiut, using and Alfredo III gear, and covered NAFO Div. 1C and 1D between the 3 nm line and the midline against Canada at depths between 400 and 1500 m. The survey was conducted with the chartered commercial vessel Helga Maria, using also Alfredo III gear, in 2019. The survey was not conducted years 2020 and 2021. In 2022, a new survey series started with a new research vessel, R/V Tarajoq and a new gear, Bacalao 476 (Nogueira and Barcia-Estévez, 2023).

Greenland halibut trawl survey in NAFO1AB offshore:

In 2001 the survey area was expanded to include NAFO Div. 1B-1A (to 74°N), and in 2004 a survey was conducted in the northern part of the Baffin Bay (73°N-77°N) (Div. 1A) at depths down to 1500 m. In 2010 was conducted a survey in Div. 1A to 75°30' where 93 successful hauls were made. There has been no deep-sea survey in the area since then.

EU-Germany survey in ICES 14b and NAFO 1:

An annual stratified random trawl survey has been conducted by Germany since 1982 in West and East Greenland from 67°N in West Greenland to 67°N in East Greenland, covering the depth zone between 0-400 m. The main purpose of the survey is to evaluate the biomass and abundance of the Atlantic cod. The survey is conducted with the German vessel Walther Herwing III.

Nuuk Fjord trawl survey in NAFO 1D inshore:

In 2015, a trawl survey was initiated in the Nuuk fjord with the GINR research vessel RV Sanna. The purpose is to evaluate the local stock of Greenland halibut, shrimp and cod. The survey continued without interruption since 2017 and has already been completed for 2023 (Nygaard et al., 2023).

c) Gillnet surveys

Cod recruitment gillnet survey

A survey using gangs of gill nets with different mesh-sizes has been developed and used since 1985

with the objective of assessing the abundance of age 2 and age 3 cod in the inshore areas of NAFO subdivisions 1B and 1D (previously NAFO subdivision 1F was also surveyed).

Greenland halibut gillnet surveys in NAFO 1A inshore:

In 2001, the longline survey in the Disko Bay was changed to a gillnet survey. The survey normally covers four transects, and each gillnet set is compiled of 4 different nets with different mesh size (46, 55, 60, 70, and 90 mm halfmesh). From 2013 to 2016, the surveys in Uummannaq and Upernavik gradually changed from longline surveys to gillnet surveys (Nygaard, 2023a, Nygaard, 2023b, Nygaard, 2023c).

d) Snow crab surveys

The annual monitoring program (trap survey) was initiated in 1997 in Disko Bay (Div. 1A) and Sisimiut (Div. 1B). Since 2001, the offshore trap survey has been conducted, yearly, in more southern areas in West Greenland (Div. 1C and 1D) but has been canceled since 2010. The trap survey in Disko Bay has been canceled since 2018 and only the management area of Sisimiut is monitored on an annually basis. Large and small meshed conical traps are used. Snow crab are enumerated by sex, carapace width and carapace condition. The chelae height is measured in males and the abdomen width in females, respectively for maturity determination. Ovary contents, clutch weight, sperm load, and egg development stage in females are determined, and females were sampled in relation fecundity studies. The objective of both monitoring programs is to assess the abundance of snow crab in inshore and offshore waters of Greenland. Results from this survey are presented in the advising documents of the GINR.

A map of GINR research vessel stations for 2022 is given in fig 1.

e) GINR Research vessels

The Greenland institute of Natural Resources operates 2 larger multipurpose research vessels.

R/V Tarajoq - OYLD

RV Tarajoq is operational in 2022. Tarajoq was commissioned in 2021. The vessel is 61m long and 16m . BRT is 2 841t. Speed up to 14 knots and pull +50t. Tarajoq has room for 12 crew members and 20 scientists. Tarajoq is equipped for bottom trawling, pelagic trawling, with A-frame, scientific Simrad EK80 echosounders and a deep water multibeam and sub-bottom profiler and has various labs and a hydrographic hangar.

R/V Sanna - OZEK

RV Sanna was commissioned in 2012 and mainly operates inshore. Sanna is 32m long and 10 m wide. BRT is 458t. Speed is up to 11 knots. Sanna is capable of both bottom and pelagic trawling and

is equipped with A-frame, scientific Simrad EK80 echosounders and hull-mounted multibeam echosounder Reson SeaBat T50-R with extended range projector to achieve 1000 m water depth range.

Besides these larger vessels GINR operates various smaller vessels and boats.

f) Marine mammals

For yearly summaries of studies of marine mammals in Greenland, see the annual reports of the North Atlantic Marine Mammal Commission (NAMMCO).

References

- ICES. 2022. Report of the North-Western Working Group (NWWG), April–3 May, 2022, ICES HQ, Copenhagen, Denmark. ICES CM 2022/ACOM:09. 733 pp.
- Nogueira A. and Estevez-Barcia, D. 2023. Survey for Greenland Halibut in NAFO Divisions 1C-1D, 2022. *NAFO SCR Doc.*, No. 23/12, Serial No. N. 7399
- Nogueira, A, Christiansen H. and Hedges K,J. 2023. Comparison of vessels used for the the 1A-F west Greenland shelf surveys. NAFO SCR 23/032.
- Nygaard, R. and Nogueira, A. 2023. Biomass and Abundance of Demersal Fish Stocks off West and East Greenland estimated from the Greenland Institute of Natural resources (GINR) Shrimp and Fish Survey (SFW), 1990-2022. NAFO SCR Doc. 23/020.
- Nygaard R., Post S., Retzel A, Zinglensen K., Heilmann L., Jeremiassen S., Jeremiassen S., Mølgaard L. and Sethsen J. 2023. Biomass and Abundance of Demersal Fish Stocks in the Nuuk fjord and Ameralik fjord derived from the GINR Shrimp and fish inshore (SFI) survey. NAFO SCR Doc. 23/013.
- Nygaard R. 2023a. Survey results from the Disko Bay Gillnet survey, NAFO Division 1A inshore. NAFO SCR Doc. 23/21
- Nygaard R. 2023b. Survey results from the Uummannaq gillnet survey in NAFO Division 1A inshore. NAFO SCR. Doc. 23/22.
- Nygaard R. 2023c. Survey results from the Upernavik Gillnet survey, NAFO Division 1A inshore. NAFO SCR Doc. 23/37
- Mortensen J., 2023 Report on hydrographic conditions off Southwest Greenland June/July 2022 *NAFO SCR Doc.*, No. 23/005, Serial No. N7387.
- Therkildsen, N.O., Hemmer-Hansen, J., Hedeholm, R.B., Wisz, M.S., Pampoulie, C., Meldrup, D., Bonanomi, S., Retzel, A., Olsen, S.M. & E. E., Nielsen. 2013. Spatiotemporal SNP analysis reveal pronounced biocomplexity at the northern range margin of Atlantic cod *Gadus morhua*. *Evolutionary Applications*. DOI 10.1111/eva. 12055

Table 1. Greenland Catches (tons) in NAFO Area in 2022 by species and Division, based on the STATLANT 21 A.

Code	Species	Common name	1A	1B	1C	1D	1E	1F	TOTAL
CAB	Anarhichas denticulatus	Northern wolffish	0.0	2.6	0.2	0.4	2.6	0.0	5.7
CAT	Anarhichas spp	Wolffishes	0.0	28.4	53.0	62.1	5.8	2.8	151.9
RNG	Coryphaenoides rupestris	Roundnose grenadier	0.0	4.8	0.4	1.7	3.9	0.0	10.9
COD	Gadus morhua	Atlantic cod	0.0	1542.7	2464.5	5643.3	4850.3	104.0	14604.7
GRC	Gadus ogac	Greenland cod	0.0	128.1	40.0	0.0	0.0	0.0	168.0
HAL	Hippoglossus hippoglossus	Atlantic halibut	0.0	0.0	0.0	6.9	1.4	0.0	8.3
CAP	Mallotus villosus	Capelin	0.0	221.9	1.7	0.9	10.9	11.7	247.1
PRA	Pandalus borealis	Northern prawn	17.0	39399.9	55876.2	5586.8	9387.2	5510.6	115777.8
SKA	Raja spp	Raja rays	0.0	27.6	1.9	1.3	3.8	0.0	34.7
GHL	Reinhardtius hippoglossoides	Greenland halibut	35100.7	64.0	585.1	2260.3	4084.1	193.9	42288.1
RED	Sebastes spp	Atlantic redfishes nei	0.0	67.5	126.2	23.1	37.5	4.6	258.9
GSK	Somniosus microcephalus	Greenland shark	0.0	53.8	2.6	7.4	36.5	0.0	100.3

Table 2. Samples and individuals sampled at the commercial fleet by species, gear, and NAFO Division in 2021.

Code	Species	Common name	Div.	Gear	Numb.ind	Sample Size
COD	Atlantic cod	Cod	1A	Trawl	44567	12
COD	Atlantic cod	Cod	1D	Fishing rods	795	4
COD	Atlantic cod	Cod	1D	Gillnet	168	1
COD	Atlantic cod	Cod	1D	Pound nets	1091	5
COD	Atlantic cod	Cod	1D	Trawl	771161	12
GHL	Reinhardtius hippoglossoides	Greenland halibut	1A	Gillnet	1352	10
GHL	Reinhardtius hippoglossoides	Greenland halibut	1A	Longline	5716	22
GHL	Reinhardtius hippoglossoides	Greenland halibut	1A	Pound nets	392	1
GHL	Reinhardtius hippoglossoides	Greenland halibut	1A	Trawl	9709	62
GHL	Reinhardtius hippoglossoides	Greenland halibut	1B	Trawl	150	1
GHL	Reinhardtius hippoglossoides	Greenland halibut	1C	Trawl	1213	7
GHL	Reinhardtius hippoglossoides	Greenland halibut	1D	Longline	168	1
GHL	Reinhardtius hippoglossoides	Greenland halibut	1D	Trawl	2748	18
GHL	Reinhardtius hippoglossoides	Greenland halibut	1F	Longline	161	1
RNG	Coryphaenoides rupestris	Roundnose grenadier	1A	Trawl	173	1
TOTAL					839564	158

Table 3. Biological samples collected in NAFO by species and Division in 2021:

Code	Species	Division	Otolith	Maturity
COD	<i>Gadus morhua</i>	1D	87	-
GHL	<i>Reinhardtius hippoglossoides</i>	1A	153	153
GHL	<i>Reinhardtius hippoglossoides</i>	1C	174	174
GHL	<i>Reinhardtius hippoglossoides</i>	1D	90	90
TOTAL			504	417

Table 4. Length distribution of Greenland halibut samples in 1AB offshore:

Length	3rdQ.	4rdQ.
62	86	70
63	62	81
64	45	54
65	40	40
66	46	53
67	39	38
68	26	19
69	34	23
70	27	21
71	18	19
72	9	17
73	10	9
74	12	10
75	11	3
76	5	6
77	9	7
78	6	8
79	4	3
80	4	4
81	4	3
82	3	4
83	1	4
84	3	2
85	1	2
86	1	3
87	1	5
88	0	1
89	0	1
90	0	1
91	0	0
92	1	0
93	1	0
94	0	1
95	0	1
96	1	0
Ind samples	3812	5532
Samples	25	36
Mean L (cm)	52.8	51.3

Table 5. Length distribution of Greenland halibut samples in 1CD offshore

Length	3rdQ.	4rdQ.
23	1	0
24	0	0
25	1	0
26	0	0
27	1	0
28	1	0
29	1	0
30	1	3
31	3	1
32	0	3
33	2	4
34	2	4
35	1	2
36	4	6
37	4	3
38	4	5
39	5	9
40	10	18
41	50	16
42	12	17
43	21	21
44	20	27
45	22	46
46	28	46
47	43	63
48	41	70
49	51	70
50	51	101
51	80	78
52	84	124
53	78	129
54	104	169
55	90	154
56	78	141
57	87	120
58	70	75
59	45	69
60	62	81

Length	3rdQ.	4rdQ.
61	54	69
62	33	67
63	34	70
64	38	54
65	27	48
66	29	57
67	17	44
68	16	34
69	18	32
70	17	33
71	15	25
72	12	15
73	6	22
74	13	16
75	5	15
76	7	9
77	6	9
78	6	12
79	2	6
80	4	7
81	5	5
82	3	15
83	0	4
84	4	6
85	3	14
86	0	8
87	2	8
88	2	7
89	2	6
90	1	2
91	4	1
92	1	2
93	1	3
94	1	3
95	1	3
96	0	3
97	0	0
98	0	3
99	1	1
Ind samples	1548	2413
Samples	10	15
Mean L (cm)	55.9	57.5




Figure 1. Distribution of stations from GINR surveys completed in Subarea 1. Stations from hydrography samples taken onboard the Royal Danish Navy vessel Hdms Knud Rasmussen are not included.