

International Commission for



the Northwest Atlantic Fisheries

Serial No. 3829
(A.c.1)

ICNAF Summ.Doc. 76/VI/7
(also ICES C.M.1976/F:6
Demersal Fish
(Northern) Committee)

ANNUAL MEETING - JUNE 1976

Report of the North-Western Working Group¹

Charlottenlund, 8-12 March 1976

The Report of the Working Group has been reproduced in part only. Chapters A, D, and E, Tables 1-11 and 22-33, and Figures 1-8 and 10 reproduced here are pertinent to the Greenland-Iceland cod stock relationship which the Commission, at its June 1974 and again at its 1975 Annual Meetings, asked NEAFC and ICES to study with a view to ensuring adequate management.

¹ This Report has not yet been approved by the International Council for the Exploration of the Sea; it has, therefore, at present the status of an internal document and does not represent an advice given on behalf of the Council. The proviso that it shall not be cited without prior reference to the Council (General Secretary, ICES, Charlottenlund Slot, DK-2920 Charlottenlund, Denmark) should be strictly observed.

Contents

| | <u>Page</u> |
|---|-------------|
| A. INTRODUCTION | 1 |
| 1. Terms of Reference | 1 |
| 2. Participants | 1 |
| 3. Previous and Present Assessments | 1 |
| B. COD IN DIVISION Va - ICELAND GROUNDS | 2 |
| 4. Nominal Catches | 2 |
| 5. Spawning and Non-Spawning Fisheries | 2 |
| 6. Effort | 2 |
| 7. Age Composition of Landings | 3 |
| 8. Mean Weight by Age | 3 |
| 9. Virtual Population Analysis | 4 |
| 10. Stock Biomass | 5 |
| 11. Recruitment | 7 |
| C. HADDOCK IN DIVISION Va - ICELAND GROUNDS | 8 |
| 13. Nominal Catches | 8 |
| 13. Effort and Catch per Unit of Effort 1970-75 | 8 |
| 15. Catch in Numbers by Age Groups | 8 |
| 16. Mean Weight by Age | 8 |
| 17. Input Data to VPA | 9 |
| 18. Results of VPA | 9 |
| 19. Stock Size | 9 |
| 20. Yield Curves | 9 |
| 21. Biomass of Stock | 10 |
| 22. Catch Predictions | 10 |
| D. COD GREENLAND | 10 |
| 23. Nominal Catch (ICES Sub-area XIV and ICNAF Divs. 1E-1F) | 10 |
| 24. Effort | 12 |
| 25. Catch in Numbers by Age Groups | 12 |
| 26. Mean Weight by Age | 13 |
| 27. Natural Mortality and Emigration | 14 |
| 28. Input Data to VPAs of Cod at Greenland | 14 |
| 29. Results of the VPAs and Predictions of Stock Size and Catches for 1976-78 | 14 |
| E. INTERRELATIONSHIP BETWEEN THE COD STOCKS AT ICELAND AND GREENLAND | 15 |
| 30. Introduction | 15 |
| 31. Migration of Adult Fish from West to East Greenland and to Iceland | 15 |
| 32. Recruitment to the West Greenland Stock of Cod originating from East Greenland and Iceland | 17 |
| 33. Management Problems for Cod at Greenland | 18 |
| TABLES 1 - 21 | 19 |
| APPENDIX I, II and III (Iceland Haddock) | 39 |
| TABLES 22 - 33 | 42 |
| FIGURES 1 - 10 | 54 |

Report of the North-Western Working Group

A. INTRODUCTION

1. Terms of Reference

At the Council's Statutory Meeting in 1975 the following resolution was adopted (C.Res.1975/2:29):

"It was decided, that

- (i) the North-Western Working Group should meet at Charlottenlund from 19-23 January 1976 (postponed to 8-12 March) under the chairmanship of Mr J Møller Christensen in order to:
 - (a) investigate the interrelationship between the cod at East and West Greenland and adjacent waters, and
 - (b) report separately on the state of the stocks of cod and haddock in Icelandic and adjacent waters.
- (ii) ICNAF should be invited to participate in the discussions under Item (a), and that
- (iii) this report be made available to the STACRES of ICNAF."

2. Participants

| | |
|------------------------------------|---------------------|
| A C Burd | U.K. (England) |
| Sv. Aa. Horsted * | Denmark |
| J Jakobsson | Iceland |
| J S Joensen | Faroe Islands |
| B W Jones | U.K. (England) |
| R Jones | U.K. (Scotland) |
| P Kannevorff | Denmark |
| J Møller Jensen | Denmark |
| J Møller Christensen (Chairman) | Denmark |
| S A Schopka | Iceland |
| A Schumacher * | Germany, Fed.Rep.of |
| <u>Ø Ulltang</u> | Norway |

* also representing ICNAF.

3. Previous and Present Assessments

At its meeting in 1970 the North-Western Working Group made assessments of the stocks of cod and haddock in ICES Division Va (Iceland Grounds). It also made an estimate of the migration of mature cod from East Greenland to Iceland.

The assessments of the cod stocks in these areas were reviewed by the Joint ICES/ICNAF Working Group on Cod Stocks in the North Atlantic in 1972.

At the present meeting the North-Western Working Group made new assessments of the cod and haddock stock at Iceland Grounds (Sections B and C). The Group also made further analyses of the interrelationship between the cod stocks at Iceland and at Greenland (Section E) and assessed the cod stock at Greenland (ICES Subarea XIV and ICNAF Divisions 1E and 1F) (Section D).

21. Biomass of stock

Table 19 shows estimates of the biomass of the stock and of the spawning stock from 1962-75. The results show that there has been a significant decline for both stock components.

22. Catch predictions

Predictions have been made of haddock catches for the period 1976-78 (Table 21). The method of computation is illustrated in Appendix III. Input data consisted of:

- 1) numbers landed at each age in 1975;
- 2) mean weights at age, based on averages for the period 1971-75 (Table 14);
- 3) a natural mortality rate (assumed $M = 0.2$);
- 4) values of F at each age. Calculations were done using two of the sets of F -at-age for 1975 given in Table 15; and
- 5) estimates of year class strengths for the 1974-76 year classes as 2 year old fish.

Values of haddock year class strengths from the VPA results are given in Table 20 and for each of the three input sets of F used in these analyses. These show that the estimates of year class strength at age 2 years were effectively independent of the input F values for the year classes 1960-70. For these year classes the mean value was 64 million fish and this value has been used for the sets of predictions in Table 21.A.

A second sets of predictions (Table 21.B) were made assuming 30 million fish for the 1974-76 year class strength, this being the lowest year class strength observed in the 1960s.

For each of the assumptions made about the F values in 1975, catches are expected to decline in 1976 and 1977. Estimates for 1978 depend on the values assessed for the strengths of the 1974-76 year classes. It should be noted that the further ahead the forecasts are made, the more depend the predictions on estimates of the recruiting year class strength. For example, a large proportion of the predictions given for 1978 in Table 21 are due to the values adopted for strengths of the 1974-76 year classes.

In view of the relatively high variability of year class strengths in practice, the confidence limits for these estimates and for the 1978 estimates in particular, are likely to be large.

D. COD - GREENLAND

23. Nominal catch (ICES Sub-area XIV and ICNAF Divs. 1E-1F)

23.1 Data used

The catches of cod in Greenland waters are reported nationally through the STATLANT system to ICNAF and ICES for West Greenland (ICNAF Subarea 1) and East Greenland (ICES Sub-area XIV), respectively. The ICNAF Subarea 1 is further split into six divisions (Divs. 1A-1F) whereas no further breakdown of the ICES Sub-area XIV exists at present.

In its present report the North-Western Working Group has as far as cod is concerned confined itself to analyses of the stocks at Iceland, at East Greenland and off the southern part of West Greenland (ICNAF Divs. 1E-1F). The inclusion in the analyses of only part of the ICNAF Subarea 1 creates some difficulties since some countries have reported part of their catch or even their total catch at West Greenland as Div. 1NK, i.e. without a breakdown on statistical divisions. It has, therefore, been necessary to allocate such unspecified

catches by divisions. The allocation here adopted is the one used by the Greenland Fisheries Institute (Horsted, unpubl.), and which is also used in analyses by ICNAF (Horsted, ICNAF Res.Doc. 75/31). The allocation is made partly on various assumptions, e.g. that unspecified catches from one country are distributed like specified catches from the same country, and partly on observations on fishing activities at Greenland. A full list of the allocations and the principles followed is available in the Greenland Fisheries Institute, but is not given here.

In order to show the magnitude of the problem, the unspecified catches (Div. LNK) are given in Table 22 together with the total amount of these catches which is allocated to Divisions 1E and 1F and added to the specified Divs. 1E-1F catches to give the best estimate of the actual nominal catch from these divisions. The figures for which a part or the total amount of catch has been based upon allocation from Division LNK are marked with an asterisk in the table. It will be seen that of the annual totals for Divisions 1E-1F cod catches up to about 40% of the total have been allocated from unspecified catches, 1974 being the only year for which all catches were reported by divisions.

The nominal catches for the fisheries at East Greenland (Sub-area XIV) are readily available in ICES "Bulletin Statistique". For 1975, members of the Working Group supplied provisional data at the meeting. Sub-area XIV covers a wide area, and although the cod fisheries in that area are known to occur between Cape Farewell and the Dohrn Bank it is not possible to break catches down by smaller units. The problem of a probable break-down of Sub-area XIV was discussed briefly by the Working Group but referred to the ICES Statistics Committee.

23.2 Trends in catches

23.2.1 Nominal catches of cod in ICNAF Divisions 1E-1F, 1960-74

As explained in Section 23.1, the nominal catches for Divisions 1E-1F as set out in Table 22 contain part of some catches reported as West Greenland unspecified (ICNAF notation: Div. LNK).

In the course of the late 1960s the cod fisheries at West Greenland (ICNAF Subarea 1) had a tendency to concentrate more on the southern Divisions (Divs. 1E-1F) than previously, and by 1970 about half the West Greenland catch was taken in those Divisions. Whereas the overall Subarea 1 cod catches reached a maximum in 1962, the Divisions 1E-1F fishery obtained its highest catch in 1968. However, since then, this part of Subarea 1 has also faced the same drastic decline as the Subarea 1 fishery as a whole, and the relative importance of the Division has dropped again to about $\frac{1}{4}$ of the total of West Greenland (Table 25). The catch in Divisions 1E-1F by 1974 was only about 12% of the catch in the peak year 1968.

Catches for 1975 are not yet known by Division, but the overall Subarea 1 catch seems to have had a further small decline from 1974.

The fishery in Subarea 1 as a whole has been under quota regulation since 1974, but neither in 1974 nor in 1975 has the total allowable catch been taken. The TAC for 1976 is 46 thousand tons.

23.2.2 Nominal catches of cod off East Greenland (ICES Sub-area XIV) 1960-75

The fishery off East Greenland is almost entirely due to trawling, with a few nations participating, primarily the Federal Republic of Germany and Iceland. The target species are cod and redfish, and although fishing can be directed to one of these species the by-catch of the other species is normally so high that it seems proper to speak of a mixed fishery of the two species. Up to 1969 redfish made up the major part of the fishery but since 1970 cod is the predominant species.

In the period 1960-72 the total catch of cod in the area (Table 23) has fluctuated between 13 and 36 thousand tons (1960-72, mean: 22 100 tons), with 1964 and 1971 as the peak years (35 600 and 31 500 tons, respectively). A drastic decline in the catches has occurred after 1972 with a provisional figure for 1975 of only 3 400 tons or 15% of the 1960-72 level. This decline is closely combined with a decline in effort seen in Section 24.

23.2.3 Nominal catches of cod at East Greenland and off Southwest Greenland as a whole (ICES Sub-area XIV and ICNAF Divisions 1E-1F), 1960-74

The cod catches in ICES Sub-area XIV and ICNAF Divisions 1E-1F mentioned in the preceding sections are combined in Table 24. For the combined area the cod catches have fluctuated between 74 and 130 thousand tons in the period 1960-71, the mean for the period being 99 thousand tons. Peak years are 1963 and 1968, both with 130 thousand tons. A drastic decline is observed after 1971, and the 1974 catch is only about 20 thousand tons or 20% of the 1960-71 level.

24. Effort

24.1 Data used

Both ICES and ICNAF request countries to report fishing effort. For East Greenland (ICES Sub-area XIV) the effort figures as set up in Table 26 were obtained from the German research reports to ICNAF (by A Meyer). This effort is an effort directed partly to cod and partly to redfish or to both species combined. The catch per unit effort as a measure of cod abundance must, therefore, be taken with great reservation.

For ICNAF Divisions 1E-1F no attempt was made by the Working Group to set up a table of an overall effort for the area. Such an exercise would, of course, also contain the same problem of allocation as with the nominal catches.

24.2 Trends in effort

Due to the complexity of the fisheries at West Greenland and the problem of allocating unspecified catches no attempt has been made recently to obtain effort-unit figures for ICNAF Divisions 1E-1F separately.

As explained in para. 24.1 some effort figures can be given for the fisheries off East Greenland (Table 26). These clearly demonstrate a decrease of effort after 1972, so that the level of effort by 1974 is 1/4 - 1/5 of the high level in the mid-1960s. The catch-per-unit of effort figures vary considerably, being highest in 1971. The c.p.u.e. level in 1974 falls within the same range as the figures in the 1960s. However, due to the mixed nature of the fisheries, no definite conclusions are drawn from these c.p.u.e. figures, nor has it been considered appropriate to use these figures to obtain an overall effort for ICES Sub-area XIV and ICNAF Divisions 1E-1F combined. However, the low catch figures for Divisions 1E-1F in recent years do suggest that effort has declined also in these Divisions and hence also in the combined Sub-area XIV-Divisions 1E-1F area.

25. Catch in Numbers by Age Groups

25.1 ICNAF Divisions 1E-1F

The numbers by age groups for the cod catches in ICNAF Divisions 1E-1F for the period 1960-75 are given in Table 27. These figures are taken from ICNAF Res.Doc.75/31 (by Sv. Aa. Horsted) for the years 1965-73, and for the years 1974-75 they are preliminary estimated by Horsted. For the years prior to 1969 (including 1960-64) the basic material is submitted by the Federal Republic of Germany (Schumacher and Meyer, unpubl.), and adjusted to the total catches for Divisions 1E-1F as they occur after allocation of unspecified West Greenland catches (see para.23.1). The German method of raising samples to catches has generally been based on the observed weight of the total sample,

whereas Horsted's figures are based on samples for which a total weight has been calculated by applying mean weights for each age group. This latter method may lead to more heavily biased figures than the former, but the method has been the only possible one since few samples with observed total weight exist for recent years. For the years 1974 and 1975 it has even been necessary to use samples from catches containing a mixture of fish from various divisions. The figures given for 1974 and 1975 are, therefore, very uncertain, although the 1968 year class has the expected very strong predominance.

25.2 East Greenland (ICES Sub-area XIV)

The numbers by age group for the cod catches off East Greenland as given in Table 28 are based on figures for the German (Fed.Rep. of) catches made available to the Working Group by A Meyer. The raising of numbers in samples to numbers in catches is based on observed total weight of the samples. The figures supplied by A Meyer have been raised to total Sub-area XIV cod catches by the Working Group. Since German catches account for the major part of the Sub-area XIV catch, the possible bias by this latter raising seems to be very small. However, due to the wide statistical area, it is not clear whether great variation in catch composition exists between the northern part (the Dohrn Bank) and the southern part (close to Cape Farewell) nor to judge whether the whole area, if fished, is covered by the sampling.

25.3 ICES Sub-area XIV plus ICNAF Divisions 1E-1F

The numbers by age group for the overall southwest and East Greenland cod catches as given in Table 29 are simple sums of figures given in Tables 27 and 28.

26. Mean Weight by Age

The mean weight by age for Greenland cod is known to vary considerably between years and between year classes. In the present analyses the following values taken from ICNAF Res.Doc.75/31 were used:

| <u>Age</u> | <u>Mean Weight (kg)</u> |
|------------|-------------------------|
| 3 | 0.65 |
| 4 | 0.99 |
| 5 | 1.68 |
| 6 | 2.77 |
| 7 | 3.84 |
| 8 | 4.72 |
| 9 | 5.34 |
| 10 | 5.34 |
| 11 | 5.48 |
| 12 | 5.39 |
| 13 | 8.70 |
| 14+ | 10.00 |

These figures were checked on the only sample available from Division 1E at present (a length sample from U.K. supplied to the ICNAF Assessment Meeting, April 1976 and broken down in age groups by means of a Danish age/length key for Divisions 1C-1E, 1975). The same sample was converted to weight by means of German length/weight data (A Meyer, ICNAF Res. Doc.66/18). This exercise showed that the weight figures as given above correspond reasonably well both with the weight obtained by German data and with the actual observed total weight for the U.K. sample.

27. Natural Mortality and Emigration

Natural mortality has been taken as $M = 0.20$, the value used throughout all previous analyses of Greenland cod. However, apart from this mortality (and the fishing mortality) the VPA analyses should also take into account the "mortality" due to emigration. The emigration has been adopted as being 25% annually for mature cod (see para. 30.2). This corresponds to a coefficient (instantaneous rate) of 0.29. Taking the age of emigration as knife-edge at age 7, the VPA analysis for the combined stocks in ICES Sub-area XIV and ICNAF Divisions 1E-1F has been made with a value of $M = 0.20$ for age groups to and including six years. From seven years onwards the M value is taken as 0.49, treating emigration as a component of the natural mortality.

28. Input Data to Virtual Population Analyses of Cod at Greenland

The basic input figures for VPA analyses are the catch in numbers and the mortality rates. Nominal catches and catch by numbers have already been considered in the previous Sections, and so have the natural mortality and the emigration parameter. For estimating forecasts, figures for mean weight by age are needed. These are also dealt with above.

The most critical input is the terminal figure for fishing mortality rate, F . In the analysis carried out it has been assumed that F in 1975 is the same for East Greenland as for ICNAF Divisions 1E-1F. At the same time it has been taken into account that catches and effort in 1975 are very much lower than in the years prior to 1974. The actual 1975 catches seem to be close to those predicted (for Divisions 1E-1F) in forecasts by an F value of 0.20 (ICNAF Res.Doc.75/31). A value of 0.22 was then chosen for the analyses, but other values of the same order might as well have been considered.

29. Results of the VPA and Predictions of Stock Size and Catches for 1976-78

The VPA analyses (Tables 30 and 31) carried out for the ICES Sub-area XIV and ICNAF Divisions 1E-1F combined show, as expected from the fisheries themselves, that there has been an overall decline in the stock over the last five years. Taking only the spawning stock, i.e. cod of age 7 and older, the numbers (in millions) at the beginning of each year are as follows:

| | | | | | | |
|------------------|-------------|-------------|-------------|-------------|-------------|-------------|
| Year | <u>1960</u> | <u>1961</u> | <u>1962</u> | <u>1963</u> | <u>1964</u> | <u>1965</u> |
| Nos. x 10^{-6} | 161.2 | 101.8 | 65.3 | 91.5 | 89.4 | 70.3 |
| Year | <u>1966</u> | <u>1967</u> | <u>1968</u> | <u>1969</u> | <u>1970</u> | <u>1971</u> |
| Nos. x 10^{-6} | 39.7 | 45.2 | 82.0 | 76.0 | 96.4 | 64.7 |
| Year | <u>1972</u> | <u>1973</u> | <u>1974</u> | <u>1975</u> | | |
| Nos. x 10^{-6} | 28.3 | 13.2 | 7.3 | 21.4 | | |

This reflects the very poor general recruitment to the stock since year class 1963 recruited. The only year class of average strength since then is the 1968 year class. The recruitment of this year class to the spawning stock may have led to some increase in spawning stock in 1975. If no good year classes enter the stock in the next few years, a further decline is to be expected.

There seems to be evidence that the 1973 year class is of some importance. Rather arbitrarily it is here judged to be somewhat stronger than other year classes since 1963 and about 1/3 of the 1963 year class, i.e. in round

figures about 70 million individuals by the age of 3. The following 1974 year class has not shown any signs of importance and is set at 10 million by age 3.

On the basis of this a prediction of stock size and catches has been made for the area considered for 1976-77, using values of F corresponding to the input F in 1975 in the VPA ($F = 0.22$) and $F_{0.1} = 0.45$ (ICNAF Res.Doc. 75/31). The results are set out in Table 32.

The predictions show that for both F values a slight improvement in the total stock could be expected from 1975 to 1978. However, this improvement is mainly due to the above-mentioned optimistic judgment of the incoming 1973 year class. Figures in brackets in the table reflect that part of the predicted catches and stock which is dependent on the incoming year classes 1973 and 1974. If the 1973 year class is overestimated, catches and stock size will remain at the present very low level.

E. INTERRELATIONSHIP BETWEEN THE COD STOCKS AT ICELAND AND AT GREENLAND

30. Introduction

Throughout the period when investigations of cod in Greenland waters have been made, i.e. since the 1920s, it has been known that part of the stock of cod at West Greenland migrates to East Greenland and Iceland when reaching maturity. This has been demonstrated mainly by tagging experiments at Greenland, but also other studies confirm this migration.

As would be expected the migration to East Greenland and Iceland has been most pronounced for cod tagged in the southernmost part of West Greenland, i.e. ICNAF Divisions 1E and 1F. Tagging off East Greenland has shown a considerable migration from these waters to Iceland but only a small-scale migration to West Greenland. Tagging at Iceland has revealed a negligible number of recaptures at Greenland thus confirming that once the cod have migrated from Greenland to Iceland they will remain at Iceland.

However, the interrelationship between the stocks is not only a matter of adult cod migrating and mixing but also a matter of recruitment of young cod to one area originating from spawning in another area. As far as this question is concerned, there seems to be some feed-back of fry from East Greenland to West Greenland and from Iceland to East Greenland, and possibly even to West Greenland.

These two separate aspects of the interrelationship between cod at Greenland and Iceland are described in further details in the following.

31. Migration of Adult Fish from West to East Greenland and to Iceland

Although it has been known that cod from Greenland waters contribute to the fisheries at Iceland no quantitative estimates of this contribution have been made until the North-Western Working Group tried to carry out such analyses at its last meeting in 1970. At that meeting the Working Group based its analyses partly on tagging experiments and partly on analyses of stock size and composition of stock and catches at Iceland and Greenland.

31.1 Estimates from tagging experiments

Based upon tagging experiments at Greenland the Working Group in 1970 concluded that the actual overall proportion of mature fish at East Greenland and in the southern part of West Greenland (ICNAF Divisions 1E-1F) emigrating to Iceland was about 25% per year.

Since then only few fish have been tagged at Greenland. Danish tagging experiments at West Greenland in the years 1966-72 were presented to the Working Group at its present meeting. They reconfirmed that from the northern divisions at West Greenland (Divisions 1B-1D) the migration to East Greenland and Iceland is insignificant, whereas tagged cod released in the

southern part of the area (Divisions 1E-1F) revealed several recaptures at East Greenland and at Iceland. Considering only fish that were 70 cm or bigger at the time of tagging, the total recaptures from the 1966-72 experiments in Divisions 1E-1F amount to 7.6% (25 recaptures, 329 fish tagged). 44% of the recaptures came from East Greenland or Iceland. The overall recapture rate from these experiments is lower than in previous experiments, but the decrease is mainly due to a lower recapture rate at West Greenland than in previous experiments, although also the recapture rate at Iceland and at East Greenland has decreased somewhat. However, the material is so limited and fishermen's reporting rate of tags so uncertain that the Working Group did not find itself in a position to change the conclusions from the meeting in 1970.

From Icelandic tagging experiments at East Greenland in the years 1971-74, only 2% has been returned, probably due to a high tagging mortality. 2/3 of the recaptures came from East Greenland and 1/3 from Icelandic waters. Again, these experiments do not allow any revision of former conclusions.

31.2 Estimate of emigrants from Greenland to Iceland

Since no new information on the number of cod of age 7 and older emigrating from Greenland to Iceland is available, the percentage of emigrants (25% annually) given in the previous report of the Working Group was used. This figure corresponds to an instantaneous emigration rate of 0.29, which was applied to the number of cod from age 7 and onwards in each year and age group derived from VPA (using the parameters outlined in Sections 27 and 28) for ICNAF Divisions 1E-1F and ICES Sub-area XIV combined. In estimating the number of cod emigrating from Greenland, F and M values have also been taken into account (see Section 27).

The annual contribution of Greenland cod to the Icelandic spawning stock (Table 33 and Figure 10) varies according to the size of the year classes and F values at Greenland, ranging from 34.7×10^6 cod in 1960 to 1.3×10^6 in 1974. From 1971 onwards there was a steady decline of emigration from Greenland from 12.1×10^6 in 1971 to 1.3×10^6 in 1974, when the very poor year classes 1965, 1966 and 1967 entered the spawning stock. In 1975, when the about average 1968 year class was expected to emigrate, the number increased slightly to 4.4×10^6 . The average over the period 1960-69 of 7 year old fish (8.0×10^6 fish) is of the same order as the estimate given in the previous report (7.3×10^6).

31.3 Some observations on the use of VPA for the Icelandic/Greenland cod stock

The Group discussed the difficulties of obtaining valid estimates of F and stock size from VPA when dealing with two stocks with interchange between them.

A VPA using only catches made at Iceland would tend to overestimate stock sizes at Iceland, especially among the younger age groups. This is because these estimates might include a proportion of fish that had commenced life in Greenland waters. A VPA using only catches made at East Greenland might underestimate stock sizes at East Greenland if no account has been taken of fish that commenced life at East Greenland but were caught at Iceland. To take account of this, the effective value of M on the older age groups could be increased to take account of an instantaneous coefficient of emigration, and the result of a trial made in this way is given in Tables 30 and 31.

A VPA using catches from Iceland and East Greenland would be useful since this should provide estimates of total stock sizes but without any indication of how this should be distributed between the two areas.

For all the VPAs it was recognised that values of F were liable to be biased. All assessments depending on VPA Fs were, therefore, regarded as provisional and subject to revisions.

It was recommended that further work be done on a simulation of the Iceland/Greenland situation with a view to obtaining better estimates of F, stock sizes and coefficient of emigration from Greenland to Iceland.

32. Recruitment to the West Greenland Stock of Cod Originating from East Greenland and Iceland

32.1 Distribution of cod at Greenland

The recruitment to the cod stock off West Greenland is dependent on fluctuations in the environment not only at West Greenland, but also at East Greenland and Iceland. These fluctuations in the environment lead to fluctuations in the strength of the cod year classes.

The distribution of cod at West Greenland depends on whether the year classes originate from West Greenland or from East Greenland-Iceland. A year class originating from West Greenland seems to come from the spawning area in the northern part of ICNAF Division 1E and Division 1D. The main nursery grounds are in ICNAF Divisions 1B-1D. Seasonal spawning/feeding migrations occur between various areas.

A year class originating from East Greenland-Iceland has a more southerly distribution at West Greenland than a West Greenland year class. A year class from East Greenland-Iceland is normally observed in ICNAF Divisions 1E and 1F at an age of one year. They grow up in this area and at an age of 7-8 years old they begin to migrate from West Greenland to the spawning grounds in ICES Sub-area XIV and Division Va. Some migration back to West Greenland may occur from the southern part of East Greenland.

The following year classes which were and some of which still are important for the fishery originate from West Greenland: 1947, 1950, 1953, 1957, 1960, 1961 and 1968. Of East Greenland origin the following were important or relatively important for the fishery at West Greenland: 1945, 1956, 1958, 1961, 1962, 1963, 1964 and 1968. The 1956 and the 1961 year classes were the most important.

32.2 Distribution of cod eggs and larvae

The ICNAF NORWESTLANT Survey 1963 showed that cod eggs in April 1963 were distributed in a continuous belt from Iceland to East Greenland, along East Greenland, round Cape Farewell and over the banks at West Greenland. Concentrations of larvae were, however, only found in two areas. One at West Greenland (ICNAF Divisions 1B-1D) which is the normal area of distribution for cod larvae of West Greenland origin. The other concentration was found from Iceland to East Greenland over the ridge. Thus, the distribution of larvae was disrupted into two parts compared to the more continuous distribution of the eggs.

If the occurrence of eggs in April 1963 reflects the general picture of distribution of eggs in April shortly after spawning, then the contribution of cod from East Greenland-Iceland to West Greenland may depend upon how successful the spawning is in the various areas off East Greenland and at Iceland, and of course upon the size of the spawning stock.

Icelandic investigations have shown that the incubation time for cod eggs off East Greenland is 20-30 days. The speed of the East Greenland Current is known to be 4.5-9.5 nautical miles per day. Thus, eggs from the South East Greenland area can be transported to South West Greenland before hatching.

In 1963 no larvae were found at South West Greenland (Divisions 1E and 1F). This indicates that there may have been a spawning failure in an area at East Greenland from Angmagssalik Bank to Cape Farewell. As the year class 1963 was relatively important for the fishery at West Greenland, and as

the West Greenland component of that year class contributed very little to the fishery, it seems likely to assume that these catches consisted of cod originating from the larvae concentrations found in July between Iceland and East Greenland.

The International O-Group Surveys in the Iceland-East Greenland area in the years 1970-74 found no O-group cod along East Greenland from 64°N to 60°N. Only in the year of 1973 was a dense concentration of O-group cod found over the Dohrn Bank. This year class was found at West Greenland in ICNAF Division 1F as 1 year old and also as 2 and 3 years old in Divisions 1E and 1D. These 1-3 year old cod from the year class 1973 may have originated from the concentration over the Dohrn Bank like the year class 1963 did.

These observations indicate that in some years not only the spawning areas off East Greenland are important to the fishery at West Greenland, but also spawning grounds rather close to Iceland.

33. Management Problems for Cod at Greenland

Apart from the problems of adequate data and parameters for analyses of the state of stocks and for forecasts of stocks and catches, management of the cod stocks round Greenland is faced with another problem.

The Working Group observed that a quota regulation is applied to the ICNAF part of the Greenland area. It is also observed that while for practical reasons the ICNAF Subarea 1 cod quota is not split up in areal sections, the analyses on which the scientific advice to ICNAF are based consider the stocks in Divisions 1A-1D and Divisions 1E-1F separately.

In recent years the ICNAF scientists have advised that due to the very low stock size and a possible danger of failure in recruitment due to low spawning stock size, fishing should be kept at the lowest practical level. In this context the scientists have also pointed out that the recruitment to West Greenland stocks is depending partly upon the spawning stock at East Greenland.

The present report confirms that there is a strong interrelationship between cod in ICNAF Divisions 1E-1F and cod at East Greenland and partly at Iceland. Although the migration of adult cod is mainly from West Greenland to East Greenland and to Iceland, the Working Group considers that the cod fisheries at West Greenland are depending to a certain degree on spawning stocks at East Greenland and possibly even at Iceland.

The Working Group also considers that for cod in ICNAF Divisions 1E-1F the interrelationship with the East Greenland cod is just as pronounced as the interrelationship with cod in Divisions 1A-1D. It therefore seems proper to consider East and West Greenland as a unit management area. If a break down for management purposes is to be considered, it may be as proper to combine Divisions 1E-1F with East Greenland (ICES Sub-area XIV) as with ICNAF Divisions 1A-1D.

Table 1. Nominal catch of Cod. ICES Division Va (Iceland Grounds). In thousand tons. 1955-75 (Bulletin Statistique).

| Species: COD Country | 1955 | 1956 | 1957 | 1958 | 1959 | 1960 | 1961 | 1962 | 1963 | 1964 | 1965 | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 | 1975* |
|-------------------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Belgium | 9.0 | 7.0 | 6.7 | 9.9 | 5.5 | 5.6 | 5.4 | 8.2 | 6.3 | 3.1 | 3.7 | 3.0 | 2.3 | 3.4 | 2.7 | 3.0 | 3.0 | 2.5 | 1.1 | 1.1 | 1.0 |
| Denmark | + | | | | | | | | | | | | | | | | | | | | |
| Faroe Isl. | 18.7 | 16.2 | 20.9 | 17.9 | 7.7 | 11.8 | 10.6 | 8.7 | 6.3 | 6.9 | 5.2 | 3.4 | 2.8 | 4.3 | 2.6 | 4.3 | 8.6 | 11.1 | 14.2 | 12.1 | 9.6 |
| France | | | | | | | 0.1 | 0.1 | | | | 0.1 | 0.4 | 0.1 | 0.1 | 1.9 | 1.5 | - | - | 0.2 | - |
| Germany (Fed.Rep.) ¹⁾ .. | 48.2 | 30.0 | 23.3 | 37.8 | 35.6 | 37.9 | 21.8 | 34.2 | 33.0 | 19.3 | 15.3 | 9.9 | 15.4 | 29.6 | 19.4 | 24.7 | 27.3 | 11.7 | 6.6 | 5.5 | 2.2 |
| German.Dem.Rep. ²⁾ | | | | | | | 0.3 | 0.5 | 0.9 | 0.5 | 0.5 | 0.3 | 0.4 | 0.9 | 0.5 | 2.7 | 0.7 | 0.7 | | | |
| Iceland | 315.4 | 292.6 | 247.1 | 284.4 | 284.3 | 295.7 | 233.9 | 221.8 | 232.8 | 273.6 | 233.5 | 224.0 | 193.4 | 227.6 | 281.7 | 302.9 | 250.3 | 225.4 | 234.9 | 238.3 | 266.8 |
| Netherlands | | | + | | | | 0.1 | 0.5 | 0.7 | 0.7 | 0.5 | 0.1 | | | + | | | | | | |
| Norway | 7.1 | 4.6 | 8.2 | 6.8 | 5.5 | 3.4 | 4.2 | 4.7 | 3.5 | 2.7 | 0.4 | 0.5 | 0.2 | 0.3 | 0.4 | 0.4 | 0.3 | 0.6 | 0.1 | 0.2 | 0.1 |
| Poland ²⁾ | | | | + | | | | | 0.2 | 0.1 | | | | | | 1.6 | 0.3 | 0.2 | | | |
| U.K.(England & Wales) | 138.7 | 127.8 | 144.3 | 150.5 | 112.7 | 109.4 | 96.5 | 105.1 | 123.2 | 122.2 | 128.1 | 109.0 | 126.6 | 111.6 | 95.4 | 125.2 | 157.7 | 144.2 | 121.3 | 115.4 | 91.0 |
| U.K.(Scotland) | 1.0 | 2.5 | 1.4 | 1.2 | 1.3 | 1.2 | 2.1 | 3.1 | 3.2 | 4.6 | 6.8 | 4.8 | 3.6 | 2.8 | 4.0 | 5.3 | 4.1 | 3.0 | 1.0 | 2.1 | 1.6 |
| U.S.S.R. ²⁾ | | | | | | | | | | | 0.2 | 2.0 | 0.3 | 1.4 | 0.2 | + | 0.1 | + | | | |
| Total | 538.1 | 480.7 | 451.9 | 508.5 | 452.6 | 465.0 | 375.0 | 386.9 | 410.1 | 433.7 | 394.2 | 357.1 | 345.0 | 382.0 | 407.0 | 472.0 | 453.9 | 399.4 | 379.2 | 374.9 | 372.3 |
| Bull.Stat. Total | 536.8 | 482.2 | 453.0 | 510.5 | 454.2 | 465.0 | 375.6 | 386.4 | 409.4 | 434.5 | 393.6 | 357.4 | 344.0 | 379.5 | 405.2 | 470.8 | 453.0 | 398.5 | 379.9 | 375.0 | |

The national statistics used in the table (see footnotes 1 and 2) differ slightly from those given in Bulletin Statistique. The order of magnitude of these discrepancies is shown by comparison of the total catches at the bottom of the table.

* Provisional.

1) From national statistics from Bundesforschungsanstalt f. Fischerei, Hamburg.

2) From national statistics.

+ = less than 0.1 thousand tons.

Note: Due to a mistake during the preparation of the table minor discrepancies (less than 2 thousand tons) occur between the total given in the table and the catch data used in the assessment for the years 1966 (2 000 tons), 1967 (300 tons), 1968 (1 400 tons), 1969 (200 tons) and 1971 (100 tons).

Table 2. Effort and catch per unit of effort 1970-75.

a) English effort

| Year | Hours trawling | | | Tons/hours trawling | Effort raised to total catches (non-spawning) |
|------|----------------|---------|---------|---------------------|---|
| | Motor | | Total | | |
| | Steam | Total | | | |
| 1970 | 59 159 | 140 365 | 199 524 | 0.628 | 491 222 (1.00) |
| 1971 | 89 237 | 211 430 | 300 667 | 0.525 | 627 835 (1.28) |
| 1972 | 98 937 | 220 673 | 319 610 | 0.451 | 606 184 (1.23) |
| 1973 | 82 913 | 194 971 | 277 884 | 0.437 | 594 369 (1.21) |
| 1974 | 68 770 | 164 612 | 233 382 | 0.495 | 545 297 (1.11) |
| 1975 | | | 212 608 | 0.428 | 640 889 (1.30) |

b) Icelandic big trawlers (>500 GRT)

| Year | Hours trawling | Cod catch (1 000 tons) | Tons/hours trawling |
|------|----------------|------------------------|-----------------------|
| 1970 | 59 941 | 25.7 | 0.429 |
| 1971 | 62 406 | 14.5 | 0.232 |
| 1972 | 61 328 | 11.8 | 0.192 |
| 1973 | 55 909 | 7.9 | 0.141 |
| 1974 | 63 752 | 20.9 ^{*)} | 0.328 ^{*)} |
| 1975 | 65 629 | (20) ^{*)} | (0.305) ^{*)} |

*) Splitting of catch between big trawler and stern-trawler estimated.

c) Icelandic multigear boats (less than 500 GRT)

| Year | Days absent when trawling | Catch (1 000 tons) | Tons/day absent |
|------|---------------------------|--------------------|-----------------|
| 1970 | 39 103 | 74.2 | 1.90 |
| 1971 | 38 669 | 58.2 | 1.49 |
| 1972 | 39 041 | 29.8 | 0.76 |
| 1973 | 28 379 | 20.3 | 0.71 |
| 1974 | 30 306 | 17.9 | 0.59 |

Table 2 (Continued)

d) Icelandic stern trawlers (300 - 500 GRT)

| Year | Hours trawling | Cod catch (1 000 tons) | Tons/hours trawling |
|------|----------------|------------------------|----------------------|
| 1970 | 1 266 | 0.4 | 0.326 |
| 1971 | 13 942 | 6.6 | 0.472 |
| 1972 | 18 939 | 8.1 | 0.431 |
| 1973 | 57 302 | 25.5 | 0.445 |
| 1974 | 111 814 | 51.5 | 0.461 |
| 1975 | 146 866 | (78.6) [⊘] | (0.535) [⊘] |

⊘) Splitting of catch between big trawler and stern trawler estimated.

e) Icelandic long lines

| Year | Days absent | Cod catch (1 000 tons) | Tons/days absent |
|------|-------------|------------------------|------------------|
| 1970 | 28 629 | 43.7 | 1.528 |
| 1971 | 30 442 | not available | |
| 1972 | 31 486 | 33.9 | 1.076 |
| 1973 | 29 831 | 34.7 | 1.162 |
| 1974 | 27 570 | 28.9 | 1.049 |
| 1975 | | 22.7 | |

f) Icelandic gill nets

| Year | Days absent | Cod catch (1 000 tons) | Tons/days absent |
|------|-------------|------------------------|------------------|
| 1970 | 20 460 | 132.5 | 6.48 |
| 1971 | 22 834 | not available | |
| 1972 | 27 801 | 114.3 | 4.11 |
| 1973 | 30 451 | 119.9 | 3.94 |
| 1974 | 28 817 | 99.9 | 3.47 |
| 1975 | | 94.4 | |

g) Icelandic hand lines

| Year | Days absent | Cod catch (1 000 tons) | Tons/days absent |
|------|-------------|------------------------|------------------|
| 1970 | 17 901 | 23.5 | 1.31 |
| 1971 | 22 143 | not available | |
| 1972 | 25 932 | 20.8 | 0.80 |
| 1973 | 23 418 | 19.7 | 0.84 |
| 1974 | 25 423 | 15.6 | 0.61 |
| 1975 | | 16.4 | |

Table 3. Iceland Cod.
Age compositions of catches 1955-75 used as input data for Virtual
Population Analysis (thousands of fish).

| Age | 1955 | 1956 | 1957 | 1958 | 1959 | 1960 | 1961 | 1962 | 1963 | 1964 |
|-----|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2 | 169 | 219 | 781 | 2 938 | 1 945 | 1 581 | 1 840 | 0 996 | 1 356 | 2 268 |
| 3 | 3 981 | 6 318 | 15 967 | 25 694 | 20 329 | 13 434 | 14 665 | 12 309 | 14 884 | 16 284 |
| 4 | 24 277 | 16 112 | 23 354 | 30 535 | 44 478 | 30 544 | 19 971 | 28 867 | 29 298 | 28 590 |
| 5 | 33 115 | 28 249 | 17 250 | 16 012 | 22 953 | 29 000 | 19 680 | 19 718 | 22 390 | 20 312 |
| 6 | 25 328 | 22 879 | 18 212 | 11 501 | 6 693 | 12 105 | 18 826 | 15 786 | 11 586 | 10 882 |
| 7 | 9 749 | 14 945 | 12 627 | 15 304 | 4 760 | 8 681 | 7 617 | 12 424 | 17 508 | 7 482 |
| 8 | 4 545 | 4 551 | 12 927 | 14 876 | 7 561 | 5 967 | 6 502 | 4 243 | 8 295 | 17 182 |
| 9 | 5 757 | 3 433 | 3 734 | 7 466 | 11 698 | 6 512 | 3 633 | 7 852 | 2 640 | 5 169 |
| 10 | 18 172 | 1 983 | 2 197 | 1 982 | 7 221 | 12 136 | 2 962 | 2 614 | 6 063 | 1 763 |
| 11 | 2 548 | 14 391 | 1 327 | 1 492 | 979 | 3 661 | 6 181 | 1 866 | 1 410 | 3 315 |
| 12 | 1 380 | 1 475 | 8 020 | 6 001 | 981 | 911 | 1 230 | 3 007 | 946 | 7 768 |
| 13 | 2 083 | 1 679 | 531 | 1 192 | 223 | 221 | 90 | 386 | 1 396 | 463 |
| 14 | 186 | 980 | 740 | 663 | 1 203 | 219 | 126 | 68 | 204 | 969 |

/...

| Age | 1965 | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 |
|-----|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| 1 | 0 | 0 | 3 | 0 | 0 | 0 | 0 | 28 | 0 | 0 | 0 |
| 2 | 2 922 | 3 852 | 5 189 | 0 772 | 0 140 | 315 | 0 820 | 3 685 | 2 406 | 2 320 | 1 445 |
| 3 | 22 039 | 16 957 | 27 444 | 11 514 | 9 828 | 10 670 | 13 303 | 8 664 | 37 583 | 14 211 | 29 522 |
| 4 | 30 535 | 30 039 | 25 937 | 49 731 | 23 168 | 50 014 | 35 932 | 27 765 | 25 602 | 58 770 | 28 786 |
| 5 | 21 562 | 19 791 | 24 063 | 22 280 | 43 262 | 24 737 | 45 939 | 30 861 | 26 338 | 22 632 | 44 057 |
| 6 | 11 002 | 12 338 | 11 953 | 16 072 | 16 968 | 27 188 | 21 275 | 23 346 | 16 048 | 15 183 | 12 421 |
| 7 | 9 050 | 6 196 | 7 807 | 17 478 | 12 826 | 15 497 | 17 443 | 11 190 | 12 011 | 9 640 | 11 118 |
| 8 | 6 228 | 7 118 | 2 838 | 5 657 | 17 411 | 12 066 | 12 334 | 10 596 | 3 607 | 6 140 | 4 202 |
| 9 | 11 670 | 2 305 | 4 142 | 1 728 | 1 881 | 14 581 | 6 885 | 11 243 | 5 893 | 1 705 | 2 212 |
| 10 | 1 694 | 5 862 | 1 279 | 3 169 | 578 | 516 | 4 710 | 4 298 | 7 853 | 3 059 | 867 |
| 11 | 974 | 526 | 2 017 | 526 | 498 | 175 | 360 | 1 281 | 1 452 | 2 162 | 1 146 |
| 12 | 587 | 281 | 95 | 598 | 101 | 99 | 108 | 83 | 261 | 293 | 466 |
| 13 | 131 | 374 | 40 | 57 | 63 | 43 | 57 | 33 | 11 | 108 | 83 |
| 14 | 246 | 54 | 153 | 53 | 29 | 18 | 18 | 3 | 1 | 31 | 19 |

Table 4. Cod.
 Division Va. Mean weight at age.
 Average of the period 1970-74.

| Age | English data | Icelandic data | | Stock |
|-----------------|--------------|----------------|----------|-------|
| | | Non-spawning | Spawning | |
| 1 | | 0.22 | | 0.22 |
| 2 | 0.69 | 0.78 | 0.43 | 0.64 |
| 3 | 0.91 | 1.19 | 1.30 | 1.12 |
| 4 | 1.32 | 1.80 | 2.78 | 1.93 |
| 5 | 1.84 | 2.63 | 4.51 | 2.92 |
| 6 | 2.73 | 3.47 | 5.40 | 3.80 |
| 7 | 3.86 | 4.12 | 6.17 | 4.65 |
| 8 | 4.69 | 4.55 | 6.60 | 5.25 |
| 9 | 4.96 | 4.82 | 6.78 | 5.48 |
| 10 | 5.55 | 5.33 | 7.30 | 6.01 |
| 11 | 6.61 | 6.72 | 8.37 | 7.18 |
| 12 | 9.69 | 7.31 | 9.68 | 8.93 |
| 13 | 11.41 | 9.29 | 12.82 | 11.14 |
| 14 | 15.40 | 12.11 | 18.10 | 15.14 |
| 15 ⁺ | 13.41 | 11.17 | 23.95 | 15.90 |

Length/weight regression parameters: $l_n w = a_l + b$

| | | |
|-------------------------|-------|---------|
| | (a) | (b) |
| Non-spawning: England : | 3.000 | 11.6183 |
| " Iceland | 2.551 | 9.7361 |
| Spawning Iceland | 3.072 | 11.8913 |

Table 5.a. Cod at Iceland.
Non-spawning fishery. Mean weight (kg) at age.

| Age/Year | 1965 | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 |
|----------|-------|-------|-------|-------|-------|------|-------|-------|-------|-------|-------|
| 1 | - | - | 0.41 | - | - | - | - | 0.22 | - | - | - |
| 2 | 1.18 | 1.06 | 1.29 | 0.75 | 0.69 | 0.58 | 0.65 | 0.63 | 1.32 | 0.74 | 1.00 |
| 3 | 1.59 | 1.61 | 1.64 | 1.40 | 1.28 | 1.04 | 1.14 | 1.12 | 1.40 | 1.26 | 1.32 |
| 4 | 2.62 | 2.33 | 2.36 | 1.57 | 1.92 | 1.60 | 1.80 | 1.81 | 1.83 | 1.94 | 1.87 |
| 5 | 2.98 | 3.34 | 3.16 | 2.56 | 2.66 | 2.59 | 2.54 | 2.59 | 2.79 | 2.62 | 2.74 |
| 6 | 3.94 | 4.10 | 4.11 | 3.28 | 3.45 | 3.47 | 3.49 | 3.36 | 3.40 | 3.61 | 3.48 |
| 7 | 4.63 | 5.09 | 4.94 | 4.25 | 4.20 | 4.10 | 4.16 | 3.99 | 4.20 | 4.14 | 4.49 |
| 8 | 5.31 | 5.77 | 6.21 | 4.37 | 4.53 | 4.31 | 4.15 | 4.54 | 4.67 | 5.09 | 4.92 |
| 9 | 5.55 | 5.18 | 6.04 | 6.39 | 5.41 | 4.78 | 4.41 | 4.85 | 5.07 | 5.03 | 5.32 |
| 10 | 7.07 | 5.57 | 7.10 | 6.77 | 5.27 | 6.45 | 4.93 | 4.59 | 5.13 | 5.56 | 6.79 |
| 11 | 8.39 | 6.67 | 6.81 | 7.74 | 6.38 | 9.09 | 7.79 | 5.47 | 5.34 | 5.89 | 7.04 |
| 12 | 7.58 | 8.94 | 9.70 | 7.21 | 9.82 | 9.44 | 8.62 | 7.27 | 5.37 | 5.87 | 7.77 |
| 13 | 13.39 | 8.43 | 9.22 | 11.41 | 10.40 | 6.71 | 9.79 | 11.90 | 11.88 | 6.16 | 8.61 |
| 14 | 10.86 | 9.92 | 12.09 | 10.32 | 9.51 | - | 13.04 | - | - | 11.17 | - |
| 15+ | 11.67 | 11.56 | 13.66 | 16.26 | 18.49 | - | 11.17 | - | - | - | 15.20 |

Table 5.b. Cod at Iceland.
Spawning fishery. Mean weight (kg) at age.

| Age/Year | 1965 | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 |
|----------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| 1 | - | - | 0.55 | 0.42 | - | 0.33 | - | - | 0.52 | 0.44 | 0.53 |
| 2 | 2.08 | 1.55 | 1.72 | 2.00 | 1.22 | 0.95 | 1.07 | 0.99 | 1.59 | 1.92 | 1.38 |
| 3 | 3.80 | 3.22 | 3.53 | 3.47 | 2.87 | 2.47 | 2.42 | 2.70 | 2.80 | 3.51 | 3.49 |
| 4 | 4.37 | 4.92 | 5.04 | 4.52 | 3.87 | 4.68 | 4.44 | 4.19 | 4.66 | 4.56 | 4.92 |
| 5 | 5.15 | 5.56 | 6.10 | 5.68 | 4.78 | 5.30 | 5.45 | 5.15 | 5.30 | 5.79 | 5.26 |
| 6 | 5.91 | 6.69 | 6.85 | 5.62 | 5.32 | 5.45 | 6.39 | 6.08 | 6.29 | 6.66 | 6.14 |
| 7 | 6.57 | 7.08 | 8.21 | 7.41 | 6.50 | 6.01 | 6.04 | 6.52 | 7.05 | 7.38 | 7.16 |
| 8 | 7.18 | 7.74 | 7.49 | 8.40 | 6.47 | 6.51 | 6.53 | 6.21 | 6.66 | 7.97 | 8.04 |
| 9 | 9.02 | 8.44 | 9.61 | 8.66 | 7.90 | 8.52 | 6.87 | 6.60 | 6.73 | 7.79 | 8.79 |
| 10 | 12.52 | 11.66 | 9.89 | 10.65 | 7.94 | 10.55 | 8.59 | 7.43 | 7.14 | 8.14 | 8.82 |
| 11 | 10.09 | 14.25 | 13.54 | 11.80 | 9.52 | 12.20 | 10.73 | 8.63 | 7.86 | 8.97 | 10.09 |
| 12 | 9.92 | 12.36 | 10.91 | 15.99 | 14.78 | 11.33 | 16.04 | 11.02 | 13.50 | 12.24 | 11.28 |
| 13 | 16.13 | 14.22 | 15.32 | 16.32 | 14.43 | 21.73 | 16.94 | 31.12 | - | 20.69 | 12.68 |
| 14 | 18.22 | 14.98 | - | - | - | 28.00 | 18.63 | 15.44 | - | - | - |
| 15+ | 21.39 | 20.76 | 20.20 | 18.34 | 21.80 | 28.00 | 29.70 | - | - | - | - |

Table 6. Cod at Iceland.
VPA input values of F for 1975

| | (A) | (B) | (C) | (D) | (E) |
|-----|--|--------------------|---------|---------|---------|
| Age | F 1970 from prelimin- ary run | Adjusted F 1970 | (B)x1.2 | (B)x1.3 | (B)x1.4 |
| 1 | 0.00 | 0.001 | 0.001 | 0.001 | 0.001 |
| 2 | 0.00 | 0.01 | 0.01 | 0.01 | 0.01 |
| 3 | 0.06 | 0.10 | 0.12 | 0.13 | 0.14 |
| 4 | 0.31 | 0.31 | 0.37 | 0.40 | 0.43 |
| 5 | 0.36 | 0.36 | 0.43 | 0.47 | 0.50 |
| 6 | 0.38 | 0.38 | 0.46 | 0.49 | 0.53 |
| 7 | 0.26 | 0.40 | 0.48 | 0.52 | 0.56 |
| 8 | 0.50 | 0.50 | 0.60 | 0.65 | 0.70 |
| 9 | 1.00 | } 0.75 | 0.90 | 0.98 | 1.05 |
| 10 | 0.55 | | | | |
| 11 | 0.67 | | | | |
| 12 | 0.83 | | | | |
| 13 | 0.73 | | | | |

Table 7. Cod at Iceland.
Derived values of F for 1970 (see Table 6).

| Age | (C) | (D) | (E) |
|-----|------|------|------|
| 3 | 0.06 | 0.06 | 0.06 |
| 4 | 0.31 | 0.31 | 0.32 |
| 5 | 0.36 | 0.36 | 0.36 |
| 6 | 0.38 | 0.38 | 0.38 |
| 7 | 0.26 | 0.26 | 0.26 |
| 8 | 0.50 | 0.50 | 0.51 |
| 9 | 1.00 | 1.00 | 1.00 |
| 10 | 0.55 | 0.56 | 0.56 |
| 11 | 0.68 | 0.68 | 0.68 |
| 12 | 0.84 | 0.84 | 0.84 |
| 13 | 0.77 | 0.75 | 0.82 |

Table 8. Iceland Cod.
Estimates of fishing mortality coefficients for 1955-75 calculated
by VPA for age and year.

| Age | 1955 | 1956 | 1957 | 1958 | 1959 | 1960 | 1961 | 1962 | 1963 | 1964 |
|-----|------|------|------|------|------|------|------|------|------|------|
| 1 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 2 | 0.00 | 0.00 | 0.00 | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 | 0.00 | 0.01 |
| 3 | 0.03 | 0.04 | 0.10 | 0.12 | 0.08 | 0.10 | 0.09 | 0.10 | 0.11 | 0.06 |
| 4 | 0.15 | 0.17 | 0.18 | 0.30 | 0.30 | 0.16 | 0.22 | 0.25 | 0.36 | 0.30 |
| 5 | 0.17 | 0.26 | 0.27 | 0.18 | 0.38 | 0.33 | 0.14 | 0.35 | 0.32 | 0.46 |
| 6 | 0.26 | 0.17 | 0.26 | 0.29 | 0.10 | 0.35 | 0.36 | 0.16 | 0.35 | 0.25 |
| 7 | 0.39 | 0.24 | 0.13 | 0.37 | 0.19 | 0.19 | 0.39 | 0.44 | 0.28 | 0.40 |
| 8 | 0.29 | 0.32 | 0.34 | 0.22 | 0.31 | 0.37 | 0.21 | 0.40 | 0.59 | 0.48 |
| 9 | 0.30 | 0.36 | 0.48 | 0.34 | 0.27 | 0.49 | 0.41 | 0.42 | 0.47 | 0.94 |
| 10 | 0.40 | 0.16 | 0.42 | 0.51 | 0.65 | 0.50 | 0.43 | 0.58 | 0.68 | 0.66 |
| 11 | 0.42 | 0.65 | 0.15 | 0.57 | 0.51 | 0.84 | 0.52 | 0.53 | 0.73 | 1.04 |
| 12 | 0.29 | 0.46 | 0.97 | 2.03 | 0.94 | 1.38 | 0.78 | 0.52 | 0.56 | 1.23 |
| 13 | 0.65 | 0.68 | 0.30 | 0.36 | 0.37 | 0.57 | 0.45 | 0.60 | 0.49 | 0.60 |
| 14 | 0.75 | 0.75 | 0.75 | 0.75 | 0.75 | 0.75 | 0.75 | 0.75 | 0.75 | 0.75 |

| Age | 1965 | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 |
|-----|------|------|------|------|------|------|------|------|------|------|------|
| 1 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 2 | 0.01 | 0.01 | 0.03 | 0.00 | 0.00 | 0.00 | 0.01 | 0.01 | 0.02 | 0.01 | 0.01 |
| 3 | 0.10 | 0.07 | 0.09 | 0.07 | 0.04 | 0.06 | 0.08 | 0.07 | 0.15 | 0.13 | 0.14 |
| 4 | 0.16 | 0.19 | 0.14 | 0.24 | 0.21 | 0.32 | 0.31 | 0.24 | 0.30 | 0.36 | 0.43 |
| 5 | 0.40 | 0.15 | 0.23 | 0.17 | 0.34 | 0.36 | 0.53 | 0.49 | 0.38 | 0.48 | 0.50 |
| 6 | 0.50 | 0.42 | 0.13 | 0.23 | 0.19 | 0.38 | 0.61 | 0.58 | 0.51 | 0.39 | 0.53 |
| 7 | 0.34 | 0.58 | 0.51 | 0.28 | 0.30 | 0.26 | 0.45 | 0.79 | 0.67 | 0.67 | 0.56 |
| 8 | 0.70 | 0.50 | 0.58 | 0.87 | 0.49 | 0.51 | 0.34 | 0.54 | 0.64 | 0.91 | 0.70 |
| 9 | 0.70 | 0.62 | 0.62 | 0.87 | 0.82 | 1.00 | 0.61 | 0.59 | 0.66 | 0.72 | 1.05 |
| 10 | 0.98 | 0.98 | 0.86 | 1.53 | 0.84 | 0.56 | 1.14 | 1.02 | 1.14 | 0.89 | 1.05 |
| 11 | 0.98 | 1.01 | 1.19 | 1.16 | 1.20 | 0.68 | 1.01 | 1.22 | 1.29 | 1.24 | 1.05 |
| 12 | 0.51 | 0.88 | 0.49 | 1.73 | 0.73 | 0.84 | 1.28 | 0.69 | 0.91 | 1.06 | 1.05 |
| 13 | 0.72 | 0.72 | 0.28 | 0.62 | 0.92 | 0.82 | 2.39 | 2.90 | 0.18 | 1.39 | 1.05 |
| 14 | 0.75 | 0.75 | 0.75 | 0.75 | 0.75 | 0.75 | 1.05 | 1.05 | 1.05 | 1.05 | 1.05 |

Table 9. Iceland Cod.
Estimates of stock size at beginning of year 1955-75 calculated by
VPA (thousands of fish).

| Age | 1955 | 1956 | 1957 | 1958 | 1959 | 1960 | 1961 | 1962 | 1963 | 1964 |
|-----|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|
| 1 | 264 785 | 386 714 | 458 889 | 229 164 | 284 501 | 214 518 | 243 737 | 438 848 | 388 094 | 437 131 |
| 2 | 246 532 | 216 787 | 316 614 | 375 706 | 187 622 | 232 929 | 175 632 | 199 554 | 359 297 | 317 744 |
| 3 | 146 263 | 201 691 | 177 292 | 258 516 | 304 949 | 151 856 | 189 278 | 142 133 | 162 481 | 292 943 |
| 4 | 193 883 | 116 155 | 159 427 | 130 756 | 188 490 | 231 330 | 112 214 | 141 740 | 105 268 | 119 607 |
| 5 | 237 875 | 136 862 | 80 586 | 109 493 | 79 606 | 114 343 | 161 878 | 73 898 | 90 080 | 59 880 |
| 6 | 120 581 | 164 924 | 86 644 | 50 465 | 75 223 | 44 572 | 67 561 | 114 799 | 42 793 | 53 632 |
| 7 | 32 788 | 75 943 | 114 418 | 54 558 | 30 977 | 55 552 | 25 622 | 38 411 | 79 768 | 24 631 |
| 8 | 20 045 | 18 095 | 48 731 | 82 296 | 30 928 | 21 076 | 37 665 | 14 142 | 20 306 | 49 566 |
| 9 | 24 630 | 12 325 | 10 726 | 28 287 | 53 990 | 18 527 | 11 898 | 24 984 | 7 771 | 9 205 |
| 10 | 60 008 | 14 991 | 7 009 | 5 435 | 16 453 | 33 683 | 9 333 | 6 482 | 13 412 | 3 996 |
| 11 | 8 129 | 32 824 | 10 487 | 3 767 | 2 675 | 7 018 | 16 705 | 4 984 | 2 968 | 5 565 |
| 12 | 6 033 | 4 370 | 14 013 | 7 390 | 1 749 | 1 313 | 2 484 | 8 141 | 2 410 | 1 171 |
| 13 | 4 746 | 3 699 | 2 256 | 4 342 | 7 797 | 559 | 270 | 936 | 3 972 | 1 126 |
| 14 | 384 | 2 024 | 1 528 | 1 369 | 2 485 | 452 | 260 | 140 | 421 | 2 001 |

| Age | 1965 | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 |
|-----|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|
| 1 | 513 056 | 270 268 | 387 258 | 284 339 | 281 503 | 211 458 | 456 807 | 191 024 | 374 214 | 195 673 | 1 104 |
| 2 | 357 892 | 420 054 | 221 276 | 317 058 | 232 796 | 230 474 | 173 127 | 374 001 | 156 372 | 306 380 | 160 203 |
| 3 | 258 098 | 290 378 | 340 432 | 176 480 | 258 888 | 190 471 | 188 412 | 141 003 | 302 878 | 125 854 | 248 747 |
| 4 | 225 147 | 191 437 | 222 440 | 253 970 | 134 102 | 203 088 | 146 316 | 142 258 | 107 627 | 214 108 | 90 231 |
| 5 | 72 230 | 156 826 | 129 686 | 158 743 | 163 190 | 88 938 | 121 331 | 87 504 | 91 490 | 65 106 | 122 523 |
| 6 | 30 818 | 39 787 | 110 565 | 84 523 | 109 897 | 94 751 | 50 605 | 58 204 | 43 988 | 51 263 | 33 022 |
| 7 | 34 122 | 15 374 | 21 506 | 79 748 | 54 740 | 74 696 | 53 169 | 22 407 | 26 764 | 21 640 | 28 344 |
| 8 | 13 452 | 19 808 | 7 043 | 10 614 | 49 576 | 33 288 | 47 217 | 27 890 | 8 367 | 11 182 | 9 104 |
| 9 | 25 182 | 5 453 | 9 840 | 3 227 | 3 652 | 24 987 | 16 445 | 27 579 | 13 347 | 3 626 | 3 691 |
| 10 | 2 938 | 10 197 | 2 403 | 4 353 | 1 104 | 1 314 | 7 500 | 7 307 | 12 522 | 5 662 | 1 447 |
| 11 | 1 696 | 900 | 3 137 | 829 | 770 | 389 | 614 | 1 964 | 2 164 | 3 288 | 1 912 |
| 12 | 1 612 | 523 | 269 | 781 | 212 | 189 | 162 | 182 | 473 | 487 | 778 |
| 13 | 279 | 794 | 178 | 135 | 114 | 83 | 67 | 37 | 75 | 155 | 138 |
| 14 | 508 | 112 | 316 | 109 | 60 | 37 | 30 | 5 | 2 | 52 | 32 |

Table 10. Cod at Iceland.
Total stock biomass and spawning stock biomass
(thousands of tons).

| Year | Total stock biomass age groups 3 and older | Spawning stock biomass age groups 7 and older |
|------|--|---|
| 1955 | 2 615 | 924 |
| 1956 | 2 429 | 952 |
| 1957 | 2 208 | 1 138 |
| 1958 | 2 089 | 1 036 |
| 1959 | 2 006 | 783 |
| 1960 | 1 868 | 748 |
| 1961 | 1 745 | 587 |
| 1962 | 1 635 | 550 |
| 1963 | 1 505 | 694 |
| 1964 | 1 480 | 543 |
| 1965 | 1 474 | 422 |
| 1966 | 1 592 | 288 |
| 1967 | 1 846 | 237 |
| 1968 | 1 959 | 487 |
| 1969 | 1 994 | 551 |
| 1970 | 1 899 | 673 |
| 1971 | 1 677 | 637 |
| 1972 | 1 371 | 462 |
| 1973 | (1 319) [*]) | 337 |
| 1974 | (1 183) [*]) | (244) [*]) |
| 1975 | | (231) [*]) |

^{*}) Values sensitive to VPA input values of F for 1975.

Table 11. Estimated year class strengths of Cod from the three VPA's (3 years old, number in 10^{-6}).

| Year Class | Iceland | E. Greenland + W. Greenland 1 E & F | E. Greenland + W. Greenland 1 E & F + Iceland |
|------------|---------|---|---|
| 1952 | 146 | | |
| 1953 | 202 | | |
| 1954 | 177 | | |
| 1955 | 259 | | |
| 1956 | 305 | | |
| 1957 | 152 | 81 | 232 |
| 1958 | 189 | 71 | 260 |
| 1959 | 142 | 16 | 158 |
| 1960 | 162 | 53 | 215 |
| 1961 | 293 | 151 | 444 |
| 1962 | 258 | 78 | 336 |
| 1963 | 290 | 135 | 425 |
| 1964 | 340 | 42 | 382 |
| 1965 | 176 | 12 | 188 |
| 1966 | 259 | 13 | 275 |
| 1967 | 190 | 8 | 200 |
| 1968 | 188 | 6 | 220 |
| 1969 | 141 | 4 | 151 |
| 1970 | 303 | 13 | 342 |

Average 1952 - 1970 year classes 220

Table 22. Nominal catch of Cod. ICNAF Divisions IE and IF in thousand tons 1960-1975 (Statistical Bulletin).

| COD | 1960 | 1961 | 1962 | 1963 | 1964 | 1965 | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 |
|--------------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|------|------|------|------|------|------|
| Faroe Islands | 13.0* | 16.3* | 28.5* | 22.7* | 17.7* | 19.3* | 22.6* | 20.5* | 12.7* | 5.1* | 2.7* | 6.4* | 2.8* | 2.1* | 2.0 |
| France | 0.1 | 0.2 | 0.2 | 0.7 | 1.0 | 0.9 | 2.0 | 1.3 | 7.7 | 3.2 | 0.5 | 0.5 | 0.3 | | |
| Germany, Fed. Rep. | 7.7 | 20.4 | 31.1 | 44.9 | 27.9 | 20.5 | 21.7 | 32.3 | 55.6 | 38.4 | 31.0 | 26.2 | 6.8 | 4.0 | 0.8 |
| German Dem. Rep. | + | | | 2.2 | 1.7 | 0.5* | 1.8 | 1.1 | 4.7 | 1.7 | 3.4* | 0.1 | | | |
| Greenland | 10.2 | 15.9 | 17.2 | 12.1 | 7.2 | 7.9 | 7.1 | 8.6 | 10.0 | 8.2* | 8.6* | 7.1* | 6.9* | 6.0 | 7.6 |
| Iceland | 2.8* | 3.6* | 0.5 | 1.7 | 1.2 | 0.7 | 0.6 | 0.1 | | | | | | | |
| Norway | 14.3* | 13.7* | 3.4* | 9.2* | 11.6* | 8.2* | 10.2* | 13.7* | 10.7* | 5.8* | 1.6* | 1.5* | 6.3* | 4.2* | 1.8 |
| Poland | + | | 0.3 | 0.2 | | + | 0.1 | + | + | 0.1 | | | | | |
| Portugal | 5.4 | 0.4 | 2.6 | 1.5 | | + | 0.2 | + | 6.4 | 5.8 | 1.4 | + | + | + | 0.4 |
| Spain | 0.1 | + | 0.4 | 0.1 | 0.2 | | + | 3.0 | 1.0 | 2.2 | 1.0 | 0.6 | 0.6 | 0.6 | + |
| U.K. (England & Wales) | 8.1 | 2.7 | 6.6 | 10.7 | 13.4 | 6.1 | 11.2 | 5.2 | 4.7 | | 2.9 | 1.4 | 0.4 | 0.5 | 0.8 |
| USSR | 0.1 | | | 1.0 | | | | | 0.5 | | 0.3 | | | + | |
| Total | 61.7 | 73.1 | 90.8 | 106.9 | 81.9 | 64.1 | 77.7 | 85.8 | 114.0 | 70.5 | 53.5 | 43.8 | 24.0 | 17.4 | 13.4 |
| Division INK ^x) | 76.2 | 88.0 | 115.9 | 99.7 | 84.3 | 99.2 | 95.1 | 95.9 | 68.6 | 35.9 | 23.0 | 26.4 | 20.1 | 1.1 | 0 |
| IE-IF Allocated ^x) | 25.2 | 26.5 | 31.0 | 29.8 | 25.9 | 26.1 | 32.2 | 30.9 | 20.3 | 10.8 | 8.2 | 9.2 | 7.3 | 0.7 | 0 |

^x) Catches reported as Division INK (West Greenland unspecified) are given two lines above. Parts of these catches have been allocated (by the Greenland Fisheries Institute) to Divisions IE-IF as given in the last line. The countries for which the catch or part of the catch was reported as Division INK are marked with an asterisk.

Table 23. Nominal catch of Cod.
ICES Sub-Area XIV in thousand tons 1960-1975 (Bulletin Statistique).

| COD | 1960 | 1961 | 1962 | 1963 | 1964 | 1965 | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 ¹⁾ |
|-------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|--------------------|
| Faroe Islands | 0.4 | 1.2 | | | | | | | | | | | 0.9 | 0.2 | 0.7 | 0.3 |
| Germany, Fed.Rep. | 19.1 | 15.0 | 14.3 | 13.9 | 30.6 | 11.0 | 7.8 | 12.1 | 8.3 | 12.6 | 13.9 | 25.6 | 21.6 | 9.3 | 2.3 | 1.5 |
| German Dem.Rep. | | | | | | | | | | | | | | | + | 0.3 |
| Greenland | 1.6 | 1.2 | 0.9 | 0.9 | 1.1 | 0.9 | 0.9 | 0.7 | 0.6 | 0.6 | 0.5 | 0.5 | 0.3 | 0.2 | + | 0.2 |
| Iceland | 2.5 | 1.4 | 0.3 | 1.8 | 2.9 | 4.7 | 4.0 | 10.5 | 6.7 | 4.5 | 5.5 | 4.6 | 3.2 | 1.4 | 3.0 | 0.8 |
| Poland | | | | | | | | | | | 0.8 | 0.4 | 0.3 | + | + | + |
| U.K. | 0.3 | 0.9 | 1.8 | 0.8 | 1.0 | 0.9 | 0.2 | 1.4 | + | | 0.1 | + | 0.2 | 0.7 | 0.5 | 0.4 |
| USSR | | | | 5.7 | | | | + | | + | + | 0.3 | 0.1 | | | |
| Total | 23.9 | 19.7 | 17.3 | 23.1 | 35.6 | 17.5 | 12.9 | 24.7 | 15.7 | 17.8 | 20.9 | 31.5 | 26.6 | 11.8 | 6.6 | 3.4 |

1) Preliminary figures based on verbal information by the Working Group.

Table 24. Nominal catch of Cod.
ICES Sub-Area XIV and ICNAF Divisions IE and IF in thousand tons 1960-1974.
(Bulletin Statistique and Statistical Bulletin).

| COD | 1960 | 1961 | 1962 | 1963 | 1964 | 1965 | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 |
|--------------------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|--------------------|--------------------|--------------------|--------------------|--------------------|------|
| Faroe Islands | 13.4 ^(*) | 17.5 ^(*) | 28.5 ^(*) | 22.7 ^(*) | 17.7 ^(*) | 19.3 ^(*) | 22.6 ^(*) | 20.5 ^(*) | 12.7 ^(*) | 5.1 ^(*) | 2.7 ^(*) | 6.4 ^(*) | 3.7 | 2.3 | 2.7 |
| France | 0.1 | 0.2 | 0.2 | 0.7 | 1.0 | 0.9 | 2.0 | 1.3 | 7.7 | 3.2 | 0.5 | 0.5 | 0.3 | | |
| Germany, Fed. Rep. | 26.8 | 35.4 | 45.4 | 58.8 | 58.5 | 31.5 | 29.5 | 44.4 | 63.9 | 51.0 | 44.9 | 51.8 | 28.4 | 13.3 | 3.1 |
| German Dem. Rep. | + | | | 2.2 | 1.7 | 0.5 ^(*) | 1.8 | 1.1 | 4.7 | 1.7 | 3.4 ^(*) | 0.1 | + | + | + |
| Greenland | 11.8 | 17.1 | 18.1 | 13.0 | 8.3 | 8.8 | 8.0 | 9.3 | 10.6 | 8.8 ^(*) | 9.1 ^(*) | 7.6 ^(*) | 7.2 ^(*) | 6.2 | 7.6 |
| Iceland | 5.5 ^(*) | 5.0 ^(*) | 0.8 | 3.5 | 4.1 | 5.4 | 4.6 | 10.6 | 6.7 | 4.5 | 5.5 | 4.6 | 3.2 | 1.4 | 3.0 |
| Norway | 14.3 ^(*) | 13.7 ^(*) | 3.4 ^(*) | 9.2 ^(*) | 11.6 ^(*) | 8.2 ^(*) | 10.2 ^(*) | 13.7 ^(*) | 10.7 ^(*) | 5.8 ^(*) | 1.6 ^(*) | 1.5 ^(*) | 6.3 ^(*) | 4.2 ^(*) | 1.8 |
| Poland | + | | 0.3 | 0.2 | | + | 0.1 | + | + | 0.1 | 0.8 | 0.4 | 0.3 | + | + |
| Portugal | 5.4 | 0.4 | 2.6 | 1.5 | | + | 0.2 | + | 6.4 | 5.8 | 1.4 | + | + | + | 0.4 |
| Spain | 0.1 | + | 0.4 | 0.1 | 0.2 | | + | 3.0 | 1.0 | 2.2 | 1.0 | 0.6 | 0.6 | 0.6 | + |
| U.K. | 8.4 | 3.6 | 8.4 | 11.5 | 14.4 | 7.0 | 11.4 | 6.6 | 4.7 | | 3.0 | 1.4 | 0.6 | 1.2 | 1.3 |
| USSR | 0.1 | | | 6.7 | | | | + | 0.5 | + | 0.3 | 0.3 | 0.1 | + | |
| Total | 85.6 | 92.8 | 108.1 | 130.0 | 117.5 | 81.6 | 90.6 | 110.5 | 129.7 | 88.3 | 74.4 | 75.3 | 50.6 | 29.2 | 20.0 |
| Division INK ^(x) | 76.2 | 88.0 | 115.9 | 99.7 | 84.3 | 99.2 | 95.1 | 95.9 | 68.6 | 35.9 | 23.0 | 26.4 | 20.1 | 1.1 | 0 |
| IE-IF Allocated ^(x) | 25.2 | 26.5 | 31.0 | 29.8 | 25.9 | 26.1 | 32.2 | 30.9 | 20.3 | 10.8 | 8.2 | 9.2 | 7.3 | 0.7 | 0 |

(x) Catches reported as Division INK (West Greenland, unspecified) are given two lines above. Parts of these catches have been allocated (by the Greenland Fisheries Institute) to Divisions IE-IF as given in the last line. The countries for which the catch or part of the catch was reported as Division INK are marked with an asterisk.

Table 25. Nominal catches of Cod in ICNAF Divisions IE-IF compared to the total catch of Cod in ICNAF Sub-Area I.

| Year | 1960 | 1961 | 1962 | 1963 | 1964 | 1965 | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 |
|---|------|------|------|-------|------|------|------|------|-------|------|------|------|------|------|------|
| Sub-Area I (tons x 10 ⁻³) | 243 | 345 | 451 | 406 | 350 | 360 | 366 | 430 | 394 | 215 | 113 | 121 | 111 | 63 | 48 |
| Divisions IE-IF (tons x 10 ⁻³) | 61.7 | 73.1 | 90.8 | 106.9 | 81.9 | 64.1 | 77.7 | 85.8 | 114.0 | 70.5 | 53.5 | 43.8 | 24.0 | 17.4 | 13.4 |
| Divisions IE-IF as % of Sub-Area I | 25.4 | 21.2 | 20.1 | 26.3 | 23.4 | 17.8 | 21.2 | 20.0 | 28.9 | 32.8 | 47.3 | 36.2 | 21.6 | 27.6 | 27.9 |

Table 26. Cod. East Greenland.
Estimates of total effort (Germany, Fed.Rep. of days fished used as unit).

| Year | Germany, Fed.Rep. catch ^{b)} | Germany, Fed.Rep. effort ^{a)} | Germany, Fed.Rep. c.p.u.e. | Total catch | Total effort |
|--------------------|---------------------------------------|--|----------------------------|-------------|--------------|
| 1962 | 14 299 | 1 660 | 8.61 | 17 295 | 2 008 |
| 1963 | 13 877 | 2 182 | 6.36 | 23 057 | 3 625 |
| 1964 | 30 623 | 3 287 | 9.32 | 35 577 | 3 819 |
| 1965 | 10 965 | 2 734 | 4.01 | 17 497 | 4 363 |
| 1966 | 7 786 | 1 827 | 4.26 | 12 870 | 3 020 |
| 1967 | 12 117 | 2 157 | 5.62 | 24 732 | 4 403 |
| 1968 | 8 323 | 1 361 | 6.12 | 15 701 | 2 567 |
| 1969 | 12 635 | 2 164 | 5.84 | 17 771 | 3 044 |
| 1970 | 13 930 | 1 532 | 9.09 | 20 907 | 2 299 |
| 1971 | 25 644 | 1 737 | 14.8 | 31 516 | 2 135 |
| 1972 | 21 592 | 1 732 | 12.5 | 26 629 | 2 136 |
| 1973 | 9 262 | 931 | 9.95 | 11 752 | 1 181 |
| 1974 | 2 309 | 312 | 7.40 | 6 553 | 885 |
| 1975 ^{c)} | 1 526 | | | 3 435 | |

a) Germany, Federal Republic of, research reports to ICNAF.

b) Bulletin Statistique Sub-Area XIV.

c) Provisional.

Table 27. Cod. ICNAF Divisions IE-IF 1960-75.
Catch in numbers per age group (1 000 fish).

| Age | 1960 | 1961 | 1962 | 1963 | 1964 | 1965 | 1966 | 1967 | 1968 | 1969 |
|-----|-------|-------|--------|--------|-------|-------|--------|--------|--------|--------|
| 3 | - | 11 | 435 | 33 | 77 | - | 1 180 | 49 | 8 | - |
| 4 | 2 214 | 283 | 2 022 | 534 | 562 | 2 447 | 1 996 | 1 070 | 994 | 142 |
| 5 | 798 | 7 745 | 4 879 | 7 710 | 1 061 | 5 336 | 19 836 | 3 211 | 10 713 | 3 167 |
| 6 | 935 | 1 860 | 11 631 | 8 201 | 8 239 | 1 889 | 4 597 | 14 391 | 9 972 | 15 355 |
| 7 | 5 233 | 1 343 | 1 415 | 11 852 | 5 550 | 5 110 | 1 588 | 5 800 | 11 520 | 6 595 |
| 8 | 1 541 | 4 741 | 1 291 | 912 | 4 823 | 3 965 | 3 018 | 583 | 2 236 | 4 662 |
| 9 | 752 | 945 | 2 676 | 248 | 542 | 1 662 | 2 232 | 369 | 182 | 731 |
| 10 | 1 469 | 604 | 475 | 996 | 245 | 223 | 707 | 917 | 123 | 43 |
| 11 | 220 | 1 203 | 308 | 178 | 733 | 158 | 79 | 55 | 314 | 75 |
| 12 | 394 | 129 | 737 | 178 | 81 | 552 | 56 | 28 | 23 | 146 |
| 13 | 1 425 | 245 | 47 | 443 | 48 | 22 | 186 | 36 | 5 | 27 |
| 14+ | 712 | 1 220 | 1 303 | 751 | 256 | 129 | 128 | 107 | 56 | 4 |

| | 1960 | 1961 | 1962 | 1963 | 1964 | 1965 | 1966 | 1967 | 1968 | 1969 |
|--|--------|--------|--------|---------|--------|--------|--------|--------|---------|--------|
| Total corresponding catch (tons) ^{x)} | 15 693 | 20 329 | 27 219 | 32 036 | 22 217 | 21 493 | 35 603 | 26 616 | 36 146 | 30 947 |
| \bar{w} (kg) | 61 705 | 73 086 | 90 789 | 106 882 | 81 942 | 64 137 | 77 661 | 85 751 | 114 001 | 70 476 |
| | 3.93 | 3.60 | 3.34 | 3.34 | 3.69 | 2.98 | 2.18 | 3.22 | 3.15 | 2.28 |

| Age | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 ^{xx)} |
|-----|-------|-------|-------|-------|-------|---------------------|
| 3 | - | - | 1 | 4 | 56 | 59 |
| 4 | 171 | 66 | 2 944 | 60 | 145 | 304 |
| 5 | 1 496 | 1 118 | 952 | 5 133 | 235 | 531 |
| 6 | 3 323 | 2 064 | 2 218 | 980 | 2 664 | 184 |
| 7 | 8 763 | 3 274 | 737 | 1 005 | 206 | 2 587 |
| 8 | 2 989 | 6 054 | 1 482 | 254 | 240 | 160 |
| 9 | 1 874 | 1 266 | 1 611 | 742 | 105 | 54 |
| 10 | 647 | 657 | 293 | 373 | 107 | 27 |
| 11 | 88 | 207 | 173 | 63 | 205 | 21 |
| 12 | 33 | 10 | 60 | 36 | 128 | 7 |
| 13 | 97 | 24 | 4 | 10 | 53 | 3 |
| 14+ | 27 | 44 | 26 | 10 | 16 | 3 |

| | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 |
|----------------------------------|--------|--------|--------|--------|--------|--------|
| Total corresponding catch (tons) | 19 508 | 14 784 | 10 501 | 8 670 | 4 160 | 3 940 |
| \bar{w} (kg) | 53 530 | 43 837 | 23 970 | 17 438 | 13 447 | 11 300 |
| | 2.74 | 2.97 | 2.28 | 2.01 | 3.23 | 2.87 |

x) Including estimates of catches reported as Division INK.
xx) Including estimates of catches for countries other than Germany, Fed. Rep. of (4 652 tons), U.K. (92) and Denmark (G) (3 186) and partly using samples from Divisions north of Divisions IE-IF. 1975 sampling very poor.

Table 28. Cod. East Greenland.
ICES Sub-Area XIV 1960-1975. Catch in numbers per age group (1 000 fish).

| Age | 1960 | 1961 | 1962 | 1963 | 1964 | 1965 | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 |
|----------------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|-------|-------|
| 3 | - | 23 | 4 | - | 1 | - | 28 | - | - | - | - | - | - | 4 | 4 | 25 |
| 4 | 78 | 87 | 64 | 61 | 26 | 131 | 21 | 145 | 104 | 31 | 66 | 25 | 27 | 25 | 63 | 25 |
| 5 | 144 | 240 | 113 | 419 | 108 | 35 | 470 | 302 | 630 | 252 | 76 | 171 | 85 | 197 | 22 | 149 |
| 6 | 255 | 203 | 974 | 743 | 933 | 91 | 89 | 2 346 | 502 | 849 | 500 | 159 | 254 | 126 | 488 | 38 |
| 7 | 1 321 | 215 | 344 | 2 555 | 2 281 | 879 | 137 | 564 | 2 505 | 770 | 1 539 | 1 051 | 295 | 250 | 176 | 344 |
| 8 | 525 | 1 080 | 151 | 419 | 3 682 | 661 | 1 071 | 210 | 238 | 2 103 | 1 060 | 3 785 | 1 299 | 82 | 185 | 68 |
| 9 | 475 | 377 | 1 050 | 70 | 383 | 1 484 | 359 | 1 292 | 62 | 170 | 1 715 | 1 580 | 3 184 | 710 | 52 | 36 |
| 10 | 1 636 | 244 | 298 | 648 | 64 | 59 | 418 | 492 | 144 | 38 | 237 | 1 326 | 818 | 959 | 329 | 9 |
| 11 | 409 | 719 | 132 | 154 | 443 | 27 | 23 | 371 | 69 | 82 | 32 | 171 | 470 | 222 | 259 | 29 |
| 12 | 60 | 184 | 362 | 96 | 74 | 139 | 3 | 37 | 27 | 68 | 63 | 19 | 136 | 72 | 65 | 23 |
| 13 | 487 | 64 | 60 | 190 | 35 | 29 | 27 | 17 | 5 | 24 | 48 | 4 | 26 | 19 | 11 | 7 |
| 14 | 16 | 192 | 15 | 23 | 146 | 41 | 18 | 49 | 10 | 7 | 16 | 9 | 22 | - | - | 2 |
| 15 | 83 | 23 | 143 | 12 | 31 | 80 | 2 | 2 | 9 | 10 | 2 | 5 | 24 | - | - | 2 |
| 16 | - | 76 | - | 71 | 8 | 1 | 5 | 2 | - | 10 | 5 | 3 | 7 | 3 | 2 | |
| 17 | - | - | 64 | 12 | 102 | 2 | 2 | 16 | - | 1 | 3 | - | - | 1 | - | |
| 18 | 39 | - | - | 18 | - | 37 | - | - | 4 | - | - | - | - | - | - | |
| 19 | - | 37 | - | - | 29 | 1 | - | - | - | 3 | - | - | - | 3 | - | |
| ≥20 | - | - | - | - | - | 16 | 9 | 12 | 2 | 5 | 1 | - | - | - | - | |
| Total | 5 528 | 3 764 | 3 774 | 5 491 | 8 346 | 3 713 | 2 682 | 5 857 | 4 311 | 4 423 | 5 363 | 8 305 | 6 647 | 2 673 | 1 656 | 755 |
| Corresponding catch (tons) | 23 914 | 18 597 | 17 295 | 23 057 | 35 577 | 17 497 | 12 870 | 24 732 | 15 701 | 17 771 | 20 907 | 31 516 | 26 629 | 11 752 | 6 553 | 3 435 |

Table 29. Cod. ICES Sub-Area XIV plus ICNAF Divisions IE-IF 1960-1975. Catch in numbers per age group (1 000 fish).

| Age | 1960 | 1961 | 1962 | 1963 | 1964 | 1965 | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 | 1975* |
|----------------------------|--------|--------|---------|---------|---------|--------|--------|---------|---------|--------|--------|--------|--------|--------|--------|--------|
| 3 | | 34 | 439 | 33 | 78 | | 1 208 | 49 | 8 | | | | 1 | 8 | 60 | 84 |
| 4 | 2 292 | 370 | 2 086 | 595 | 588 | 2 578 | 2 017 | 1 215 | 1 098 | 173 | 237 | 91 | 2 971 | 85 | 208 | 329 |
| 5 | 942 | 7 985 | 4 992 | 8 129 | 1 169 | 5 371 | 20 306 | 3 513 | 11 343 | 3 419 | 1 572 | 1 289 | 1 037 | 5 330 | 257 | 680 |
| 6 | 1 190 | 2 063 | 12 605 | 8 944 | 9 172 | 1 980 | 4 686 | 16 737 | 10 474 | 16 204 | 3 823 | 2 223 | 2 472 | 1 106 | 3 152 | 222 |
| 7 | 6 554 | 1 558 | 1 759 | 14 407 | 7 831 | 5 989 | 1 725 | 6 364 | 14 025 | 7 365 | 10 302 | 4 325 | 1 032 | 1 255 | 382 | 2 931 |
| 8 | 2 066 | 5 821 | 1 442 | 1 331 | 8 505 | 4 626 | 4 089 | 793 | 2 474 | 6 765 | 4 049 | 9 839 | 2 781 | 336 | 425 | 228 |
| 9 | 1 227 | 1 322 | 3 726 | 318 | 925 | 3 146 | 2 591 | 1 661 | 244 | 901 | 3 589 | 2 846 | 4 795 | 1 452 | 157 | 90 |
| 10 | 3 105 | 848 | 773 | 1 644 | 309 | 282 | 1 125 | 1 409 | 267 | 81 | 884 | 1 983 | 1 111 | 1 332 | 436 | 36 |
| 11 | 629 | 1 922 | 440 | 332 | 1 176 | 185 | 102 | 426 | 383 | 157 | 120 | 378 | 643 | 285 | 464 | 50 |
| 12 | 454 | 313 | 1 099 | 274 | 155 | 691 | 59 | 65 | 50 | 214 | 96 | 29 | 196 | 108 | 193 | 30 |
| 13 | 1 912 | 309 | 107 | 633 | 83 | 51 | 213 | 53 | 10 | 51 | 145 | 28 | 30 | 29 | 64 | 10 |
| ≥14 | 850 | 1 548 | 1 525 | 887 | 572 | 307 | 164 | 188 | 81 | 40 | 54 | 58 | 79 | 17 | 18 | 5 |
| Total | 21 221 | 24 093 | 30 993 | 37 527 | 30 563 | 25 206 | 38 285 | 32 473 | 45 457 | 35 370 | 24 871 | 23 089 | 17 148 | 11 343 | 5 816 | 4 695 |
| Corresponding catch (tons) | 85 619 | 91 683 | 108 084 | 129 939 | 117 519 | 81 634 | 90 531 | 110 483 | 129 702 | 89 247 | 74 437 | 75 353 | 50 599 | 29 190 | 20 000 | 14 735 |

*) Provisional figures.

Table 30. Cod. ICES Sub-Area XIV plus ICNAP Divisions IE-IF.
Fishing mortalities by year and age.

| Age | 1960 | 1961 | 1962 | 1963 | 1964 | 1965 | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 |
|-----------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| 3 | 0.00 | 0.00 | 0.02 | 0.00 | 0.00 | 0.00 | 0.01 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 4 | 0.02 | 0.00 | 0.03 | 0.03 | 0.01 | 0.02 | 0.02 | 0.01 | 0.02 | 0.02 | 0.02 | 0.01 | 0.07 | 0.03 | 0.02 | 0.02 |
| 5 | 0.04 | 0.08 | 0.06 | 0.14 | 0.08 | 0.11 | 0.18 | 0.05 | 0.10 | 0.10 | 0.19 | 0.14 | 0.18 | 0.17 | 0.12 | 0.09 |
| 6 | 0.06 | 0.12 | 0.17 | 0.15 | 0.24 | 0.20 | 0.14 | 0.22 | 0.22 | 0.21 | 0.16 | 0.45 | 0.45 | 0.30 | 0.14 | 0.15 |
| 7 | 0.11 | 0.13 | 0.17 | 0.35 | 0.22 | 0.28 | 0.30 | 0.32 | 0.33 | 0.27 | 0.23 | 0.31 | 0.46 | 0.50 | 0.18 | 0.22 |
| 8 | 0.16 | 0.19 | 0.23 | 0.26 | 0.52 | 0.28 | 0.43 | 0.31 | 0.28 | 0.36 | 0.32 | 0.52 | 0.47 | 0.37 | 0.45 | 0.22 |
| 9 | 0.12 | 0.19 | 0.24 | 0.10 | 0.41 | 0.52 | 0.34 | 0.45 | 0.20 | 0.21 | 0.47 | 0.56 | 0.78 | 0.71 | 0.42 | 0.22 |
| 10 | 0.19 | 0.15 | 0.23 | 0.22 | 0.18 | 0.29 | 0.52 | 0.45 | 0.16 | 0.13 | 0.47 | 0.77 | 0.66 | 0.76 | 0.71 | 0.22 |
| 11 | 0.09 | 0.23 | 0.15 | 0.20 | 0.33 | 0.21 | 0.22 | 0.54 | 0.29 | 0.18 | 0.40 | 0.55 | 0.95 | 0.50 | 1.03 | 0.22 |
| 12 | 0.06 | 0.08 | 0.28 | 0.18 | 0.18 | 0.47 | 0.13 | 0.30 | 0.15 | 0.37 | 0.23 | 0.22 | 0.94 | 0.58 | 1.18 | 0.22 |
| 13 | 0.39 | 0.08 | 0.05 | 0.36 | 0.10 | 0.11 | 0.36 | 0.23 | 0.09 | 0.31 | 0.67 | 0.13 | 0.52 | 0.48 | 1.36 | 0.22 |
| 14 | 0.50 | 0.50 | 0.50 | 0.50 | 0.50 | 0.50 | 0.50 | 0.50 | 0.50 | 0.50 | 0.50 | 0.50 | 0.50 | 0.50 | 0.50 | 0.22 |
| Mean $F \geq 7$ | 0.14 | 0.18 | 0.22 | 0.31 | 0.32 | 0.32 | 0.38 | 0.36 | 0.31 | 0.30 | 0.29 | 0.48 | 0.63 | 0.61 | 0.50 | 0.22 |

The last group is a plus group.

Table 31. Cod. ICES Sub-Area XIV plus ICNAF Divisions IE-IF.
Stock in numbers at beginning of year.

| Age | 1960 | 1961 | 1962 | 1963 | 1964 | 1965 | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 |
|-----|---------|---------|--------|--------|---------|---------|---------|---------|---------|--------|--------|--------|--------|--------|--------|--------|
| 3 | 135 469 | 103 633 | 25 548 | 82 807 | 210 123 | 116 737 | 191 570 | 60 170 | 15 000 | 16 137 | 10 362 | 60 481 | 3 786 | 13 246 | 25 472 | 42 172 |
| 4 | 144 361 | 110 912 | 84 817 | 20 521 | 67 767 | 171 963 | 95 576 | 155 753 | 49 219 | 12 273 | 13 212 | 8 484 | 49 518 | 3 099 | 10 838 | 20 801 |
| 5 | 24 790 | 116 123 | 90 473 | 67 558 | 16 264 | 54 952 | 138 464 | 76 430 | 126 423 | 39 305 | 9 892 | 10 603 | 6 864 | 37 861 | 2 461 | 8 686 |
| 6 | 21 321 | 19 446 | 87 870 | 69 568 | 47 986 | 12 261 | 40 148 | 95 075 | 59 405 | 93 277 | 29 097 | 6 684 | 7 520 | 4 686 | 26 197 | 1 783 |
| 7 | 78 457 | 16 382 | 14 062 | 60 588 | 48 899 | 31 035 | 8 256 | 28 647 | 62 776 | 39 209 | 61 783 | 20 378 | 3 479 | 3 940 | 2 842 | 18 607 |
| 8 | 17 966 | 43 009 | 8 835 | 7 261 | 26 111 | 23 943 | 14 423 | 3 737 | 12 683 | 27 732 | 18 375 | 29 942 | 9 174 | 1 346 | 1 461 | 1 447 |
| 9 | 14 042 | 9 416 | 21 871 | 4 305 | 3 428 | 9 541 | 11 123 | 5 722 | 1 683 | 5 874 | 11 823 | 8 160 | 10 877 | 3 506 | 568 | 571 |
| 10 | 22 965 | 7 656 | 4 752 | 10 540 | 2 392 | 1 395 | 3 458 | 4 834 | 2 242 | 843 | 2 906 | 4 515 | 2 843 | 3 062 | 1 054 | 229 |
| 11 | 9 435 | 11 681 | 4 037 | 2 318 | 5 194 | 1 228 | 638 | 1 265 | 1 889 | 1 168 | 454 | 1 109 | 1 276 | 903 | 874 | 317 |
| 12 | 9 254 | 5 295 | 5 681 | 2 134 | 1 165 | 2 283 | 610 | 313 | 452 | 864 | 595 | 187 | 393 | 302 | 337 | 190 |
| 13 | 7 384 | 5 318 | 3 002 | 2 638 | 1 097 | 594 | 873 | 328 | 142 | 238 | 366 | 291 | 92 | 94 | 103 | 63 |
| 14 | 1 683 | 3 065 | 3 020 | 1 756 | 1 133 | 608 | 325 | 372 | 160 | 79 | 107 | 115 | 156 | 34 | 36 | 16 |

Table 32. Prediction of catch and biomass for Cod in ICES Sub-Area XIV and ICNAF Divisions IE-IF.

| 1975 | 1976 | | | 1977 | | | 1978 |
|------|---|---------|---|---|---------|---|---|
| | Biomass ^x Age ≥ 4 (1 000 tons) | F (≥ 7) | Predicted ^x catch (1 000 tons) | Biomass ^x Age ≥ 4 (1 000 tons) | F (≥ 7) | Predicted ^x catch (1 000 tons) | Biomass ^x Age ≥ 4 (1 000 tons) |
| | | | | | 0.22 | 16.5 (0.9) | 203 (85) |
| | | 0.22 | 13.9 (0.08) | 187 (57) | 0.45 | 30.5 (0.9) | 185 (83) |
| 126 | 136 | 0.45 | 26.2 (0.2) | 174 (56) | 0.22 | 14.7 (0.9) | 196 (85) |
| | | | | | 0.45 | 27.1 (0.9) | 178 (83) |

x) The biomass is given by 1 January and therefore includes only fish 4 years and older at that time. During the year 3-year-old fish will recruit, and some of these are included in the catch figures.

The figures in brackets reflect that part of the predicted catches and stock which is dependent on the incoming year classes.

Table 33. Numbers of Cod emigrating from Greenland (ICES Sub-Area XIV and ICNAF Divisions IE-IF) to Iceland (ICES Division Va) in Nos. x 10⁻⁶.

| Age Years | 1960 | 1961 | 1962 | 1963 | 1964 | 1965 | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 |
|-----------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| 7 | 17.1 | 3.5 | 3.0 | 11.9 | 10.2 | 6.3 | 1.7 | 5.7 | 12.4 | 8.0 | 12.8 | 4.1 | 0.7 | 0.7 | 0.6 | 3.9 |
| 8 | 3.8 | 9.0 | 1.8 | 1.5 | 4.8 | 4.8 | 2.7 | 0.7 | 2.6 | 5.4 | 3.7 | 5.5 | 1.7 | 0.3 | 0.3 | 0.3 |
| 9 | 3.1 | 2.0 | 4.6 | 0.9 | 0.7 | 1.7 | 2.2 | 1.1 | 0.4 | 1.2 | 2.2 | 1.5 | 1.8 | 0.6 | 0.1 | 0.1 |
| 10 | 4.8 | 1.6 | 1.0 | 2.2 | 0.5 | 0.3 | 0.6 | 1.0 | 0.5 | 0.2 | 0.5 | 0.7 | 0.5 | 0.5 | 0.2 | - |
| 11 | 2.1 | 2.4 | 0.9 | 0.5 | 1.0 | 0.3 | 0.1 | 0.2 | 0.4 | 0.2 | 0.1 | 0.2 | 0.2 | 0.2 | 0.1 | 0.1 |
| 12 | 2.1 | 1.2 | 1.1 | 0.5 | 0.2 | 0.4 | 0.1 | 0.1 | 0.1 | 0.2 | 0.1 | - | 0.1 | 0.1 | - | - |
| 13 | 1.4 | 1.2 | 0.7 | 0.5 | 0.2 | 0.1 | 0.2 | - | - | - | 0.1 | 0.1 | - | - | - | - |
| 14 | 0.3 | 0.6 | 0.6 | 0.3 | 0.2 | 0.1 | 0.1 | 0.1 | - | - | - | - | - | - | - | - |
| Total | 34.7 | 21.5 | 13.6 | 18.3 | 17.8 | 14.0 | 7.7 | 8.9 | 16.4 | 14.2 | 19.5 | 12.1 | 5.0 | 2.4 | 1.3 | 4.4 |


C ∞

| Age/Year Classes | 1953 | 1954 | 1955 | 1956 | 1957 | 1958 | 1959 | 1960 | 1961 | 1962 | 1963 | 1964 | 1965 | 1966 | 1967 | 1968 |
|------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|-------|
| 7 | 17.1 | 3.5 | 3.0 | 11.9 | 10.2 | 6.3 | 1.7 | 5.7 | 12.4 | 8.0 | 12.8 | 4.1 | 0.7 | 0.7 | 0.6 | 3.9 |
| 8 | 9.0 | 1.8 | 1.5 | 4.8 | 4.8 | 2.7 | 0.7 | 2.6 | 5.4 | 3.7 | 5.5 | 1.7 | 0.3 | 0.3 | 0.3 | - |
| 9 | 4.6 | 0.9 | 0.7 | 1.7 | 2.2 | 1.1 | 0.4 | 1.2 | 2.2 | 1.5 | 1.8 | 1.6 | 0.1 | 0.1 | - | - |
| 10 | 2.2 | 0.5 | 0.3 | 0.6 | 1.0 | 0.5 | 0.2 | 0.5 | 0.7 | 0.5 | 0.5 | 0.2 | - | - | - | - |
| 11 | 1.0 | 0.3 | 0.1 | 0.2 | 0.4 | 0.2 | 0.1 | 0.2 | 0.2 | 0.2 | 0.1 | 0.1 | - | - | - | - |
| 12 | 0.4 | 0.1 | 0.1 | 0.1 | 0.2 | 0.1 | - | 0.1 | 0.1 | - | - | - | - | - | - | - |
| 13 | 0.2 | - | - | - | 0.1 | 0.1 | - | - | - | - | - | - | - | - | - | - |
| 14 | 0.1 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Total | 34.6 | 7.1 | 5.7 | 19.3 | 18.9 | 11.0 | 3.1 | 10.3 | 21.0 | 13.9 | 20.7 | 6.7 | 1.1 | 1.1 | 0.9 | (3.9) |

Figure 1.

Iceland Cod.

Trends in estimates of fishing effort and of weighted F values (age groups 4-6) from two VPAs, all expressed relative to the 1970 value.



| | Z | F | | Z | \hat{F} |
|-----------|------|------|------|------|-----------|
| 1961 | .570 | .224 | 1973 | .880 | 0.52 |
| 1962 | .550 | .269 | 1974 | .930 | 0.55 |
| 1963 | .805 | .311 | | | |
| 1964 | .755 | .368 | | | |
| 1965 | .770 | .420 | | | |
| 1966 | .690 | .231 | | | |
| 1967 | .285 | .211 | | | |
| 1968 | .345 | .213 | | | |
| 1969 | .540 | .283 | | | |
| 1970 | .610 | .338 | | | |
| 1971 | .730 | .529 | | | |
| 1972 | .910 | .560 | | | |
| \bar{X} | .630 | .312 | | | |




Figure 2. Iceland Cod.
The relation between fishing mortality from VPA and total mortality based on English trawler catch per effort.




Figure 3. Iceland Cod, Spawning stock biomass (age groups 7+) estimated from VPA.

Figure 4. Iceland Cod.
Change in biomass with age in an unexploited year class.




Figure 5. Iceland Cod.
Yield per recruit and spawning stock
per recruit under the present exploitation
pattern.




Figure 6. Iceland Cod.
Relationship between stock size of 3
year olds and catch per effort of 3
year olds by English trawlers.




Figure 7. Iceland Cod.
 Relationship between stock size of 4 year olds
 and catch per effort of 4 year olds by English
 trawlers.

| Year class | x | y | \hat{R}_3 |
|------------|------|-----|-------------|
| 1957 | 4.1 | 112 | |
| 1958 | 5.2 | 142 | |
| 1959 | 4.8 | 105 | |
| 1960 | 4.2 | 120 | |
| 1961 | 5.6 | 225 | 12.8 |
| 1962 | 6.4 | 192 | |
| 1963 | 5.4 | 222 | |
| 1964 | 11.1 | 254 | 282 |
| 1965 | 5.7 | 134 | |
| 1966 | 9.5 | 203 | 146 |
| 1967 | 5.9 | 146 | |
| 1968 | 5.6 | 142 | |
| 1969 | 5.5 | 108 | |

b 17.8783
 a 53.2780
 Sxy 37.4695

Si 5.3913

\bar{x} 6.08
 Σx 79.0
 $S(x-\bar{x})^2$ 48.3031

\bar{y} 162
 Σy 2105

$S(y-\bar{y})^2$ 3.0883

r = .7071 p < .001

n = 13




Figure 8. Iceland Cod.
Relationship between English catch per effort
of 3 year old cod and estimates from inter-
national 0-group surveys.




Figure 10. The dependence of the estimated fishing mortalities at Iceland (4-6 years old) on the numbers of 7-year old fish at Greenland (B) and emigration numbers of emigrants from Greenland (A).

• A Emigrants from Greenland (age 7 and older)
 $y = -.001 x + .416$; $r = .86$

* B Number of 7-year old fish at Greenland
 $y = -.0041 x + .431$; $r = .91$

